

ROLE OF CINNAMOMUM TAMALA (LAURACEAE) AND NIGELLA SATIVA (RANUNCULACEAE) VOLATILE OILS IN MAIZE WEEVIL, SITOPHILUS ZEAMAI, MOTSCHULSKY MANAGEMENT

 Mukesh Kumar Chaubey^{1,*},  Namita Kumar²

¹National Post Graduate College, Department of Zoology, Barahalganj, Gorakhpur, India

²Mahatma Gandhi Post Graduate College, Department of Zoology, Gorakhpur, India

*Corresponding Author:

E-mail: mgpgc9839427296@gmail.com

(Received 22nd August 2023; accepted 05th October 2023)

ABSTRACT. Synthetic insecticides used indiscriminately in insect pest management programme results in carcinogenicity, mutagenesis, neurotoxicity and teratogenicity in non-target animals and development of resistance in target animals. These issues have diverted the researches aiming insect pest management towards the use of plant volatiles. In this study, insecticidal properties of *Cinnamomum tamala* (Lauraceae) and *Nigella sativa* (Ranunculaceae) volatile oils have been evaluated against maize weevil, *Sitophilus zeamais* (Coleoptera: Curculionidae). Volatile oils were isolated and tested for repellent, toxic, oviposition inhibitory, developmental inhibitory and feeding inhibitory properties against *S. zeamais*. In toxicity assay, median lethal concentrations of *C. tamala* and *N. sativa* oils were 0.396 and 0.334 μcm^{-3} ; and 0.369 and 0.328 μcm^{-3} air respectively when *S. zeamais* adults were fumigated for 24 and 48h. In contact toxicity assay, lethal concentrations of *C. tamala* and *N. sativa* oils were 0.287 and 0.205 μcm^{-2} ; and 0.246 and 0.195 μcm^{-2} area for 24 and 48h respectively when *S. zeamais* adults were exposed. These two volatile oils used reduced acetylcholine esterase activity in adults when fumigated with sub-lethal concentrations. Both volatile oils significantly reduced oviposition, progeny production and feeding, but, increased developmental period in *S. zeamais*. Therefore, it can be concluded that these two oils viz. *C. tamala* and *N. sativa* oils can be used in preparation of volatile oil based formulations in insect pest management.

Keywords: Volatile oil, insecticides, *Sitophilus zeamais*, oviposition inhibition, acetylcholine esterase.

INTRODUCTION

Insect pest especially of grains under storage damage grains both qualitatively and quantitatively causing huge economic loss annually. To mitigate such losses, synthetic insecticides have been applied variously. But continuous and unlimited uses of these insecticides damage human health and environment greatly including depletion of ozone layer, neurotoxic, carcinogenic, teratogenic and mutagenic effects in non-target animals as well as cross- and multi resistance in target/non-target insects [1-3]. Synthetic insecticides are killing over thousands of people per year unintentionally [4]. Besides, insecticides of synthetic origin have entered in our ecosystems affecting food chain [5]. Among these insecticides, organochlorines are persistent and accumulate at population level causing mass killings of bees, fishes, amphibians, birds and mammals [6,7]. Thus,

scientific communities have diverted their researches towards the use of plant derived volatile oils as alternative of insect pest management. These volatile oils are secondary metabolites in plant families like *Alliaceae*, *Apiaceae*, *Asteraceae*, *Cupressaceae*, *Lamiaceae*, *Lauraceae*, *Myrtaceae*, *Piperaceae*, *Poaceae*, *Rutaceae* and *Zingiberaceae*. Volatile oils are complex mixtures of compounds in different proportions depending on plant parts used for extraction, method of extraction, phenological stage of plant, season of harvesting, age of plant, genotype of plant, nature of soil and conditions of environment [8-10]. Most of these oils possess larvicidal, adulticidal, antifeedant, oviposition, developmental and adult emergence inhibitory properties for a number of insects [11-15].

Cinnamomum tamala, Indian bay leaf also known as tejpat or tejapatta is a tree of *Lauraceae* family. It is native to India, Bangladesh, Bhutan, China and Nepal. Its leaves have a clove like smell. It is used mainly for culinary and medicines [16]. The volatile oil from fresh leaves contains eugenol as major constituent followed by spathulenol and eugenyl acetate while volatile oil from dried leaves contains eugenol as main constituents followed by spathulenol, methyl eugenol, β -humulene and caryophyllene [17]. *C. tamala* has known for antibacterial, antioxidant, antiulcer, anticancer and antimicrobial properties [18-24]. *C. tamala* leaves have several biological properties and act as diuretic, diaphoretic, lactagogue and hypoglycemic agents [25,26]. *C. tamala* oil is effective as hypoglycaemic and hypolipidemic agents [27].

Nigella sativa also known as black caraway, black cumin, kalonji or siyahdaneh is a herb of family *Ranunculaceae*. It is native to eastern European countries like Bulgaria and Romania and Asian countries like Cyprus, Iran, Iraq, Pakistan and Turkey. It is an important spice and drug in Indian traditional system of medicine [28]. *N. sativa* is used in lowering blood pressure, triglycerides cholesterol level [29]. It has anti-bacterial, anti-cancer, anti-diabetic, anti-convulsant activity and anti-oxidant properties [30-34]. The *nigella* seeds contain volatile oil (0.4-0.45 %) which has constituents like anethol, carvacrol, d-citronellol, *p*-cymene, d-limonene, nigellone, α -pinene, β -pinene, 4-terpineol, thymoquinone, thymohydroquinone, dithymoquinone and thymol and longifoline [35,36].

Sitophilus zeamais, Motschulsky (Coleoptera: *Curculionidae*) is a serious pest of maize, wheat, rice, sorghum, oats, barley, rye, buckwheat, peas and cottonseed [37]. It is commonly found in humid tropical areas of the world where maize is cultivated. Adults attack whole grains and larvae cryptically feed and develop inside grains [38]. The present study aims to evaluate repellent, insecticidal, acetylcholine esterase inhibitory, oviposition inhibitory, developmental inhibitory and antifeedant activities of *C. tamala* and *N. sativa* volatile oils against maize weevil, *S. zeamais*.

MATERIALS AND METHODS

Essential oils

Dried leaves of *C. tamala* and seeds of *N. sativa* were purchased from local market of Gorakhpur. These were grounded and hydrodistilled in Clevenger apparatus to isolate volatile oils. Oils were stored in Eppendorff tubes at 4 °C till further use.

Insects

Maize weevil, *S. zeamais* was cultured on whole maize grain at $28\pm4^{\circ}\text{C}$, $50\pm5\%$ relative humidity (RH) and photoperiod of 10:14 (Light:Dark) hours to evaluate insecticidal properties of *C. tamala* and *N. sativa* volatile oils.

Repellent activity

Repellent activity of *C. tamala* and *N. sativa* volatile oils was assayed by filter paper disc method. Experimental solutions of *C. tamala* and *N. sativa* oils were prepared in acetone. Whatman filter papers were cut into two halves: one half was treated with test solution and the other half was treated with solvent only. Oil treated and solvent treated halves were dried to evaporate solvent, attached with cellophane tape and placed in each glass petri dish (diameter 8.5 cm, height 1.2 cm). Now, forty *S. zeamais* adults were released in petri dish, covered and kept in dark. In the experiment six replicates were set for each concentration of oil. After 4 hours of exposure, adults in treated and untreated halves were counted and percent repellency (PR) was calculated using formula: $\text{PR} = [(C-T)/(C+T)] \times 100$, C = Number of insects in the untreated half of the filter paper and T = number of insect in treated half of the filter paper. Preference index (PI) was calculated using formula: $\text{PI} = (\text{percentage of insects in treated halves} - \text{percentage of insects in untreated halves}) / (\text{percentage of insects in treated halves} + \text{percentage of insects in untreated halves})$.

Fumigant toxicity

Experimental solutions of *C. tamala* and *N. sativa* oils were made in acetone. Ten adults taken from laboratory culture were placed in glass petri dish and added 2 gm of maize grains. Experimental oil formulation was applied in a filter paper strip (2 cm diameter) and was left for few minutes to evaporate solvent. Now, treated paper was pasted on the inner surface of the petri dish cover, air tightened with parafilm and kept in rearing conditions in dark. For each concentration of oil and control, six replicates were set. Mortality in adults was recorded after 24 and 48h of exposure.

Contact toxicity

Experimental solutions of *C. tamala* and *N. sativa* oils were made in acetone. Experimental oil solution was applied on the inner surface of glass petri dish (diameter 8.5 cm, height 1.2 cm). Petri dish was left open for few minutes to evaporate solvent completely. Now, ten *S. zeamais* adults were released in petri dish, covered and kept in rear conditions in dark. For each concentration of oil and control, six replicates were set. Mortality in adults was recorded after 24 and 48h of exposure.

Acetylcholine esterase activity

S. zeamais adults were fumigated with 40 and 80% of 24h-LC₅₀ of *C. tamala* and *N. sativa* oils for 24 hours. After fumigation, adult insects were collected and used for acetylcholine esterase enzyme activity estimation [39]. The experimental adult insects were homogenized in phosphate buffer saline (50mM, pH8), centrifuged and supernatant collected was used as enzyme source for estimation of acetylcholine esterase activity. To 0.1ml of enzyme source, 0.1ml acetylthiocholine iodide (0.5 mM), 0.05 ml, 5,5-Dithio-bis 2-nitrobenzoic acid (0.33 mM) and 1.45 ml phosphate

buffer (50 mM, pH 8) were added. The reaction mixture was incubated for 3 min at 25°C. Now, changes in absorbance of reaction mixture were measured at 412nm for the determination of acetylcholine esterase activity. Acetylcholine esterase activity was estimated in terms of mmol of 'SH' hydrolysed min⁻¹mg⁻¹ protein.

Oviposition inhibition

In this assay, *S. zeamais* adults of mixed sex were taken from the laboratory culture and fumigated with 40 and 80% of 24h-LC₅₀ and 48h-LC₅₀ of *C. tamala* and *N. sativa* oils for 24 hours and 48 hours respectively. After fumigation, the treated adults were reared on maize grain. After 10 days, discarded adults and counted the number of F₁ progeny after 45 days. Six replicates were set for each concentration of oil as well as control. The percentage oviposition deterrence (POD) was calculated using the formula [40]: Percentage oviposition deterrence (POD) = [(E_C-E_T)/E_C] × 100, where E_C = number of adults emerged in control and E_T = number of adults emerged in test.

Developmental inhibition

Ten *S. zeamais* adults of mixed sex were placed in a 250 ml plastic container with 20 gm of maize grains and allowed to mate and lay eggs for seven days in condition applied for rearing of insects. Now, adults were removed and a filter paper strip (2 cm diameter) impregnated with *C. tamala* and *N. sativa* oil was pasted on undercover of the container. The eggs and juveniles were fumigated with three concentrations (0.2, 0.4 and 0.6 µlcm⁻³) of *C. tamala* and *N. sativa* oils. The duration from the end of fumigation till the emergence of F₁ adults and the number of adults emerged in control as well as in test was recorded. Six replicates were set for each concentration of oil and control. Inhibition rate (IR) was calculated using formula [41]: Inhibition rate (IR) = [(C_n-T_n)/C_n] × 100, where C_n = number of adults emerged in control and T_n = number of adults emerged in test.

Antifeedant activity

For the determination of antifeedant activity of *C. tamala* and *N. sativa* oils, flour discs were prepared by mixing 10 gm of maize flour with 50 ml water until completely suspended. Maize flour suspension was pipetted out (200 µl) onto a plastic sheet, held for 24 hours at room temperature and then dried in an oven for one hour at 60°C. Each flour disc was treated with 40 and 80% of 96h-LC₅₀ of *C. tamala* and *N. sativa* oil, weighed, placed in glass petri dish and twenty-five *S. zeamais* adults were introduced into it and allowed to feed. After four days of feeding, flour discs were taken and reweighed to calculate antifeedant activity (AFA) using formula [42]: AFA = [C-T/C] × 100, where C = consumption of food in control group and T = consumption of food in treated group. Six replicates were set for each concentration of oil and control.

Statistical analysis

Median lethal concentration was estimated by POLO programme [43]. One-way analysis of variance and correlation and regression were conducted to study concentration-response relationship [44]. Tukey's test was performed to detect the significant difference between means [45].

RESULTS AND DISCUSSION

Repellent activity

Percent Repellency (PR) and Preference Index (PI) were increased with increase in the concentration of *C. tamala* and *N. sativa* oils and were recorded maximum at 0.8% concentrations (Table1).

Table 1. Repellent activity of *C. tamala* and *N. sativa* volatile oils against *S. zeamais* adults

Oil	Concentration (%)	Percent Repellency (PR)* Mean±SD	Preference Index (PI)**
<i>C. tamala</i>	0.1	22.50±1.04	-0.22
	0.2	48.52±0.93	-0.48
	0.4	72.50±1.09	-0.72
	0.8	98.35±0.08	-0.98
<i>N. sativa</i>	0.1	15.80±0.98	-0.15
	0.2	35.80±0.75	-0.35
	0.4	64.25±0.84	-0.64
	0.8	90.80±0.07	-0.90

*Percent repellency (PR) was calculated using formula: $PR = [(C-T)/(C+T)] \times 100$, C = number of insects in the untreated halves and T = number of insect in treated halves; **Preference index (PI) was calculated using formula: $PI = (\text{percentage of insects in treated halves} - \text{percentage of insects in untreated halves}) / (\text{percentage of insects in treated halves} + \text{percentage of insects in untreated halves})$. PI value between -1.0 to -0.1 indicates repellent oil, -0.1 to +0.1 neutral oil and +0.1 to +1.0 attractant oil.

Fumigant toxicity

Median lethal concentrations were recorded 0.396 and 0.334 $\mu\text{lc m}^{-3}$ air for *C. tamala* oil after 24 and 48 hours exposure period respectively (Table 2). On the other hand, LC₅₀ values were 0.369 and 0.328 $\mu\text{lc m}^{-3}$ air for *N. sativa* oil after 24 and 48 hours exposure period respectively (Table 2). The index of significance of potency estimation, g-value indicates that the mean value is within the limits of all probability levels ($P < 0.1$, 0.5 and 0.01) as it is less than 0.5. Values of t-ratio greater than 1.6 indicated that regression was significant. Values of heterogeneity factor less than 1.0 denotes that model fits the data adequate. Regression analysis showed concentration-dependent mortality in *S. zeamais* adults as lethality was found to increase with increase in concentration of oils (Table 2).

Contact toxicity

Median lethal concentrations were 0.287 and 0.205 $\mu\text{lc m}^{-2}$; and 0.246 and 0.195 $\mu\text{lc m}^{-2}$ area for *C. tamala* and *N. sativa* essential oils after 24 and 48 hours exposure period respectively (Table 2). Regression analysis showed concentration-dependent mortality in *S. zeamais* adults by *C. tamala* and *N. sativa* oil (Table 2).

Table 2. Fumigant and contact toxicity of *C. tamala* and *N. sativa* volatile oils against *S. zeamais* adults

Oil	Toxicity	Exposure period (h)	LC ₅₀ *	g-value	Heterogeneity	t-ratio	Regression Equation	Correlation Coefficient
<i>C. tamala</i>	Fumigant	24	0.396	0.20	0.33	3.78	Y = - 3.87+5.87X	0.98
	toxicity	48	0.334	0.15	0.35	4.67	Y = 5.23+2.87X	0.99
	Contact	24	0.287	0.28	0.34	4.19	Y = - 4.05+2.86X	0.98
	toxicity	48	0.205	0.23	0.33	3.87	Y = 9.08+5.31X	0.99
<i>N. sativa</i>	Fumigant	24	0.369	0.29	0.35	3.93	Y = - 7.46+7.91X	0.99
	toxicity	48	0.328	0.27	0.34	4.18	Y = 6.84+3.54X	0.98
	Contact	24	0.246	0.24	0.32	4.79	Y = - 7.98+6.54X	0.99
	toxicity	48	0.195	0.26	0.35	3.83	Y = 7.43+6.43X	0.99

* $\mu\text{lc m}^{-3}$ for fumigant toxicity and $\mu\text{lc m}^{-2}$ for contact toxicity

Acetylcholine esterase enzyme activity

Fumigation of *S. zeamais* adults with 40 and 80% of 24h-LC₅₀ of *C. tamala* oil and *N. sativa* oil reduced acetylcholine esterase activity to 82.38 and 55.97%; and 74.73 and 50.83% of control respectively (Table 3). Both *C. tamala* and *N. sativa* oils significantly inhibited activity of acetylcholine esterase enzyme in *S. zeamais* adults (For *C. tamala* oil F=106.43; For *N. sativa* oil F = 120.00; P<0.05; Table 3). Post Hoc Tukey's Honestly Significant Difference (HSD) analysis shows significant (P<0.05) difference between all the treated and control groups (Table 4).

Table 3. Effect of *C. tamala* and *N. sativa* volatile oils on acetylcholine esterase activity in *S. zeamais*

Oil	Concentration	Enzyme activity* (Mean±SD)	F-value (df = 2,15)
<i>C. tamala</i>	Control	0.0954±0.0029(100)	106.43**
	40% of 24h-LC ₅₀	0.0768±0.00024(82.38)	
	80% of 24h-LC ₅₀	0.0534±0.0016(55.97)	
<i>N. sativa</i>	Control	0.0954±0.0029 (100)	120.00**
	40% of 24h-LC ₅₀	0.0713±0.0022(74.73)	
	80% of 24h-LC ₅₀	0.0485±0.0015(50.83)	

*mmol of 'SH' hydrolysed $\text{min}^{-1}\text{mg}^{-1}$ protein; Values in parentheses indicate per cent change with respect to control taken as 100%; **Significant at P<0.05 (df = 2,15)

Table 4. Post Hoc Tukey's Honestly Significant Difference (HSD) analysis of effect of *C. tamala* and *N. sativa* volatile oils on acetylcholine esterase activity in *S. zeamais*

Oil	Pair wise comparison		HSD0.05 = 0.0074 HSD0.01 = 0.0098	Q0.05 = 3.6734 Q0.01 = 4.8359
<i>C. tamala</i>	T1:T2	M1 = 0.10 M2 = 0.07	0.02	Q = 11.37*
	T1:T3	M1 = 0.10 M3 = 0.06	0.04	Q = 20.61*
	T2:T3	M2 = 0.07 M3 = 0.06	0.02	Q = 9.24*
	T1:T2	M1 = 0.10 M2 = 0.07	0.03	Q = 13.43*
	T1:T3	M1 = 0.10 M3 = 0.07	0.05	Q = 21.70*
	T2:T3	M2 = 0.07 M3 = 0.05	0.02	Q = 8.27*

*Significant ($P < 0.05$) difference between all the treated and control groups.

Oviposition inhibition

Fumigation of *S. zeamais* adults with *C. tamala* and *N. sativa* oils reduced oviposition capacity of insects (Table 5). Oviposition was reduced to 63.47 and 58.56% of control respectively when *S. zeamais* adults were fumigated with 80% of 24h-LC₅₀ of *C. tamala* and *N. sativa* oils (For *C. tamala* oil $F = 186.95$; For *N. sativa* oil $F = 173.62$; $P < 0.05$; Table 5). Percent oviposition deterrence was recorded 40.33 and 41.14% when *S. zeamais* adults were fumigated with 80% of 24h-LC₅₀ of *C. tamala* and *N. sativa* oils respectively (Table 5).

Table 5. Oviposition inhibitory activities of *C. tamala* and *N. sativa* volatile oils in *S. zeamais*

Oil	Conc.	No. of progeny emerged (Mean±SD)	POD*	F-value	Conc.	No. of progeny emerged (Mean±SD)	POD*	F-value
<i>C. tamala</i>	Control	94.00±3.52 (100%)	-		Control	94.00±3.52 (100%)	-	
	40% of 24h-LC ₅₀	72.67±2.34 (78.37)	22.63	186.95**	40% of 48h-LC ₅₀	55.00±2.13 (58.51)	41.49	163.26**
	80% of 24h-LC ₅₀	59.67±2.23 (63.47)	40.33		80% of 48h-LC ₅₀	39.33±1.20 (41.84)	58.16	
<i>N. sativa</i>	Control	94.00±3.52 (100%)	-		Control	94.00±3.52 (100%)	-	
	40% of 24h-LC ₅₀	69.33±2.12 (73.75)	26.25	173.62**	40% of 48h-LC ₅₀	53.00±2.08 (56.38)	43.62	177.28**
	80% of 24h-LC ₅₀	55.33±2.17 (58.86)	41.14		80% of 48h-LC ₅₀	39.50±1.07 (42.02)	57.98	

Values in parentheses indicate per cent change with respect to control taken as 100%; *Percentage of oviposition deterrence (POD) = $[(EC-ET)/EC] \times 100$; EC = number of adults emerged in control and ET = number of adults emerged in test; **Significant at $P < 0.05$ (df = 2,15)

Similarly, reduction in oviposition was 41.84 and 42.02% of control when *S. zeamais* adults were fumigated with 80% of 48h-LC₅₀ of *C. tamala* and *N. sativa* oils respectively (For *C. tamala* oil F = 163.26; For *N. sativa* F = 177.28) (Table 5). Post Hoc Tukey's Honestly Significant Difference (HSD) analysis shows significant (P<0.05) difference between all the treated and control groups (Table 6).

Table 6. Post Hoc Tukey's Honestly Significant Difference (HSD) analysis of oviposition inhibitory activities of *C. tamala* and *N. sativa* volatile oils in *S. zeamais*

Oil	Exposure period 24h			Exposure period 48h		
	Pair wise comparison T ₁ :T ₂	HSD _{0.05} = 6.6693 HSD _{0.01} = 8.7799 21.33	Q _{0.05} = 3.6734 Q _{0.01} = 4.8359 Q = 11.75*	Pair wise comparison T ₁ :T ₂	HSD _{0.05} = 7.3195 HSD _{0.01} = 9.6359 39.00	Q _{0.05} = 3.6734 Q _{0.01} = 4.8359 Q = 19.57*
<i>C. tamala</i>	M ₁ = 94.00 M ₂ = 72.67			M ₁ = 94.00 M ₂ = 55.00		
	T ₁ :T ₃	34.33	Q = 18.91*	T ₁ :T ₃	54.67	Q = 27.44*
	M ₃ = 59.67			M ₃ = 39.33		
	T ₂ :T ₃	13.00	Q = 7.16*	T ₂ :T ₃	15.67	Q = 7.86*
<i>N. sativa</i>	M ₃ = 59.67			M ₃ = 39.33		
	Pair wise comparison	HSD _{0.05} = 8.1099 HSD _{0.01} = 10.6764 24.67	Q _{0.05} = 3.6734 Q _{0.01} = 4.8359 Q = 11.17*	Pair wise comparison	HSD _{0.05} = 7.9684 HSD _{0.01} = 10.4902 41.00	Q _{0.05} = 3.6734 Q _{0.01} = 4.8359 Q = 18.90*
	T ₁ :T ₂	M ₁ = 94.00 M ₂ = 69.33		T ₁ :T ₂	M ₁ = 94.00 M ₂ = 53.00	
	T ₁ :T ₃	M ₁ = 94.00 M ₃ = 55.33	Q = 17.51*	T ₁ :T ₃	54.50	Q = 25.12*
	T ₂ :T ₃	M ₂ = 69.33 M ₃ = 55.33	Q = 6.34*	T ₂ :T ₃	13.50	Q = 6.22*

*Significant (P<0.05) difference between all the treated and control groups

Developmental inhibition

Progeny production was reduced to 51.27 and 48.43% as compared to the control when fumigated with $0.6 \mu\text{cm}^{-3}$ of *C. tamala* and *N. sativa* oils respectively (Table 7). Both *C. tamala* and *N. sativa* oils slow down the developmental progression of *S. zeamais* larva and pupa (Table 7). The duration of larval phase to adult emergence was increased significantly to 120.25% and 128.27% as compared to the control when fumigated with $0.6 \mu\text{cm}^{-3}$ of *C. tamala* and *N. sativa* essential oils respectively (For *C. tamala* oil $F = 34.65$; For *N. sativa* 48.79; $P < 0.01$; Table 7). Post Hoc Tukey's Honestly Significant Difference (HSD) analysis shows significant ($P < 0.05$) difference between all the treated and control groups (Table 8).

Table 7. Effect of *C. tamala* and *N. sativa* volatile oils on development of *S. zeamais*

Oil	Conc.	No. of progeny emerged (Mean±SD)	PR*	F-value	Duration of developmental period in days (Mean±SD)	F-value
<i>C. tamala</i>	Control	85.17±6.01 (100)	-		24.83±0.54 (100)	
	0.2 μcm^{-3}	69.83±3.18 (81.98)	18.02		26.50±0.96 (106.59)	
	0.4 μcm^{-3}	56.67±2.68 (66.53)	33.47		28.33±0.76 (114.09)	34.65**
	0.6 μcm^{-3}	43.67±1.47 (51.27)	48.73	59.11**	29.86±0.47 (120.25)	
<i>N. sativa</i>	Control	85.17±6.01 (100)	-		24.83±0.54 (100)	
	0.2 μcm^{-3}	67.00±3.03 (78.66)	21.34		27.00±0.69 (108.73)	48.79**
	0.4 μcm^{-3}	53.88±2.26 (63.26)	36.74		29.00±0.57 (116.65)	
	0.6 μcm^{-3}	41.00±2.05 (48.13)	51.87	65.82**	32.00±0.34 (128.87)	

Values in parentheses indicate per cent change with respect to control taken as 100%; *Inhibition rate (IR) = $[(Cn - Tn) / Cn] \times 100$, Cn = number of adults emerged in control and Tn = number of adults emerged in test; **Significant at $P < 0.05$ (df = 3,20)

Table 8. Post Hoc Tukey's Honestly Significant Difference (HSD) analysis of effect of *C. tamala* and *N. sativa* volatile oils on development of *S. zeamais*

Oil	Progeny emergence				Developmental period			
	Pair wise comparison	HSD _{0.05} = 9.1580 HSD _{0.01} = 11.6097	Q _{0.05} = 3.9583 Q _{0.01} = 5.0180		Pair wise comparison	HSD _{0.05} = 1.4842 HSD _{0.01} = 1.8773	Q _{0.05} = 3.9419 Q _{0.01} = 4.9859	
<i>C. tamala</i>	T ₁ :T ₂ M ₁ = 85.17 M ₂ = 69.83	15.33	Q = 6.63*		T ₁ :T ₂ M ₁ = 24.83 M ₂ = 26.50	1.67	Q = 4.43*	
	T ₁ :T ₃ M ₁ = 85.17 M ₃ = 56.67	28.50	Q = 12.32*		T ₁ :T ₃ M ₁ = 24.83 M ₃ = 28.33	3.50	Q = 9.30*	
	T ₁ :T ₄ M ₁ = 85.17 M ₄ = 43.67	41.50	Q = 17.94*		T ₁ :T ₄ M ₁ = 24.83 M ₄ = 29.86	5.02	Q = 13.34*	
	T ₂ :T ₃ M ₂ = 69.83 M ₃ = 56.67	13.17	Q = 5.69*		T ₂ :T ₃ M ₂ = 26.50 M ₃ = 28.33	1.83	Q = 4.87*	
	T ₂ :T ₄ M ₂ = 69.83 M ₄ = 43.67	26.17	Q = 11.31*		T ₂ :T ₄ M ₂ = 26.50 M ₄ = 29.86	3.36	Q = 8.92*	
	T ₃ :T ₄ M ₃ = 56.67 M ₄ = 43.67	13.00	Q = 5.62*		T ₃ :T ₄ M ₃ = 28.33 M ₄ = 29.86	1.52	Q = 4.05*	

*Significant ($P < 0.05$) difference between all the treated and control groups.

Table 8. Continues

Oil	Progeny emergence				Developmental period		
	Pair wise comparison	HSD _{0.05} = 9.9424 HSD _{0.01} = 12.6359	Q _{0.05} = 3.9766 Q _{0.01} = 5.0539	Pair wise comparison	HSD _{0.05} = 1.7266 HSD _{0.01} = 2.1889	Q _{0.05} = 3.9583 Q _{0.01} = 5.0180	
N. sativa	T ₁ :T ₂ M ₁ = 85.17 M ₂ = 68.33	16.80	Q = 6.73*	T ₁ :T ₂ M ₁ = 24.83 M ₂ = 27.00	2.17	Q = 4.97*	
	T ₁ :T ₃ M ₁ = 85.17 M ₃ = 54.33	30.83	Q = 12.33*	T ₁ :T ₃ M ₁ = 24.83 M ₃ = 29.00	4.17	Q = 9.55*	
	T ₁ :T ₄ M ₁ = 85.17 M ₄ = 42.60	42.57	Q = 17.03*	T ₁ :T ₄ M ₁ = 24.83 M ₄ = 32.00	7.17	Q = 16.43*	
	T ₂ :T ₃ M ₂ = 68.33 M ₃ = 54.33	14.00	Q = 5.60*	T ₂ :T ₃ M ₂ = 27.00 M ₃ = 29.00	2.00	Q = 4.58*	
	T ₂ :T ₄ M ₂ = 68.33 M ₄ = 42.60	25.73	Q = 10.29*	T ₂ :T ₄ M ₂ = 27.00 M ₄ = 32.00	5.00	Q = 11.46*	
	T ₃ :T ₄ M ₃ = 54.33 M ₄ = 42.60	11.73	Q = 4.69*	T ₃ :T ₄ M ₃ = 29.00 M ₄ = 32.00	3.00	Q = 6.88*	

*Significant ($P < 0.05$) difference between all the treated and control groups.

Antifeedant activity

Both *C. tamala* and *N. sativa* oils inhibited feeding in *S. zeamais* adults (Table 9). Both oils decreased consumption of flour disc by *S. zeamais* adults. Antifeedant activity was reduced to 75.47 and 79.36% with respect to control at 80% of 96h-LC₅₀ of *C. tamala* and *N. sativa* oil respectively (For *C. tamala* oil $F = 261.48$; For *N. sativa* $F = 703.46$; $P < 0.01$; Table 9) Post Hoc Tukey's Honestly Significant Difference (HSD) analysis shows significant ($P < 0.05$) difference between all the treated and control groups (Table 10).

Table 9. Antifeedant activity of *C. tamala* and *N. sativa* volatile oils against *S. zeamais*

Concentration	<i>C. tamala</i>		<i>N. sativa</i>	
	Consumption of flour disc (mg) (Mean±SD)	AFA*	Consumption of flour disc (mg) (Mean±SD)	AFA*
Control	11.29±0.15 (100)	-	11.29±0.15 (100)	-
40% of 96h-LC ₅₀	6.49±0.24 (57.48)	42.51	5.60±0.23 (49.61)	50.39
80% of 96h-LC ₅₀	2.77±0.12 (24.53)	75.46	2.33±0.12 (20.64)	79.36
	$F = 261.48^{**}$		$F = 703.46^{**}$	

Values in parentheses indicate per cent change with respect to control taken as 100%;

*Antifeedant activity was calculated using $AFA = [C-T/C] \times 100$; Where, C = consumption of flour disc in control group, and T = consumption of flour disc in treated group. Six replicates were set for each concentration of oil and control; **Significant at $P < 0.05$ ($df = 2, 15$)

Table 10. Post Hoc Tukey's Honestly Significant Difference (HSD) analysis of antifeedant activity of *C. tamala* and *N. sativa* volatile oils against *S. zeamais*

Oil	Pair wise comparison		HSD _{0.05} = 0.9700	Q _{0.05} = 3.6734
			HSD _{0.01} = 1.2769	Q _{0.01} = 4.8359
<i>C. tamala</i>	T ₁ :T ₂	M ₁ = 11.29	4.80	Q = 18.18*
		M ₂ = 6.49		
	T ₁ :T ₃	M ₁ = 11.29	8.52	Q = 32.25*
		M ₃ = 2.77		
	T ₂ :T ₃	M ₂ = 6.49	3.72	Q = 14.07*
		M ₃ = 2.77		
<i>N. sativa</i>	T ₁ :T ₂	M ₁ = 11.29	5.69	Q = 33.29*
		M ₂ = 5.60		
	T ₁ :T ₃	M ₁ = 11.29	8.96	Q = 52.41*
		M ₃ = 2.33		
	T ₂ :T ₃	M ₂ = 5.60	3.27	Q = 19.12*
		M ₃ = 2.33		

*Significant ($P < 0.05$) difference between all the treated and control groups.

Volatile oils of plant origin have been well known for their insecticidal properties against several insect pests including those of stored grains [46-52]. These volatile oils show repellent, contact toxicity, fumigant toxicity, oviposition inhibitory and developmental inhibitory activities against a variety of coleopteran insect of stored grains including rice weevil, *S. zeamais*. These volatile oils cause mortality in insects as these have been reported to inhibit activity of acetylcholine esterase [53,54]. Besides crude and complete oils, its individual constituents also cause repellency, contact toxicity, fumigant toxicity, oviposition inhibition and developmental inhibition in insects. Linalool, linalyl acetate, menthol, methonene, limonene, α -pipene, β -pipene, β -caryophyllene and linalool have been shown to cause toxicity in *S. zeamais* [46,55].

In this study, repellent, toxic, oviposition inhibitory, developmental inhibitory and feeding inhibitory activities of *C. tamala* and *N. sativa* oils were studied in *S. zeamais*. Both *C. tamala* and *N. sativa* oils repel and cause mortality in *S. zeamais* adults. The rapid mortality caused shows neurotoxic mode of action of the two oils under investigation. Similar results have also been obtained with *Piper cubeba*, *Zingiber officinale* and other oils in *S. zeamais* adults [56]. Several other volatile oils and its pure compounds have also been reported to inhibit acetylcholine esterase activity causing paralysis and death in insects [46,53,54]. These oils interfere with neuromodulator octopamine or GABA-gated chloride channels [57,58]. The octopaminergic system of insects uses octopamine as neurotransmitter, neurohormone and neurohormone-neuromodulator. Disruption in its activity breaks down the nervous system in insects [59].

Oviposition and progeny production was reduced *S. zeamais* when fumigated with *C. tamala* and *N. sativa*. Volatile oils reduce oviposition potency of insects probably by disrupting mating and sexual communication in *S. zeamais* adults. These two volatile oils inhibit development of juvenile phases and increase developmental period in *S. zeamais*. Reduced adult emergence may be due to the death of egg and larval phases while delay in development may be due to inhibition of metabolic processes or disturbances in hormonal networking and tissue responsiveness. Both *C. tamala* and *N. sativa* oils reduce feeding potency in *S. zeamais*. This reduction may be due to repellent behaviour of these oils. Similar results have been obtained with *Allium sativum*, *Anthum graveolense*, *P. cubeb* and *Z. officinale* oils in *S. zeamais* [56,60]. The rapid mode of action of the two

oils under investigation shows their low persistence in the environment. The persistence of the insecticidal activity depends on the chemical composition of volatile oils and nature and position of the functional groups in oil constituents [61]. Volatile oils possessing high content of hydrogenated compound loss their activity as compared to those containing high content of oxygenated compounds [62,63].

Further studies have also been carried out to explore insecticidal activity of oil's constituents and its role in antagonistic and synergistic relationship [64,65]. It must be kept in mind that oils/constituents should be effective against targets only but not against non-targets including humans. Issues like risk associated to users, mode of exposure, degradation in the environment and chronic toxicity should also be considered and evaluated for compatible use of volatile oils for insect pest management.

CONCLUSION

The present study reports that *C. tamala* and *N. sativa* oils possess repellent, toxic as well as oviposition, developmental and feeding inhibitory activities against *S. zeamais*. These oils act at multiple target sites which minimize the possibility of resistance development. Since *C. tamala* and *N. sativa* are the parts of our food, these oils are safe for humans. For these reasons, *C. tamala* and *N. sativa* oils can be considered as a natural alternative in the management of stored grains insects.

Conflict of Interest. The authors declared that there is no conflict of interest.

Authorship Contributions. Concept: M.K.C., N.K., Design: M.K.C., N.K., Data Collection or Processing: M.K.C., N.K., Analysis or Interpretation: M.K.C., N.K., Literature Search: M.K.C., N.K.

Financial Disclosure. Author is highly thankful to Department of Higher Education, Uttar Pradesh, India for financial assistance [No. 46/2021/603/70-4-2021-4(56)/2020] to carry out this research.

REFERENCES

- [1] World Meteorological Organization (WMO): Scientific assessment of ozone depletion. Geneva (Switzerland). 1991; Report No. 25.
- [2] Lu, F.C. (1995): A review of the acceptable daily intakes of pesticides assessed by the World Health Organization. *Regulatory Toxicology and Pharmacology* 21: 351-364.
- [3] United Nations Environment Programme (UNEP): The Montreal Protocol on substances that deplete the ozone layer. Nairobi (Kenya). 2000.
- [4] Alavanja, M.C.R., Bonner, M.R. (2012): Occupational pesticide exposures and cancer risk: a review. *Journal of Toxicology and Environmental Health B* 15, 238-263.
- [5] EEA; Late lessons from early warnings: science, precaution, innovation. European Environment Agency, Report No 1/2013. EEA, Copenhagen, 2013.
- [6] Köhler, H.R., Triebkorn, R. (2013): Wildlife ecotoxicology of pesticides: can we track effects to the population level and beyond? *Science* 341, 759-765.
- [7] World Health Organization (WHO): Agrochemicals, health and environment: directory of resources, 2017.
- [8] Atti-Santos, A.C., Panseira, M.R., Paroul, N., Atti-Serafini, L., Moyna, P.P. (2004): Seasonal variation of essential oil yield and composition of *Thymus vulgaris* L. (Lamiaceae) from south Brazil. *Journal of Essential Oil Research* 16, 294-295.
- [9] Angioni, A., Barra, A., Coroneo, V., Dessi, S., Cabras, P. (2006): Chemical composition, seasonal variability and antifungal activity of *Lavandula stoechas* L. ssp. *stoechas* essential oils from stem/ leaves and flowers. *Journal of Agricultural Food Chemistry* 54, 4364-4370.

- [10] Verma, R.S., Verma, R.K., Yadav, A.K. (2011): Seasonal variation in essential oil content and composition of Thyme, *Thymus serpyllum* L. cultivated in Uttarakhand hills. Indian Journal of Pharmacological Sciences 341, 233-235.
- [11] Caballero-Gallardo, K., Olivero-Verbel, J., Stashenko, E.E. (2011): Repellent activity of essential oils and some of their individual constituents against *Tribolium castaneum* Herbst. Journal of Agricultural Food Chemistry 59, 1690-1696.
- [12] Isman, M.B., Miresmaili, S., Machial, C. (2011): Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. Phytochemical Review 10, 197-204.
- [13] Liu, Z.L., Chu, S.S., Jiang, G.H. (2011): Insecticidal activity and composition of essential oil of *Ostericum sieboldii* (Apiaceae) against *Sitophilus zeamais* and *Tribolium castaneum*. Records of Natural Products 5, 74-81.
- [14] Stefanazzi, N., Stadler, T.A., Ferrero, A. (2011): Composition and toxic, repellent and feeding deterrent activity of essential oils against the stored-grain pests *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae). Pest Management Science 67, 639-646.
- [15] Chaubey, M.K. (2022): Study of insecticidal activities of *Aegle marmelos* (Rutaceae) essential oil against pulse beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). Biopesticide International 18(2):145-154.
- [16] Quattrocchi U. (2016): *CRC World Dictionary of Medicinal and poisonous plants: Common names, Scientific names, Eponyms and Etymology (5 Volume set)*. CRC Press. pp. 959. ISBN 978-1-4822-5064-0.
- [17] Chowdhury J.U., Shaha, G.C., Begum, F., Bhuiyan, M.N.H., Rahim, M. (2013): Essential oil composition from fresh and dried leaves of *Cinnamomum tamala*. Bangladesh Journal of Scientific and Industrial Research 48(2), 151-154.
- [18] Osawa, T., Namaki, M. (1983): A novel type antioxidant isolated from leaf wax of Eucalyptus leaves. Agricultural Biological Chemistry 45, 735-739.
- [19] Miller, N.J., Rice-Evans, C.A. (1997): Factors affecting the antioxidant activity determined by ABTS⁺ radical cation assay. Free Radicals Research 26, 195-199.
- [20] Vardar Unlu, G., Sokmen, A., Domez Eand Tepe, B. (2003): Antimicrobial activity of essential oil from *Cinnamomum tamala* (Lauraceae). Journal of Agricultural Food Chemistry 51: 63-67.
- [21] Rao, C.V., Vijayakumar, M., Sairam K, Kumar V. (2008): Antidiarrhoeal activity of the standardised extract of *Cinnamomum tamala* in experimental rats. Journal of Natural Medicine 62(4):396-402.
- [22] Lima, Z.P., Severi, J.A., Pellizzon, C.H., Brito, A.R., Solis, P.N. (2010): Gastroprotective activity of *Cinnamomum tamala* leaves on experimental gastric ulcers in rats. Journal of Ethnopharmacology 128(2010): 537-540.
- [23] Shahwar, D., Ullah, S., Khan, M.A., Ahmad, N., Saeed, A., Ullah S. (2015): Anticancer activity of *Cinnamon tamala* leaf constituents towards human ovarian cancer cells. Pakistan Journal of Pharmacological Science 28(3), 969-972.
- [24] Taha, A.M., Eldahshan, O.A. (2017): Chemical characteristics, antimicrobial and cytotoxic activities of the essential oil of Egyptian *Cinnamomum glanduliferum* Bark. Chemical Biodiversity 14(5).
- [25] Chopra, R.N., Nayer, S.L., Chopra, I.C. (1956): Glossary of Indian Medicinal Plants, Council of Scientific and Industrial Research, New Delhi. 65-66.
- [26] Yusuf, M., Haque, N., Begum, J., Chowdhury, J.U. (1994): Medicinal Plants of Bangladesh. BCSIR Laboratories, Chittagong-4220, Bangladesh. 164.
- [27] Sharma, S.R., Dwivedi, S.K., Swarup, D. (1996): Hypoglycaemic and Hypolipidaemic Effects of *Cinnamomum tamala* Nees Leaves. Journal of Experimental Biology 34, 372.
- [28] Sharma, P.C., Yelne, M.B., Dennis, T.J. (2005): *Database on medicinal plants used in Ayurveda*. New Delhi, pp. 420-440.
- [29] Sahebkar, A., Beccuti, G., Simental-Mendía, L.E., Nobili, V., Bo, S. (2016): *Nigella sativa* (black seed) effects on plasma lipid concentrations in humans: A systematic review and meta-analysis of randomized placebo-controlled trials. Pharmacological Research 106, 37-50.

- [30] Abdelmeguid, N.E., Fakhoury, R., Kamal, S.M., Al Wafai, R.J. (2010): Effects of *Nigella sativa* and thymoquinone on biochemical and subcellular changes in pancreatic β -cells of streptozotocin-induced diabetic rats. *Journal of Diabetes* 2(4), 256-266.
- [31] Bakathir, H.A., Abbas, N.A. (2011): Detection of the antibacterial effect of *Nigella sativa* ground seeds with water. *African Journal of Traditional, Complementary and Alternative Medicines* 8(2), 159-164.
- [32] Ezz, H.S., Khadrawy, Y.A., Noor, N.A. (2011): The neuroprotective effect of curcumin and *Nigella sativa* oil against oxidative stress in the pilocarpine model of epilepsy: a comparison with valproate. *Neurochemical Research* 36(11):2195-2204.
- [33] Mahmoud, S.S., Torchilin, V.P. (2012): Hormetic/cytotoxic effects of *Nigella sativa* seed alcoholic and aqueous extracts on MCF-7 breast cancer cells alone or in combination with doxorubicin. *Cell Biochemistry and Biophysics* 25(7), 1392-1398.
- [34] Umar, S., Zargan, J., Umar, K., Ahmad, S., Katiyar, C.K., Khan, H.A. (2012): Modulation of the oxidative stress and inflammatory cytokine response by thymoquinone in the collagen induced arthritis in Wistar rats. *Chemico-Biological Interactions* 197(1), 40-46.
- [35] Enomoto, S., Asano, R., Iwahori, Y., Narui, T., Okada, Y., Singab, A.N. (2001): Hematological studies on black cumin oil from the seeds of *Nigella sativa* L. *Biological and Pharmaceutical Bulletin* 24, 307-310.
- [36] Tembhurne, S.V., Feroz, S., Sakarkar, D.M. (2014): A review on therapeutic potential of *Nigella sativa* (kalonji) seeds. *Journal of Medicinal Plants Research* 8, 166-167.
- [37] Demissie, G., Tefera, T., Tadesse, A. (2008): Efficacy of SilicoSec, filter cake and wood ash against the maize weevil, *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae) on three maize genotypes. *Journal of Stored Product Research* 44, 227-231.
- [38] Ileleji, K.E., Maier, D.E., Woloshuk, C.P. (2007): Evaluation of different temperature management strategies for suppression of *Sitophilus zeamais* (Motschulsky) in stored maize. *Journal of Stored Product Research* 43, 480-488.
- [39] Ellman, G.L., Courtney, K.D., Andres, Jr. V., Featherstone, R.M. (1961). A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 1961, 7, 88-95.
- [40] Vanmathi, J.S., Padmalatha, C., Singh, A.J.A.R., Charman, K. (2012). Effect of chosen botanicals on the oviposition deterrence and adult emergence of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *Elixir Biological Technology* 51A, 11120-11123.
- [41] Tapondju, L.A., Alder, A., Fontem, H., Fontem, D.A. (2002): Efficacy of powder and essential oil from the *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six stored products beetles. *Journal of Stored Product Research* 38, 395-402.
- [42] Sithisut, D., Fields, P.G., Chandrapathya, A. (2011): Contact toxicity, feeding reduction and repellency of essential oils from three plants from the ginger family (Zingiberaceae) and their major components against *Sitophilus zeamais* and *Tribolium castaneum*. *Journal of Stored Product Research* 104, 1445-54.
- [43] Russel, R.M., Robertson, J.L., Savin, S.A. (1977): POLO: A new computer programme for probit analysis. *Bulletin of Entomological Research* 23, 209-213.
- [44] Sokal, R.R., Rohlf, F.J. (1973): Introduction to biostatistics. W.H. Freeman and Co, San Francisco, CA, USA, 185-207.
- [45] <https://www.socscistatistics.com/tests/anova/default2.aspx>.
- [46] Chaubey, M.K. (2012a): Fumigant toxicity of essential oils and pure compounds against *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *Biological Agriculture Horticulture* 28, 111-119.
- [47] Chaubey, M.K. (2012b): Responses of *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae) against essential oils and pure compounds. *Herba Polonica* 58, 33-45.
- [48] Chaubey, M.K. (2012c): Biological effects of essential oils against Rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *Journal of Essential Oil Bearing Plants*, 15, 809-815.

- [49] Chaubey, M.K. (2013): Insecticidal effects of *Allium sativum* (Alliaceae) essential oil against *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Journal of Biologically Active Products from Nature* 3, 248-258.
- [50] Chaubey, M.K. (2014): Biological activities of *Allium sativum* essential oil against pulse beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Herba Polonica* 60, 41-55.
- [51] Chaubey, M.K. (2016a): Insecticidal activities of *Cinnamomum tamala* (Lauraceae) essential oil against *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *International Journal of Entomological Research* 4, 91-98.
- [52] Chaubey, M.K. (2016b): Fumigant toxicity of *Allium sativum* (Alliaceae) essential oil against *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *Entomology Applied Science Letters* 3, 43-48.
- [53] Chaubey, M.K. (2017a): Evaluation of insecticidal properties of *Piper nigrum* and *Cuminum cyminum* essential oils against *Sitophilus zeamais*. *Journal of Entomology* 14, 148-154.
- [54] Chaubey, M.K. (2017b): Study of insecticidal properties of *garlic*, *Allium sativum* (Alliaceae) and *bel*, *Aegle marmelos* (Rutaceae) essential oils against *Sitophilus zeamais* L. (Coleoptera: Curculionidae). *Journal of Entomology* 14, 191-198.
- [55] Ogendo, J.O., Kostyukovsky, M., Ravid, U., Matasyoh, J.C., Deng, A.L., Omolo, E.O., Kariuki, S.T., Shaaya, E. (2008): Bioactivity of *Ocimum gratissimum* L. oil and two constituents against five insect pests attacking stored food products. *Journal of Stored Product Research* 44, 328-334.
- [56] Chaubey, M.K.; Kumar, N. (2021). Role of *Piper cubeba* and *Zingiber officinale* volatile oils in maize weevil, *Sitophilus zeamais* management. *International Journal of Green and Herbal Chemistry* 10(4):404-417.
- [57] Enan, E.E. (2005): Molecular and pharmacological analysis of an octopamine receptor from American cockroach and fruit fly in response to plant essential oils. *Archives of Insect Biochemistry and Physiology* 159, 161-171.
- [58] Tong, F., Coats, J.R. (2012): Quantitative structure-activity relationship of monoterpenoid binding activities to the house flies GABA receptor. *Pest Management Science* 68, 1122-1129.
- [59] Hollingworth, R.M., Johnstone, E.M., Wright, N. (1984): Pesticide Synthesis through Rational Approaches. ACS Symposium Series No. 255, American Chemical Society, Washington, DC. 103-125.
- [60] Chaubey, M.K., Kumar, N. (2022): Insecticidal properties of *Allium sativum* and *Anethum graveolens* essential oils against maize weevil, *Sitophilus zeamais* (Motschulsky). *International Journal of Green and Herbal Chemistry* 11(1):60-74.
- [61] Kumbhar, P.P., Dewang, P.M. (2001): Monoterpenoids: The natural pest management agents. *Fragrance and Flavour Association of India* 3, 49-56.
- [62] Huang, Y., Ho, S.H. (1998): Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Product Research* 34, 11-17.
- [63] Regnault-Roger, C., Philigene, B. J., Vincent, C. (2000): *Biopesticides d'origine vegetales*. Paris, 337.
- [64] Kordali, S., Aslan, I., Almasur, O.C., Cakir, A. (2006): Toxicity of essential oils isolated from three *Artemisia* species and some of their major components to granary weevil, *Sitophilus granarius* (L.) (Coleoptera: Curculionidae). *Industrial Crops and Products* 23, 162-170.
- [65] Fields, P., Woods, S.M., Taylor, W. (2010): Triterpenoid saponins synergize insecticidal pea peptides: effect on feeding and survival of the rice weevil, *Sitophilus oryzae*. *Canadian Entomology* 142, 501-512.