

‘Formulation and Evaluation of Carica Papaya Leaf Extract Herbal Syrup For The Treatment of Dengue,,

Himanshu patidar,Mamta ahirwar, Satyaendra Shrivastava

Parijat college of pharmacy, Indore (M.P) India

Abstract:-

The present study deals with the formulation and evaluation of herbal syrup prepared from *Carica papaya* leaf extract for dengue management. Dengue is a mosquito-borne viral disease that leads to a decrease in platelet count. Papaya leaf extract is known for its platelet enhancing activity. The prepared syrup was evaluated for organoleptic properties, pH, viscosity and stability. The results showed that the formulation was stable, safe and suitable for oral use.

Keywords: Carica papaya, Dengue, Herbal syrup, Platelet count

Introduction :-

Dengue is a viral disease transmitted by *Aedes aegypti* mosquito. It is one of the major public health problems in tropical countries. The disease is characterized by high fever, headache, muscle pain and decreased platelet count.

Herbal medicines are gaining importance due to their safety and effectiveness. *Carica papaya* leaves contain phytochemicals such as flavonoids, alkaloids and phenolic compounds which help in increasing platelet count and boosting immunity.

The present study aims to formulate and evaluate a herbal syrup using papaya leaf extract for dengue treatment.

Aim:-

To formulate and evaluate a herbal syrup of *Carica papaya* leaf extract for the supportive treatment of dengue.

Carica papaya :-

Carica papaya, commonly known as papaya, is a well-known medicinal plant belonging to the family Caricaceae. The leaves of the plant are widely used in traditional and herbal medicine due to their significant therapeutic properties. The dried and powdered form of papaya leaves, commonly referred to as Cerica Papaya Powder, is used as an active ingredient in various herbal formulations.

Papaya leaf powder contains several important bioactive constituents such as alkaloids (carpaine), flavonoids, phenolic compounds, enzymes (papain and chymopapain), and vitamins. These constituents are responsible for its pharmacological activities including antioxidant, anti-inflammatory, and immunomodulatory effects.

2. MATERIALS AND METHODS

2.1Experimental Requirements

Experimental requirement Papaya Powder were obtained from marotiya bazar .

2.2 List of Equipment

Ostwald viscometer, sieve, pH meter, mixer grinder, and heating mantle were used.

2.3 Preparation of Formulation

Extraction (Decoction Method):

The powder was taken and boiled with distilled water. After boiling, it was cooled and filtered to get a clear extract.



Fig. 1: Dry Powder of carica papaya



Fig.2 Decotion method

Formulation of Syrup

The extraction of Carica papaya leaves is carried out to obtain the active phytoconstituents likeFlavonoids, alkaloids (especially carpaine), and other therapeutic compounds that are useful in theTreatment of dengue and related ailments. Fresh or shade-dried papaya leaves are collected aWashed thoroughly to remove dust and impurities. The leaves are then either crushed directly (inThe case of juice extraction) or ground into a coarse powder if dried. In aqueous extraction, about50 grams of powdered leaves are added to 500 ml of distilled water and heated on a water bath at60–70°C for about 2 hours with occasional stirring. The mixture is then cooled and filtered usingMuslin cloth or Whatman filter paper to obtain a clear green extract. The filtrate may beConcentrated by gentle heating or left to evaporate naturally under shade. In ethanolic extraction, the powdered leaves are soaked in 70% ethanol for 48–72 hours in a closed container withOccasional shaking. After maceration, the extract is filtered and the ethanol is evaporated using aRotary evaporator or gentle heating to yield a concentrated extract. The final extract is stored in anAmber-colored airtight container in a cool and dry place to preserve its medicinal properties. Thisextract can be further used for pharmacological testing or formulation into syrups, tablets,

Table No.1: Formulation of Convolvulus pluricaulisAsparagus racemosus Herbal syrup

Sr. No.	Ingredients	BatchA	BatchB	BatchC
1	Papaya leafextract	5 ml	10 ml	15 ml
2	Simple syrup	5 ml	10 ml	15 ml
3	Glycerin	1.5 ml	1.5 ml	1.5 ml
4	Honey	3 ml	5 ml	8 ml
5	Sodium Benzoate	0.1%	0.1%	0.1%
6	Guar Gum	0.5 gm	0.5 gm	0.5 gm
7	Lemon grass Oil	2-3 drops	2-3 drops	2-3 drops
8	Distilled Water	q.s	q.s	q.s

2.4.1 Organoleptic Property

Prepared syrup of *Asparagus racemosus* and *Convolvulus pluricaulis* was observed for its colour, appearance, odour and taste. The syrup was light brown in colour, clear in appearance, having characteristic odour and sweet taste.

2.4.2 Flavonoids Test

1 ml of the herbal extract was taken in a test tube and 2 ml of 1% sodium hydroxide solution was added. Formation of yellow colour indicated the presence of flavonoids.

2.4.3 Determination of pH

The pH of the prepared syrup was determined using a digital pH meter. The readings were taken three times and average value was noted.

2.4.4 Solubility

The prepared syrup was checked for its miscibility with water. It was found to be completely miscible without any separation.

2.4.5 Determination of Density

Density of the syrup was determined using a specific gravity bottle. The bottle was cleaned, dried and weighed. Then it was filled with syrup and weighed again to calculate density.

2.4.6 Viscosity

Viscosity of the syrup was determined using an Ostwald viscometer. The instrument was cleaned properly before use and flow time of the syrup was recorded.

2.4.7 Stability Study

The prepared syrup was stored at $40 \pm 2^{\circ}\text{C}$ and $75 \pm 5\%$ RH. The samples were checked at different intervals (0, 7, 14 and 21 days) for changes in colour, odour and taste. No significant changes were observed.

RESULTS AND DISCUSSION Evaluation parameters

Features	F1	F2	F3
Colour	Light Brown	Light Brown	Light Brown
Odor	Characteristics	Characteristics	Characteristics
Test	Sweet	Sweet	Sweet

Table No.2. Organoleptic characteristics features of syrup



Fig.No.3:Estimation of Ph



Fig.No.4: Determination of viscosity



Fig.No.5: Determination of density



Fig.No.6: Solubility test

Features	F1	F2	F3
Ph	5.8	5.9	5.7
Solubility	Soluble in methanol and water	Soluble in methanol and water	Soluble in methanol and water
Density	0.82g	0.84g	0.79g
Viscosity	3.64cp	3.62cp	3.66cp

Table No.3. Estimation of pH, solubility, Density & Viscosity

Summary

The present study focuses on the formulation and evaluation of a herbal syrup containing Carica papaya leaf extract for the treatment of dengue. Carica papaya leaves are well known for their platelet-enhancing and immunomodulatory properties, which are beneficial in managing dengue fever. The syrup was prepared using papaya leaf extract along with suitable excipients such as sucrose, glycerin, honey, and preservatives to improve stability, taste, and patient compliance.

The formulated syrup was evaluated for various physicochemical parameters like pH, viscosity, color, taste, and stability. The results indicated that the formulation was stable, palatable, and within acceptable limits. The herbal constituents present in the formulation contribute to increasing platelet count and reducing symptoms associated with

Conclusion

The developed Carica papaya leaf extract herbal syrup showed satisfactory results in terms of formulation and evaluation parameters. It can be concluded that the syrup is safe, effective, and suitable for use as a supportive treatment in dengue management. The natural herbal formulation offers advantages such as fewer side effects, better patient acceptability, and cost-effectiveness.

Further clinical studies are recommended to confirm its therapeutic efficacy and to establish it as a standard herbal treatment for dengue

Uses (Extended)

1. Carica papaya (Papaya Leaf)

- Increases platelet count (anti-thrombocytopenic)
- Supports immune system function
- Helps in recovery from dengue fever
- Reduces body pain and inflammation
- Improves digestion and liver function
- Acts as antioxidant (reduces oxidative stress)

References

1. J.K.K. Asamoah, E. Okyere, et al, Optimal control & comprehensive Cost- effective analysis For COVID-19 (2022) 105177 .
2. M.A. Mamun, J.M. Misti, M.D. Griffiths, D. Gozal, et al, The dengue epidemic in Bangladesh: Risk factors and actionable items, *Lancet* 394 (10215) (2019) 2149– 2150.
3. B. Gakkhar S, Aziz- Mishra A, Ambrosio Alaoui MA. Et al, A network mode for control of Dengue epidemic using sterile insect technique. *Math Biosci Eng* 2018
4. Boonchutima S, Kachentawa K, Limpavithayakul M, Prachansri A. Et al Longitudinal study of Thai people media exposure, knowledge, and behavior on dengue fever prevention and control. *J Infect Public Health* 2017
5. S. Khaleda, Q.A. Mowla, Y. Murayama, Dhaka et al, metropolitan, in: *Urban Development in Asia and Africa Urban Book Series*, Springer, Singapore, 2017
6. Singhai A, Juneja V, Abbas S., Jha R.K et al, The effect of Carica papaya leaves extract capsules On platelets count and hematocrit levels in acute febrile illness with thrombocytopenia patient, *International Journal of Medical Research & Health Sciences*, 2016, Vol. 5(1), 254-257
7. Krzywinska E, Kokoza V, Morris M, de la Casa-Esperon E, Raikhel AS, Krzywinski et al, The sex locus is tightly linked to factors conferring sex-specific lethal effects in the mosquito *Aedes Aegypti*. *Heredity (Edinb)*. 2016
8. Mukherjee K Pulok, Nema K Neelesh, Bhadra Santanu, Mukhrjee D, et al, Braga C Fernao, Matsabisa G Motiale, Immunomodulatory leads from medicinal plants, *Indian Journal of Traditional Knowledge*, Vol. 13 (2), April 2014, 235-256
9. Massonnet Bruneel B, Corre-Catelin N, Lacroix R, Lees RS, Hoang KP, Nimmo et al. Fitness Of transgenic mosquito *Aedes aegypti* males carrying a dominant lethal genetic system *One* 2013
10. Kokate K. C., Purohit P.A., Gokhale B.S, et al, *Book of Pharmacognosy*, Nirali Prakashan Publisher, 48th edition, January 2013, 10.5
11. Kunle, Folashade O., Omoregie H. And Ahmadu, Egharevba, Ochogu P., et al Standardization Of herbal medicines – A review, *International Journal of Biodiversity and Conservation*, March 2012, Vol. 4(3), 101-112
12. Romasi F., Karina J., Parhusip N. J. A et al, antibacterial activity of papaya leaf extracts Against pathogenic bacteria, *Makara Journal Of Technology*, November 2011, Vol.15 (2), 173-177 (2014): 281508.
- 13 .Revised and Expanded et al, New Delhi: WHO; 2011. World Health Organization. Comprehensive Guidelines for Prevention and Control of Dengue and Dengue Haemorrhagic Fever.