



An Integrated Approach to Harness the Anti-inflammatory Potential of *Achyranthes aspera* Linn

Devi Datt Joshi^{1,*}, Bharat G. Somkuwar², Lokesh Deb³, Virendra Singh Rana⁴

¹Amity Herbal Consortium, Amity University Uttar Pradesh, Noida (UP), India.

²Institute of Bioresources and Sustainable Development, Node Mizoram A-1, C/o P. Lalthangzauva Building, Chawnga Road, Nursery Veng, Aizawl (Mizoram), India.

³Pharmacologist, Institute of Bioresources and Sustainable Development-Sikkim Centre, Department of Biotechnology, Government of India, 5th mile, Near Metro point, Tadong, Gangtok, Sikkim, India.

⁴Division of Agricultural Chemicals, Indian Agricultural Research Institute (ICAR), Pusa Campus, New Delhi. India.

How to cite this paper: Devi Datt Joshi, Bharat G. Somkuwar, Lokesh Deb, Virendra Singh Rana. (2023) An Integrated Approach to Harness the Anti-inflammatory Potential of *Achyranthes aspera* Linn. *International Journal of Food Science and Agriculture*, 7(1), 112-121. DOI: 10.26855/ijfsa.2023.03.016

Received: February 28, 2023

Accepted: March 26, 2023

Published: April 25, 2023

***Corresponding author:** Devi Datt Joshi, Amity Herbal Consortium, Amity University Uttar Pradesh, Noida (UP), India.

Abstract

Plants have been used to overcome pathologic conditions and body ailments since the pre-historic period. A few plants e.g. *Borago officinalis*, *Curcuma longa*, *Rosmarinus officinalis*, *Zingiber officinale*, etc. are frequently used for treating inflammatory disorders, globally. Inflammation is a complex biological response of vascular tissue to different harmful stimuli and results in discomfort to the body. There is not a single medicine to treat all the inflammation at the time. In recent years plant-based medicines are evaluated for efficacy through phytopharmacology, molecular mechanisms, and clinical trials. The therapeutic ingredients of polyherbal formulations generally contain different chemical classes, and their efficacy is based on synergism. *Achyranthes aspera* is an abundantly available herb, globally, has been in use in Ayurveda, Siddha, and Unani system of medicine to cure many ailments like arthritis, asthma, cardiac disease, dermatological diseases, gynecological disorder, kidney stone, leprosy, wound, snakebite, malaria, pneumonia, dysentery, rabies, toothache, etc. The applications of *A. aspera* in pathological investigations have scientifically justified the traditional utilization. Further, phytopharmacological studies have highlighted its synergistic mechanism of action and revealed its vital role in modulating inflammatory reactions for fast recovery with the least side effects, though there are a few investigations.

Keywords

Anti-inflammatory herbs, Traditional uses, Phytopharmacology, *Achyranthes aspera*, Synergistic combination

1. Introduction

Inflammatory diseases are the major cause of morbidity and mortality [1]. Inflammation is a complex response of the vascularized connective tissues to the exogenous and endogenous stimuli as a protective response to injury caused by microorganisms, noxious chemicals, or physical trauma to destroy or inactivate the foreign matter, and its elimination [2]. The process usually subsides on completion of healing but sometimes it turns severe, even if it may be worse than the disease and even maybe fatal [3, 4]. It starts with initiation, chemoattraction of inflammatory cells, and activation to release inflammatory mediators. Increased temperature, pain, redness, swelling, loss of activeness, etc. are the classical sign of inflammation [5]. Different inflammatory mediators such as cytokines, digestive enzymes, eicosanoids (prostaglandins and leukotriene B₄, C₄, D₄, E₄), histamine, kinins (bradykinin), narcosis factor, neuropeptides, nitric oxide,

platelet-activating factors, serotonin, tumors, biological metabolites, complement proteins, and adhesion factors play important role in the pathogenesis of inflammation [6]. Different cells like basophils, connective tissue fibroblasts, eosinophils, lymphocytes, mast cells, monocytes, neutrophils, resident macrophages, etc., are involved in the pathogenesis of inflammation [7]. Inflammations based on the severity of cases have been categorized as acute, and chronic [8-10].

Traditional herbal medicines have derived from the rich scientific heritage of different traditional healing practices. The utilization and pattern of application are different for different diseases [11]. Further, these medicinal herbs are in use with a successful history of thousands of years [12, 13].

Currently, the market of herbal drugs to treat inflammatory diseases constitutes 83%, globally. It is expected to be more than 95% in the forthcoming years as the increased acceptability of these preparations [14-17]. Generally, herbs are used to promote health and restore the body's ability to regulate the various functions such as self-healing, and protection, but safety is always a matter of concern. Drew and Myers (1997) have reported that about 40% of the anticancer drugs of plant origin have successfully treated cancer, but they still have serious side effects [18].

The social priority to use herbal medicine is because of the cultural acceptability, lesser side effects, and better compatibility with the human body [11]. Different chemical classes of herbal origin such as alkaloids, and rographolide, coumarin, flavonoids, xanthone, sterols, terpenoids, withaferin-A, etc., have been proved effective to treat the inflammations [1, 19]. Anti-inflammatory drugs of synthetic origin are classified as steroidal and nonsteroidal types, which have severe adverse effects on patients with other body ailments [20]. Currently, both doctors and patients are looking for herbal remedies to treat inflammation and low immunity problems. A study on different folklores and traditional practices for better and safe treatment, even in combination, using *A. aspera* is to be availed.

2. Material and Methods

The computerized database search engines, such as PubMed, ScienceDirect, Google Scholar, and MeSH were used to search the research publications on 'anti-inflammatory herbs'. The keywords for searching the scientific publications in these databases were 'anti-inflammatory herbs', 'herbals', 'folklore', 'traditional healing practices', and 'polyherbal formulation'. Search hits that appeared were further segregated using key terms like 'traditional applications', 'market demand', and 'OTC product'. All the retrieved references were further manually scanned to ensure a full research article. The relevant articles were short-listed, and key findings were retrieved, analysed, and systematically compiled in the form of a current review article.

3. Results and Discussion

The literature search on anti-inflammatory traditional medicinal plants has revealed the scientific heritage of different healing systems, and the usage of different plant species to overcome different inflammations. Further, it has been revealed that the current market of herbal drugs to treat inflammatory diseases is expected to increase by 12% in forthcoming years [14-17]. The use of different plants to treat inflammation, promote health, and restore the body's ability has been recorded (Table 1). More recently, a trend to investigate anti-inflammatory herb for analgesic properties, isolation of active ingredient, dose determination, toxic properties, etc. have been in practice for safe uses [11]. Molecular investigations have deciphered that different class of phytochemicals have a role in treating different types of inflammation [1].

3.1 *Achyranthes aspera* Linn

Achyranthes aspera Linn. (Apamarg), fam. Amaranthaceae has been used for its medicinal properties in Allopathic, Ayurvedic, Homeopathic, Siddha, Unani-Tibbi, Naturopathy, and many more folklore remedies [79]. Traditional healers claim that the addition of *A. aspera* would enhance the efficacy of any drug of plant origin [80]. In Ayurvedic toxicology, *A. aspera* plant is used for the treatment of poisoning which destroys it or is eliminated from the body [81]. Traditionally, it is used for the treatment of asthma, diabetes, dysentery, fever, hypertension, and malarial fever. Its root extract has a potent insect hormonal molting activity. The whole plant decoction has diuretic properties, and the aqueous extract is used to treat pneumonia. *A. aspera* dried herb is used to treat colic in children, and dry leaves are used to treat asthma. Different extracts of leaves have been reported for antiperoxidative, hypoglycemic, and thyroid-stimulating properties. In ancient Indian literature, it is claimed as a significant abortifacient. *A. aspera* is a primary constituent of an herbal formulation namely, Naayuruvi Kuzhi Thailum, a formulation of Siddha system of medicine. It is quite effective in the management of asthma. In different folklore, the inflorescence, fruits, seeds, and roots of *A. aspera* are used to treat hydrophobia [82]. In the present article, we want to generate innovative ideas based on the pool of evidence to develop an anti-inflammatory modern herbal medicine using the *A. aspera*.

3.2 Phyto-constituents

A. aspera has alkaloids (betaine, achyranthine), amino acids, flavonoids, phenolics saponins, steroids, and triterpeno-

ids, the different chemical classes known for anti-inflammatory properties [83, 84]. Achyranthine is a water-soluble alkaloid isolated from *A. aspera*. It decreases the blood pressure and heart rate, dilated blood vessels, and increased the rate and amplitude of respiration in dogs and frogs [85]. The traditional claims of anti-inflammatory potential of *A. aspera*, current knowledge of its chemical composition, and standardization of therapeutic parameters have an array for new modern green medicine.

Table 1. Traditional anti-inflammatory herbs and properties

S. No.	Description of plant	Traditional uses	Phyto-pharmacology	
			Active ingredients	Pharmacological properties
1.	<i>Borago officinalis</i> L. (Borage) Family.: Boraginaceae	Seeds oil is a dietary supplement [21-23], used to treat arthritis, autoimmune disorders eczema, diabetes, multiple sclerosis, heart diseases, and cancer [24, 25]	Seeds have 28% γ -linolenic acid (GLA), and it is the highest of all the borage species [26]	GLA is a potent food supplement [24, 26], do not have any side effects. It is prescribed as an anti-inflammatory drug, but its seeds have a miscarriage risk during pregnancy [27]
2.	<i>Curcuma longa</i> L. (Turmeric) Family: Zingiberaceae	Its rhizome is used to treat rheumatoid arthritis (RA) [28]	Curcuminoids [29]	The rhizome is used to treat irritable bowel syndrome (IBS) (Bundy et al., 2004), maintain the amelioration of ulcerative colitis [30], and psoriasis [31].
3.	<i>Elaeagnus angustifolia</i> L. (Oleaster) Family: Elaeagnaceae	It has analgesic effects. Fruits are used to cure asthma, fever, jaundice, tetanus, and rheumatoid arthritis [32]	Fruits are rich in proteins, sugar, vitamins, and minerals [32]	Aqueous extract of fruits is used to treat oral lichen planus (OLP) lesions [32], and it has anti-inflammatory properties [33, 34].
4	<i>Harpagophytum procumbens</i> (Burch.) DC. Ex Meisn (Devil's Claw) Family: Pedaliaceae	Roots are used to reduce swelling, pain, and inflammation [35].	Iridoid glycosides, harpagoside, and harpagide are the main ingredients [35].	To treat degenerative rheumatoid arthritis, osteoarthritis, tendonitis, kidney inflammation, and heart diseases [35-37]. Patients with ulcers, gallstone, and diabetes have been reported gastrointestinal upset, the side effect of <i>H. procumbens</i> [38].
5.	<i>Persea americana</i> Mill (Avocado) Family: Lauraceae	Fruits have excellent nutritional and medical value, and seed oil is edible [39].	Fats, fibres, proteins, vitamins, and minerals, but low sugar content. [39].	The use of avocado and soybean extracts together has been reported effective to treat mild to moderate vulvar lichen sclerosis (VLS) [20,40].
6.	<i>Oenothera biennis</i> L. (Evening Primrose) Family: Onagraceae	Seeds oil is used to cure rheumatic and arthritic problems, atopic dermatitis, psoriasis, premenstrual, and menopausal syndromes, and diabetic neuropathy [41].	The seed oil has ferulic, γ -linolenic acids, and linear aliphatic alcohols [41].	The seed oil has protective roles against pro-inflammatory agents [41], because of these reasons, it has more varied applications as an anti-inflammatory agent than the Borage oil [41].
7.	<i>Olea europaea</i> L. (Olive) Family: Oleaceae	Its oil is known as extra virgin olive oil (EVOO) [42]. It is a main source of fat in the Mediterranean diet.	Mono-unsaturated fatty acids, moderate amount of saturated and poly-unsaturated fatty acids, traces of phenols, sitosterol, triterpenes, and pigments [43]	The oil is used to modulate postprandial plasma lipopolysaccharide, proinflammatory cytokines, TXB2 and LTB4, and reduce the risk of coronary heart disease and metabolic syndrome in patients [42, 43]. Its oral use accelerates the wound healing process even in second-degree and more burn wound patients [44].
8.	<i>Ribes nigrum</i> L. (Black Currant) Family: Grossulariaceae	Fruits and leaves are used in the Eurasian traditional medicine to treat a variety of body ailments and are popularly used to prepare 'cassis liqueur', a traditional alcoholic beverage in France [45]	The seed oil has omega-6-PUFA, γ -linoleic, and α -linoleic acids [45].	Oil is used to treat morning stiffness and reduce pro-inflammatory mediators [46]. Exocarp extract reduces heat shock proteins [47].
9.	<i>Rosa canina</i> L. (Dog Rose) Family: Rosaceae	Used to treat dog bites [48] asthma, bronchitis, cough, flu, gallbladder diseases, burns, vitamin C deficiency, colic, and lower urinary tract disorders [49-51]	Fruits have flavonoids [52, 53], GOPO,PUFAs, linoleic, and α -linolenic acids [54-56]	It has diuretic, anti-arthritis, and anti-rheumatic properties. Its other uses are to treat diarrhea, diabetes, and inadequate peripheral circulation [49-51, 57, 58].

10.	<i>Rosmarinus officinalis</i> L. (Rosemary) Family: Lamiaceae	Rosemary, a sacred plant symbolizes love and death. It is used in weddings and funerals as a symbol of enduring love and eternal bond by Greeks and Romans. Arabians believe that it maintains brain health and memory [59].	Volatile, and phenolic fractions are responsible for therapeutic properties [60].	Essential oil of rosemary is used to treat inflammation [59], and gastric ulcers, and is considered better than Omeprazole [60].
11.	<i>Salvia officinalis</i> L. (Sage) Family: Lamiaceae	Its tea is used to treat digestive and circulation disturbances, asthma, angina, bronchitis, cough, excessive sweating, mouth and throat inflammations, depression, skin diseases, etc. [61, 62]	Carnosol, carnosic acid [63], and essential oil [64].	The plant is reported for anti-cancer, and anti-inflammatory properties [63, 65], but its essential oil does not have any immunomodulatory effects [64].
12.	<i>Spirulina platensis</i> (Nordstedt) Geitler (Blue-green algae) Family: Oscillatoriaceae	It is used for centuries by humans as a food supplement [66]	Crude protein (5-70 %), vitamins, and minerals [66]	<i>S. platensis</i> has analgesic and antiarthritic properties [67]. It helps to control body weight, and no toxic effects have been reported [66].
13.	<i>Urtica dioica</i> L. (Stinging Nettle) Family: Urticaceae	Traditionally used as an aphrodisiac, diuretic, and anti-inflammatory herb [68]	Minerals and trace elements [68].	The lipophilic fraction has more anti-inflammatory potential than its traditional application [68].
14.	<i>Uncaria tomentosa</i> (Willd. ex Schult.) DC (Cat's Claw) Family: Rubiaceae	Traditionally, it is used to treat asthma, abscesses, fever, urinary tract infections, viral infections, and wounds [69]	Alkaloids, sterols, glycosides, organic acids, and triterpenes [69]	Osteoarthritis of the knee remiss after 4 weeks by inhibiting TNF- α and diminishing PGE2 production [70]
15.	<i>Vaccinium myrtillus</i> L. (Bilberry) Family: Ericaceae	<i>V. myrtillus</i> leaves were in use to treat diabetes mellitus before the discovery of insulin [71]	Phenolics [71, 72]	Phytopharmacological studies and clinical trials have not been supported the traditional application [71, 72]
16.	<i>Zingiber officinale</i> Roscoe (Ginger) Family: Zingiberaceae	The rhizome is used to treat pain, muscular aches, fever, sore throats, indigestion, and vomiting [73, 74]	Phenolics, vitamin C, β -carotene, flavonoids, tannins, and minerals [75].	The therapeutic effects depend on the rhizome quantity consumption [73]. It increases the serum corticosterone level and reduces proinflammatory markers [73, 74]. Its extract in osteoarthritis (OA) patients enhances the function of ibuprofen and indomethacin [75-78]

3.3 Anti-inflammatory properties

Bhosale et al. (2012) have studied the anti-inflammatory properties of aqueous extract of *A. aspera*. Two types of extract namely aqueous extracts of *A. aspera* leaves (AEAL) and aqueous extracts of *A. aspera* whole plant (AEAW) were studied in albino mice using carrageenan-induced hind paw edema along with the phytochemical analysis and acute toxicity of extracts. Toxicological studies have revealed the toxic dose of AEAL to be more than 2,000 mg/kg. AEAL reduced the carrageenan-induced edema by 35.71-54.76% compared to the untreated control group, whereas Diclofenac sodium (10 mg/kg) has inhibited the edema volume by 42.85%. The results indicated that the AEAL (800 mg/kg body weight) has more significant anti-inflammatory activity when compared with the standard, and untreated control, respectively. Both extracts of *A. aspera* exhibit promising anti-inflammatory activity, which is rich in alkaloids, flavonoids, saponins, and triterpenoids [84].

Further, the alcoholic extract of the roots of *A. aspera* exhibits anti-inflammatory activity in Wistar rats in carrageenan-induced paw edema and cotton pellet granuloma test [80].

3.4 Anti-HSV activity

Mukherjee et al. (2013) have studied the herpes simplex virus (HSV) with 95% methanolic extract of *A. aspera* roots, and oleanolic acid extracted from *A. aspera*, and reported that both methanolic extract and oleanolic acid inhibit HSV and their plaque formation in a dose-dependent manner [86].

3.5 Anti-HIV activity

HIV can spread in the body via either a cell-free (virus floating free in plasma) mode or a cell in the associated mode (virus particles that remain attached to or within the host cell after replication) involving direct cell-cell contact. *A. aspera* plant extract has been reported as an anti-HIV agent [87] and reported that the methanolic extracts of *A. aspera* and *Rosa centifolia* have anti-HIV activity [88].

3.6 Molecular docking

Dutta et al. (2021) have conducted a molecular docking investigation on the inhibitory efficiency of eight anti-HIV compounds from *A. aspera* extract against severe acute respiratory syndrome coronavirus (SARS-Cov-2). The binding energies and inhibition constant values of these eight compounds (with ACE2 protein of the human body, Mpro, and RdRp proteins of SARS-CoV-2) were compared with the three FDA-approved drugs (Chloroquine, Hydroxychloroquine, and Remdesivir). Most of the compounds were found with good binding energies and inhibition constant values compared to drugs. Further, the absorption, distribution, metabolism, excretion, and toxicity (ADMET) profile of these compounds reveals the safety of the human body. The highest inhibitory activity was achieved by oleanolic acid (OA) with binding energy values of -8.9, -7.6, and -7.4 Kcal/mol against ACE2, Mpro, and RdRp, respectively, reveals that OA has the highest binding affinity with the ACE2 receptor. Hence, OA is a potential therapeutic option for drug development to combat SARS-CoV-2, it has been concluded in the study [89].

3.7 Nephroprotective, anti-arthritic, and antioxidant activity

A. aspera whole plant has been reported for nephroprotective activity against lead acetate induced nephrotoxicity [90], and hypolipidemic activity [91]. Roots of the *A. aspera* are antioxidant, spermicidal activity [92]. Leaves of the plant have been reported for antioxidant, anti-fertility, anti-depressant, anti-cancer, and anti-microbial properties [92-95]. The whole aerial parts are reported as hepatoprotective and the seeds of the plant have anti-microbial activity [96].

3.8 Cardiovascular activity

Achyranthine, a water-soluble alkaloid isolated from *A. aspera* decreased blood pressure and heart rate, dilated blood vessels, and increased the rate and amplitude of respiration in dogs and frogs. The contractile effect of achyranthine at 0.5 mg/ml on frog rectus muscles has been reported less than that of the acetylcholine (0.1 mg/ml), and its spasmogenic effect was not blocked by tubocurarine [80].

3.9 Synergistic behavior

Natural products and related structures are sources of new pharmaceuticals, because of the immense variety of functionally relevant secondary metabolites of herbals [97]. Polyherbal therapy has often produced a promising effect in the treatment of diseases over a single herb. Polyherbal ingredients organized into certain formulas have been deciphered to have potential interaction effects such as mutual enhancement, mutual assistance, mutual restraint, and mutual antagonism [98, 99]. In the Ayurvedic and Unani systems of medicine polyherbal formulation is a common practice [100]. Polyherbal formulations are more effective in numerous diseases with safe high doses. Synergism has followed either pharmacodynamics or pharmacokinetics [101]. The capacity of herbs to ease the absorption, distribution, metabolism, and elimination of the other herbs is focused on pharmacokinetic synergism. Pharmacodynamics synergism includes the effect of active constituents with a similar therapeutic activity that is targeted by the diverse mechanism of action.

The combination of herbals may act on multiple targets at the same time with better relief which lacks in a single herbal formulation. These benefits have resulted in the popularity of polyherbal formulations compared to single herbal formulations [102]. In polyherbal formulation different molecules cure a disease by different mechanisms against a disease condition [103] in E-721B, a polyherbal formulation, which has *A. aspera* as an ingredient, is used to treat acute hypersensitivity in asthma patients. The process is supposed due to the synergistic effect of *A. aspera* with other ingredients [104]. *A. aspera* has a weak anti-viral activity [86], but has good antioxidant and anti-inflammatory properties [105]. The richness in ascorbic acid of *A. aspera* confers the antioxidant activity that boosts the immunity against infections like SARS-coronavirus [105, 106], the most desirable properties for alleviating inflammation due to SARS-coronavirus [83, 107], and utilization of *A. aspera* in the management of COVID-19.

4. Conclusions

The plant-based anti-inflammatory modern green medicines are the standardized plant extracts, and the abundant availability of *A. aspera* is an opportunity as an additional therapeutic armamentarium for chronic inflammatory diseases such as arthritis. It is proved to be a multipurpose medicinal agent that paves the way for further attention and research to identify the synergistic combinations that require more studies to refine the use and improvement of the efficacy of this valuable medicinal plant.

As nature always stands for symbiosis, the study of various phytoconstituents and the discovery of useful herbal combinations to produce a synergistically desirable effect is another thirst area of the present article. Although polyherbal formulation has various phytoconstituents and synergistically produces desirable effects may be supported by the *in-vivo* studies and safety evaluations.

Many more investigations have been conducted on *A. aspera* over many years, and current studies with molecular know-how are only a few but do not reveal any adverse reaction as medicine. Future studies to explore possible synergistic and adverse effects of *A. aspera* should be combined with current recommended pharmaceutical therapies and other adjunct treatments.

Funding

This paper was supported by the Core Grant of Amity University Uttar Pradesh, Noida-201313 (UP), India.

Compliance with ethical standards

This article does not contain any studies with human participants or animals performed by any of the authors.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Abbreviations

GLA = gamma-linolenic acid
 IBS = irritable bowel syndrome
 OLP= oral lichen planus
 EVOO = extra virgin olive oil
 GOPO = galactolipid compound
 PUFAs = polyunsaturated fats
 RA = rheumatoid arthritis
 TNF = Tumor necrosis factor
 AEAAL = aqueous extracts of *A. aspera* leaves
 AEAALW= aqueous extracts of *A. aspera* whole plant
 HSV = herpes simplex virus
 HIV = human immunodeficiency virus
 COVID-19 = *Coronavirus* disease

References

- [1] Shah B.N., Nayak, B.S., Seth A.K., et al. (2006). Medicinal plants as a source of anti-inflammatory and anti-arthritic agents - A review. *Pharmacognosy Magazine*, 2 (6), 77–86.
- [2] Young, A., Thomson, A. J., Ledingham, M., Jordan, F., Greer, I.A., Norman, J. E. (2002). Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term1. *Biology of Reproduction*, 66 (2), 445–449.
- [3] Richard, F., Luigi, X., Cubeddu-Michelle, A. C. (2009). Lippincott's Illustrated Reviews Bundle, Lippincott's Illustrated Reviews: Biochemistry, 4th Ed. Lippincott's. Lippincott Williams & Wilkins.
- [4] Rang, H.P., Dale, M.M., Lamb, P., Ritter, J.M., Moore, P. (2003). *Pharmacology*. Churchill Livingstone.
- [5] Craig, C.R., Stitzel, R.E. (2003). *Modern pharmacology: clinical applications*. Philadelphia, Pa.; London: Lippincott Williams & Wilkins.
- [6] Kumar, V., Abbas, A., Aster, J. (2010). Robbins and Cotran's Pathologic Basis of Disease, 7th Edition. American Journal of Surgical Pathology, 34, 132.
- [7] Richardson, K.T. (1971). Pharmacology and pathophysiology of inflammation. *Archives of Ophthalmology*, 86 (6), 706–724.
- [8] Kumar, R., Clermont, G., Vodovotz, Y., Chow, C.C. (2004). The dynamics of acute inflammation. *Journal of Theoretical Biology*, 230 (2), 145–155.
- [9] Cottam, D.R., Mattar, S.G., Barinas-Mitchell, E., et al. (2004). The chronic inflammatory hypothesis for the morbidity associated with morbid obesity: Implications and effect of weight loss. *Obesity Surgery*, 14 (5), 589–600. doi: 10.1381/096089204323093345.
- [10] Wu, J.T., Wu, L. L. (2007). Acute and chronic inflammation: Effect of the risk factor(s) on the progression of the early inflammatory response to the oxidative and nitrosative stress. *Measurement*, 19 (3), 71–75.
- [11] Kamboj, V.P. (2000). Herbal medicine. *Current Science*, 78 (1), 35–39.
- [12] Wang, M.Y., West, B.J., Jensen, C.J., et al. (2002). *Morindacitrifolia* (Noni): A literature review and recent advances in Noni research. *Acta Pharmacologica Sinica*, 23 (12), 1127–1141.

- [13] Verma, S., Singh, S.P. (2008). Current and future status of herbal medicines. *Veterinary World*, 2(2), 347-350.
- [14] Association, B.M. (1993). *Complementary Medicine: New Approaches to Good Practice*. Oxford; New York, Oxford University Press.
- [15] Boullata, J.I., Nace, A.M. (2000). Safety issues with herbal medicine. *Pharmacotherapy* 20, 357-69. doi: 10.1592/phco.20.4.257.34886.
- [16] Bent, S., Ko, R. (2004). Commonly used herbal medicines in the United States: A review. *American Journal of Medicine*, 116 (7), 478-485. doi: 10.1016/j.amjmed.2003.10.036.
- [17] Sherman, K. J. (2005). Complementary and alternative medicine in the United States. *American College of Physicians. Annals of Internal Medicine*; Philadelphia, 143 (9), e-ISSN: 15393704.
- [18] Drew, A.K., Myers, S.P. (1997). Safety issues in herbal medicine: Implications for the health professions. *Medical Journal of Australia*, 166 (10), 538-541.
- [19] Singh, A., Malhotra, S., Subban, R. (2008). Anti-inflammatory and analgesic agents from Indian medicinal plants. *International Journal of Integrative Biology*, 3 (1), 57-72.
- [20] Ghasemian, M., Owlia, S., Owlia, M.B. (2016). Review of anti-inflammatory herbal medicines. *Advances in Pharmacological Sciences*, 2016: 9130979. doi: 10.1155/2016/9130979.
- [21] Dharmananda, S. (2001). Safety issues affecting herbs: Pyrrolizidine alkaloids. <http://www.itmonline.org/arts/pas.htm>. As accessed on 18th April 2022.
- [22] Bianco, V.V., Santamaria, P., Elia, A. (1998). Nutritional value and nitrate content in edible wild species used in southern Italy. *Acta Horticulturae*, 467, 71-87.
- [23] Larson, K.M., Roby, M.R., Stermitz, F.R. (1984). Unsaturated pyrrolizidines from borage (*Borago officinalis*), A common garden herb. *Journal of Natural Products*, 47(4), 747-748. <http://doi.org/10.1021/np50034a045>.
- [24] Gupta, M., Singh, S. (2010). *Borago officinalis* Linn. An important medicinal plant of Mediterranean region: Review. *International Journal of Pharmaceutical Sciences Review and Research*, 5, 27-34.
- [25] Al-Khamees, W.A., Schwartz, M.D., Alrashdi, S., Algren, A.D., Morgan, B.W. (2011). Status epilepticus associated with Borage Oil ingestion. *Journal of Medical Toxicology*, 7,154-157. doi: 10.1007/s13181-011-0135-9.
- [26] Asadi-Samani, M., Bahmani, M., Rafieian-Kopaei, M. (2014). The chemical composition, botanical characteristic and biological activities of *Borago officinalis*: A review. *Asian Pac J Trop Med*, 7, S22-S28.
- [27] Kast RE. (2001). Borage oil reduction of rheumatoid arthritis activity may be mediated by increased cAMP that suppresses tumor necrosis factor-alpha. *International Immunopharmacology*, 1(12):2197-9. doi: 10.1016/s1567-5769(01)00146-1.
- [28] Deodhar, S.D., Sethi, R., Srimal, R.C. (1980). Preliminary study on antirheumatic activity of curcumin (diferuloyl methane). *Indian Journal of Medical Research*, 71(4), 632-634.
- [29] Bundy, R., Walker, A.F., Middleton, R.W., Booth, J. (2004). Turmeric extract may improve irritable bowel syndrome symptomatology in otherwise healthy adults: A pilot study. *Journal of Alternative and Complementary Medicine*, 10 (6), 1015-1018.
- [30] Hanai, H., Iida, T., Takeuchi, K., et al. (2006). Curcumin maintenance therapy for ulcerative colitis: Randomized, multicenter, double-blind, placebo-controlled trial. *Clinical Gastroenterology and Hepatology*, 4 (12), 1502-1506.
- [31] Taheri, B. J., Anbari, F., Maleki, Z., et al. (2010). Efficacy of *Elaeagnus angustifolia* tropical gel in the treatment of symptomatic oral Lichen Planus. *Journal of Dental Research, Dental Clinics, Dental Prospects*, 4(1), 29-32.
- [32] Ahmadiani, A., Hosseiny, J., Semnani, S., et al. (2000). Antinociceptive and anti-inflammatory effects of *Elaeagnus angustifolia* fruit extract. *Journal of Ethnopharmacology*, 72 (1-2), 287-292. doi: 10.1016/s0378-8741(00)00222-1.
- [33] Farahbakhsh, S., Arbabian, S., Emami, F., et al. (2011). Inhibition of cyclooxygenase type 1 and 2 enzymes by aqueous extract of *Elaeagnus angustifolia* in mice. *Basic and Clinical Neuroscience*, 2(2), 31-37.
- [34] Gyurkovska, V., Alipieva, K., Maciuk, A., et al. (2011). Anti-inflammatory activity of Devil's claw in vitro systems and their active constituents. *Food Chemistry*, 125 (1), 175-178.
- [35] Fiebich, B.L., Fiebich, B.L., Heinrich, M., Hiller, K.O., Kammerer, N. (2001). Inhibition of TNF- α synthesis in LPS-stimulated primary human monocytes by *Harpagophytum* extract SteiHan 69. *Phytomedicine*, 8 (1), 28-30.
- [36] Loew, D., Mollerfeld, J., Schrodter, A., Puttkammer, S., Kaszkin, M. (2001). Investigations on the pharmacokinetic properties of *Harpagophytum* extracts and their effects on eicosanoid biosynthesis *in-vitro* and *ex-vivo*. *Clinical Pharmacology and Therapeutics*, 69 (5), 356-364. doi.org/10.1067/mcp.2001.115445.
- [37] Loew, D., Mollerfeld, J., Schrodter, A., Puttkammer, S., Kaszkin, M. (2001). Investigations on the pharmacokinetic properties of *Harpagophytum* extracts and their effects on eicosanoid biosynthesis *in-vitro* and *ex-vivo*. *Clinical Pharmacology and Therapeutics*, 69 (5), 356-364. doi.org/10.1067/mcp.2001.115445.
- [38] Setty, A.R., Sigal, L.H. (2005). Herbal medications commonly used in the practice of rheumatology: Mechanisms of action, efficacy, and side effects. *Seminars in Arthritis and Rheumatism*, 34(6), 773-784. doi: 10.1016/j.semarthrit.2005.01.011.
- [39] Adeyemi, O. O., Okpo, S. O., Ogunti, O. O. (2002). Analgesic and anti-inflammatory effects of the aqueous extract of leaves of

Persea americana Mill (Lauraceae). *Fitoterapia*, 73 (5), 375–380.

- [40] Borghi, A., Corazza, M., Minghetti, S., Toni, G., Virgili, A. (2015). Avocado and soybean extracts as active principles in the treatment of mild-to-moderate vulvar lichen sclerosus: Results of efficacy and tolerability. *J Eur Acad Dermatology Venereol*, 29, 1225-1230.
- [41] Montserrat-De, L. P. S., Fernandez-Arche, M.A., Angel-Martin, M., Garca-Gimenez, M.D. (2014). Phytochemical characterization of potential nutraceutical ingredients from evening primrose oil (*Oenothera biennis* L.). *Phytochemistry Letters*, 8(1), 158-162.
- [42] Bogani, P., Galli, C., Villa, M., Visioli, F. (2007). Postprandial anti-inflammatory and antioxidant effects of extra virgin olive oil. *Atherosclerosis*, 190 (1), 181-186.
- [43] Camargo, A., Rangel-Zuniga, O.A., Haro, C., et al. (2014). Olive oil phenolic compounds decrease the postprandial inflammatory response by reducing postprandial plasma lipopolysaccharide levels. *Food Chemistry*, 162, 161-171.
- [44] Najmi, M., Shariatpanahi, Z.V., Tolouei, M., Amiri, Z. (2015). Effect of oral olive oil on healing of 10-20% total body surface area burn wounds in hospitalized patients. *Burns*, 41(3), 493-496.
- [45] Gopalan, A., Reuben, S. C., Ahmed, S., Darvesh, A. S., Hohmann, J., Bishayee, A. (2012). The health benefits of blackcurrants. *Food Funct*, 3, 795-809. doi: 10.1039/c2fo30058.
- [46] Watson, J., Byars, M.L., McGill, P., Kelman, A.W. (1993). Cytokine and prostaglandin production by monocytes of volunteers and rheumatoid arthritis patients treated with dietary supplements of blackcurrant seed oil. *Rheumatology*, 32(12), 1055-1058.
- [47] Bishayee, A., Thoppil, R.J., Mandal, A., et al. (2013). Black currant phytoconstituents exert chemoprevention of diethyl nitrosamine-initiated hepatocarcinogenesis by suppression of the inflammatory response. *Molecular Carcinogenesis*, 52(4), 303-317.
- [48] Winther, K., Campbell-Tofte, J., Vinther-Hansen, A.S. (2016). Bioactive ingredients of rose hips (*Rosa canina* L.) with special reference to antioxidative and anti-inflammatory properties: *in-vitro* studies. *Botanics: Targets and Therapy*, 2016 (6), 11-23.
- [49] Kultur, S. (2007). Medicinal plants used in Kirklareli Province (Turkey). *Journal of ethnopharmacology*, 111 (2), 341-364.
- [50] Yeşilada, E., Ustün, O., Sezik, E., Takaishi, Y., Ono, Y., Honda, G. (1997). Inhibitory effects of Turkish folk remedies on inflammatory cytokines: Interleukin-1 α , interleukin-1 β and tumor necrosis factor α . *Journal of Ethnopharmacology*, 58(1), 59-73.
- [51] Carle, R., Gomma, K. (1991). Medicinal uses of *Matricariae* Flos. *British Journal of Phytotherapy*, 2 (4), 147–153.
- [52] Ninomiya, K., Matsuda, H., Kubo, M., Morikawa, T., Nishida, N., Yoshikawa, M. (2007). Potent anti-obese principle from *Rosa canina*: Structural requirements and mode of action of trans-tiliroside. *Bioorganic and Medicinal Chemistry Letters*, 17(11), 3059–3064.
- [53] Nagatomo, A., Nishida, N., Fukuhara, I., et al. (2015). Daily intake of rosehip extract decreases abdominal visceral fat in pre-obese subjects: A randomized, double-blind, placebo-controlled clinical trial. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 2015 (8), 147-156.
- [54] Larsen, E., Kharazmi, A., Christensen, L. P., Christensen, S.B. (2003). An anti-inflammatory galactolipid from rose hip (*Rosa canina*) that inhibits chemotaxis of human peripheral blood neutrophils in vitro. *Journal of Natural Products*, 66(7), 994-995.
- [55] Jager, A. K., Eldeen, I.M.S., Van-Staden, J. (2007). COX-1 and -2 activity of rose hip. *Phytotherapy Research*, 21(12), 1251-1252.
- [56] Jager, A.K., Petersen, K.N., Thomasen, G., Christensen, S.B. (2008). Isolation of linoleic and α -linolenic acids as COX-1 and -2 inhibitors in rose hip. *Phytotherapy Research*, 22 (7), 982-984. doi: 10.1002/ptr.2446.
- [57] Marstrand, C., Warholm, L., Kharazmi, A., Winther, K. (2013). The anti-inflammatory capacity of Rose-hip is strongly dependent on the seeds - A comparison of animal and human studies. *Osteoarthritis and Cartilage*, 21, S216-S217.
- [58] Winther, K. (2014). Shells from rose-hip (*Rosa canina*) do not reduce symptom scores or improve anti-inflammatory property in patients with osteoarthritis-a double-blind, placebo-controlled, randomized study. *Osteoarthritis and Cartilage*, 22, S321. doi:10.1016/j.joca.2014.02.594.
- [59] Borges, R. S., Ortiz, B.L.S., Pereira, A.C.M., et al. (2019). *Rosmarinus officinalis* essential oil: A review of its phytochemistry, anti-inflammatory activity, and mechanisms of action involved. *Journal of Ethnopharmacology*, 229, 29-45.
- [60] Amaral, G.P., de-Carvalho, N.R., Barcelos, R.P., et al. (2013). Protective action of ethanolic extract of *Rosmarinus officinalis* L. in gastric ulcer prevention induced by ethanol in rats. *Food and Chemical Toxicology*, 55, 48-55. doi: 10.1016/j.fct.2012.12.038.
- [61] Hamidpour, M., Hamidpour, R., Hamidpour, S., Shahlari, M. (2014). Chemistry, pharmacology, and medicinal property of sage (salvia) to prevent and cure illnesses such as obesity, diabetes, depression, dementia, lupus, autism, heart disease, and cancer. *Journal of Traditional and Complementary Medicine*, 4 (2), 82-88. doi: 10.4103/2225-4110.130373.
- [62] Fawzi, M., Kamel, Z., Farhan, S. (2017). Anti-inflammatory effect of Sage (*Salvia officinalis*) extracts; Abstract on Oral Health. *Iraqi Dent J*, 39 (1), 1. doi:10.26477/idx.v39i1.111.
- [63] Poeckel, D., Greiner, C., Verhoff, M., et al. (2008). Carnosic acid and carnosol potently inhibit human 5-lipoxygenase and suppress pro-inflammatory responses of stimulated human polymorphonuclear leukocytes. *Biochemical Pharmacology*, 76 (1), 91-97.

- [64] Carrasco, F.R., Schmidt, G., Romero, A.L., et al. (2009). Immunomodulatory activity of *Zingiber officinale* Roscoe, *Salvia officinalis* L. and *Syzygium aromaticum* L. essential oils: Evidence for humor-and cell-mediated responses. *Journal of Pharmacy and Pharmacology*, 61(7), 961-967.
- [65] Rodrigues, M.R.A., Kanazawa, L.K.S., Neves., et al. (2012). Antinociceptive and anti-inflammatory potential of extract and isolated compounds from the leaves of *Salvia officinalis* in mice. *Journal of Ethnopharmacology*, 139 (2), 519-126.
- [66] Rasool, M., Sabina, E. P., Lavanya, B. (2006). Anti-inflammatory effect of *Spirulina fusiformis* on adjuvant-induced arthritis in mice. *Biological and Pharmaceutical Bulletin*, 29 (12), 2483-2487.
- [67] Beg, S., Swain, S., Hasan, H., et al. (2011). Systematic review of herbals as potential anti-inflammatory agents: Recent advances, current clinical status and future perspectives. *Pharmacogn Rev*, 5 (10), 120–137.
- [68] Johnson, T.A., Sohn, J., Inman, W.D., Bjeldanes, L. F., Rayburn, K. (2013). Lipophilic stinging nettle extracts possess potent anti-inflammatory activity, are not cytotoxic and may be superior to traditional tinctures for treating inflammatory disorders. *Phytomedicine*, 20 (2), 143–147.
- [69] Batiha, G. E., Beshbishy, A. M., Wasef, L., et al. (2010). *Uncaria tomentosa* (Willd. ex Schult.) DC.: A review on chemical constituents and biological activities. *Appl Sci* 10, 2668. doi:10.3390/app10082668.
- [70] Piscoya, J., Rodriguez, Z., Bustamante, S. A., et al. (2001). Efficacy and safety of freeze-dried cat's claw in osteoarthritis of the knee: Mechanisms of action of the species *Uncaria guianensis*. *Inflammation Research*, 50 (9), 442–448.
- [71] Kolehmainen, M., Mykkanen, O., Kirjavainen, P.V., et al. (2012). Bilberries reduce low-grade inflammation in individuals with features of metabolic syndrome. *Molecular Nutrition and Food Research*, 56 (10), 1501-1510.
- [72] Hoggard, N., Cruickshank, M., Moar, K.M., et al. (2013). A single supplement of a standardized bilberry (*Vaccinium myrtillus* L.) extract (36 % wet weight anthocyanins) modifies glycaemic response in individuals with type 2 diabetes controlled by diet and lifestyle. *Journal of Nutritional Science*, 2: e22. doi: 10.1017/jns.2013.16.
- [73] Ueda, H., Ippoushi, K., Takeuchi, A. (2010). Repeated oral administration of a squeezed ginger (*Zingiber officinale*) extract augmented the serum corticosterone level and had anti-inflammatory properties. *Bioscience, Biotechnology and Biochemistry*, 74(11), 2248–2252.
- [74] Mahluji, S., Ostadrahimi, A., Mobasseri, M., Attari, V.E., Payahoo, L. (2013). Anti-inflammatory effects of *Zingiber officinale* in type 2 diabetic patients. *Advanced Pharmaceutical Bulletin*, 3 (2), 273-276.
- [75] Shirin, A. P. R., Prakash, J. (2010). Chemical composition and antioxidant properties of ginger root (*Zingiber officinale*). *J Med Plant Res*, 4 (24), 2674-2679. doi: 10.5897/JMPR09.464.
- [76] Haghighi, A., Tavalaei, N., Owlia, M.B. (2006). Effects of ginger on primary knee osteoarthritis. *Indian Journal of Rheumatology*, 1(1), 3-7.
- [77] Haghighi, M., Khalvat, A., Toliati, T., Jallaei, S. (2005). Comparing the effects of ginger (*Zingiber officinale*) extract and ibuprofen on patients with osteoarthritis. *Archives of Iranian Medicine*, 8(4), 267-271.
- [78] Srivastava, K.C., Mustafa, T. (1992). Ginger (*Zingiber officinale*) in rheumatism and musculoskeletal disorders. *Medical Hypotheses*, 39 (4), 25-28.
- [79] Dhale, A. D., Bhoi, S.S. (2013). Pharmacognostic characterization and phytochemical screening of *Achyranthes aspera* Linn. *Current Agriculture Research Journal*, 1(1), 51-57. <http://dx.doi.org/10.12944/CARJ.1.1.07>.
- [80] Hasan, S. (2014). Pharmacological and Medicinal Uses of *Achyranthes Aspera*. *Int J Sci Environ Technol*, 3, 123-129.
- [81] Chougule., Bhupal, S., Patil, A. B., et al. (2021). A Review of Apamarg (*Achyranthes aspera*) with reference to Visha Chikitsa (anti-poisonous effect). *International Journal of Scientific Research*, 10, 45–47.
- [82] Gupta, V., Krishna, C.M., Bansal, P., Kumar, S., Prasad, G.P., Ravi-Krishna, D.V. (2010). Phytochemistry and Pharmacological Potential of *Achyranthes aspera*- A review. *Int. J. Ayurvedic Med*, 1, 1–11. doi:10.47552/ijam.v1i1.
- [83] Vetrichelvan T, Jegadeesan M. (2003). Effect of alcohol extract of *Achyranthes aspera* Linn. on acute and subacute inflammation. *Phytotherapy Research*, 17 (1): 77-79.
- [84] Bhosale, U., Pophale, P., Somani, R., Yegnanarayan, R. (2012). Effect of aqueous extracts of *Achyranthes aspera* Linn. on experimental animal model for inflammation. *Anc Sci Life*, 31 (4), 202-206.
- [85] Sohani, S. (2020). Ethno-medicinal uses and chemical constituents of Indian flora: *Achyranthes aspera* (Latjeera). *International Journal of Science and Research*, 9 (4), 1778-1780.
- [86] Mukherjee, H., Ojha, D., Bag, P., Chandel, H. S., et al. (2013). Anti-herpes virus activities of *Achyranthes aspera*: An Indian ethnomedicine, and its triterpene acid. *Microbiol Res*, 168 (4), 238-244.
- [87] Mothes, W., Sherer, N.M., Jin, J., Zhong, P. (2010). Virus cell-to-cell transmission. *Journal of Virology*, 84 (17), 8360-8368. <https://doi.org/10.1128/JVI.00443-10>.
- [88] Palshetkar, A., Pathare, N., Jadhav, N., et al. (2020). *in-vitro* anti-HIV activity of some Indian medicinal plant extracts. *BMC Complementary Medicine and Therapies*, 20 (69). <https://doi.org/10.1186/s12906-020-2816-x>.
- [89] Dutta, T., Ghorai, S., Khan, A. A., Baildya, N., Ghosh, N.N. (2021). Screening of potential anti-HIV compounds from *Achy-*

- ranthes aspera* extracts for SARS-CoV-2: An insight from molecular docking study. Journal of Physics: Conference Series. 1797 (2021) 012042.
- [90] Jayakumar, T., Sridhar, M.P., Bharathprasad T.R., Ilayaraja, M., Govindasamy, S., Balasubramanian, M.P. (2009). Experimental studies of *Achyranthes aspera* (L) preventing nephrotoxicity induced by lead in albino rats. Journal of Health Science, 55 (5), 701–708.
- [91] Girach-Aminuddin, R.D., Khan, S.A (1992). Ethnomedicinal uses of *Achyranthes aspera* L. in Orissa (India). International Journal of Pharmacognosy, 30 (2), 113–115.
- [92] Sebastian, M.K., Bhandari, M. M. (1984). Medico-ethnobotany of Mount Abu, Rajasthan, India. Journal of Ethnopharmacology, 12(2), 223-230.
- [93] Anis, M., Iqbal, M. (1994). Medicinal plant-lore of Aligarh, India. Pharmaceutical Biology, 32 (1), 59-64. <https://doi.org/10.3109/13880209409082973>.
- [94] Reddy, M.B., Reddy, K. R., Reddy, M. N. (1988). A survey of medicinal plants of Chenchu tribes of Andhra Pradesh, India. Pharmaceutical Biology, 26 (4), 189-196.
- [95] Pal, D. C., Jain, S. K. (1989). Notes on Lodha medicine in Midnapur District, West Bengal, India. Economic Botany, 43(4), 464-470.
- [96] Neogi, N., Rathor, R., Shrestha, A., Banerjee, D. (1969). Studies on the anti-inflammatory and anti-arthritic activity of achyranthine. Indian Journal of Pharmacology, 1 (3), 37–48.
- [97] Ngo, L.T., Okogun, J.I., Folk, W.R. (2013). 21st Century natural product research and drug development and traditional medicines. Natural Product Reports, 30 (4), 584-592.
- [98] Ramaiah, M., Chakravathi, G., Yasaswini, K. (2013). *In-vitro* biological standardization, formulation and evaluation of directly compressed polyherbal anthelmintic tablets. Pharmacogn, J 5(3), 130-134.
- [99] Karole, S., Shrivastava, S., Thomas, S., et al. (2019). Polyherbal Formulation concept for synergic action: A review. J Drug Deliv Ther, 9, 453–66.
- [100] Ajazuddin., Saraf, S. (2010). Evaluation of physicochemical and phytochemical properties of Safoof-E-Sana, a Unani polyherbal formulation. Pharmacognosy Res, 2 (5), 318-22. doi: 10.4103/0974-8490.72332.
- [101] Spinella, M. (2002). The importance of pharmacological synergy in psychoactive herbal medicines. Altern Med Rev, 7 (2), 130-137.
- [102] Parasuraman, S., Thing, G.S., Dhanaraj, S.A. (2014). Polyherbal formulation: Concept of Ayurveda. Pharmacognosy Reviews, 8 (16), 73-80.
- [103] Sarwar, M., Attitall, I.H., Abdollahi, M. (2011). A review on the recent advances in pharmacological studies on medicinal plants; animal studies are done but clinical studies need completing. Asian Journal of Animal and Veterinary Advances, 6 (8), 867-883. doi: 10.3923/ajava.2011.867.883.
- [104] Mitra, S.K., Gopumadhavan, S., Venkataranganna, M. V., et al. (1999). Anti-asthmatic and anti-anaphylactic effect of E-721B, An herbal formulation. Indian Journal of Pharmacology, 31(2), 133-137.
- [105] Adeleye, O. A., Femi-Oyewo, M. N., Bamiro, O.A., et al. (2021). Ethnomedicinal herbs in African traditional medicine with potential activity for the prevention, treatment, and management of coronavirus disease. Futur J Pharm Sci, 7 (1). doi: 10.1186/s43094-021-00223-5.
- [106] Carr, A. C., Maggini, S. (2017). Vitamin C and immune function. Nutrients, 9 (11), 1211. doi: 10.3390/nu9111211.
- [107] Sharma, V., Chaudhary, U. (2015). An overview on indigenous knowledge of *Achyranthes aspera*. J Crit Rev, 2 (1), 7-19.