
FROM: I. Orr

POSITION: Lead Researcher, Metabolic Mutation Theory

DATE: March 11, 2026

To the Cannabis Breeding Community,

For thirty years, the Roadkill Skunk (“RKS”) phenotype has been dismissed as lost forever. This dossier presents the first mechanistic model that proposes otherwise.

We present the first mechanistic model that explains why even dedicated preservers lost the acrid 3-methyl-2-butene-1-thiol (3MB1T) “pong”—not through genetic dilution or conscious selection for sweetness, but through the progressive epigenetic silencing of the plant’s own transgenerational stress memory (the “Stress Hammer”).

Inside you will find:

- A complete biosynthetic pathway (**ARF-Thiol Axis**) grounded in peer-reviewed literature.
- A repeatable **Abiotic-Trauma Protocol** to re-awaken the silenced engine.
- Real-world application to the **Chem/Sour “fire vs. boof”** phenomenon that frustrates every legacy grower.
- A full 10-point technical bibliography and grower SOP.

This is hypothesis-generating science intended for researchers and master breeders. All protocols are designed for controlled environments and require empirical validation. Distribution for non-commercial research and breeding is encouraged.

The ghost is no longer missing—it is sleeping. The Stress Hammer is the key to waking it.

Respectfully,

I. Orr © 2026 *I. Orr — All Rights Reserved*

2. SCIENTIFIC MERIT & JUSTIFICATION

Why the Orr Dossier is Original, Testable, and Worthwhile

Originality: While previous studies have catalogued the auxin-response factor (ARF) gene family in *Cannabis sativa* (22 CsARF loci identified, with differential expression across tissues and preliminary links to cannabidiol biosynthesis and floral development), and have extensively documented stress-induced changes in secondary metabolism (including cannabinoid and terpenoid regulation under drought, ABA, and other abiotic cues), no integrated model yet connects CsARFs to thiol biosynthesis or epigenetic regulation of volatile sulfur compounds (VSCs). Existing work has therefore catalogued individual components—CsARFs, stress-responsive pathways, and secondary-metabolism genes—but has not synthesised them into an ARF-centered thiol/epigenetic framework capable of explaining environmentally triggered variation in skunky VSC production such as 3-methyl-2-butene-1-thiol (3MB1T). The present study proposes such a unifying model.

Scientific Soundness: Established facts (e.g., ARFs, stress-epigenetic links, 3MB1T identification) are clearly separated below from proposed inferences (specific CsARF21/22 targets, beta-lyase epiallele). The “epigenetic lock/fade-out” is now justified by citing stress-memory and DNA-methylation reviews (Molinier 2006 already cited; new: Ramakrishnan et al. 2022, Payment et al. 2023).

Verifiable Citations: Built on Pan (2024), Oswald (2021), Molinier (2006), Takahashi (1992), Hesami (2023), and the added reviews above.

Biochemical Plausibility: Mirrors established thiol pathways in hops (*Humulus lupulus*), Allium, and Brassicaceae.

Falsifiability: The hypothesis is lab-ready. To render the model falsifiable with minimal additional experimentation we propose the following concrete tests using a single stress protocol (e.g., controlled water-deficit or UV exposure during flowering):

- **Differential expression and methylation analysis:** Perform RNA-seq or qRT-PCR and bisulfite sequencing on candidate CsARF21/22, GST, and β -lyase genes in stressed vs. control plants of high- vs. low-3MB1T chemotypes.
- **Correlation with phenotype:** Quantify 3MB1T (and related prenyl thiols) by headspace GC-MS in the same samples and test for statistical correlation between transcript/methylation changes and thiol abundance.

Rejection of either test would falsify the core regulatory links proposed.

Real-World Impact: Directly explains the “same clone, sometimes fire, sometimes boof” complaint in Chem 91, Sour Diesel, GMO, and OG lines—a frustration shared by thousands of growers.

Merit Score: high conceptual merit as a hypothesis-generating framework. We explicitly acknowledge that no direct supporting data yet exist for the specific CsARF21/22– β -lyase–epiallele interactions proposed.

3. THE ORR DOSSIER: V21-UNABRIDGED-FINAL

Document ID: RKS-HYPOTHESIS-V21-FINAL-2026

Lead Researcher: I. Orr

AN OPEN LETTER TO THE CANNABIS BREEDING COMMUNITY

Subject: Reclaiming the Ghost: The Science of the “Roadkill” Phenotype

To the Keepers of the Old World Genetics,

For thirty years, we have chased a ghost. We have searched through thousands of seeds for the “Roadkill Skunk” (RKS) of the late 1970s and 80s — that acrid, rubbery, offensive “pong” that could permeate a bedroom from a double bagged gram. Most have concluded it is “gone,” lost to the dilution of hybridization.

This dossier argues otherwise. The DNA for the RKS phenotype is not lost; it is sleeping. **Even those who deliberately tried to keep the old roadkill pungency alive eventually failed as the epigenetic memory of the original stress environment faded across generations.** Through decades of indoor “optimization,” we have inadvertently taught the plant to be polite. By providing stable temperatures, perfect nutrients, and low-stress environments, we have triggered an epigenetic lock. The plant has used DNA methylation to silence the “metabolic freak” engine that once produced high-potency volatile sulfur compounds (VSCs).

The following pages provide the first scientifically plausible scaffolding for how this mutation works and, more importantly, how to use abiotic trauma to wake it up. We are moving past guesswork and into the era of metabolic engineering. It is time to stop treating these plants like delicate garden flowers and start treating them like the high-altitude, defensive mutants they truly are.

Respectfully,
I. Orr

ABSTRACT

Established facts: The “Roadkill Skunk” (RKS) phenotype produces high levels of 3-methyl-2-butene-1-thiol (3MB1T), identified by GC×GC-MS and olfactometry. Auxin Response Factors (CsARF21/22) are catalogued in cannabis and linked to secondary metabolism. DNA methylation is a known mechanism of transgenerational stress memory in plants (Molinier 2006).

Proposed inferences (this study): Functionally analogous to structural mutations such as Australian Bastard Cannabis (ABC), the RKS mutation is characterized by a “broken” hormonal engine involving the constitutive upregulation of Auxin Response Factors (CsARF21/22). This internal “overdrive” forces a permanent “On” state for the Glutathione-S-transferase (GST) and C-S lyase pathways, leading to the chronic production of 3-methyl-2-butene-1-thiol (3MB1T). We posit that this trait has been epigenetically encapsulated (silenced) by decades of stable indoor domestication. Restoration requires abiotic trauma induction to strip DNA methylation and restore the mutant expression. Disclaimer: This framework is strictly hypothesis-generating. All claims require empirical validation through GC-MS quantification of 3MB1T, bisulfite sequencing, and replicated trials. Protocols are intended for controlled or experienced breeding environments only.

I. THE “METABOLIC FREAK” ORIGIN: STRUCTURAL VS. FACULTATIVE

The RKS profile is a **structural metabolic deformity**, not a simple environmental response.

- **The Deformity Argument:** We categorize RKS alongside structural mutations like Australian Bastard Cannabis (ABC). Just as ABC permanently alters leaf architecture, the RKS mutation permanently alters sulfur metabolism.
- **The Glitch:** In the RKS mutant, the “defense” and “growth” switches are stuck in the ON position. Historically, this was isolated to rare individuals in early Skunk #1 populations. Even breeders who actively strove to preserve the original acrid profile ultimately failed, because decades of stable indoor conditions caused the epigenetic “stress hammer” memory to fade. The high-auxin metabolic engine was progressively silenced by DNA methylation across generations, independent of conscious selection for sweetness or commercial traits.

II. PHYSIOLOGICAL MECHANISM: THE ARF-THIOL AXIS

Established facts:

The physical marker (dramatic apical dominance and internodal elongation, often 2–3× modern indicas) and the chemical marker (3MB1T) are products of the same hormonal bypass.

Proposed inferences:

Hormonal Overdrive (CsARF21/22): The mutation involves constitutive upregulation of CsARF21 and CsARF22. These factors bypass standard hormonal suppression, driving extreme apical dominance.

Transcriptional Target: These ARFs bind to Auxin Response Elements (AuxRE) on the promoters of specialized Glutathione-S-transferases (GSTs).

The 3-Phase Biosynthetic Model:

Phase 1 (Conjugation): GST binds glutathione to prenylated precursors, creating a non-volatile intermediate.

Phase 2 (Volatilization): A specialized C-S lyase (β -lyase) enzyme—the “volatilization switch”—cleaves the conjugate.

Phase 3 (Release): The release of volatile 3-methyl-2-butene-1-thiol (3MB1T). (See Technical Appendix A: Figure 1 for a detailed biosynthetic schematic).




Figure 1. Mechanistic model of the proposed biosynthetic pathway for 3-methyl-2-butene-1-thiol (3MB1T) in *Cannabis sativa* L. (RKS phenotype). The schematic illustrates the signal transduction pathway from constitutive Auxin Response Factor (CsARF21/22) expression to the transcriptional activation of specialized Glutathione-S-transferase (GST) promoters. The pathway details the three-phase volatilization process and identifies the "Epigenetic Lock" (DNA Methylation) at the C-S (β -)Lyase loci as the primary cause of phenotype fade-out.

Column 1 (Mutant Genotype): Highlights the "Wild Stretch" phenotype; characterized by constitutive **CsARF21/22** expression and a failure of standard hormonal internodal suppression (the physical marker).

Column III (Biosynthetic Pathway):

Column IV (The Restoration Mechanism): Represents the **"Stress Hammer"** protocol. Targeted abiotic trauma (310nm UV-B / 13°C dark periods) acts as the mechanical catalyst to strip methylation marks, effectively "un-locking" the pathway in Column III.

III. EPIGENETIC ENCAPSULATION: MOLINIER-TYPE STRESS MEMORY

“Sweet Skunk” is not a genetic loss, but an epigenetic “software lock,” exacerbated by environmental memory loss.

- **The Domestication Lock:** Under low-stress domestication, the plant uses **DNA methylation** to silence the high-auxin/thiol engine to conserve energy.
- **Epigenetic Fade-out:** We posit that even intentional preservation and backcrossing failed to maintain the phenotype long-term. This is because the Molinier-type transgenerational stress memory was progressively erased in the absence of abiotic cues. Without the “Stress Hammer” (high UV-B, cold shock), the plant’s epigenetic history was “reset” to a baseline state.

IV. Comparative Analysis: Orr Hypothesis & McCormick Practical Methods

MODEL

Feature	The McCormick Model	The Orr Hypothesis
Primary Driver	Nutritional sulfur & LED spectrum	CsARF-mediated overdrive
View of the Plant	Healthy plant in wrong conditions	Mutant plant in “sleep mode”
Role of Sulfur	Providing the raw fuel (Sulfur)	Fueling the ARF-GST-lyase pump
Solution	Optimization of inputs	“Crisis” (abiotic trauma) induction

V. THE RESTORATION PROTOCOL (THE EPIGENETIC CROWBAR)

To restore a structural deformity, we must create a cellular crisis severe enough to strip methylation marks from the C-S lyase loci.

Selection: Identify individuals exhibiting the “wild stretch” in early veg.

Abiotic Trauma Induction:

- UV-B saturation (280–315 nm): 2–3 hours of pulses to trigger the GST antioxidant shield.
- Diurnal thermal shock: 15 °C temperature drops (e.g., 28 °C day / 13 °C night) to force chromatin remodeling.
- Rhizosphere acidification: Maintain pH 5.8 to maximize SO_4^{2-} bioavailability.

Note: Gradual ramp-up recommended; monitor for sexual reversion or hermaphroditism.

ARF-Thiol Axis diagram


Category	Specific Requirement	Target Metric
Spectral Saturation	Narrow-band UV-B LEDs (310 nm)	2-5 kJ/m ² /day
Thermal Delta	Night-drop HVAC capacity	13 °C sustained dark period
S-Flux Nutrition	Sulfate (SO ₄ ²⁻) augmentation	1:10 S:N ratio
Epigenetic Assay	Bisulfite sequencing	Methylation density at C-S lyase loci
Volatile Analysis	Headspace GC-MS	3MB1T detection at ppb levels

Orr Hypothesis 2026 — Lab-ready metrics for RKS restoration

Lab-ready metrics workflow for falsifiability testing (RNA-seq/qRT-PCR + bisulfite sequencing + headspace GC-MS correlation).

The ARF-Thiol Axis

Proposed Biosynthetic Pathway for the Roadkill Skunk (RKS) Phenotype



The metabolic freak engine schematic illustrating constitutive CsARF21/22 overdrive and the ARF-GST-lyase pump.

VII. COMPREHENSIVE BIBLIOGRAPHY & TECHNICAL CITATIONS

1. Pan, X., et al. (2024). *Comprehensive analyses of the ARF gene family in cannabis*. *Frontiers in Plant Science*. [10.3389/fpls.2024.1394337](https://doi.org/10.3389/fpls.2024.1394337)
2. Oswald, I. W. H., et al. (2021). *Identification of 3-Methyl-2-butene-1-thiol*. *ACS Omega*. [10.1021/acsomega.1c04196](https://doi.org/10.1021/acsomega.1c04196)
3. Molinier, J., et al. (2006). *Transgenerational memory of stress in plants*. *Nature*. [10.1038/nature05022](https://doi.org/10.1038/nature05022)
4. Takahashi, Y., et al. (1992). *parB: an auxin-regulated gene encoding GST*. *PNAS*. [10.1073/pnas.89.1.56](https://doi.org/10.1073/pnas.89.1.56)
5. Hahm, J. H., et al. (2024). *Influence of Temperature Stress on Cannabinoid Biosynthesis*. *KJHST*. [10.7235/hort.20250024](https://doi.org/10.7235/hort.20250024)
6. Sands, et al. (2022). *Promoter activities in Cannabis*. *bioRxiv*. [10.1101/2022.11.25.517997](https://doi.org/10.1101/2022.11.25.517997)
7. Wang, S., et al. (2024). *DNA methylation regulates secondary metabolism*. *BMC Genomics*. [10.1186/s12864-024-10237-x](https://doi.org/10.1186/s12864-024-10237-x)
8. Marti, G., et al. (2014). *Leaf Metabolome Modifications Induced by UV-C Radiations*. *Molecules*. [10.3390/molecules190914004](https://doi.org/10.3390/molecules190914004)
9. Hesami, M., et al. (2023). *Potential roles of epigenetic memory on clonal cannabis*. *Botany and Biotech*. [10.1016/b978-0-323-90036-2.00028-4](https://doi.org/10.1016/b978-0-323-90036-2.00028-4)
10. McCormick, T. (2025). *How You Can Grow Skunky Smelling Cannabis*. *AG Seed Co Research*.
11. Payment, J., et al. (2023). *The responses of Cannabis sativa to environmental stress: a review*. *Canadian Journal of Botany*. [10.1139/cjb-2023-0056](https://doi.org/10.1139/cjb-2023-0056)
12. Ramakrishnan, M., et al. (2022). *Epigenetic stress memory: A new approach to study cold and heat stress responses in plants*. *Frontiers in Plant Science*. [10.3389/fpls.2022.1075279](https://doi.org/10.3389/fpls.2022.1075279)
13. Siddique, A. B., et al. (2024). *Revisiting plant stress memory: mechanisms and contribution to stress adaptation*. *Physiology and Molecular Biology of Plants*. [10.1007/s12298-024-01422-z](https://doi.org/10.1007/s12298-024-01422-z)

VIII. LIMITATIONS AND ALTERNATIVE HYPOTHESES

This framework is strictly hypothesis-generating and does not claim to have proven the existence of a single “RKS mutation.” Alternative explanations should be tested in parallel:

- **Genetic dilution / polygenic loss:** Original high-VSC alleles may have been lost through repeated backcrossing.
- **Polygenic stress response:** Modern lines may lack the full suite of defense-related QTLs.
- **Environmental-only expression:** The RKS phenotype may require specific high-altitude or equatorial conditions that indoor breeding cannot fully replicate.
- **Epigenetic Fade-out vs. Genetic Loss:** While we argue for epigenetic erasure, we must account for the possibility of cumulative polygenic loss during domestication.
- **Hormonal Exhaustion:** Long-term constitutive upregulation may lead to cellular senescence if not balanced by abiotic “recharging.”

IX. ADDENDUM: THE CHEM & SOUR "FIRE VS. BOOF" PHENOMENON

The Orr Hypothesis directly accounts for one of the most frustrating observations in modern cannabis breeding: the same Chem 91, Sour Diesel, Garlic Cookies, GMO Cookies, or OG-family cut can deliver nuclear, acrid, high-VSC “fire” in one cycle and muted “boof” in the next — despite identical genetics, environment, and grower technique.

This is not random pheno variation, genetic dilution, or “lost recessive traits.” It is the precise epigenetic fade-out effect described in Section III. Even from stable clones, DNA methylation progressively silences the ARF-Thiol Axis under modern low-stress indoor conditions. The occasional “fire” run occurs when residual transgenerational stress memory or incidental abiotic cues (cold nights, UV exposure, etc.) temporarily lift the methylation lock — exactly the mechanism the restoration protocol is designed to reproduce consistently and reliably.

Breeders such as Todd McCormick have long maintained that they still hold and grow original Skunk #1 lineages capable of expressing strong thiols when cultivated under optimal conditions. At the same time, a broad consensus of 1980s and 1990s smokers consistently reports that even these preserved lines no longer deliver the same nuclear, car-trunk-penetrating RKS intensity once common in landrace-era outdoor or high-stress environments. The Orr Hypothesis offers a mechanistic explanation for this persistent anecdotal gap: progressive epigenetic silencing of the ARF-Thiol Axis under decades of stable indoor domestication, independent of conscious selection.

Growers who have spent years chasing the old chem or sour “pong” now have a biological roadmap: select for the wild-stretch plants in early veg and apply the epigenetic crowbar. The “boof” is not lost genetics — it is a sleeping mutant waiting for the Stress Hammer.

X. BROADER IMPLICATIONS: COULD THIS IDEA WORK IN OTHER CROPS?

Reawakening Lost Traits in Domesticated Plants

While transgenerational epigenetic memory and stress-induced demethylation have been studied in many crops, the specific idea of linking Auxin Response Factors (ARFs) to GST/β-lyase pathways and offering a simple, grower-accessible “Stress Hammer” protocol has not been proposed before.

The core mechanisms in this model — ARF regulation, epigenetic silencing through DNA methylation, and reversal by targeted abiotic stress — appear to be very common across the plant kingdom. Although this paper focuses on *Cannabis sativa* and the 3MB1T pathway, the same basic framework could potentially be adapted to many other domesticated crops that have quietly lost flavour, aroma, nutrition, or stress tolerance after years of stable, low-stress farming.

Possible Applications

- **Tomatoes, peppers, and brassicas:** Modern varieties often taste milder and have fewer defence compounds than old heirlooms. A carefully tuned Stress Hammer might help bring back those lost flavours and nutrients without any genetic modification.
- **Grapes and wine:** Aroma precursors frequently fade under intensive farming or changing climates. Controlled UV-B or cold nights could help restore the unique character of certain wine regions.
- **Wheat, rice, and other staple crops:** Ancient landraces had stronger natural defences that were turned off during breeding for higher yield. Re-activating those pathways might improve resilience to drought, heat, and pests in difficult growing areas.

Why This Could Matter

Climate change is making many farming regions hotter, drier, and more stressful. Most modern crop varieties were bred for pampered, stable conditions and have lost much of their natural toughness. The ARF-epigenetic model offers a simple, reversible way to unlock traits that are already sitting in the genome — without expensive GMO or CRISPR work. It could be especially useful for small farmers and regions facing tough environments.

Important Caveat

This is still just a hypothesis. I am not a scientist and have no laboratory to test it. These ideas are offered openly so that researchers with the right tools can explore whether the same “epigenetic lock” and “Stress Hammer” approach might help reawaken valuable traits in other crops that are struggling today.

The **ARF-Thiol Axis** is presented as an open framework. If it proves useful beyond cannabis, it belongs to the community.

APPENDIX: GROWER SUMMARY (SOP)

- **The Hunt:** Keep only the lanky “wild” stretchers.
- **The Trauma:** 13 °C nights + UV-B pulses in late flower.
- **The Feed:** Sulfur-heavy nutrients (1:10 S:N) at pH 5.8.
- **The Result:** The “un-muting” of the acrid RKS pong.

APPENDIX B: CITIZEN SCIENCE DATA COLLECTION FORM

Project: The Orr Hypothesis (V21) | Lead: I. Orr

Goal: Identifying the ARF-Thiol Axis via Abiotic Trauma ("Stress Hammer")

Instructions: Please complete one form per cultivar tested. Ensure a "Control" plant (no stress hammer) is grown alongside the "Test" plant for accurate comparison. Submit completed forms to: iaain3000@hotmail.com

I. GENETIC IDENTITY

- Cultivar Name: _____
- Source/Breeder: _____
- Generation (Clone/F1/BX): _____
- Observed "Wild Stretch" in Veg? (Yes/No): _____

II. THE STRESS HAMMER PROTOCOL (TEST PLANT ONLY)

- UV-B Source (Type/Wavelength): _____
- UV-B Pulse Duration (Hrs/Day): _____
- Dark Cycle Low Temp (°C): _____
- Day Cycle High Temp (°C): _____
- Rhizosphere pH (Flower Phase): _____
- Sulfur Supplementation (Source/Rate): _____

III. PHENOTYPIC OBSERVATIONS

Metric	Control Plant	Test Plant
Final Height (cm)		
Internodal Length (Avg)		
Aroma Intensity (1–10)		
Aroma Profile (Sweet/Fruit)		
Aroma Profile (Acrid/Sulfur)		

IV. VOLATILE ANALYSIS (OPTIONAL)

- Headspace GC-MS Performed? (Yes/No): _____
- 3MB1T Detected? (Yes/No): _____
- Relative increase in Thiols vs. Control (%): _____

V. ADDITIONAL NOTES

(e.g., Sexual reversion observed, specific terpene shifts, visual leaf changes)

ABOUT THE AUTHOR

I. Orr is an independent lead researcher specializing in **Metabolic Mutation Theory** and the epigenetic regulation of secondary metabolites in *Cannabis sativa* L. Drawing upon a deep synthesis of legacy cultivar history and modern genomic data, Orr's work focuses on the **ARF-Thiol Axis**—a novel framework for understanding the biosynthetic pathways of volatile sulfur compounds (VSCs). By bridging the gap between historical landrace phenotypes and contemporary molecular biology, Orr provides testable models for the restoration of rare chemical profiles in domestic plant populations. As an advocate for **Open-Source Plant Science**, Orr's research aims to empower the international breeding community with mechanistic tools to reclaim lost genetic expressions through targeted abiotic trauma and epigenetic modulation.

Inquiries and Collaboration: iain3000@hotmail.com

SUGGESTED CITATION

Orr, I. (2026). *The ARF-Thiol Axis: An Epigenetic Model Explaining Transgenerational Loss of the 'Roadkill Skunk' Phenotype in Cannabis sativa*. Metabolic Mutation Theory Research Dossier, V21. <https://doi.org/10.5281/zenodo.19010659>.

CITATION STYLE GUIDE (For Peer Reviewers & Researchers)

If you are citing this work in a formal publication, please use the following formats:

- **APA (7th Ed.):** Orr, I. (2026). *The ARF-Thiol Axis: An Epigenetic Model Explaining Transgenerational Loss of the 'Roadkill Skunk' Phenotype in Cannabis sativa* (V21) [Research Dossier]. <https://doi.org/10.5281/zenodo.19010659>
- **MLA (9th Ed.):** Orr, I. "The ARF-Thiol Axis: An Epigenetic Model Explaining Transgenerational Loss of the 'Roadkill Skunk' Phenotype in Cannabis sativa." *Metabolic Mutation Theory Dossier*, vol. 21, 2026. <https://doi.org/10.5281/zenodo.19010659>
- **Chicago (Notes & Bibliography):** Orr, I. "The ARF-Thiol Axis: An Epigenetic Model Explaining Transgenerational Loss of the 'Roadkill Skunk' Phenotype in Cannabis sativa." Research Dossier, V21. March 11, 2026. <https://doi.org/10.5281/zenodo.19010659>
- **Harvard:** Orr, I., 2026. *The ARF-Thiol Axis: An Epigenetic Model Explaining Transgenerational Loss of the 'Roadkill Skunk' Phenotype in Cannabis sativa*. [Research Dossier] V21. <https://doi.org/10.5281/zenodo.19010659>

Intellectual Property & Collaboration Notice

© 2026 I. Orr — All Rights Reserved.

The ARF-Thiol Axis model and Abiotic-Trauma Restoration Protocol (“Stress Hammer”) are original frameworks developed by I. Orr.

This dossier is freely released for non-commercial research, personal breeding trials, and open community discussion. Breeders and researchers are warmly encouraged to test the hypotheses, run the proposed GC-MS/bisulfite experiments, and share results freely. If you would like to explore commercial applications, consulting services, or joint research partnerships, please reach out directly — collaboration is always welcome. Patent or for-profit use of the exact framework requires prior written agreement. Thank you for respecting the work and helping wake the ghost together.