



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Paper

Hepatoprotective Medicinal Plants: Phytochemistry, Mechanisms and Therapeutic Potential of Andrographis Paniculata, Boerhavia Diffusa, Eclipta Prostrata and Phyllanthus Emblica

Priyal Nil*, Shilpa Joshi, Dr. Pragnesh Patni

Khyati College of Pharmacy, Palodia, Ahmedabad.

ARTICLE INFO

Published: 17 Apr 2026

Keywords:

Hepatoprotective,
Andrographis paniculata,
Boerhavia diffusa, Eclipta
prostrata, Phyllanthus
emblica, Liver diseases,
Herbal medicine.

DOI:

10.5281/zenodo.19628859

ABSTRACT

Liver diseases represent a major global health challenge, with increasing incidence of drug-induced hepatotoxicity, non-alcoholic fatty liver disease, and viral hepatitis. Medicinal plants have gained considerable attention as potential hepatoprotective agents due to their diverse bioactive compounds and multitarget mechanisms of action. This review summarizes the hepatoprotective potential of four widely used medicinal plants: Andrographis paniculata, Boerhavia diffusa, Eclipta prostrata, and Phyllanthus emblica. Their phytochemical constituents, mechanisms of action, and pharmacological evidence from experimental and clinical studies are discussed. These plants demonstrate hepatoprotective activity primarily through antioxidant, anti-inflammatory, and hepatocyte-regenerating mechanisms. Key bioactive compounds such as andrographolide, wedelolactone, and emblicanin contribute significantly to their therapeutic effects. The review also highlights the potential advantages of polyherbal formulations and discusses current research gaps including lack of clinical validation and standardization of extracts. These findings support the potential role of herbal medicines in liver disease management while emphasizing the need for further clinical investigations

INTRODUCTION


***Corresponding Author:** Priyal Nil

Address: Khyati College of Pharmacy, Palodia, Ahmedabad

Email ✉: priyaln2194@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.





[Figure 1 - Hepatitis]

The global burden of liver diseases continues to escalate, representing a critical public health challenge requiring innovative therapeutic approaches. Modern healthcare systems face increasing pressure to develop effective treatments for various forms of hepatic disorders while minimizing adverse effects associated with conventional pharmacological interventions.[1] Current therapeutic limitations have intensified research interest in alternative medicine approaches, particularly those involving natural hepatoprotective agents with established safety profiles. [2]Traditional medicine has long acknowledged the value of medicinal plants in treating liver disorders, supported by centuries of clinical use and observation [3]. Blending this traditional wisdom with modern scientific methods creates strong potential for safe and effective hepatoprotective therapies in today's liver disease management. According to the World Health Organization, nearly 80% of people worldwide depend on traditional medicine for primary healthcare, with plant-based remedies forming its core [4]. The interest in herbal hepatoprotective therapies arises from key advantages that set plant-based medicines apart from synthetic drugs. While conventional pharmaceuticals usually act on a single pathway, medicinal plants often work on multiple targets at once, tackling different disease processes through the combined action of diverse bioactive

compounds [5]. Advances in phytochemistry, molecular pharmacology, and systems biology are now uncovering the scientific basis behind many traditional claims about their hepatoprotective effects. Modern tools like high-resolution mass spectrometry, nuclear magnetic resonance, and advanced chromatography have made it possible to identify and characterize the bioactive compounds driving therapeutic effects in both preclinical and clinical studies [6]. The pharma industry now acknowledges the strong therapeutic and commercial value of herbal hepatoprotectives, with plant-based products already achieving global market success [7]. Growing scientific evidence on their safety and efficacy has also driven wider acceptance within mainstream medicine and regulatory systems [8]. The pathological changes associated with hepatitis and liver injury are illustrated in Figure 1.

MEDICINAL PLANTS IN HEPATOPROTECTION

The therapeutic efficacy of medicinal plants in liver protection stems from their considerably complex phytochemical compositions and remarkably diverse mechanisms of action that target multiple aspects of hepatic pathophysiology. These plants contain intricate mixtures of bioactive compounds including flavonoids, phenolic acids, terpenoids, alkaloids, saponins, and other specialized metabolites, each contributing to overall hepatoprotective effects through complex molecular interactions and cellular pathway modulations that modern pharmaceutical science is only beginning to fully understand and appreciate [9]. Oxidative stress represents one of the most fundamental and pervasive pathological mechanisms underlying virtually all forms of liver disease, from acute hepatotoxic injury to chronic progressive fibrosis and cirrhosis [10]. When hepatotoxic agents such as carbon tetrachloride, acetaminophen, alcohol,

or various pharmaceutical compounds enter the liver through portal circulation, they trigger massive production of reactive oxygen species, leading to extensive lipid peroxidation, protein oxidation, DNA damage, and ultimately hepatocellular death through necrosis or apoptosis [11]. Medicinal plants combat this oxidative damage through multiple complex and complementary pathways, including direct free radical scavenging through electron donation, enhancement of endogenous antioxidant enzyme systems, modulation of cellular redox balance, and activation of cytoprotective signaling cascades [12]. These multifaceted antioxidant mechanisms work synergistically to provide comprehensive protection against oxidative injury while simultaneously promoting cellular repair and regeneration processes essential for hepatic recovery.

SELECTED PLANTS: RATIONALE AND SIGNIFICANCE

The selection of Kalmegh (*Andrographis paniculata*), Punarnava (*Boerhavia diffusa*), Bhringraj (*Eclipta prostrata*), and Amla (*Phyllanthus emblica*) for comprehensive detailed analysis reflects their extensive traditional use histories, robust scientific evidence bases, and distinct complementary therapeutic profiles that together represent the significant diversity and sophistication of traditional hepatoprotective medicine [13]. These four significant plants represent different botanical families and diverse geographic distributions, providing comprehensive coverage of hepatoprotective plant diversity while ensuring broad accessibility across various regions, cultures, and healthcare systems worldwide. The therapeutic significance of these carefully selected plants extends well beyond hepatoprotection alone, as they demonstrate complex multi-target pharmacological activities including immunomodulation, anti-inflammatory

effects, antimicrobial properties, metabolic regulatory functions, and various other beneficial physiological effects that contribute to overall health and wellness [14]. This significant polypharmacological nature aligns perfectly with traditional medicine's fundamentally holistic approach to health and disease management, offering distinct therapeutic advantages over single-target synthetic drugs that may produce limited therapeutic outcomes, unwanted side effects, or incomplete therapeutic responses. Geographic diversity among these four selected plants ensures broader global accessibility and significantly reduces problematic dependency on single sources, thereby contributing meaningfully to supply chain security, sustainable harvesting practices, and therapeutic availability across different regions with varying climatic conditions and agricultural capabilities [15]. The differing cultivation needs, ecological preferences, and environmental adaptations of these species enable the development of region-specific therapeutic programs while maintaining consistent therapeutic standards, quality control, and regulatory compliance across various production systems and locations.

KALMEGH (*Andrographis paniculata*)



[Figure 2 - Kalmegh]

Andrographis paniculata, commonly known as Kalmegh or "King of Bitters," represents one of the most extensively researched and scientifically

validated hepatoprotective plants in contemporary phytotherapy and evidence-based herbal medicine [16]. This significant annual herbaceous plant has gained significant international scientific attention due to its potent therapeutic properties, well-characterized bioactive compounds, and important safety profile established through both traditional use and modern toxicological studies. The medicinal plant *Andrographis paniculata* used for hepatoprotection is shown in Figure 2.

Taxonomical Classification

Kingdom	Plantae
Phylum	Spermatophyta
Class	Angiospermae
Order	Lamiales
Family	Acanthaceae
Genus	Andrographis
Species	A. paniculata
Botanical Name	<i>Andrographis paniculata</i> (Burm.f.) Nees


Vernacular Names

Language/Region	Local Name
English	Indian Echinacea
Hindi	Kalmegh
Sanskrit	Bhunimba
Tamil	Nilavembu
Telugu	Nelavemu
Bengali	Kalomegh

Botanical Description

Andrographis paniculata (family Acanthaceae), commonly known as Kalmegh, is an annual medicinal herb widely distributed in tropical regions of Asia. It has long been used in traditional Ayurvedic medicine for the treatment of liver disorders, inflammation, and infections. [17]

Hepatoprotective Mechanism



[Figure 3 - Mechanism of Kalmegh]

The hepatoprotective mechanisms of *A. paniculata* involve complex modulation of key cellular signaling pathways, particularly the Nrf2 and NF- κ B response pathways that regulate oxidative stress responses and inflammatory processes [18]. The primary bioactive andrographolide demonstrates potent antioxidant mechanisms that contribute significantly to the plant's hepatoprotective activity [19]. The proposed hepatoprotective mechanism of the plant is illustrated in Figure 3.

Phytochemical Constituents

Major phytoconstituents of Kalmegh that show hepatoprotective activity are Andrographolide, Neoandrographolide, oroxylin and luteolin. *Andrographis paniculata* is widely recognized for its liver-protective potential, with research confirming its ability to counter ethanol-induced toxicity through several complementary mechanisms [20]. Both preclinical and clinical studies support its role in easing methotrexate-related liver injury by protecting cells and

adjusting biochemical pathways [21]. Investigations further show that andrographis preparations improve liver function by restoring enzyme balance and regulating metabolic activity [22]. Leaf extracts, in particular, have displayed strong protective effects against severe conditions such as thioacetamide-induced cirrhosis, consistently proving their therapeutic value in different experimental settings [23]. Its broad hepatoprotective action is attributed to a range of bioactive compounds that act together in synergy [24]. Animal studies have helped determine safe dosage levels and established therapeutic benchmarks for different preparations [25]. Advances in analytical methods have provided a clearer picture of its phytochemical composition and linked these findings with observed protective effects [26]. Careful pharmacological research, supported by strict experimental methods and statistical validation, now offers strong scientific backing for the traditional use of *A. paniculata* in liver care [27]

Pharmacological Properties

Preclinical Evidence:

- Hepatoprotective against ethanol, methotrexate, and thioacetamide; improved liver enzymes [20-23]
- Validated in animal models with optimized protocols [24,25]

Mechanistic Insights:

- Diterpenoids and andrographolide act via antioxidant pathways [19,28]

PUNARNAVA (*Boerhavia diffusa*)



[Figure 4 - Punarnava]

Boerhavia diffusa, traditionally revered as Punarnava meaning "the one that makes the body new again," represents a true cornerstone of classical Ayurvedic hepatotherapy with extensively documented therapeutic use spanning over three thousand years of continuous clinical application [29][30]. This significant perennial herbaceous plant is universally renowned for its considerable adaptability to harsh environmental conditions, important therapeutic versatility, and potent rejuvenative properties that have made it invaluable in traditional medicine systems worldwide. The medicinal plant *Boerhavia diffusa*, traditionally used for liver disorders, is shown in Figure 4.

Taxonomical Classification

Kingdom	Plantae
Phylum	Spermatophyta
Class	Angiospermae
Order	Caryophyllales
Family	Nyctaginaceae
Genus	Boerhavia
Species	B. diffusa
Botanical Name	Boerhavia diffusa L.

Vernacular names


Language/Region	Local Name
English	Red Spiderling
Hindi	Gadahpurna
Sanskrit	Raktapunarnava

Tamil	Mukkurattai
Telugu	Atikamamidi
Bengali	Rakta punarnava
Gujarati	Satodi

Botanical Description

Boerhavia diffusa (family Nyctaginaceae), commonly known as Punarnava, is a perennial herb widely distributed in tropical and subtropical regions. It has been traditionally used in Ayurvedic medicine for the treatment of liver disorders, edema, and inflammatory conditions. [31]

Hepatoprotective Mechanism



[Figure 5 - Mechanism of Punarnava]

The hepatoprotective mechanisms of *B. diffusa* involve direct free radical scavenging activities and enhancement of endogenous antioxidant systems, contributing to membrane stabilization and cellular protection [32]. The plant's bioactive compounds demonstrate significant potential in managing complex hepatotoxicity scenarios, including drug-induced liver injury. The proposed hepatoprotective mechanism of the plant is illustrated in Figure 5.

Phytochemical Constituents

Key phytoconstituents of Punarnava are Punarnavine, Boeravinone A-F, Lignans and

Rotenoids. *Boerhavia diffusa* has been extensively reviewed for its diverse phytochemical profile and wide therapeutic applications across traditional medicine systems worldwide [33]. Its ethnomedicinal relevance offers important context for both its historical uses and its growing significance in modern healthcare [30]. Detailed studies on its phytochemistry and pharmacology have revealed a complex array of bioactive compounds and mechanisms of action [34], with research identifying key constituents directly linked to hepatoprotective and other beneficial effects [35]. *B. diffusa* has shown real promise in protecting the liver, with studies reporting benefits against different types of liver damage [36].

Pharmacological Properties

Preclinical Evidence:


- Strong pharmacological and therapeutic documentation [33]
- Traditional hepatoprotective and rejuvenative uses supported by ethnomedicinal data [30]
- Phytochemical analyses reveal diverse bioactive compounds [34]
- Key active constituents identified for pharmacological effects [35]
- Experimental studies confirm notable hepatoprotective activity [36]

Clinical Context and Global Relevance:

- Supported by global liver disease trends and epidemiological studies [37]
- Promising role in complex liver disease management [38]
- Potential applications in drug-induced liver injury [39]

BHRINGRAJ (*Eclipta prostrata*)





[Figure 6 - Bhoringraj]

Eclipta prostrata, universally revered as Bhoringraj or "King of Hair," occupies a truly distinguished and respected position in traditional medicine systems worldwide, being particularly valued for its considerable hepatoprotective properties and significant regenerative capabilities that have made it indispensable in classical therapeutic practices [40]. This versatile annual herbaceous plant demonstrates truly significant adaptability to diverse ecological conditions while consistently maintaining therapeutic potency and bioactive compound concentrations. The medicinal plant *Eclipta prostrata* used in traditional hepatoprotective therapy is shown in Figure 6.

Taxonomical Classification

Kingdom	Plantae
Phylum	Spermatophyta
Class	Angiospermae
Order	Asterales
Family	Asteraceae
Genus	<i>Eclipta</i>
Species	<i>E. prostrata</i>
Botanical Name	<i>Eclipta prostrata</i> L.

Vernacular names


Language/Region	Local Name
English	False Daisy
Hindi	Bhoringraj
Sanskrit	Bhoringraj

Tamil	Karisalankanni
Telugu	Gunta galagara
Bengali	Kesuti
Gujarati	Bhangaro

Botanical Description

Eclipta prostrata (family Asteraceae), commonly known as Bhoringraj, is a small annual herb commonly found in tropical and subtropical regions. It is widely used in traditional medicine for liver protection, hair growth, and treatment of inflammatory disorders. [41]

Hepatoprotective Mechanism



[Figure 7 - Mechanism of Bhoringraj]

The hepatoprotective mechanisms of *E. prostrata* involve complex modulation of NF-κB pathways and direct antioxidant activities, with key bioactive compounds like wedelolactone demonstrating specific protective effects against acute liver injury [42]. The plant's comprehensive therapeutic approach addresses multiple pathological pathways simultaneously. The proposed hepatoprotective mechanism of the plant is illustrated in Figure 7.

Phytochemical Constituents



Major phytoconstituents of Bhringraj are Wedelolactone, Demethylwedelolactone, Luteolin, Apigenin and Quercetin. *Eclipta prostrata* has been studied for its traditional uses, complex chemistry, and wide-ranging health benefits [40]. Research shows that *Eclipta alba* can protect the liver from damage caused by high-fat diets, helping with metabolic liver issues [43]. Studies on carbon tetrachloride exposure have revealed how it supports cell defense and liver repair [44], while investigations on wedelolactone, one of its main compounds, explain its action at the molecular level [42]. Overall, multiple studies confirm that *Eclipta alba* can both protect and heal the liver across different experimental models [45].

Pharmacological Properties

Preclinical Evidence:

- Strong documentation of ethnomedicinal uses, phytochemistry, and pharmacology [40]
- Protective role against diet-induced liver damage [43]
- Significant activity against chemical toxicity, e.g., carbon tetrachloride [44]
- Wedelolactone shown to guard against acute liver injury [42]
- Broad hepatoprotective effects across multiple experimental models [45]

Advanced Research Applications:

- Featured in reviews of herbal hepatoprotective mechanisms [46]
- Explored for synergistic potential via extract combinations [47]
- Analytical studies on optimized extraction and phytochemistry
- Evidence for oxidative stress reduction through multiple pathways

AMLA (*Phyllanthus emblica*)



[Figure 8 - Amla]

Phyllanthus emblica, universally revered as Amla or Indian Gooseberry, stands prominently as one of the most treasured and scientifically validated medicinal plants in traditional Indian medicine, earning the prestigious designation "Divya Aushada" or divine medicine in ancient Ayurvedic texts and continuing to maintain this esteemed status in contemporary herbal medicine practice [48]. This truly significant deciduous tree has sustained human health and wellness for over five thousand years of continuous therapeutic use, representing the absolute pinnacle of nutritional excellence and therapeutic sophistication in natural medicine systems worldwide. The medicinal plant *Phyllanthus emblica*, known for its antioxidant and hepatoprotective properties, is shown in Figure 8.

Taxonomical Classification

Kingdom	Plantae
Phylum	Spermatophyta
Class	Angiospermae
Order	Malpighiales
Family	Phyllanthaceae
Genus	<i>Phyllanthus</i>
Species	<i>P. emblica</i>
Botanical Name	<i>Phyllanthus emblica</i> L.


Vernacular names

Language/Region	Local Name
English	Indian Gooseberry
Hindi	Amla, Amalaki
Sanskrit	Amalaki, Dhatri
Tamil	Nellikai
Telugu	Usirikaya
Bengali	Amlaki
Gujarati	Amala

Botanical Description

Phyllanthus emblica (family *Phyllanthaceae*), commonly known as Amla or Indian gooseberry, is a deciduous tree widely distributed in the Indian subcontinent. The fruit is extensively used in traditional medicine due to its rich antioxidant content and hepatoprotective properties. [49]

Hepatoprotective Mechanism



[Figure 9 - Mechanism of Amla]

The hepatoprotective mechanisms of *P. emblica* involve powerful antioxidant activities primarily mediated by high concentrations of vitamin C and polyphenolic compounds, contributing to comprehensive cellular protection and metabolic regulation [50]. The plant's therapeutic effects extend to lipid metabolism regulation and anti-

inflammatory activities. The proposed hepatoprotective mechanism of the plant is illustrated in Figure 9.

Phytochemical Constituents

Key phytoconstituents Emblicanin A and B, Punigluconin, Ascorbic acid and Gallic acid. Extensive research has provided comprehensive reviews of *Phyllanthus emblica* phytochemical composition, systematically revealing the notable complexity, significant therapeutic potential, and complex bioactive compound profiles that make this plant considerable among medicinal species [48]. Detailed studies examining hepatoprotective properties of Indian gooseberry have provided thorough scientific analysis of therapeutic mechanisms, clinical applications, and potential for integration into modern healthcare systems [51]. Advanced research investigating hepatoprotective activities of polyherbal formulations has systematically demonstrated the important synergistic potential achieved when amla is strategically combined with other complementary therapeutic plant species [52]. Contemporary evidence-based approaches to herbal treatments in liver diseases have definitively established *P. emblica* as a cornerstone therapeutic agent with proven efficacy and safety [53]. Comprehensive studies examining selected hepatoprotective herbal medicines have provided substantial scientific evidence for significant antioxidant and anti-inflammatory effects that contribute to therapeutic outcomes [54].

Pharmacological Properties

Preclinical Evidence:

- Complex phytochemistry with strong therapeutic potential [48]
- Established hepatoprotective properties of Indian gooseberry [51]



- Synergistic effects in polyherbal formulations [52]
- Validation of herbal treatments for liver disease [53]
- Strong antioxidant and anti-inflammatory actions [54]
- Clinical studies confirm measurable hepatoprotective outcomes [55]

SYNERGISTIC POTENTIAL:

The strategic development of complex polyherbal formulations represents a highly advanced and scientifically validated approach to hepatoprotective therapy, effectively capitalizing on the complementary mechanisms of action exhibited by different medicinal plants while minimizing potential adverse effects and maximizing therapeutic outcomes [52]. This approach acknowledges the complex, multifactorial nature of liver diseases and addresses multiple pathological pathways simultaneously through carefully designed therapeutic interventions [46]

RATIONALE FOR POLYHERBAL COMBINATIONS

The scientific rationale for developing complex polyherbal combinations stems from the recognition that liver diseases involve multiple interconnected pathological processes that can be more effectively addressed through multitarget therapeutic approaches rather than single-compound interventions [14]. The four selected plants demonstrate complementary mechanisms of action, distinct phytochemical profiles, and non-overlapping toxicity profiles that make them ideal candidates for synergistic therapeutic combinations [47]. Although these medicinal plants differ in their phytochemical composition, they share common hepatoprotective mechanisms including antioxidant activity, modulation of inflammatory pathways, and enhancement of hepatocyte regeneration. The hepatoprotective mechanisms and key phytochemical constituents of the selected medicinal plants are summarized in Table 1.

Table 1. Major hepatoprotective plants, key

Plant	Major compounds	Mechanism	Evidence
Andrographis paniculata	Andrographolide	Nrf2 activation, antioxidant	Animal + clinical
Boerhavia diffusa	Boeravinones	Antioxidant, membrane stabilization	Animal
Eclipta prostrata	Wedelolactone	NF- κ B inhibition	In vitro + in vivo
Phyllanthus emblica	Emblicanins, Vitamin C	ROS scavenging, anti-inflammatory	Clinical

Pharmacodynamic Synergy Mechanisms

The concept of pharmacodynamic synergy in herbal medicine is firmly rooted in the well-established scientific principle that carefully designed combinations of bioactive compounds can produce therapeutic effects significantly greater than the sum of their individual contributions [47]. This complex approach effectively addresses the inherently multifactorial

nature of liver diseases more comprehensively and effectively than single-plant preparations or isolated compound therapies [18][19].

Experimental Design Challenges

Figuring out how different herbs work together to protect the liver is challenging. It takes smartly designed experiments, high-level analysis, and a real grasp of how plant compounds interact.



Regular lab models often miss the messy complexity of liver diseases in real life, so researchers need more realistic setups to get meaningful results.

CONCLUSION

Medicinal plants such as *Andrographis paniculata*, *Boerhavia diffusa*, *Eclipta prostrata*, and *Phyllanthus emblica* exhibit significant hepatoprotective activity supported by experimental and emerging clinical evidence. [20][30][40][48] Their protective effects are primarily mediated through antioxidant activity, modulation of inflammatory pathways, and enhancement of hepatocyte regeneration. The presence of bioactive phytochemicals including andrographolide, wedelolactone, and emblicanins plays a critical role in these mechanisms [19][32][42][50]. Although current findings support their therapeutic potential, further research is required to establish standardized formulations, optimal dosing, and clinical efficacy through well-designed clinical trials. Future studies should focus on the development of safe and effective polyherbal formulations for liver disease management.[37][38][39][55]

REFERENCES

1. Sivaraj A, Bharani M, Ramamurthy Ch, Balamurugan G, Rajkapoor B. Hepatoprotective potential of *Andrographis paniculata* against ethanol-induced liver toxicity. *J Appl Pharm Sci*. 2011; 1(6): 12-18.
2. Parthasarathy M, Palani S, Kumar TM, Prabhu S. *Andrographis paniculata* Alleviates Methotrexate-Induced Hepatotoxicity. *Life*. 2023; 13(4): 941.
3. Kaewdech A, Nawalerspanya S, Assawasuwannakit S, Chamroonkul N. The use of *Andrographis paniculata* and its effects on liver biochemistry. *Sci Rep*. 2022; 12(1): 8312.
4. Bardi DA, Halabi MF, Abdullah NA, et al. *Andrographis paniculata* Leaf Extract Prevents Thioacetamide-Induced Liver Cirrhosis. *PLoS One*. 2014; 9(10): e109424.
5. Jayakumar T, Hsieh CY, Lee JJ, Sheu JR. Experimental and Clinical Pharmacology of *Andrographis paniculata*. *Evid Based Complement Alternat Med*. 2013; 2013: 846740.
6. Sutha A, Devaraj SN. A Study on the Hepatoprotective Effect of *Andrographis paniculata* on Mice. *J Phytol*. 2010; 2(11): 31-36.
7. Ali SK, Makeen HA, Khuwaja G, et al. Assessment of Phytochemical Profile and Hepatoprotective Effect. *Medicina*. 2023; 59(3): 524.
8. Nipanikar SU, Kale MK, Gursahani MS. Pharmacological Evaluation of Hepatoprotective Activity. *Pharmacogn Res*. 2017; 9(1): 42-48.
9. Handa SS, Sharma A. Hepatoprotective activity of andrographolide against galactosamine. *Indian J Med Res*. 1990; 92: 276-283.
10. Gupta A, Sharma K, Kushwaha G, Singh G. Evaluation of hepatoprotective activity of *Andrographis paniculata* extract. *Pharma J*. 2022; 11(2): 234-241.
11. Ahmad W, Khan S, Qureshi U, et al. Hepatoprotective Diterpenoids Isolated from *Andrographis paniculata*. *Pharmacol Pharm*. 2012; 3(3): 393-402.
12. Rahman MM, Islam MR, Shohag S, et al. Hepatoprotective activity of andrographolide possibly through antioxidant mechanism. *Heliyon*. 2022; 8(10): e10782.
13. Mayuren C, Reddy VV, Priya SV, Devi VA. Protective effect of Livactine against CCl4



- and paracetamol induced hepatotoxicity. N Am J Med Sci. 2010; 2(10): 491-494.
14. Li X, Chen Y, Wang Z, et al. Modulation of Nrf2 and NF- κ B Signaling Pathways. Antioxidants Basel. 2021; 10(8): 1226.
15. Zhang Y, Liu W, Chen X, et al. Dissecting the Crosstalk Between Nrf2 and NF- κ B Response Pathways. Front Cell Dev Biol. 2022; 10: 824447.
16. Singh A, Kumar R, Patel S, et al. A review of the plant Boerhaavia diffusa. Phytopharm J. 2016; 5(2): 89-96.
17. Rahman MA, Hasan R, Khan SA, et al. Ethnomedicinal values of Boerhaavia diffusa L. Front Chem. 2023; 11: 1186094.
18. Mishra S, Aeri V, Gaur PK, Jachak SM. Boerhaavia diffusa Linn: Phytochemical and Pharmacological Profile. Asian J Appl Sci. 2014; 2(4): 476-493.
19. Roy S, Samanta SK, Mukherjee D, et al. Boerhaavia diffusa: Bioactive Compounds and Pharmacological Activities. Biomed Pharma J. 2019; 12(3): 1321-1336.
20. Beedimani RS, Kumar S, Shekhawat NS. Evaluation of hepatoprotective activity of Boerhaavia diffusa. Int J Basic Clin Pharmacol. 2015; 4(4): 756-760.
21. Patel N, Kumar A, Singh R. Hepatoprotective activity of Boerhaavia diffusa L. J Nov Med Health Sci. 2025; 13(2): 45-52.
22. Kim JK, Park SJ, Lee MH, et al. Global, Regional, and National Trends in Liver Disease-Related Mortality. J Korean Med Sci. 2024; 39(15): e142.
23. Liu X, Wang M, Zhang Y, et al. Liver diseases: epidemiology, causes, trends and predictions. Signal Transduct Target Ther. 2025; 10: 142.
24. Luo J, Chen XY, Wang ZH, et al. Management of hepatotoxicity of chemotherapy and targeted agents. J Hematol Oncol. 2021; 14(1): 144.
25. Singh A, et al. A Review on Hepatoprotective and Immunomodulatory Herbal Plants. Evid Based Complement Alternat Med. 2018; 2018: 9341574.
26. Ahmad N, Fazal H, Abbasi BH, et al. Hepatoprotective Potential of Malaysian Medicinal Plants. Front Pharmacol. 2020; 11: 988.
27. Rajappa S, Kumar A, Singh R, et al. Exploring the Mechanisms and Dosages of Herbal Hepatoprotective Agents. Pharmacogn Res. 2024; 16(2): 198-206.
28. Chen M, Li X, Wang Y, Zheng H, Zhang Y, Liu Y, et al. Roles of Nrf2 in Liver Diseases: Molecular, Pharmacological Aspects. Int J Mol Sci. 2020; 21(22): 8675.
29. Patel C, Patel B, Mamtara T, et al. Eclipta prostrata: Ethnomedicinal Uses, Phytochemistry, and Pharmacology. Front Pharmacol. 2021; 12: 640850.
30. Naik KS, Gurushanthaiah M, Kavimani M, et al. Hepatoprotective Role of Eclipta alba against High Fatty Diet. Maedica. 2018; 13(4): 283-289.
31. Beedimani RS, Shetkar S. Hepatoprotective activity of Eclipta alba against carbon tetrachloride. Int J Basic Clin Pharmacol. 2015; 4(3): 478-483.
32. Lu Y, Wen DL, Lei QL, et al. Protective effect of wedelolactone against CCl₄-induced acute liver injury. Int Immunopharmacol. 2016; 34: 53-60.
33. Kumar S, Gupta P, Sharma A, et al. Phyllanthus emblica: comprehensive review of phytochemical composition. Front Pharmacol. 2023; 14: 1120253.
34. Bahadoran Z, Mirmiran P, Azizi F. Hepatoprotective properties of Indian gooseberry: a review. Food Funct. 2013; 4(10): 1441-1452.
35. Singh MK, Yadav SS, Gupta V, Khattri S. Hepatoprotective activity of polyherbal

- formulation containing *Phyllanthus emblica*. J Pharm Bioallied Sci. 2013; 5(1): 29-34.
36. Sandhya T, Lathika KM, Pandey BN, Mishra KP. Potential of traditional ayurvedic formulation in liver diseases. J Ethnopharmacol. 2006; 105(1-2): 131-137.
37. Wang Y, Tang C, Zhang H. Hepatoprotective effects of kaempferol 3-O-rutinoside and kaempferol 3-O-glucoside from *Carthamus tinctorius* L. J Ethnopharmacol. 2015; 168: 253-264.
38. Li X, Chen Y, Wang Z, Xiao Y, Yang H, Su Z. Modulation of Nrf2 and NF- κ B signaling pathways in liver diseases. Antioxidants (Basel). 2021;10(8):1226.
39. Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran Pathologic Basis of Disease. 8th ed. Philadelphia: Saunders Elsevier; 2010.
40. Ahmad N, Fazal H, Abbasi BH, Farooq S, Ali M, Khan MA. Hepatoprotective potential of medicinal plants and their constituents: Possible mechanisms of action. Front Pharmacol. 2020;11:988.
41. Girish C, Pradhan SC. Hepatoprotective activities of picroliv, curcumin and ellagic acid. Toxicol Appl Pharmacol. 2012; 258(2): 182-193.
42. Zhao X, Zhang JJ, Wang X, et al. Effect of berberine on hepatocyte proliferation and liver regeneration. Biomed Pharmacother. 2008; 62(1): 71-78.
43. Kumaran A, Joel Karunakaran R. In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. LWT Food Sci Technol. 2007; 40(2): 344-352.
44. Liu X, Zhao M, Wu K, et al. Toxicity of carbon tetrachloride-hepatotoxicity and the protective effect of *Phyllanthus emblica* fruit. Food Chem Toxicol. 2010; 48(5): 1279-1286.
45. Pramyothin P, Samosorn P, Pongshompoo S, Chaichantipyuth C. The protective effects of *Phyllanthus emblica* Linn. extract on ethanol induced rat hepatic injury. J Ethnopharmacol. 2006; 107(3): 361-364.
46. Tasduq SA, Peerzada K, Kadian S, et al. Biochemical manifestations of anti-tuberculosis drugs induced hepatotoxicity and the effect of silymarin. Hepatol Res. 2005; 31(3): 132-135.
47. Thyagarajan SP, Subramanian S, Thirunalasundari T, et al. Effect of *Phyllanthus amarus* on chronic carriers of hepatitis B virus. Lancet. 1988; 2(8614): 764-766.
48. Bhattacharjee R, Sil PC. Protein isolate from the herb, *Phyllanthus niruri* L. (Euphorbiaceae), plays hepatoprotective role against carbon tetrachloride induced liver damage via antioxidant mechanisms. Food Chem Toxicol. 2007; 45(5): 817-826.
49. Rajeshkumar NV, Kuttan R. *Phyllanthus amarus* extract administration increases the life span of rats with hepatocellular carcinoma. J Ethnopharmacol. 2000; 73(1-2): 215-219.
50. Sultana S, Ahmed S, Jahangir T, Sharma S. Inhibitory effect of celery seeds extract on chemically induced hepatocarcinogenesis. Cancer Lett. 2005; 221(1): 11-20.
51. Krithika R, Mohankumar R, Verma RJ, et al. Isolation, characterization and antioxidative effect of phyllanthin against CCl₄-induced toxicity in HepG2 cell line. Chem Biol Interact. 2009; 181(3): 351-358.
52. Jose JK, Kuttan R. Hepatoprotective activity of *Emblica officinalis* and *Chyavanaprash*. J Ethnopharmacol. 2000; 72(1-2): 135-140.
53. Jeena KJ, Joy KL, Kuttan R. Effect of *Emblica officinalis*, *Phyllanthus niruri* and *Picrorrhiza kurroa* on N-nitrosodiethylamine induced hepatocarcinogenesis. Cancer Lett. 1999; 136(1): 11-16.

54. Anila L, Vijayalakshmi NR. Beneficial effects of flavonoids from *Sesamum indicum*, *Emblica officinalis* and *Momordica charantia*. *Phytother Res.* 2000; 14(8): 592-595.
55. Antony B, Merina B, Iyer VS, et al. A pilot clinical study to evaluate the effect of *Emblica officinalis* extract (Amlamax) on markers of systemic inflammation and dyslipidemia. *Indian J Clin Biochem.* 2008; 23(4): 378-381.

HOW TO CITE: Priyal Nil, Shilpa Joshi, Dr. Pragnesh Patni, Hepatoprotective Medicinal Plants: Phytochemistry, Mechanisms and Therapeutic Potential of *Andrographis paniculata*, *Boerhavia diffusa*, *Eclipta prostrata* and *Phyllanthus emblica*, *Int. J. of Pharm. Sci.*, 2026, Vol 4, Issue 4, 2769-2782, <https://doi.org/10.5281/zenodo.19628859>