HiLL PUBLISHING

International Journal of Food Science and Agriculture, 2025, 9(4), 464-469
https://www.hillpublisher.com/journals/jsfa/
ISSN Online: 2578-3475
ISSN Print: 2578-3467

Extraction of Oil from Safou Pulp (*Dacryodes edulis* (G. Don) H. J. Lam) by Hot Pressure: Yield and AG Composition

Feueltgaldah Christian Bopoundza[1,3,*], Bob Wilfrid Loumouamou[1,2,3], Jean Bruno Bassiloua[1,3,4], Mignon Prince Exaucé Taty[1]

[1]Multidisciplinary Food and Nutritional Research Team, Regional Center of Excellence in Food and Nutrition, Faculty of Sciences and Technology, Marien Ngouabi University, Brazzaville, BP 69, Republic of the Congo.
[2]Process Engineering Laboratory, UNESCO-ENSP Chair, Marien Ngouabi University, Brazzaville, BP 69, Republic of the Congo.
[3]CR2IE, Chemistry and Technology Platform, Higher School of Cataract Technology (EPrES), Brazzaville, BP 69, Republic of the Congo.
[4]Non Wood Forest Products Valorization Center, CVPFNL, BP 5700 Pointe-Noire, Republic of Congo.

How to cite this paper: Feueltgaldah Christian Bopoundza, Bob Wilfrid Loumouamou, Jean Bruno Bassiloua, Mignon Prince Exaucé Taty. (2025). Extraction of Oil from Safou Pulp (*Dacryodes edulis* (G. Don) H. J. Lam) by Hot Pressure: Yield and AG Composition. *International Journal of Food Science and Agriculture*, 9(4), 464-469.
DOI: 10.26855/ijfsa.2025.12.020

Received: November 30, 2025
Accepted: December 29, 2025
Published: January 28, 2026

Corresponding author: Feueltgaldah Christian Bopoundza, Multidisciplinary Food and Nutritional Research Team, Regional Center of Excellence in Food and Nutrition, Faculty of Sciences and Technology, Marien Ngouabi University, Brazzaville, BP 69, Republic of the Congo; CR2IE, Chemistry and Technology Platform, Higher School of Cataract Technology (EPrES), Brazzaville, BP 69, Republic of the Congo.

Abstract

The actual launch of industrial or semi-industrial production of oil from safou pulp can only be effective if it is profitable enough to compete with the species currently being exploited. Studies on oil extraction by mechanical pressing, which is the main industrial processing technique for oilseeds, are outdated and need to be reevaluated using modern presses to better assess oil extraction yields. The present study, carried out using a screw press (KOMET type CA 59 G) on the pulp studied, gave an extraction yield of 81.18% of the total oil, more or less exceeding those reported in the literature. In addition, as the oil from the kernel has a similar fatty acid composition to that of the pulp, using the whole fruit could increase the fat content and eliminate the need for pitting, thereby reducing processing costs. The extracted oil has a fatty acid profile of C16:0>C18:2>C18:1>C18:0>C18:3, which has already been reported in the literature. Mechanical pressing has no influence on the fatty acid composition of the oil obtained.

Keywords

Safou pulp; oil; press extraction; yield; AG

1. Introduction

The ever-increasing growth of the world population, estimated at 9.8 billion by 2050 by the UN [1], is forcing decision-makers to focus on developing food supply strategies to ensure food security for populations. The demand for fats, which generally account for 30 to 40% of total energy intake (WHO, 2018; ANSES, 2021), corresponding to approximately 70 to 100 g of fat per day per person, demonstrates the importance attached to this class of metabolites [2]. Vegetable oils remain the most widely consumed and recommended fat in terms of nutritional quality and cardiovascular disease prevention (FAO) compared to animal fats.

Safou (*Dacryodes edulis*) is an unconventional oilseed crop that is promising and potentially nutritious [3-11], domesticated and highly prized in sub-Saharan Africa [12-17], and could help meet the demand for fat, especially in Africa, where increasing food insecurity is a growing concern [18]. This contribution can only be effective if cost-effective processing techniques are mastered, the effective implementation of which could also help reduce post-harvest losses of this species, which amount to more than 50% [9, 19-21].

Knowledge of yield and quality in finished products using mechanical presses, which offer automation, intelligent control, and energy savings, making the process accessible to small producers at the industrial level and constituting the main technique for industrial processing of oilseeds, is relatively old [5, 22, 23] and was not always one of the authors' main objectives [5, 22]. Updating knowledge in line with scientific advances, including the development of modern presses, is essential for a better appreciation of the qualities of this species.

2. Materials and Methods

2.1 Plant material

The safous studied were purchased at the Total market in Brazzaville (Republic of Congo). The only selection criterion was the ripeness of the fruit (dark blue or black color). According to the seller, these safous came from Cameroon.

2.2 Study methods

After purchase, the safous were transported to the laboratory. They were cleaned before being processed.

2.2.1 Morphological and physicochemical characteristics

The masses of the fruits and pulps studied were determined using a CONSTANT PN-Model precision balance: 14192-005R; Max: 620 g; d: 0.01 g, morphological dimensions (length, width) using a caliper, and water and oil content of the pulp using formula 1 after drying part of the pulp halves in an oven (T = 103°C, duration = 24 h) and by formula 2 after Soxhlet extraction with hexane for 3 h of these oven-dried pulp halves [24].

$$\%(water) = \frac{m_0 - m_1}{m_0} x100 \quad (1)$$

$$\%(oil) = \frac{m_1 - m_2}{m_1} x100 \quad (2)$$

avec:
m_0: initial mass of the half-pulp.
m_1: mass of the half-pulp after drying in an oven (or of the powder before extraction)
m_2: mass of the half-pulp cake after extraction.

2.2.2 Mechanical extraction of oil from pulp

The remaining pulp halves used to extract oil by pressing were dried in an oven at 60°C until the mass stabilized. Extraction (3 trials) was carried out using a KOMET CA 59 G press (IBG Montors Oekotec GmbH & Co. KG, Mönchengladbach, Germany) with a capacity of 3-5 kg/h and a power of 1.1 kW (Figure 1). In each test, 500 g of half-pulps were roasted at 80°C for 10 min. During extraction, the press speed was set to 2.4 AU, and a 7 mm diameter die was used.



Figure 1. Extraction of oil from safou pulp using the KOMET CA 59 G press.

2.2.3 Major fatty acid compositions of extracted oils

Fatty acid methyl ester (FAME) preparation and gas chromatography (GC) analysis for FA composition were performed according to the NF EN ISO 12966-2 (AFNOR, 2017). FAME was analyzed by GC (FOCUS GC chromatograph (Thermo Fisher Scientific, Massachusetts, USA)) equipped with a CP-Sil 88 column (50 m x 0.25 mm x 0.2 µm; Agilent, Santa Clara, USA) and a split injector (1:20 ratio). Carrier gas: helium, Flow rate: 1.0 mL/min. Injector temperature: 250°C. FID detector temperature: 270°C. The oven was heated from 185 to 225°C at a speed of 5°C/min and maintained at 225°C for 10 min. FAMEs were identified by comparing the retention times of each peak to those of the FAME mix standard and quantified as a relative percentage of total FA using Chromcard 2.3.32005 software (Thermo Fisher Scientific, Massachusetts, USA).

3. Results and Discussions

3.1 Morphological and physicochemical characteristics of the safous studied

The morphological and physicochemical characteristics of the safous studied (Table 1) show that they belong to size III, category III [25, 26] and class III [27].

Table 1. Characteristics of the safous studied

	Mean ± standard deviation
Characteristics	87
Length (cm)	8.41 ± 0.78
Width (cm)	4.93 ± 0.43
Thickness (cm)	0.98 ± 0.14
Fruit weight (g)	78.67 ± 7.57
Pulp mass (g)	56.74 ± 5.86
Water content (%)	62.44 ± 5.43
Oil content (%)	44.89 ± 3.72

The "*edulis*" variety [28], with a sphericity index (width/length) of ½, is the most widely grown and marketed variety in the sub-region [11].

3.2 Oil yield and fatty acid composition

The experimental results are shown in Table 2 below.

Table 2. Oil yield and major fatty acid composition

	Test 1	Test 2	Test 3	**Mean**	**Standard deviation**
Yield	36.38	36.45	36.48	36.44	0.05
C16:0	45.21	45.21	45.20	45.21	0.01
C18:0	5.74	5.71	5.69	5.71	0.02
C18:1n9c	19.03	19.06	19.02	19.04	0.02
C18:2n6c	28.96	28.86	29.02	28.95	0.08
C18:3n3	0.77	0.87	0.77	0.81	0.06
C20:0	0.29	0.29	0.29	0.29	0.00

3.2.1 Oil yield

The extraction yield was 36.38% for test 1, 36.48% for test 2, and 36.44% for test 3, with an average of 36.44 ± 0.05. This average yield, which represents 81.18% of the total oil contained in the pulps studied, slightly higher than those recently obtained using the same press on the dried pulps of *Raphia sese* [29], is higher than those obtained by [5] using an ALTECH electric press equipped with a BE 100 4/4 (2.2 kW) motor, varying between 53.80 and

60.49% of total oil. This yield is also very high compared to those obtained using traditional methods [23]. In a study on the "problems of safou pulp oil production in Cameroon," the study mentions a yield of 6.75 L of oil extracted using a traditional press, compared to 15 L of oil extracted using a solvent, which is equivalent to 45% of the total oil. [9] reports this yield to be between 25 and 49%. If we rely on the yield mentioned by the same author during chemical extraction, between 40 and 65%, we obtain a margin of oil extracted by press between 62.50 and 75.38% of the total oil, which still appears to be lower than the yield obtained in the present study.

Although this process differs from that used in the production of palm oil (*Elaeis guineensis*), the most consumed and produced oil in the world [30], whose extraction yield is determined based on fresh fruit containing nuts, this yield appears to be slightly higher than that of this species, which yields an average of 20 to 23% of fresh fruit bunches. This yield, which can reach up to 31% from fresh fruit using modern presses, remains lower than that obtained from safou pulp in the present study. [31] report a yield of 36.14% oil extracted from fresh palm fruit using a Dékanmé press. However, if the oil extraction yield from safou pulp is compared to that from whole fruit, i.e., taking into account, as in the case of palm oil, the mass of the kernels, which represents more than 25% of the fruit mass (Table 2), this yield could approach 28%. Furthermore, safou pulp has the advantage of producing oil with a fatty acid composition similar to that of the kernel [32, 33]. Using the whole fruit could increase the fat content [32] and eliminate the pitting stage, thereby reducing processing costs.

This study shows that safou pulp offers satisfactory potential for industrial or semi-industrial exploitation. The total oil yield obtained using this screw press will be able to compete with that of species currently being exploited.

3.2.2 AG composition

The fatty acid composition of the oil extracted from safou pulp using the KOMET press has a profile of C16:0>C18:2>C18:1>C18:0>C18:3, which has already been reported in the literature [11] for safou from Brazzaville. This fatty acid profile, obtained from hexane extraction and representing 32.65% of the 50 trees studied, comes after the C16:0>C18:1>C18:2>C18:0>C18:3 profile, which is the most commonly found profile in the literature [5, 8, 9, 34, 35, 36, 37, 38] and before the C16:0>C18:2>C18:1>C18:3>C18:0 profile representing 2.04%. Another profile, C18:1>C18:2>C16:0>C18:0>C20:0, rarely encountered in the literature, was obtained in Ibadan in Oyo State, Nigeria [33]. Thus, these observations show that the fatty acid composition of safou pulp oil can vary from one tree to another and from one ecological zone to another, with a common profile of C16:0>C18:1>C18:2>C18:0>C18:3, and that the type of extraction has no influence on the fatty acid composition of safou pulp oil.

The differences in FA content obtained and found in the literature remain genotypic in nature. Furthermore, the overall saturated (≃ 50%) and unsaturated (≃ 48%) FA contents are similar to those of oils found in the literature, extracted by pressing [3] and solvent extraction [39, 8, 36, 37], except for samples from [37], which had saturated FA (C16:0, C18:0, and C20:0) contents of 34.12% and unsaturated FA (C18:1 and C18:2) contents of 63.62% (Table 3).

Table 3. Fatty acid composition of safou pulp oil extracted by pressing and solvent extraction, as reported in the literature

Extraction type	**Press**			**Solvent (Soxhlet)**							
Author	[5] Boko-Rep. Congo			[39]	[3] Brazzaville-Rep. Congo			[8] Cameroun		[34] Brazzaville-Rep. Congo	[33] Nigeria	[37] Nigeria
Press/Solvent	Altech BE 100 4/4 (2,2kW)			-	Petroleum ether (harvest date: 01/30/87))			Hexane		Metha-nol/Chloro-form (2/1)	Hexane	Petro-leum ether
	E1 *	E2	E3		A1**	A2	A3	C1***	C2			
C16:0	46.1	44.9	44.8	47.89	47.5	52.1	49.3	39.0	44.0	43.2 ± 0.4	17.18	44.31
C18:0	2.5	2.6	2.4	2.13	3.3	2.7	3.7	3.7	2.3	2.3 ± 0.1	14.84	8.07
C18:1	29.8	32.0	31.4	31.25	23.2	27.1	39.5	31.0	34.2	28.3 ± 0.1	40.45	42.45
C18:2	19.8	18.6	19.8	17.5	26.0	14.2	7.5	24.9	18.6	25.0 ± 0.1	23.17	5.17
C18:3	0.7	0.6	0.7	0.29	-	3.9	-	1.3	0.7	1.1 ± 0.9	-	-
C20:0	0.8	-	-	0.21	-	-	-	-	-		2.10	-

Notes. *Sample 1, **Tree 1, ***Cultivar 1

4. Conclusion

The pulp of the safou studied has an average oil content of 44.89%. The average extraction yield obtained by mechanical hot pressing was 36.44% for three trials, with a low standard deviation of 0.05%. This yield, which represents 81.18% of total oil, is very satisfactory for a screw press and exceeds those of previous studies, making safou a competitive oilseed compared to species currently in production.

The FA composition has a C16:0>C18:2>C18:1>C18:0>C18:3 profile already found in the literature, with saturated and unsaturated FA contents of around 50% and 48%, respectively, similar to those of oils found in the literature, extracted by pressing and solvent extraction. Mechanical pressure extraction has no influence on the fatty acid composition of safou pulp oil.

Acknowledgements

We thank the CIRAD team, UMR Qualisud, Montpellier, France, for the oil extractions and the fatty acid analyses.

References

[1] United Nations Department of Economic and Social Affairs, Population Division. World population prospects 2024. New York: United Nations; 2024.

[2] Odile M, Xavier PX-P. Vegetable oil and fats: Functional resources and nutritional interest. OCL. 2012;19(2):63-75.

[3] Kiakouama S, Silou T. Safou or atanga: A regional fruit in the Gulf of Guinea, physico-chemical analysis. Muntu. CICIBA Rev. 1987;6:137-51.

[4] Tchendji C, Severin M, Wathelet JP, Deroanne C. Composition of Dacryodes edulisfat. Rev Fr Corps Gras. 1987;28:123-5.

[5] Silou T, Goma Manouigui J, Boungou P, Ouamba JM. Study of the drying of safou pulp: Preliminary results. Tropicultura. 1991;9(2):61-8.

[6] Silou T. The safou tree (Dacryodes edulis): A little-known tree. Fruits. 1996;51(1):47-60.

[7] Kinkela T, Bezard J. Study of the structure of triacylglycerols in s1afou pulp oil (Dacryodes edulis). Rev Fr Corps Gras. 1993;11(12):373-82.

[8] Kapseu C, Tchiegang C. Composition of the oil from the fruits of two safou cultivars in Cameroon. Fruits. 1996;51(3):185-91.

[9] Kengue J. Safou (Dacryodes edulis). Monograph. Southampton: International Centre for Underutilised Crops; 2002.

[10] Silou T. Unconventional fats from the Congo Basin: Characterisation, biodiversity and quality. OCL. 2014;21(2):D209. doi:10.1051/ocl/2013044.

[11] Bopoundza FC, Loumouamoui BW, Ngounga Kiminou C, Binaki AF, Biassala EL, Mampouya D, et al. Morphological and physicochemical characterisation of Safous from Congo. Res J Appl Sci. 2020;15(10):296-304.

[12] Okorie PE. Use of Dacryodes edulisin erosion control in agriculture lands in Nigeria. In: Kengue J, ed. Proceedings of the second international workshop on the valorisation of Dacyodes edulisand others non-conventional oil crops. Ngaoundéré, Cameroon: Presses Universitaires de Yaoudé; 1998. p.35-44.

[13] International Centre for Underutilised Crops. Fruits for the future: Safou. Factsheet 3. Southampton: ICUC, University of Southampton; 2001.

[14] Verheij EWM. Dacryodes edulis (G. Don) H.J. Lam. In: PROTA. Plant Resources of Tropical Africa. Wageningen, Netherlands: PROTA; 2002.

[15] Mpungi Buyungu. Contribution of safou (Dacryodes edulis) to agricultural diversification in West Africa. In: Proceedings of the 3rd International Seminar on the Development of Safou Tree and Other Unconventional Oilseeds. Yaoundé, Cameroon: Presses Universitaires d'Afrique; 2002. p.40-9.

[16] Icart-Vernière C, Zougrana S, Nkoudou Ze N, Mouquet-Rivier C. Le safou, un fruit atypique dans l'alimentation traditionnelle au Cameroun. Cah Agric. 2023;32(21). doi:10.1051/cagri/2023013.

[17] Bopoundza FC, Bassiloua JB, Binaki AF, Loumouamoui BW, Biassala EL, Boukouhola A, et al. Modelling of local processes for preserving safou pulp (Dacryodes edulis (G. Dom) H. J. Lam) in Brazzaville, Congo. Int J Food Sci Agric. 2025;9(4):298-314. doi:10.26855/ijfsa.2025.12.006.

[18] Food and Agriculture Organization of the United Nations, Economic Commission for Africa, African Union Commission, World Food Programme. Afrique – Aperçu régional de la sécurité alimentaire et de la nutrition 2023: Statistiques et tendances. Accra: FAO; 2023. doi:10.4060/cc8743fr.

[19] Nwufo MI, Anyim CO. Post-harvest handling and storage of African pear (Dacryodes edulis) in south-eastern Nigeria. In: Kengue

J, ed. Proceedings of the second international workshop on the valorisation of Dacyodes edulisand others non-conventional oil crops. Ngaoundéré, Cameroon: Presses Universitaires de Yaoudé; 1998. p.217-25.

[20] Silou T, Massamba D, Goma Maniongui J, Maloumbi MG, Biyoko SR. Post-harvest losses by natural softening of safou pulp (Dacryodes edulis) in Congo Brazzaville. J Food Eng. 2007;79(2):392-400.

[21] Mayele Kipoy D. Morphological biodiversity and biochemistry of safou (Dacryodes edulis) from Kinshasa and Bas-Congo in the Democratic Republic of Congo [PhD thesis]. Kinshasa: University of Kinshasa; 2010.

[22] Silou Th. Détermination de la teneur en huile de la pulpe de safou (Dacryodes edulis) par refractométrie. Fruits. 1991;46(3):93-100.

[23] Kenmegne Kamdem AT, Ali A, Tchiengang C, Kapseu C. Problématique de la production d'huile de safou au Cameroun. Fruits. 1997;52:325-30.

[24] Mampouya D, Kama Niamayoua R, Goteni S, Loumouamou AN, Kinkela T, Silou T. Optimisation of the Soxhlet extraction of oil from safou pulp (Dacryodes edulis). Adv J Food Sci Technol. 2013;5(3):230-5.

[25] Silou T, Rocquelin G, Gallon G, Molangui T. Contribution à la caractérisation des safou (Dacryodes edulis) d'Afrique centrale. Note II- Composition chimique et caractéristiques nutritionnelles des safous du district de Boko (Congo-Brazzaville). Variation inter-arbre. Riv Ital Sostanze Grasse. 2000;77:85-9.

[26] Silou Th, Rocquelin G, Mouaragadja IG, Gallon G. Contribution of the characterization of safous (Dacryodes edulis) of Central Africa III. Chemical composition and nutritional characteristics of safou of Cameroun, the Congo-Brazzaville, the Congo-Kinshasa and Gabon. Riv Ital Sostanze Grasse. 2002;79:177-82.

[27] Ondo Azi AS, Ella Missang C, Silou T. Classification of Dacryodes edulis (G. Dom) H.J. Lam by using morphological and physical characteristics of the fruits: Statistical approach. For Trees Livelihoods. 2009;19(2):99-109.

[28] Okafor JC. Varietal delimitation in Dacryodes edulis (G. Don) H.J. Lam (Burseraceae). Int Tree Crops J. 1983;2:255-65.

[29] Taty MPE, Loumouamou BW, Elenga M, Baréa B, Barouh N, Danflous JP, et al. Raphia seseunconventional oil from the Congo Basin: Comparison between chemical and screw press extraction. J Am Oil Chem Soc. 2024;101:1-4.

[30] United Nations Conference on Trade and Development. Huile de palme – Un profil de produit de base par INFOCOMM. New York and Geneva: United Nations; 2016.

[31] Adégbola PY, Singbo AG, Ahouansou R, Savi MC. Analyse socio-économique de la presse manuelle à huile de palme "Dé-kanmé". Bull Rech Agron Bénin. 2003;40:1-12.

[32] Silou Th. Le safou (Dacryodes edulis H. J. Lam) : Oléagineux à vocation régionale dans le Golfe de Guinée : Synthèse des données physico-chimiques et technologiques. In: Kengue J, Nya Ngatchou J, eds. Actes du séminaire régional sur la valorisation du safoutier. Douala, Cameroon; 1994. p.123-42.

[33] Ajahid IA, Adesanwo O. Comparative study of the mineral element and fatty acid composition of Dacryodes edulispulp and seed. World J Agric Sci. 2009;5(3):279-83.

[34] Nzondo-Gadet M, Nzikou JM, Kimbonguila A, Linder M, Desobry S. Solvent and enzymatic extraction of Safou and Kolo oils. Eur J Lipid Sci Technol. 2004;106:289-93. doi:10.1002/ejlt.200300919.

[35] Nzondo-Gadet M, Nzikou JM, Matouba E, Etoumongo A, Linder M, Desobry S. Characterisation and nutritional interest of Safou pulp oil. Process Biochem. 2005;40:307-12. doi:10.1016/j.procbio.2004.01.003.

[36] Silou T. Oils and fats for the future. A case study: Safou (Dacryodes edulis) from the Congo Basin countries in Africa. New York: Nova Science Publishers, Inc.; 2012.

[37] Ibanga OI. Extraction, characterization of African Pear (Dacyodes edulis) oil and its application in synthesis and evaluation of surface coating driers. Int J Adv Res Chem Sci. 2014;1(4):14-22.

[38] Noumi GB, Djounja T, Ngameni E, Kapseu C. Influence of the storage time on the fats and oil composition of safou (Dacryodes edulis), dried pulp. Int Food Res J. 2014;21(5):1837-41.

[39] Omoti O, Okiy DA. Characteristics and composition of the pulp oil and cake of the African pear, Dacryodes edulis (G. Don) H.J Lam. J Sci Food Agric. 1987;38(1):67-72.