

Evaluation of a car-based method for detecting the invasive *Lupinus polyphyllus* Lindl in road infrastructure

Juliana Daniel-Ferreira

`juliana.daniel.f@slu.se`

Swedish University of Agricultural Sciences, Swedish Biodiversity Centre

Jan Olof Helldin

Swedish University of Agricultural Sciences, Swedish Biodiversity Centre

Tommy Lennartsson

Swedish University of Agricultural Sciences, Swedish Biodiversity Centre

Article

Keywords: invasive alien plants, detection probability, road verge management, environmental monitoring

Posted Date: March 3rd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-8828049/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on June 25th, 2026. See the published version at <https://doi.org/10.1038/s41598-026-58979-w>.

Abstract

Growing recognition of the negative impacts of invasive alien plants on native species diversity and ecosystem function has led to greater focus on effective management strategies. Monitoring is crucial for both prevention of establishment and planning of eradication of established IAPs. However, monitoring is often constrained by resource limitations, and less time-consuming methods risk providing too low levels of detection.

Lupinus polyphyllus is a highly problematic non-native plant species in Europe. We evaluated the accuracy of a car-based survey method for finding *L. polyphyllus* in road verges and how the physical characteristics of the stands influenced their detection from the car. At the 200 m scale the accuracy of detecting *L. polyphyllus* by car was 85%, while at the scale of individual stands the accuracy was lower (72%). Results indicated that detection probability was influenced by stand density and position in the road verge. Dense stands had the highest detection probability, while single plants were less likely to be detected. Stands present in side and back slopes were more likely to be detected than those only in one slope, with the side slope stands having the lowest detection probability.

Our study shows that the car-based survey is a reliable method for detecting *L. polyphyllus* and estimating its abundance in road verges. Such information is useful for designing management strategies to hinder seed production and dispersal of the invasive. Since the accuracy of detection of single plants was lower, more detailed surveys at local scales are recommended for early eradication.

Introduction

Invasive alien species (IAS) are species that have been introduced beyond their native range and have become problematic in the novel ecosystems due to rapid expansion, often linked to lack of natural enemies. IAS therefore pose a threat to ecosystems worldwide (Pyšek et al., 2020; Rai & Singh, 2020). Among them, invasive alien plants (IAP) have been shown to reduce native species abundance and diversity, and can even alter ecosystem function (Vilà et al., 2011). Recent projections indicate that the rate of new introductions is rising, underlining the urgency of effective management strategies to mitigate this global challenge (Seebens et al., 2018). This exerts a growing pressure on governmental agencies and environmental managers to tackle a wide array of challenges that emerge at the different stages of invasion process (Hulme, 2006). Although prevention (i.e. early detection, rapid response, and eradication) is the main strategy for control of IAS (Maxwell et al., 2009; Simberloff et al., 2013), other management strategies need to be applied for the control of species that have already established and are at the late stages of the invasion process (Simberloff et al., 2013). At these stages, monitoring the current state of the invasion and the efficacy of management measures, as well as anticipating where the species will spread becomes increasingly important (Maxwell et al., 2009; Shackleton et al., 2020). Monitoring is also essential for keeping track of potential IAP species.

Extensive monitoring requires considerable resources, and financial constraints can hinder comprehensive monitoring efforts, potentially leading to gaps in data collection (Epanchin-Niell et al.,

2012). Furthermore, probability of detection during field surveys is often imperfect and varies depending on several factors. For plants, detection variability during surveys could originate from the observers and their level of experience (Milberg et al., 2008), size and growth form of the stand (Chen et al., 2009), conditions specific to the type of survey and environment (Chen et al., 2009), and interactions among these factors (Hauser et al., 2022). Despite these challenges, effective long-term monitoring of IAP remains essential for informed decisions in ecosystem management (Schulze-Brüninghoff et al., 2021). Failure to detect IAP results in the spread of the species and increases costs because implemented management strategies risk to miss their targets (Hauser et al., 2022). Therefore, it is important to explore new survey methods to establish their cost-effectiveness and the variation in the probability of detecting the target species.

Road verges have been shown to be an important conduit for the expansion of IAP (Christen & Matlack, 2006; Lázaro-Lobo & Ervin, 2019; Lemke et al., 2021). Seeds can disperse over long distances along the road network through the movement of vehicles (Von Der Lippe et al., 2013) and through road building and maintenance (Rauschert et al., 2017). To monitor the status and spread of IAP along road verges, researchers and practitioners have employed different approaches such as remote sensing using satellite (e.g. Lourenço et al. 2021), or aerial (Müllerová et al., 2005) imagery, georeferenced imagery, e.g. Google Street View (Kotowska et al. 2021), as well as regular field surveys (e.g., Joly et al. 2011). While each method provides distinct advantages over the others, they often also have limitations and are thus not appropriate in all situations. For instance, remote sensing images are easy to access and have broad spatial coverage, but the images are often too coarse and their analysis can be challenging due to similarity among species (Zweig & Newman, 2015). Furthermore, the detection of the plant species in question is facilitated by performing the survey at the right time, preferably when the plants are distinguishable through phenologically prominent features such as flowering stage (Schulze-Brüninghoff et al., 2021), making it challenging to use images collected for other purposes.

The Swedish Transport Administration (STA) routinely surveys selected road verges for IAP. The goal of the surveys is to detect new occurrences for eradication and also to monitor the spread and distribution of the species in the road network. However, the survey method varies among regions in the country. Here, we present and evaluate a method that could potentially serve as standard. The method could be used both by practitioners and researchers to achieve a more seamless collaboration between the two.

Lupinus polyphyllus (hereafter lupine) is one of the most problematic non-native species in Europe due to its environmental and economic impacts (Eckstein et al., 2023). Despite it not being included in the European Union's list of invasive alien species of conservation concern, the species is considered invasive and the Swedish Environmental Protection Agency is striving to reduce its spread (Naturvårdsverket, 2023; Nentwig et al., 2018). Given that lupine is the most common IAP in the Swedish road network, we developed a method to detect the species along road infrastructure. Since the method is designed to be used to survey the entire road network in a region, it needs to be fairly fast without losing too much accuracy. Therefore, a car-based method was designed, which both has the potential to fulfil these requirements, and aligns to the routines for road monitoring already in use by the STA. We

evaluated the detection accuracy of the method, as well as which characteristics of the lupine stands can affect their detection probability.

We hypothesize that the factors likely affecting the detection rate of lupines in our survey method are the density of plants in a stand, the location of the stand in the road verge, and whether the stand occurs both in the road verge and outside the road area or not.

Methods

Survey method and areas of study

To perform the survey, all accessible road verges within the selected area were scanned for lupines from a car with two people. One person drove the car while the other looked for the invasive along the road verges and registered the surveyed roads and occurrences of the species in a digital map and protocol. The recommended speed ranged between 20 and 50 km/h, depending on the density of stands and the width of the road. The speed was reduced as the density of lupines in the road verge increased or the visibility decreased, and if necessary the car could stop completely next to a stand (e.g. in order to assess stand density and to acquire as accurate positioning as possible). If the road was sufficiently narrow to allow good visibility of both road verges, the observer was advised to register the stands present in both sides of the road whilst driving through the road only one time. However, if the road was wide and visibility of the road verge on the other side was impeded, the survey was performed for each side of the road separately. The method is described in detail in Helldin et al. (2024).

The goal of the survey method was to collect data on both the distribution and the abundance of lupine in the selected regions. The abundance was measured by the total number of occurrences, their size, and also the density of plants within a stand. Using the ESRI app Field Maps (earlier Collector), each stand was marked on a basemap with a polyline object that marked its beginning and end. Through the Field Maps protocol, each line object also contained information about the density, position, and width of the stand. It was also registered whether the stand continued outside of the road verge into the neighbouring habitat or not. The density was selected by the observer as one of the following categories: individual plant(s), sparse stand (the IAP covered less than 50% of the stand surface), dense stand (the IAP covered approximately between 50 and 90% of the stand surface) and dominating (the IAP covered more than 90% of the stand surface). Stand density can be assumed to be important for the lupine's effects on plant communities (Dániel-Ferreira et al., 2024; Goossens et al., 2024), and the chosen density categories are justified by what can easily be assessed in the field. Plant density is difficult to assess (Morrison, 2016) and can easily become subjective. However, the boundary above/below 50% is usually discernible, as well as very high coverage. The position of the stand could be one of three choices: side slope (i.e. the road embankment), back slope, or both side and back slopes (Fig. 1).

For simplicity, the width of the stand was an approximation given by the observer as an integer starting at 1 m. The width of the stand could subsequently be used to create polygons to acquire a more comprehensive representation of the abundance of the species, as well as an estimate of the area in

need of control measures. A new line object was created if there was a break in the occurrences or if there was a pronounced change of stand width. The length of the gap needed to register a new stand depended on the circumstances. With a low number of plants, a 10 m gap was applied. If it became too time consuming, the occurrences were registered as a single line object with patchy occurrences (i.e. a yes / no question in the protocol related to the patchiness of the stand). A new line could also be registered if there was a clear change in density. This was only relevant for larger stands with reasonably homogeneous density, for example where there was a large dense core stand that transitioned into sparser or varied stands. In the case of smaller occurrences with different densities (which are <10m apart), a line with varied density was registered.

The survey was conducted in four areas during the peak season of lupine flowering, which is early June in the southern study areas, and late July in the northern study area (Fig. 2).

All observed occurrences of lupines, both flowering and vegetative individuals, were registered during the survey. Three of the areas were located in the boreo-nemoral lowland in south-central Sweden, to the west of the city of Uppsala, and the fourth was located in a subalpine area further north, around Funäsdalen in the province of Härjedalen. The areas were selected to represent a variation in overall lupine abundance and to include different landscape types, subalpine forest, lowland forest and lowland agricultural. Two of the lowland areas (Heby and Uppsala) were surveyed in 2021. In the summer of 2022, the Heby area was extended, and some areas were surveyed again. The Funäsdalen area was surveyed in 2022. During 2023, the third lowland area, Enköping, was surveyed (Fig. S1–S4).

Evaluation of the survey method

To assess the type and proportion of lupine stands that were detected during the survey by car, we selected 131 verification stretches, i.e. shorter road stretches (200 m) where a more detailed survey was conducted by foot instead of by car (~26.2 km in total). During these more detailed inspections, the surveyors walked the length of the stretches along both sides of the road at a slow pace while searching for lupines on the road verge from a closer distance and without obstructed view. We selected the verification stretches at random points along road sections distributed over the respective study area (Fig. S1 – S4). Verification stretches were of two types: i) *non-empty stretches*: where lupines were recorded during the survey by car, and ii) *empty stretches*: stretches that were considered empty during the survey by car. For the survey by foot, the observers started by parking the car at the predetermined given points and then proceeded to survey the road verges on both sides 100 m in opposite directions from the given point (i.e., 200 m road stretch and 400 m roadside in total). The observers recorded lupine occurrences using the same Field Maps protocol as during the survey by car (see above).

Occurrences (line objects) recorded during the survey by foot were compared with those registered during the survey by car in ArcGIS Pro 3.1. An examiner performed a visual desktop assessment of each line object to determine whether it had been missed during the survey by car or not. In general, if two registrations of stands (one from the survey by car, the other from the survey by foot) overlapped, the stand was marked as 'detected during the survey by car'. To allow for some uncertainty due to the

varying accuracy of the GPS, objects that were within a 10 m distance from each other were also counted as 'detected' during the survey by car. If the examiner was unsure during inspection as to whether the object had been missed or not, a closer inspection of the characteristics of the stand was performed. This allowed the examiner to determine whether a long stand identified during the survey by car as having varying densities, was broken up into shorter stands during the survey by foot.

Statistical analysis

Analyses were performed in R 4.1.2 (R Core Team, 2021). A generalized linear mixed model (GLMM) with a binomial distribution and a logit link was used to investigate which variables influenced the probability of detecting or missing a lupine stand from the car. The response variable was binary: 'Yes' was given for a stand that was detected both from the car and by foot, and 'No' was indicated for a stand that was detected only during the survey by foot (i.e., the object was not detected during the survey by car). The explanatory variables were the characteristics of the stand as described by the survey by foot: stand location within the road verge (factor), stand density (factor), and whether the stand continued outside of the road verge (factor). In the analysis, the density category 'dominant' was removed because there were only three observations along the verification stretches and all of them were detected from the car (Fig. 3). The stand location was of interest because we assumed that plants present in the side slope could be more easily missed when viewed from the road. The density of the stand was of particular interest as we assumed that denser stands (i.e. with more individuals per m²) are easier to detect from the car. A factor indicating whether the stand continued outside of the road verge was added to the model given that this could also be an indicator of the size of the stand. The density was assessed for each stand parallel to the road, while the presence of the stand outside the road verge was an indicator of the size perpendicular to the road, which could also influence the visibility from the car. Finally, the study area was added as a random effect to account for the variability emerging from the differences in the distribution of lupines among areas, differences that may arise from the different observers that performed the survey, and from the survey being conducted in the three different years. We did not consider any interactions among the variables of interest to be relevant for this analysis. We also assumed weather and observer to be of importance but due to lack of variation in the data collected these variables could not be examined (most of the data was collected in sunny weather).

After the model was fitted, it was visually validated using the package *DHARMa* (Hartig, 2020). To investigate exactly what characteristics of the stand were related to a lower detection probability during the car inventory, a post-hoc analysis was performed using the *emmeans* package (Lenth, 2020). To adjust for multiple comparisons, the Tukey method for comparing a family of three estimates was applied. Finally, we compared the fitted model to a null model using Akaike's information criterion corrected for small sample sizes (AICc).

Finally, given that the speed during the survey by car varied according to the density of lupines in the road verges, we also performed a separate analysis to evaluate how speed influenced the detection of lupines from the car. The speed was only recorded during the first two years of the survey along the

verification stretches, and therefore a separate analysis that used only the data from 2021 and 2022 was performed. Vehicle speeds along the verification stretches were recorded in 10 km/h intervals and ranged from 20 to 50 km/h. The highest speed occurred only along one stretch and was therefore excluded from the analysis. We evaluated how an interaction between the explanatory variables explained above (i.e. stand location within the road verge, stand density, and whether the stand continued outside of the road verge) influenced whether the stand was detected or missed during the survey by car. We added the interacting effect to each of the explanatory variables in separate models and calculated the AICc to detect the most parsimonious models.

Results

There was a total of 2,103 lupine stands found along 2,933 km within the four areas (Table 1-2, Fig. 4). Of the 131 verification stretches, 61 stretches were identified as empty from the car. Among the 61 stretches classified as empty from the car, lupines were found in nine (14.75%) of these stretches during the survey by foot, meaning that lupines had been missed during the survey by car. Along non-empty stretches, 223 stands of flowering lupine were observed during the survey by foot, and 27.8% ($n = 62$) of them were missed during the survey by car (Fig. 3). In 2021, 46 stands were observed in Uppsala and Heby, and 11 (23.91%) were missed during the survey by car. In 2022, 83 stands were observed in Uppsala, Heby, and Funäsdalen and 36 (43.37%) were missed during the survey by car. In 2023, 94 lupine stands were observed in Enköping and 15 (15.96%) were missed during the survey by car (Table S1). No commission errors were recorded during the survey by car (i.e. no stretches were falsely identified as containing lupines).

The results from the analysis of factors influencing probability of detection shown below are based on non-empty stretches only ($n = 79$). In general, the density and position of the lupine stands influenced the probability of detection from the car (Table 3, Table S2). Single plants were harder to detect from the car, but there were no differences in detection probability between single plants and sparse stands. Dense stands had the highest probability of detection (Fig. 5a). Stands that only occurred on one of the slopes (i.e., either side slope or back slope) had lower probability of detection than stands that occurred on both slopes (Fig. 5b). The continuation of the stand outside of the road verge did not influence the detection probability (Fig. 5c). The candidate model had an AICc = 198.12 and the null model had an AICc = 254.82 indicating that the candidate model was better than the null model. There was no evidence of collinearity among explanatory variables (Table 3).

The speed of the car influenced the detection of lupine stands during the survey by car. In general, the probability of detecting a stand was higher at lower speeds and steadily decreased with increasing speed (Fig. 6). This pattern was most noticeable for single plants (Fig. 6a) and for stands that were present only on the side slope (Fig. 6b). The probability of detecting dense, i.e. more conspicuous, stands remained high at higher speeds (Fig. 6a). The most parsimonious models were the ones in which the interaction effect was modeled separately for each variable ($\Delta\text{AICc} < 3$; Table S3-S6). Therefore, the figures show the results of three separate models in which the interacting effect is added to each

explanatory variable separately. Using a subset of the data did not change the results mentioned above, although the p-values increased (Table S7).

Discussion

We showed that the survey by car method is a promising method for finding and estimating the abundance of lupines along longer stretches of road verges or in a network of road verges. We found that the detection probability of lupine stands depended on the speed of the car and the density and position of the stand in the road verges but was independent of whether the stand continued also outside of the road verge. The accuracy was higher on a larger spatial scale (ca. 85% accuracy of detecting lupine occurrence at 200 m scale as compared to ca. 72 % accuracy on the level of individual stands). The survey by car can thus offer road managers a general idea of which road segments require control measures. Given that *L. polyphyllus* is well established in the country, complete eradication at the national level is no longer feasible. In such cases, the survey by car method would be primarily focused on supporting the control and containment of populations to prevent further spread. Identifying areas where eradication is still possible are keys to achieve this goal.

As predicted, the density of plants in a stand affected the probability of detecting the stand from the car such that the probability increased with the size of the stand. There was a higher, but not significantly so, probability of detecting sparse stands compared to single plants. (Fig. 5a). However, we suspect that the probability of detection of single plants is even lower than the one we report here, given that stands marked as 'single plants' could either be actual single plants or longer line objects with dispersed single plants (which we assume increased their detection probability). Nevertheless, detecting a single plant from the car was less likely in comparison, meaning that the accuracy of the method will vary depending on the proportion of the total lupine population in an area that consists of single plants. The position of the lupine stands in the road verge also affected their detection probability such that stands that were present on both side and back slopes had a significantly higher detection probability compared to stands that were only present in one of the slopes. Given that the detection probability is assumed to be mostly affected by the size of the individual plants and of the stands, target areas should be surveyed regularly (ideally every year), to allow for single plants or smaller stands to be more easily detected as they grow.

The accuracy of the survey by car increased at lower speeds, with 20 km/h providing the highest probability of detecting lupine stands in the road verge. As the speed of the vehicle increased, the probability of detecting single plants and plants located in the side slope decreased. The reduction of accuracy was most pronounced for single plants, with a drop from ~80% probability of detection at 20 km/h to less than 25% at 40 km/h. These results have important implications for planning surveys by car, and survey objectives and availability of resources should guide the choice of driving speed. If the goal is to produce a coarse distribution map of established lupine stands across a broader road network, higher speeds may be acceptable. In contrast, when the objective is early detection of newly established plants, lower speeds are recommended to maximize the probability of detection. Since the cost of eradication of IAP is lowest at an early stage of invasion, there is thus a trade-off between cost of the survey (higher

cost at lower speed due to higher time consumption) and cost of eradication (lower cost with earlier detection following lower speed). Nevertheless, our results suggest that detecting single plants is challenging because the probability of detection drops rapidly when the speed exceeds 20 km/h.

Other possible factors influencing detection probability of lupine stands that were not considered in this study are the background vegetation and observer error. For instance, Hauser et al. (2022) found that detection rate of *Pilosella* plants was determined by the interaction between target and background characteristics. In this case, the target and background were the same, that is, lupines and road verges, but variation at a more detailed level might have been of importance. For example, vegetation height surrounding the lupines could have impacted the probability of detection. Furthermore, lupine flowers can adopt a variety of colours such as white, pink, and purple, which could potentially affect their detection probability, and there could be possible differences in detection probability between lupine stands in a road verge next to a forest compared to a road verge next to an agricultural field. Observer error has been repeatedly shown to influence detection probability in plant surveys (Morrison, 2016). We did not explore how the observer influenced detection probability of lupine stands because the identity of the surveyor was not clear in the database. Another reason is that we believe that blooming individuals of lupines are among the easiest species to identify in the field, and even observers with a low degree of experience in plant surveys should be able to identify them. Consistent with this expectation, we did not record any commission errors during the survey by car. This is supported by previous studies, showing that there were no remarkable differences in detection probability between observers when the species are conspicuous and grow tall (e.g. Kéry & Gregg 2003; Chen et al. 2009). When there is no need to differentiate among species, the observer bias should be lower, and detection probability becomes more dependent on the characteristics of the types of stands and the environment they are in. However, estimation of plant abundance (in this case stand density) is highly subjective and has often very low accuracy (Morrison, 2016), which might have influenced our results and could have been partly controlled for if we had added the observer to the model.

Even though the proposed survey by car method was designed with the species *L. polyphyllus* in mind, the method could potentially be adapted to surveying other IAP in the road network. Conspicuous species that are easy to identify even for inexperienced observers such as *Solidago canadensis* (Canadian goldenrod), *Impatiens glandulifera* (Himalayan balsam) or *Heracleum mantegazzianum* (giant hogweed) could potentially be surveyed using this method, although with adapted categories of stand density. Due to differing flowering phenology, however, not all of those species can be covered in the same survey. Moreover, IAP with less distinctive morphology may have a higher risk of false positives, which needs to be taken into account when adapting the method to other taxa.

Conclusions

Road verges have been shown to often have a greater abundance and richness of exotic species and therefore need continuous and cost-effective monitoring of IAP. The suggested method will correctly

identify 85% of the 200 m sections with lupines, and 72% of the lupine stands. In addition, the method correctly identified 100% of the 200 m sections without lupines. If repeated, the method also shows changes in the distribution and abundance of lupines in an area. Based on a distribution map, eradication strategies can be developed. However, given the accuracy limitations of the survey by car method, eradication measures should be preceded by more detailed surveys and inventories of the stands in and around the chosen areas for eradication in order to find all plants, also early establishments.

The substantial geographical scope allowed by the survey by car method will enable land managers to identify areas to prioritize for eradication of lupine populations. The strategy applied to prevent further spread can vary depending on the type of stands and their locations, as well as on the availability of resources. For instance, small and isolated stands could be prioritized for eradication in order to prevent further establishment into new areas. Since eradication measures may not be 100% effective, and due to some IAP having a seed bank, most eradication measures require continuous monitoring and, if needed, repeated eradication for some years to ensure that the control measure is successful. Alternatively, road stretches with high densities of the invasive could be isolated from the rest of the road network by implementing adaptive management, e.g. cutting before seed-set or by making sure that the machinery used is lupine-free either by cleaning the grass mowers or by using specific grass mowers for lupine-free areas and others for areas with high lupine densities. This strategy requires continuous monitoring of the IAP-free parts of the road network.

Declarations

All authors have read, understood, and have complied as applicable with the statement on “Ethical responsibilities of Authors” as found in the Instructions for Authors. There were no human participants involved in this study.

Conflict of interest:

The authors declare no competing interests.

Acknowledgements

We are particularly grateful to Jörgen Wissman who was instrumental in the early project stages, including Funding acquisition, Conceptualization, Methodology, Supervision and Data curation. Izabel Eriksson Reuterborg, Andreas Ekman, Hanna Karlsson, Eleonore Larsson, Michael Kinsella, Sonja Reimer, Samuel Persson, Karl Soler Kinnerbäck, and Erica Johnson assisted with the data collection. The study was funded by the Swedish Transport Administration (Trafikverket).

Author contributions

Juliana Dániel-Ferreira: Methodology, Data curation, Formal analysis, Writing – original draft, Visualization **Jan-Olof Helldin:** Conceptualization, Funding acquisition, Methodology, Writing – review &

editing **Tommy Lennartsson**: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing – review & editing

Data availability

The dataset generated and/or analyzed during the current study are available in FigShare at <https://doi.org/10.6084/m9.figshare.31347463>.

Funding:

Swedish Transport Administration, research programme TRIIAS

References

1. Chen, G., Kéry, M., Zhang, J., & Ma, K. (2009). Factors affecting detection probability in plant distribution studies. *Journal of Ecology*, 97(6), 1383–1389. <https://doi.org/10.1111/j.1365-2745.2009.01560.x>
2. Christen, D., & Matlack, G. (2006). The Role of Roadsides in Plant Invasions: A Demographic Approach. *Conservation Biology*, 20(2), 385–391.
3. Dániel-Ferreira, J., Lennartsson, T., Wissman, J., Knudsen, C., & Eckstein, R. L. (2024). Vascular plant diversity in Swedish road verges of high conservation value is threatened by the invasive alien herb *Lupinus polyphyllus* Lindley. *Nordic Journal of Botany*, e04438. <https://doi.org/10.1111/njb.04438>
4. Eckstein, R. L., Welk, E., Klinger, Y. P., Lennartsson, T., Wissman, J., Ludewig, K., Hansen, W., & Ramula, S. (2023). Biological Flora of Central Europe–*Lupinus polyphyllus* Lindl. *Perspectives in Plant Ecology, Evolution and Systematics*, 125715. <https://doi.org/10.1016/j.ppees.2022.125715>
5. Epanchin-Niell, R. S., Haight, R. G., Berec, L., Kean, J. M., & Liebhold, A. M. (2012). Optimal surveillance and eradication of invasive species in heterogeneous landscapes. *Ecology Letters*, 15(8), 803–812. <https://doi.org/10.1111/j.1461-0248.2012.01800.x>
6. Goossens, E. P., Mertens, W., & Olde Venterink, H. (2024). *Solidago gigantea* invasion homogenizes soil properties and native plant communities. *Biological Invasions*, 26(10), 3315–3327. <https://doi.org/10.1007/s10530-024-03374-7>
7. Hartig, F. (2020). DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. *R Package Version 0.3.3.0*. <https://CRAN.R-project.org/package=DHARMA>
8. Hauser, C. E., Giljohann, K. M., McCarthy, M. A., Garrard, G. E., Robinson, A. P., Williams, N. S. G., & Moore, J. L. (2022). A field experiment characterizing variable detection rates during plant surveys. *Conservation Biology*, 36(3). <https://doi.org/10.1111/cobi.13888>
9. Helldin, J.-O., Dániel-Ferreira, J., Wissman, J., & Lennartsson, T. (2024). *Inventeringsmetod för blomsterlupin i vägkanter*. <https://triokol.se/project/inventeringsmetod-for-blomsterlupin/>

10. Hulme, P. E. (2006). Beyond control: Wider implications for the management of biological invasions. *Journal of Applied Ecology*, 43(5), 835–847. <https://doi.org/10.1111/j.1365-2664.2006.01227.x>
11. Joly, M., Bertrand, P., Gbangou, R. Y., White, M.-C., Dube, J., & Lavoie, C. (2011). Paving the Way for Invasive Species: Road Type and the Spread of Common Ragweed (*Ambrosia artemisiifolia*). *ENVIRONMENTAL MANAGEMENT*, 48(3), 514–522. <https://doi.org/10.1007/s00267-011-9711-7>
12. Kéry, M., & Gregg, K. B. (2003). Effects of life-state on detectability in a demographic study of the terrestrial orchid *Cleistes bifaria*. *Journal of Ecology*, 91(2), 265–273. <https://doi.org/10.1046/j.1365-2745.2003.00759.x>
13. Kotowska, D., Pärt, T., & Žmihorski, M. (2021). Evaluating Google Street View for tracking invasive alien plants along roads. *Ecological Indicators*, 121, 107020. <https://doi.org/10.1016/j.ecolind.2020.107020>
14. Lázaro-Lobo, A., & Ervin, G. N. (2019). A global examination on the differential impacts of roadsides on native vs. Exotic and weedy plant species. *Global Ecology and Conservation*, 17, e00555. <https://doi.org/10.1016/j.gecco.2019.e00555>
15. Lemke, A., Buchholz, S., Kowarik, I., Starfinger, U., & Von Der Lippe, M. (2021). Interaction of traffic intensity and habitat features shape invasion dynamics of an invasive alien species (*Ambrosia artemisiifolia*) in a regional road network. *NeoBiota*, 64, 155–175. <https://doi.org/10.3897/neobiota.64.58775>
16. Lenth, R. (2020). Emmeans: Estimated Marginal Means, aka Least-Squares Means. *R Package Version 1.5.2-1*. <https://CRAN.R-project.org/package=emmeans>
17. Lourenço, P., Teodoro, A. C., Gonçalves, J. A., Honrado, J. P., Cunha, M., & Sillero, N. (2021). Assessing the performance of different OBIA software approaches for mapping invasive alien plants along roads with remote sensing data. *International Journal of Applied Earth Observation and Geoinformation*, 95, 102263. <https://doi.org/10.1016/j.jag.2020.102263>
18. Maxwell, B. D., Lehnhoff, E., & Rew, L. J. (2009). The Rationale for Monitoring Invasive Plant Populations as a Crucial Step for Management. *Invasive Plant Science and Management*, 2(1), 1–9. <https://doi.org/10.1614/IPSM-07-054.1>
19. Milberg, P., Bergstedt, J., Fridman, J., Odell, G., & Westerberg, L. (2008). Observer bias and random variation in vegetation monitoring data. *Journal of Vegetation Science*, 19(5), 633–644. <https://doi.org/10.3170/2008-8-18423>
20. Morrison, L. W. (2016). Observer error in vegetation surveys: A review. *Journal of Plant Ecology*, 9(4), 367–379. <https://doi.org/10.1093/jpe/rtv077>
21. Müllerová, J., Pyšek, P., Jarošík, V., & Pergl, J. (2005). Aerial photographs as a tool for assessing the regional dynamics of the invasive plant species *Heracleum mantegazzianum*. *Journal of Applied Ecology*, 42(6), 1042–1053. <https://doi.org/10.1111/j.1365-2664.2005.01092.x>
22. Naturvårdsverket. (2023). *Blomsterlupin*. <https://www.naturvardsverket.se/amnesomraden/invasiva-frammande-arter/Arter/arter-som-ej-omfattas-av-regler/blomsterlupin/>

23. Nentwig, W., Bacher, S., Kumschick, S., Pyšek, P., & Vilà, M. (2018). More than “100 worst” alien species in Europe. *Biological Invasions*, 20(6), 1611–1621. <https://doi.org/10.1007/s10530-017-1651-6>
24. Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T., Dawson, W., Essl, F., Foxcroft, L. C., Genovesi, P., Jeschke, J. M., Kühn, I., Liebhold, A. M., Mandrak, N. E., Meyerson, L. A., Pauchard, A., Pergl, J., Roy, H. E., Seebens, H., ... Richardson, D. M. (2020). Scientists’ warning on invasive alien species. *Biological Reviews*, 95(6), 1511–1534. <https://doi.org/10.1111/brv.12627>
25. R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing [Computer software]. <https://www.R-project.org/>
26. Rai, P. K., & Singh, J. S. (2020). Invasive alien plant species: Their impact on environment, ecosystem services and human health. *Ecological Indicators*, 111(106020), 106020. <https://doi.org/10.1016/j.ecolind.2019.106020>
27. Rauschert, E. S. J., Mortensen, D. A., & Bloser, S. M. (2017). Human-mediated dispersal via rural road maintenance can move invasive propagules. *Biological Invasions*, 19(7), 2047–2058. <https://doi.org/10.1007/s10530-017-1416-2>
28. Schulze-Brüninghoff, D., Wachendorf, M., & Astor, T. (2021). Potentials and limitations of worldview-3 data for the detection of invasive lupinus polyphyllus lindl. In semi-natural grasslands. *Remote Sensing*, 13(21). Scopus. <https://doi.org/10.3390/rs13214333>
29. Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., Pagad, S., Pyšek, P., van Kleunen, M., Winter, M., Ansong, M., Arianoutsou, M., Bacher, S., Blasius, B., Brockerhoff, E. G., Brundu, G., Capinha, C., Causton, C. E., Celesti-Grapow, L., ... Essl, F. (2018). Global rise in emerging alien species results from increased accessibility of new source pools. *Proceedings of the National Academy of Sciences*, 115(10). <https://doi.org/10.1073/pnas.1719429115>
30. Shackleton, R. T., Petitpierre, B., Pajkovic, M., Dessimoz, F., Brönnimann, O., Cattin, L., Čejková, Š., Kull, C. A., Pergl, J., Pyšek, P., Yoccoz, N., & Guisan, A. (2020). Integrated Methods for Monitoring the Invasive Potential and Management of *Heracleum mantegazzianum* (giant hogweed) in Switzerland. *Environmental Management*, 65(6), 829–842. <https://doi.org/10.1007/s00267-020-01282-9>
31. Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E., & Vilà, M. (2013). Impacts of biological invasions: What’s what and the way forward. *Trends in Ecology & Evolution*, 28(1), 58–66. <https://doi.org/10.1016/j.tree.2012.07.013>
32. Vilà, M., Espinar, J. L., Hejda, M., Hulme, P. E., Jarošík, V., Maron, J. L., Pergl, J., Schaffner, U., Sun, Y., & Pyšek, P. (2011). Ecological impacts of invasive alien plants: A meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters*, 14(7), 702–708. <https://doi.org/10.1111/j.1461-0248.2011.01628.x>
33. Von Der Lippe, M., Bullock, J. M., Kowarik, I., Knopp, T., & Wichmann, M. (2013). Human-Mediated Dispersal of Seeds by the Airflow of Vehicles. *PLoS ONE*, 8(1), e52733. <https://doi.org/10.1371/journal.pone.0052733>

34. Zweig, C. L., & Newman, S. (2015). Using landscape context to map invasive species with medium-resolution satellite imagery: Using landscape context to map invasives. *Restoration Ecology*, 23(5), 524–530. <https://doi.org/10.1111/rec.12214>

Tables

Table 1. Number of verification stretches (surveyed by foot) per survey area, separated by year and output from the preceded survey by car.

Area	Year of survey	Number of non-empty stretches	Number of empty stretches
Heby	2021	10	10
Heby	2022	20	11
Uppsala	2021	10	10
Uppsala	2022	2	8
Funäsdalen	2022	7	3
Enköping	2023	21	19

Table 2. Number of surveyed kilometres per area and year, and the total number of lupine stands registered for each area. Note that the number of lupine stands per kilometre is a rough estimate of the density of the species in the road network within each area.

Area	Km 2021	Km 2022	Km 2023	Total km	Total lupine stands	Lupine stands / km
Heby	667.5	383.9	-	1051.5	1145	1.1
Enköping	-	-	395.4	395.4	564	1.4
Uppsala	723.5	230.6	-	954.1	169	0.2
Funäsdalen	-	532.1	-	532.1	225	0.4

Table 3. Results from the generalized linear model exploring which characteristics of the lupine stands are important for detection from a moving vehicle. Shown are the Chi-square values (χ^2), degrees of freedom (df), p-values (p), and variance inflation factors (GVIF^{(1/(2*df))}). P-values in bold are significant at the 0.05 level or lower. The estimates, standard errors, z-values, and p-values can be found in Table S1. Marginal R² = 0.44, conditional R² = 0.52.

	χ^2	<i>df</i>	<i>p</i>	<i>GVIF</i> ^{(1/(2*<i>df</i>))}
Stand location	16.44	2	<0.001	1.10
Density of the stand	8.24	2	<0.01	1.06
Stand continues outside the road verge	0.00	1	0.99	1.20

Figures




Figure 1

Illustration of a cross section of a road and the road verge. Plants of *L. polyphyllus* could be observed in the side slope, the back slope, or both. Stands could also continue outside of the road verge




Figure 2

Map of the four areas (Heby, Enköping, Uppsala, and Funäsdalen) surveyed using the survey by car method and verified by the survey by foot method




Figure 3

Description of the type of lupine stands, their location in the road verge, and whether they were detected or missed during the survey by car




Figure 4

Total number of lupine stands and their plant density in the four surveyed areas in Sweden




Figure 5

Result of the GLM exploring which characteristics of a lupine stand affect their probability of detection. a) Stand density, b) Stand location in the road verge, and c) Whether the stand continues outside of the road verge (Yes) or not (No). The letters represent the results from the post-hoc analysis. Comparisons that do not share a letter are significantly different from each other. The error lines represent the standard error




Figure 6

Influence of speed during the survey by car on the probability of detection lupine stands that differ in a) density, b) position in the road verge, and c) whether the stand continues outside of the road verge (Yes) or not (No). The shaded areas represent the 95% confidence intervals. Points represent the individual observations

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterial.docx](#)