

Biochemical and agronomic responses of guar to nitrogen, *Bradyrhizobium* inoculation and deficit irrigation under arid and semi-arid conditions

Received: 22 October 2025

Accepted: 5 May 2026

Published online: 11 May 2026

Cite this article as: Akbari G., Norouzi S., Alahdadi I. *et al.* Biochemical and agronomic responses of guar to nitrogen, *Bradyrhizobium* inoculation and deficit irrigation under arid and semi-arid conditions. *Sci Rep* (2026). <https://doi.org/10.1038/s41598-026-52375-0>

Gholamali Akbari, Saeed Norouzi, Iraj Alahdadi, Elias Soltani & Mohammadali Norouzian

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

Biochemical and agronomic responses of guar to nitrogen, *Bradyrhizobium* inoculation and deficit irrigation under arid and semi-arid conditions

Gholamali Akbari*¹, Saeed Norouzi*¹, Iraj Alahdadi¹, Elias Soltani¹, Mohammadali Norouzian³

1. Department of Agronomy and Plant Breeding Sciences, Aburaihan Campus, University of Tehran, Tehran, Pakdasht, Iran

2. Department of Animal and Poultry Sciences, Aburaihan Campus, University of Tehran, Tehran, Pakdasht, Iran

*Corresponding Authors: GAkbari@ut.ac.ir
Saeed_Norouzi2001@yahoo.com

Abstract

The present study aimed to investigate the effects of various levels of chemical nitrogen fertilizer and inoculation with different strains of *Bradyrhizobium japonicum* on several phenological, quantitative, qualitative, and fodder-related characteristics of guar (*Cyamopsis tetragonoloba* L.) under varying water-deficit regimes. A two-year field experiment (2021–2022) was conducted as a split-factorial arrangement based on a randomized complete block design with three replications to evaluate the interactive effects of nitrogen fertilization (0, 50, and 100 kg N ha⁻¹) and seed inoculation with strains RS-150 and RS-153 under different irrigation regimes. Deficit irrigation significantly reduced grain yield, while full irrigation × 100 kg N ha⁻¹ yielded the highest GY (3658.2 kg ha⁻¹) and severe deficit × no N application the lowest (1017.8 kg ha⁻¹). Application of 100 kg N ha⁻¹ increased GY by 30.99% compared with the control. Inoculation with RS-153 enhanced grain number per m² and 1000-grain weight by 14.41% and 14.54%, respectively, relative to non-inoculated plants. Full irrigation also increased grain protein content by up to 22.19% compared with severe deficit. Soluble sugars accumulated under water stress, reaching 490.16 µg g⁻¹ DW under deficit irrigation × 100 kg N ha⁻¹ × RS-150, indicating osmotic adjustment. Crude protein, metabolizable energy, and dry matter digestibility were further improved under full irrigation and higher nitrogen supply. Overall, integrated nitrogen management and rhizobial inoculation partially mitigated drought-induced yield reductions and improved the forage quality of guar under semi-arid conditions.

Key words: *Bradyrhizobium japonicum*, Drought stress, Enzymatic/non-enzymatic activities, Forage quality, Grain protein, Grain yield, Metabolizable energy

Abbreviations

N: Nitrogen; **BRh:** *Bradyrhizobium* inoculum; **D50F:** Days to 50% flowering; **D50P:** Days to 50% Podding; **DH:** Days to Harvesting; **CAT:** Catalase; **GN:** Grain number; **TGW:** 1000-grain weight; **GY:** Grain yield; **Chl:** Chlorophyll; **CAR:** Carotenoid; **GPr:** Grain protein; **SS:** Soluble sugars; **PC:** Proline content; **CrPr:** Crude protein; **ADF:** Acid detergent fiber; **NDF:** Neutral detergent fiber; **DMD:** Dry matter digestibility; **ME:** Metabolizable energy; **IR:** Irrigation regimes; **NA:** Nitrogen absorption; **NF:** Nitrogen fertilizer; **FW:** Fresh weight; **DW:** Dry weight; **SOD:** Superoxide dismutase; **CV:** Coefficient of Variation.

1. Introduction

Agricultural production has long served as a primary source of livelihood and a key contributor to feeding the growing global population [1,2]. However, farmland degradation remains a major constraint on crop productivity, emphasizing the importance of appropriate management practices to sustain soil fertility and yield [3,4]. Effective resource and nutrient management, particularly of nitrogen (N), one of the most limiting nutrients, plays a crucial role in determining crop growth and yield quality [5,6,7]. In Iran, where about 88% of the

land area is arid or semi-arid, soils are typically poor in organic matter and N content [8,9]. Although chemical N fertilizers can enhance plant growth by improving root development, leaf area, and nutrient uptake [10,11], their excessive use accelerates soil organic matter depletion, acidification, soil erosion, and environmental pollution [12,13]. In general, integrating organic manures with conventional fertilization improves soil carbon and nitrogen stocks while sustaining yield [14]. In this regard, legumes, as efficient atmospheric N fixers, enrich soil fertility and reduce dependence on chemical fertilizers, representing a sustainable approach to improving soil nutrient status [15,16,17,18]. Guar (*Cyamopsis tetragonoloba* L.), an annual drought- and heat-tolerant legume native to India and Pakistan, thrives in sandy and well-drained clay soils and serves diverse purposes, including fodder, food, and industrial use [19,20,21]. Its rich composition of galactomannan, phenolic, and flavonoid compounds imparts antibacterial and medicinal properties, with traditional applications for treating diabetes, asthma, anemia, and inflammation [22,23,24,25,26].

Another approach toward the sustainable intensification of agriculture to improve soil fertility and plant growth is the use of plant growth-promoting rhizobacteria (PGPRs) or bio-fertilizers [27,28]. PGPRs are eco-friendly alternatives to chemical fertilizers, and their application began about 60 years ago [29,30]. These microorganisms enhance vegetative growth, nutrient uptake, nutrient solubility, and siderophore production, as well as synthesize phytohormones, enzymes, and metabolites involved in plant growth and stress tolerance, particularly under drought conditions [30,31,32,33,34]. Well-known PGPRs include *Rhizobium*, *Bradyrhizobium*, *Bacillus*, *Azospirillum*, *Azotobacter*, *Acinetobacter*, *Pseudomonas*, *Mycorrhiza*, and *Trichoderma* [35,36,37]. Inoculation of legumes with *Rhizobium* strains has been reported to improve growth traits such as shoot biomass, plant height, branch numbers, nodules, pods, grain weight, yield, and protein content [38]. Although these bacteria generally promote plant growth and yield, their symbiotic efficiency in N fixation is influenced by environmental factors such as water availability, soil salinity, pH, nutrient deficiency, ion toxicity, and temperature extremes [39,40,41,42], which can exert either beneficial or adverse effects on plants [43]. Furthermore, bio-fertilizers can reduce the need for chemical N fertilizers, convert N compounds into amino acids, improve nutrient availability, root development, seed germination, and overall plant growth [44,45,46].

Plants are constantly exposed to various biotic and abiotic stresses under diverse environmental conditions [47,48]. Among these, water availability is a major determinant of plant growth and productivity [49]. Drought stress, particularly in arid and semi-arid regions such as Iran, is one of the most limiting factors for plant performance and yield [50,51,52]. Developing efficient irrigation systems remains a major challenge, as water requirements vary with growth stage, plant sensitivity, climatic conditions, and irrigation intervals [53,54,55]. Water stress restricts evapotranspiration via stomatal closure, limits carbon assimilation, biomass accumulation, photosynthetic activities, and ultimately reduces grain yield [56,57,58,59]. It also promotes excessive formation of reactive oxygen species (ROS), including hydrogen peroxide and superoxide radicals, leading to oxidative damage, reduced photosynthetic and water-use efficiency, lipid peroxidation, and cell death [60,61,62]. To counter these effects, plants enhance their defense mechanisms through osmolyte accumulation (e.g., proline and soluble sugars), antioxidant enzyme activation, and synthesis of secondary metabolites (e.g., flavonoids and phenolics), increased abscisic acid, and maintenance of leaf water status [63]. In guar, supplemental irrigation improves plant height, pod number, biomass, and grain yield, while full irrigation enhances protein and galactomannan content [64]. Conversely, drought cycles markedly reduce yield components and water use efficiency [65,66,67]. Despite these insights, limited information exists on how integrated management of chemical and biological nitrogen sources under deficit irrigation can improve guar performance. Furthermore, water-deficit stress, besides its negative impact on crop yield, induces significant metabolic changes and accumulation of secondary metabolites in plants, while nitrogen application can enhance drought tolerance and water

use efficiency [59,68], partially mitigating these stress-induced metabolic effects. Researchers also indicated that combined application of high nitrogen levels and *Bradyrhizobium* further strengthened photosynthetic efficiency and drought resilience, highlighting the synergy between nutrient management and microbial inoculation in maintaining biochemical stability under water deficit [69,70]. Therefore, this study investigated the effects of nitrogen fertilizer levels and *Bradyrhizobium* inoculation on some phenological, biochemical, qualitative, and quantitative traits of guar (*Cyamopsis tetragonoloba* L.) under varying irrigation regimes in arid and semi-arid conditions.

2. Materials and methods

2.1. Plant materials

In this study, the effects of three levels of chemical nitrogen fertilizer and symbiotic bacteria on several quantitative and qualitative traits of guar (*Cyamopsis tetragonoloba* L.) under different water-deficit regimes were investigated using a split-factorial arrangement in a randomized complete block design (RCBD) with three replications. Guar seeds (RGC-1031 from India) used in this study were obtained from the University of Birjand, Iran, as institutional research material. No specific collection permit or license was required because the plant material was sourced from institutional resources rather than wild populations. Formal taxonomic identification was confirmed by Dr. Hamidreza Ramazani, faculty member at the University of Birjand. At the time of the study, a voucher specimen had not been deposited in a public herbarium and no deposition number was available.

The experiment was conducted at the research farms of the Aburaihan Campus, University of Tehran, Iran (with GPS coordinates of 35°28'29" N and 51°36'35" E) in early June 2021 and early June 2022. Experimental treatments included irrigation regimes as main plots (full irrigation, deficit irrigation up to 50% of the flowering stage, and deficit irrigation up to 50% of the podding stage), and nitrogen fertilizer levels (control, 50, and 100 kg N ha⁻¹) combined with seed inoculation using *Bradyrhizobium japonicum* strains (non-inoculation, RS-150, and RS-153) as subplots. The bacterial inoculants were liquid formulations supplied by Green Biotech Company (Qom Province, Iran) and guar seeds were soaked in the inoculant solutions for 12-24 hours before sowing, following the manufacturer's instructions. *Bradyrhizobium japonicum*, previously reported as effective symbionts for guar, were used to evaluate their potential effects on plant growth, yield, and forage quality.

Before sowing, a composite soil sample was collected from the experimental field to determine soil physicochemical properties and fertilizer requirements (Table 1). Based on soil test results, phosphorus (triple superphosphate at 50 kg ha⁻¹), potassium (potassium sulfate at 170 kg ha⁻¹), nitrogen (75 kg ha⁻¹ urea 46%), and sulfur (220 kg ha⁻¹) were uniformly applied to all plots following primary tillage. One-third of the nitrogen fertilizer was applied at planting, with the remainder top-dressed at the beginning of rapid growth and early podding stages. The pedigree, origin, and key agronomic characteristics of the guar cultivar used are presented in Table 2.

Table 1. Physico-chemical properties of the studied soil

Soil texture	pH	Ec (dS.m ⁻¹)	Organic Carbon (%)	Total Nitrogen (%)	P (mg.kg ⁻¹)	K (mg.kg ⁻¹)	Cu (ppm)	Fe (ppm)	Zn (ppm)	Mn (ppm)
Loam-Clay	8.19	7.61	1.45	0.14	79	732	0.87	4.32	2.32	7.88
Fertilizer recommendations based on soil test										
		N (Kg ha ⁻¹)		P (Kg ha ⁻¹)		K (Kg ha ⁻¹)	S (Kg ha ⁻¹)			
		75		50		170	220			

Table 2. Pedigree and key characteristics of guar (*Cyamopsis tetragonoloba*) cultivar RGC-1031

Genotype / Cultivar	Type	Origin	Pedigree	Key characteristics / Adaptation
RGC-1031 (Guar Kranti)	Cluster bean (guar)	ARS Durgapura, Rajasthan, India	RGC-936 × (RGC-986 / P-10)	Dual-purpose (grain + fodder) cultivar; moderately tall; 90-100 days to maturity; tolerant to drought and high temperature; well adapted to semi-arid, rain-fed regions

Planting rows were spaced 50 cm apart with 5 cm between plants within rows, in plots measuring 3 × 2 m, corresponding to approximately 400,000 plants ha⁻¹. Field irrigation was applied via a drip system, with full irrigation every four days and deficit irrigation every eight days, corresponding to 50% water reduction at flowering and podding stages. Weeding was also carried out manually three times during the rapid growth, flowering, and podding stages. At the end of the experiment (in both years), various phenological, qualitative, and quantitative traits were measured. Biochemical analyses were performed at specific growth stages as indicated in Table 3 (BBCH scale) to capture peak metabolic activity and optimal forage quality. Photosynthetic pigments (Chl a, Chl b, Chl T, and CAR), leaf proline content, soluble sugars, antioxidant enzyme activities (CAT and SOD), grain protein, and forage quality parameters (crude protein, ADF, NDF, dry matter digestibility, and metabolizable energy) were quantified according to the BBCH stages listed in Table 3 [71].

Table 3. Growth stages of guar (BBCH scale) for biochemical analyses

Parameter / Trait	Growth Stage (Phenophase)	BBCH Code	Description
Photosynthetic pigments (Chl a, Chl b, T Chl, and CARs)	60-69	Flowering	Full flower opening; leaves fully expanded; photosynthetic and metabolic activity at maximum. Carotenoids act as accessory pigments and protect against oxidative stress.
Non-enzymatic antioxidants (Leaf proline content and soluble sugars)	60-69	Flowering	Leaves were fully developed; active osmotic adjustment and antioxidative response under stress conditions.
Enzymatic activities (CAT and SOD)	60-69	Flowering	Peak antioxidant enzyme activity corresponds to high metabolic and oxidative demand during flowering.
Grain protein	90-99	Physiological maturity	Fully mature grains with maximum accumulation of storage compounds (proteins).
Crude protein (Cr Pr)	71-75	Early to mid-milk stage (beginning of grain filling; milky endosperm)	Protein content begins to decline rapidly after BBCH 75 due to dilution by structural carbohydrates.
Acid detergent fiber (ADF)	75-83	Late milk to early dough stage	ADF starts to rise sharply after BBCH 83 as lignin deposition increases.
Neutral detergent fiber (NDF)	75-83	Late milk to early dough stage	NDF increases gradually, reaching moderate levels before senescence.
Dry matter digestibility (DMD)	75-83	Late milk to early dough stage	DMD is highest during early grain filling and starts declining as structural tissues mature.
Metabolizable energy (ME)	75-83	Late milk to early dough stage	ME peaks when digestibility is high and lignification is still moderate.

2.2. Phenological traits and yield-related traits

Phenological traits included days to 50% flowering (D50F), days to 50% podding (D50P), and days to physiological maturity (DH). These traits were recorded in the field for each plot, based on the number of days from sowing to the corresponding stage: flowers observed in at least 50% of plants, pods observed in at least 50% of plants, and 95% of pods fully matured, respectively. Also, Grain yield and yield components (number of grains per pod, 1000-grain weight, and grain yield) were determined by harvesting all plants within a 1-m² area from the central part of each plot to avoid edge effects. Measurements were conducted at physiological maturity, and all values were expressed per hectare.

2.3. Qualitative and fodder-related attributes

In addition, photosynthetic pigments (chlorophyll a, chlorophyll b, and total chlorophyll), carotenoids (CAR), grain protein (GPr), soluble sugars (SS), and proline content (PC) were determined. Furthermore, to evaluate the effects of treatments on forage quality and to compare among treatments, forage samples were collected at the 50% podding stage from the full irrigation and deficit irrigation (up to 50% podding) treatments, and forage quality parameters, including crude protein (CPro), acid detergent fiber (ADF), neutral detergent fiber (NDF), ash percentage, dry matter digestibility (DMD), and metabolizable energy (ME), were analyzed. Forage quality analysis was restricted to the full irrigation and deficit irrigation (up to 50% podding) treatments, representing the best and worst water regimes. This selection was made to illustrate the most contrasting responses while keeping the experiment manageable in terms of laboratory analysis.

2.3.1. Photosynthetic pigments (Chl a, Chl b, Chl T, and CAR)

Chl a, Chl b, Chl T, and CAR traits were quantified following the procedures described by Arnon [72] and Lichtenthaler [73]. For each sample, 0.5 g of fresh leaf blade tissue was weighed and homogenized in a mortar using 80% aqueous acetone. The homogenate was filtered through filter paper into a volumetric flask, and the residue was re-homogenized and filtered again to ensure complete pigment extraction. The combined filtrate was then adjusted to a final volume of 10 ml with 80% acetone. Pigment concentrations were determined spectrophotometrically using a Jenway 6300 spectrophotometer at 645, 663, and 470 nm. 80% aqueous acetone served as the blank. Chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents were subsequently calculated using the standard equations provided by the cited references.

$$\text{Equation (1) Chlorophyll a (mg.g}^{-1}\text{ FW)} = \frac{[(12.7 \times D_{663}) - (2.69 \times D_{645})] \times V}{1000 \times W}$$

$$\text{Equation (2) Chlorophyll b (mg.g}^{-1}\text{ FW)} = \frac{[(22.9 \times D_{645}) - (4.93 \times D_{663})] \times V}{1000 \times W}$$

$$\text{Equation (3) Total Chlorophyll (mg.g}^{-1}\text{ FW)} = \frac{[(20.2 \times D_{645}) - (8.02 \times D_{663})] \times V}{1000 \times W}$$

$$\text{Equation (4) Carotenoids (mg.g}^{-1}\text{ FW)} = \frac{[(1000 \times D_{470}) - (1.82 \times \text{Chl.a}) - (85.02 \times \text{Chl.b})]}{198}$$

Where V represents the final volume of the 80% acetone extract (mL), W is the fresh leaf tissue weight (g), and D is the absorbance measured at the specified wavelengths.

2.3.2. Proline content (PC)

Proline content was quantified following the method described by Bates et al. [74]. For each sample, 0.5 g of fresh leaf blade tissue was weighed and homogenized in 10 ml of 3% aqueous sulfosalicylic acid. The resulting homogenate was filtered through Whatman filter paper to obtain a clear extract. Subsequently, 2 ml of acid-ninhydrin solution and 2 ml of glacial acetic acid were added to 2 ml of the filtrate, and the mixture was heated at 100°C for 60 minutes. After incubation, the tubes were immediately cooled in an ice bath for 30 minutes. Thereafter, 4 ml of toluene was added, and the solution was thoroughly mixed for 15–20 seconds. Two distinct phases were formed, and the upper toluene layer was carefully collected. Absorbance was measured at 520 nm using a spectrophotometer, and proline concentration was calculated from a standard curve prepared with 0, 50, 100, 200, and 250 µM proline. Final values were expressed as µM per gram of fresh leaf tissue.

$$\text{Equation (5) Proline content (}\mu\text{M.g}^{-1}\text{FW)} = \frac{\left[\frac{(\mu\text{g.ml}^{-1} \text{ proline} \times \text{ml toluene})}{115.5 \mu\text{g.}\mu\text{mole}^{-1}} \right]}{\left[\frac{(\text{g sample})}{5} \right]}$$

2.3.3. Soluble sugars (SS)

Soluble sugar (SS) content was determined following the method of Yemm and Willis (75) with minor modifications. For each sample, 0.1 g of dried leaf tissue was weighed and hydrolyzed in 5 ml of 2.5 N HCl by boiling for three hours. After cooling to room temperature, the hydrolysate was neutralized with solid sodium carbonate until effervescence ceased. The resulting solution was then diluted to 100 ml with distilled water and centrifuged at 3500 rpm for 10 minutes. Subsequently, 0.5 ml of the supernatant was collected and further diluted to 1 ml with distilled water. Next, 4 ml of anthrone reagent was added to each tube, and the mixture was heated in a boiling water bath for eight minutes. Absorbance was measured at 630 nm using a spectrophotometer, with a glucose-free solution serving as the blank. Soluble sugar concentrations were calculated from a standard curve prepared with known concentrations of glucose (0, 50, 100, 150, 200, and 250 $\mu\text{g ml}^{-1}$), and final values were expressed as μg per gram of dry weight ($\mu\text{g g}^{-1}$ DW).

2.3.4. Enzymatic activities of catalase (CAT) and superoxide dismutase (SOD)

2.3.4.1. Catalase (CAT) Activity

Catalase activity was measured following the method of Bailly et al. [76] with minor modifications. Leaf tissues (1 g) were homogenized in 3 mL of 50 mM potassium phosphate buffer (pH 7.2) containing 1 mM EDTA, 1 mM PMSF, and 1% PVP. The homogenate was centrifuged at $14,000 \times g$ for 15 minutes at 4°C , and the supernatant was used for the assay. The reaction mixture contained 25 mM potassium phosphate buffer (pH 7.0), 10 mM H_2O_2 , and enzyme extract. Catalase activity was determined based on the rate of H_2O_2 decomposition at 240 nm using an extinction coefficient of $\epsilon = 0.036 \text{ Mm}^{-1} \text{ cm}^{-1}$. One unit of CAT activity was defined as the amount of enzyme that decomposes 1 μmol of H_2O_2 per minute per mg of protein.

2.3.4.2. Superoxide Dismutase (SOD) Activity

SOD activity was assayed according to Giannopolitis and Ries [77] with minor modifications. At first, fresh leaf tissue (1 g) was homogenized in 50 mM phosphate buffer containing 0.013 mM methionine, 0.1 μM EDTA, and 2 μM riboflavin under complete darkness. After adding riboflavin, 3 mL of the reaction mixture was transferred to a test tube, and 100 μL of the enzyme extract was added. Samples were irradiated under a light source at 30 cm for 16 minutes, and absorbance was read at 560 nm against a blank. SOD activity was expressed as units per mg protein, with one unit defined as the amount of enzyme required to inhibit 50% of superoxide-mediated reactions.

2.3.5. Grain protein (GPr) percentage

To measure grain protein and crude protein, the amount of N was first determined in the studied samples using the Kjeldahl method and calculated using the following equation [78,79].

$$\text{Equation (6) Protein percentage (\%)} = \% \text{ Kjeldahl N} \times F$$

Where N represents the nitrogen content of the grain or the whole plant, as determined by the Kjeldahl method, and F is the conversion factor used to calculate protein, set at 6.25 for legumes.

2.3.6. Ash percentage

The percentage of ash was measured based on the method proposed by Thiex et al. [79] with slight modifications. Therefore, 2 g of each sample was first weighed in dried glass crucibles and burned to ash in a furnace at 550°C for 3 hours. Then, the crucibles were quickly cooled in a desiccator, and the ash percentage for each treatment was calculated as follows.

$$\text{Equation (7) Ash (\%)} = \frac{Z-X}{Y-X} \times 100$$

In which X=weight of empty crucible; Y=weight of crucible + sample; Z=weight of crucible + ash

2.3.7. Neutral detergent fiber (NDF), Acid detergent fiber (ADF), and Dry matter digestibility (DMD)

Neutral detergent fiber (NDF) and acid detergent fiber (ADF) were determined to assess forage quality following the procedures described by Van Soest [80] with minor modifications. For NDF, samples were treated with a neutral detergent solution to remove soluble cell contents, leaving the cell wall components (cellulose, hemicellulose, and lignin) as residue, which was then weighed and expressed as a percentage of dry matter. ADF was measured by treating samples with an acid detergent solution, which removes hemicellulose and soluble components, leaving cellulose and lignin as residue; the remaining material was then weighed and expressed as a percentage of dry matter. Dry matter digestibility (DMD) was subsequently estimated using the relationship between ADF and N% according to Oddy et al. [81].

$$\text{Equation (8) DMD (\%)} = 83.58 - 0.824 \text{ ADF\%} + 2.262 \text{ N\%}$$

2.3.8. Metabolizable energy (ME)

Metabolizable energy (ME) was calculated using the equations provided by the Standing Committee on Agriculture [82] and Williams et al. [83] as follows (Equation 8):

$$\text{Equation (9) ME (MJ.kg}^{-1}\text{)} = 0.17\text{DMD\%} - 2$$

2.4. Quantitative traits (Yield and yield components)

In addition, grain numbers (GN), 1000-grain weight (TGW), and grain yield (GY) were calculated based on crop yields per m² at the physiological maturity stage. Yield and its components were evaluated based on the plants of each plot (according to the harvest area) after determining edge effects.

3. Statistical analysis

The data were analyzed using SAS software (version 9.4), and treatment means were compared using the LSD test at $p \leq 0.05$. Homogeneity of variances between both years was assessed using Bartlett's test prior to analysis. Since significant heterogeneity of variances was detected, the assumption of homogeneity required for combined analysis was not satisfied; therefore, the data were analyzed separately for each experimental year. Graphs were also prepared using Microsoft Excel (2013).

4. Results

4.1. Phenological traits (Days to 50% flowering, days to 50% podding, and days to harvesting)

The ANOVA results for phenological traits (D50F, D50P, and DH) across the two experimental years are presented in Table 4. Irrigation regime (IR) significantly affected all traits in both years ($p < 0.01$). In addition, nitrogen fertilizer (NF) significantly influenced D50F in both years ($p < 0.05$), whereas its effects on D50P and DH were significant only in the first year. In contrast, *Bradyrhizobium* inoculation and all interaction effects were not significant for any of the studied traits.


Table 4. Effects of IR, NF levels, and BRh inoculation conditions on days to 50% flowering, days to 50% podding, and days to harvesting of guar

Source of variation	df	D50F		D50P		DH	
		First year	Second year	First year	Second year	First year	Second year
Block	2	29.05 ns	4.61 ns	20.04 ns	3.79 ns	7.7 ns	2.04 ns
Irrigation regimes (IR)	2	109.12*	487.46**	147.82**	759.46**	551.44**	988.59**
Ea	4	17.29	14.14	30.41	5.86	13.59	12.07
Nitrogen fertilizer (NF) levels	2	112.75*	110.42*	122.11*	53.09 ns	180.26*	203.26 ns

<i>Bradyrhizobium</i> Inoculation (BRh) conditions	2	7.46 ns	7.75 ns	11.15 ns	2.68 ns	40.11 ns	30.26 ns
Irrigation × NF levels (IR × NF)	4	3.94 ns	4.24 ns	10.93 ns	7.94 ns	6.2 ns	32.74 ns
IR × BRh	4	0.20 ns	0.22 ns	2.35 ns	3.59 ns	1.17 ns	3.24 ns
NF × BRh	4	0.61 ns	0.51 ns	2.37 ns	2.61 ns	8.15 ns	10.19 ns
IR × NF × BRh	8	0.82 ns	0.88 ns	2.02 ns	1.79 ns	2.4 ns	7.92 ns
<i>Eb</i>	48	23.32	27.92	24.26	30.7	55.88	78.12
Coefficient of Variation (CV %)		9.43	9.79	7.94	8.15	7.19	8.25

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

Mean comparisons (Fig. 1a) showed that full irrigation regime consistently resulted in the longest D50F in both years, with 53.52 days in the first year and 58.82 days in the second year, which were significantly longer than the corresponding deficit irrigation treatments of 50.07 and 50 days in the first year and 52.07 and 50.96 days in the second year of the experiment, respectively. In addition, increasing nitrogen levels significantly reduced D50F (Fig. 1b). Results indicated that the application of 100 kg N ha⁻¹ produced the shortest flowering period with 48.85 days in the first year of the experiment and 51.63 days in the second year compared with 52.59 and 55.33 days for the control and 52.15 and 54.89 days for the 50 kg N ha⁻¹ treatment, respectively.







Fig. 1. Effects of (a) irrigation regimes and (b) nitrogen fertilizer levels on days to 50% flowering (D50F) in the guar plant
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Considering the significant effects of IR on D50P (Table 4), deficit irrigation accelerated podding compared to full IR. The shortest D50P was observed under deficit IR up to 50% podding (60.19 and 63.11 days according to the first and second-year data, respectively), whereas full IR resulted in the longest D50P (64.67 and 73.63 days; Fig. 2a). In addition, the application of NF significantly affected D50P only in the first year, where 100 kg N ha⁻¹ shortened the podding period to 59.78 days compared to 64 days in the control and 61.43 days at 50 kg N ha⁻¹ (reductions of 7.06% and 2.68%), indicating that NF accelerated podding onset under these conditions. No significant NF effects were observed in the second year (Fig. 2b).







Fig. 2. Effects of (a) irrigation regimes and (b) nitrogen fertilizer levels on days to 50% podding (D50P) in the guar plant
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

ANOVA results (Table 4) indicated that IR significantly affected DH in both years, while NF influenced DH only in the first year. DH was consistently longer under full IR (109 days in the first year and 113.67 days in the second year) than under deficit IR, which shortened the growth period (101.33 days for deficit IR up to 50% flowering and 100.67 days for deficit IR up to 50% podding in the first year; 105.33 and 102.33 days, respectively, in the second year; Fig. 3a), reflecting accelerated development under water deficit as a drought-escape strategy. In the first year, NF application slightly prolonged DH, with 100 kg N ha⁻¹ resulting in 106.67 days, compared to 101.33 days in the control and 104.33 days at 50 kg N ha⁻¹ (Fig. 3b), indicating that adequate nitrogen extended vegetative and reproductive growth. This effect likely reflects the indeterminate growth habit of guar and the role of nitrogen in delaying leaf senescence and maintaining photosynthetic activity, allowing plants to sustain growth before reaching physiological maturity.






Fig. 3. Effects of (a) irrigation regimes and (b) nitrogen fertilizer levels on days to harvesting (physiological maturity; DH) in the guar plant
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

4.2. Biochemical indices

4.2.1. Photosynthetic pigments (Chl a, Chl b, Chl T, and CAR)

Analysis of variance of the treatments on biochemical indices (Table 5) showed that Chl a content was significantly affected by IR, NF, and BRh inoculation in both years ($p < 0.01$), with a significant NF $\times$ BRh interaction in the first year ($p < 0.05$). In addition, Chl b was influenced by IR, NF, and IR $\times$ NF, IR $\times$ BRh interactions ($p < 0.01$), as well as NF $\times$ BRh in the first year of the experiment ($p < 0.05$). According to the second-year data, IR, NF, and the interactions IR $\times$ BRh and NF $\times$ BRh were significant ($p < 0.01$) on the Chl b content. Also, Chl T was strongly affected by IR, NF, BRh inoculation, and the IR $\times$ BRh interaction in both years, with NF $\times$ BRh significant in the first year ($p < 0.01$). Furthermore, the CAR content was significantly influenced by IR and NF in both years ($p < 0.01$) and by BRh inoculation ($p < 0.05$), whereas all investigated interactions had no significant effects on CAR levels in either year (Table 5).

Table 5. Effects of irrigation regimes, NF levels, and inoculation/non-inoculation conditions on photosynthetic pigments

Source of variation	df	Chl a		Chl b		Chl T		CAR	
		First year	Second year	First year	Second year	First year	Second year	First year	Second year
Block	2	0.82ns	1.61ns	0.16*	0.24*	0.26ns	0.96ns	1.64ns	0.16ns
Irrigation regimes (IR)	2	9.4**	114.89**	0.62**	0.67**	5.82**	98.45**	17.5**	16.51**
Ea	4	2.16	2.26	0.04	0.06	1.76	1.72	0.88	0.56
NF levels	2	36.98**	84.47**	3.36**	4.8**	61.09**	128.4**	15.3**	16.08**
Inoculation conditions	2	21.87**	31.93**	0.06ns	0.07ns	21.05**	31.96**	2.72*	3.12*
IR $\times$ NF	4	0.11ns	0.66ns	0.33**	0.1ns	0.33ns	0.32ns	0.1ns	0.81ns
IR $\times$ BRh	4	1.05ns	3.55*	0.74**	0.64**	2.29**	3.58**	0.13ns	0.25ns
NF $\times$ BRh	4	1.88*	1.74ns	0.14*	0.22**	2.46**	2.19ns	0.25ns	0.31ns
IR $\times$ NF $\times$ BRh	8	0.08ns	0.91ns	0.05ns	0.09ns	0.17ns	1.46ns	0.39ns	0.17ns
Eb	48	0.72	1.02	0.03	0.05	0.61	0.9	0.65	0.61
CV (%)		12.71	16.37	9.54	13.82	9.16	12.06	17.3	20.28

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

The first-year mean comparisons for Chl a content under different irrigation regimes (Fig. 4a) indicated that drought stress significantly reduced Chl a. The highest and lowest values (7.24 and 6.06 mg g⁻¹ FW) were observed under full irrigation and deficit irrigation up to 50% podding, respectively. No significant difference was found between the two deficit irrigation treatments (6.69 mg g⁻¹ FW for deficit irrigation up to 50% flowering and 6.06 mg g⁻¹ FW for up to 50% podding). Overall, deficit irrigation up to 50% podding reduced Chl a content by 16.30% relative to full irrigation and by 9.42% relative to deficit irrigation up to 50% flowering. Regarding the NF $\times$ BRh interaction (first-year data; Fig. 4b), the highest

Chl a values were recorded for 100 kg N N ha⁻¹ × RS-153 (8.50 mg g⁻¹ FW) and 50 kg N N ha⁻¹ × RS-153 (8.25 mg g⁻¹ FW) interactions. In contrast, the lowest value (4.89 mg g⁻¹ FW) was observed under the treatment without N fertilizer × no seed inoculation.




Fig. 4. Effects of (a) IR and (b) NF × BRh interaction on the content of Chl a (adapted from first-year data)
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Given the significance effects of IR × BRh interaction on the Chl a content in the second year of the research, compare means (Fig. 5) indicated that the highest values for Chl a (9.63 and 9.15 mg g⁻¹ FW), which had significant increases compared to the other interactions, were obtained for the interactions of full irrigation × seed inoculation with the RS-150 strain and full irrigation × seed inoculation with the RS-153 strain, respectively. In contrast, the significant lowest Chl a value, Chl a, which was significantly reduced compared to the others, was recorded for the interaction of deficit irrigation up to 50% podding × non-inoculation conditions with the equivalent of 3.56 mg g⁻¹ FW (Fig. 5).




Fig. 5. Effects of the IR $\times$ NF interaction on Chl a (adapted from second-year data)
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Considering the significant effects of IR $\times$ NF, IR $\times$ BRh, and NF $\times$ BRh interactions on the content of Chl b (adopted from the first-year data; Table 5), mean squares of the IR $\times$ NF interaction (Fig. 5) revealed that the highest Chl b content (2.52 mg g⁻¹ FW) was recorded under deficit irrigation up to 50% flowering combined with 100 kg N ha⁻¹, showing a significant increase compared with other treatments. In contrast, the lowest Chl b content (1.38 mg g⁻¹ FW) was observed under full irrigation $\times$ no application of chemical N fertilizer (Fig. 6). This pattern suggests that moderate water deficit combined with adequate nitrogen supply might have stimulated Chl biosynthesis and improved N assimilation efficiency, possibly due to enhanced photosynthetic adaptation and delayed leaf senescence in guar.




Fig. 6. Effects of the interaction of IR $\times$ NF on the content of Chl b
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

At same time, mean comparisons related to the IR $\times$ BRh interaction obtained from the first year of the experiment (Table 6) showed that the highest Chl b content (2.19 mg g⁻¹ FW) was recorded for the deficit irrigation up to 50% podding $\times$ RS-150 interaction, with no

significant difference from deficit irrigation up to 50% podding $\times$ RS-153 (2.09 mg g⁻¹ FW). In contrast, the lowest Chl b value (1.42 mg g⁻¹ FW) was observed under deficit irrigation up to 50% flowering $\times$ RS-153.

Table 6. Effects of IR $\times$ BRh interaction on the contents of Chl a, Chl b, and Chl T

Source of variation		Chl b (mg g ⁻¹ FW)		Chl T (mg g ⁻¹ FW)	
		First year	Second year	First year	Second year
Irrigation regimes (IR)	Inoculation conditions				
Full irrigation	Non-Inoculation (Control)	1.66 cd	1.64 bcd	7.51 ef	8.21 b
	RS-150	1.72 ef	1.46 cde	9.47 ab	11.09 a
	RS-153	2.02 bcd	1.67 bc	10.13 a	10.81 a
Deficit irrigation up to 50% flowering	Non-Inoculation (Control)	2.06 abc	1.42 de	8.01 de	6.09 d
	RS-150	1.88 de	1.71 b	8.67cd	7.39 bc
	RS-153	1.42 g	1.98 a	8.74 bcd	8.67 b
Deficit irrigation up to 50% podding	Non-Inoculation (Control)	1.88 f	1.35 e	7.14 f	4.91 e
	RS-150	2.19 a	2 a	8.18 de	6.95 cd
	RS-153	2.09 ab	1.85 ab	9.03 bc	6.56 cd
LSD		0.17	0.23	0.79	0.93

Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Regarding the NF $\times$ BRh interaction effects (Table 7; first-year data), the highest Chl b value (2.32 mg g⁻¹ FW) was observed under 100 kg N ha⁻¹ $\times$ RS-150. The lowest Chl b content (1.39 mg g⁻¹ FW) occurred under control (no N fertilizer) $\times$ RS-153, which was not significantly different from control $\times$ RS-150 (1.55 mg g⁻¹ FW). Moreover, the second-year data revealed a similar pattern, where the highest and lowest Chl b values were obtained under deficit irrigation up to 50% podding $\times$ RS-150 (2.00 mg g⁻¹ FW) and deficit irrigation up to 50% podding $\times$ non-inoculated treatment (1.35 mg g⁻¹ FW), respectively. Similar trends were also observed for Chl T (Table 7), as discussed in the following section.

Table 7. Effects of NF $\times$ BRh interaction on the contents of Chl a, Chl b, and Chl T (The first year of data)

Source of variation		Chl b (mg g ⁻¹ FW)		Chl T (mg g ⁻¹ FW)	SS (μ g g ⁻¹ DW)
		First year	Second year	First year	First year
NF levels (Kg ha ⁻¹)	Inoculation conditions				
Control	Non-Inoculation (Control)	1.55 ef	1.33 f	6.44 e	163.65 e
	RS-150	1.63 e	1.36 f	7.09 de	160.59 e
	RS-153	1.39 f	1.1 g	7.02 de	164.75 e
50	Non-Inoculation (Control)	1.81 d	1.55 ef	7.62 d	179.23 d
	RS-150	1.83 d	1.61 de	9.29 bc	200.38 c
	RS-153	2.01 c	1.81 cd	10.25 a	210.84 ab
100	Non-Inoculation (Control)	2.24 ab	1.96 bc	8.61 c	185.29 d
	RS-150	2.32 a	2.21 a	9.94 ab	207.67 bc
	RS-153	2.12 bc	2.15 ab	10.62 a	219.07 a
LSD		0.17	0.23	0.79	10.26

Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Mean comparisons related to the IR $\times$ BRh interaction on Chl T showed that the highest and lowest values (10.13 and 7.14 mg g⁻¹ FW) were recorded for the interaction of full irrigation $\times$ RS-153 and deficit irrigation up to 50% podding $\times$ non-inoculation, respectively (Table 6). Similarly, according to the NF $\times$ BRh interaction (Table 7), the maximum and minimum Chl T (10.62 and 6.44 mg g⁻¹ FW) were obtained under 100 kg N ha⁻¹ $\times$ RS-153

and no application of NF (control) $\times$ non-inoculation conditions (control), respectively. According to the second-year data, the interaction of IR $\times$ BRh also showed a significant effect on Chl T. The highest Chl T value (11.09 mg g⁻¹ FW) was observed under full irrigation $\times$ seed inoculation with RS-150, which was not significantly different from full irrigation $\times$ seed inoculation with RS-153. Conversely, the lowest Chl T content (4.91 mg g⁻¹ FW) was recorded under deficit irrigation up to 50% podding $\times$ non-inoculation (Table 6).

For CAR contents, the first-year data (Table 8) indicated that full irrigation resulted in the highest carotenoid concentration (5.37 mg g⁻¹ FW), which was not significantly different from deficit irrigation up to 50% flowering (4.78 mg g⁻¹ FW), while the lowest CAR content (3.77 mg g⁻¹ FW) occurred under deficit irrigation up to 50% podding. Nitrogen fertilization increased CAR, with 100 kg N ha⁻¹ producing 4.98 mg g⁻¹ FW compared to 50 kg N ha⁻¹ (4.28 mg g⁻¹ FW) and the control (3.78 mg g⁻¹ FW). Seed inoculation with RS-153 and RS-150 also significantly enhanced CAR content (4.85 and 4.80 mg g⁻¹ FW, respectively) relative to non-inoculated plants. Although CAR content decreased slightly in the second year, the trend among treatments remained consistent: full irrigation (4.66 mg g⁻¹ FW) increased CAR content by 24.6% and 50.3% relative to deficit irrigation up to 50% flowering and deficit irrigation up to 50% podding, respectively, while 100 kg N ha⁻¹ enhanced CAR by 50.5% and 11.9% compared with the control and 50 kg N ha⁻¹ treatments, respectively. Seed inoculation with RS-150 also resulted in significant and non-significant increases of 19.5% and 7.5% compared with non-inoculated plants and those inoculated with RS-153, respectively (Table 8; second-year data).

Table 8. Effects of IR, NF levels, and BRh inoculation conditions on CAR, PC, and GPr traits

Source of variation	CAR (mg g ⁻¹ FW)		PC (μ g g ⁻¹ FW)	GPr (%)		CAT (μ mol H ₂ O ₂ mg ⁻¹ protein min ⁻¹)		SOD (mg protein min ⁻¹)	
	First year	Second year	First year	First year	Second year	First year	Second year	First year	Second year
Irrigation Regimes (IR)									
Full irrigation	5.37 _a	4.66 _a	4.4 _c	29.27 _a	26.92 _a	23.98 _c	31.52 _c	45.27 _c	56.17 _c
Deficit irrigation up to 50% flowering	4.78 _a	3.74 _b	4.97 _b	26.16 _b	26.60 _a	31.33 _b	43.82 _b	54.23 _b	73.98 _b
Deficit irrigation up to 50% podding	3.77 _b	3.10 _c	5.66 _a	23.96 _c	24.27 _b	35.70 _a	52.20 _a	61.11 _a	90.22 _a
LSD	0.71	0.57	0.44	0.59	1.27	1.66	3.14	4.07	4.59
NF levels (Kg ha ⁻¹)									
Control	3.78 _b	2.99 _c	3.49 _b	24.19 _c	24.29 _b	34.81 _a	50.90 _a	56.37 _a	89.27 _a
50	4.97 _a	4.02 _b	5.77 _a	25.87 _b	26.61 _b	28.49 _b	41.87 _b	52.78 _b	69.33 _b
100	5.18 _a	4.50 _a	5.77 _a	29.34 _a	29.89 _a	27.72 _b	34.76 _c	51.45 _b	61.75 _c
Inoculation conditions									
Non-Inoculation (Control)	4.28 _b	3.48 _b	5.31 _a	24.01 _b	25.21 _b	32.14 _a	48.52 _a	57.90 _a	79.09 _a
RS-150	4.80 _a	4.16 _a	4.94 _b	27.24 _a	27.52 _a	30.69 _a	42.20 _b	52.70 _b	69.14 _b
RS-153	4.85 _a	3.87 _{ab}	4.77 _b	28.15 _a	28.06 _a	28.18 _b	36.81 _c	50.01 _c	72.12 _b
LSD	0.43	0.42	0.27	1.58	1.14	1.83	3.53	2.49	4.21

Means followed by the same letter within each column do not differ significantly at $p < 0.05$
 CAR: Carotenoids; PC: Proline content; GPr: Grain protein; CAT: Catalase; SOD: Superoxide dismutase

4.2.2. Soluble sugars (SS), proline content (PC), and grain protein (GPr)

In addition to the previous findings, the ANOVA results (Table 9) indicated that SS was significantly affected by IR, NF, BRh inoculation, and interaction between NF and BRh (NF

× BRh) during the first year of the experiment ($p < 0.01$). In the second year, the SS content was also significantly influenced by IR, NF, both interaction of IR × NF and IR × BRh ($p < 0.01$), as well as by BRh inoculation and the three-way interaction of IR × NF × BRh ($p < 0.05$). Moreover, IR, NF, and BRh inoculation had significant effects on PC and GPr percentage in both years ($p < 0.01$). In addition, significant differences in GPr percentage were observed among IR, NF, and BRh inoculation treatments in both experimental years ($p < 0.01$).

Table 9. Effects of irrigation regimes, NF levels, and inoculation/non-inoculation conditions on attributes of SS, PC, GPr, and enzymatic activities of CAT and SOD

Source of variation	df	SS		PC		GPr		CAT		SOD	
		First year	Second year	First year	Second year	First year	Second year	First year	Second year	First year	Second year
Block	2	67.22ns	85.4ns	0.5ns	0.46ns	35.19*	26.9**	5.58ns	37.89ns	14.46ns	96.59ns
Irrigation regimes (IR)	2	21055.2**	541258**	10.74**	171.72**	192.22**	192.16**	947.35**	2920.32**	1704.44**	7823.71**
Ea	4	25.37	24.77	0.34	0.69	0.61	2.82	4.83	17.22	29.05	36.93
Nitrogen Fertilizer (NF) levels	2	12949.14**	2379.08**	46.83**	83.52**	186.12**	214.2**	408.03**	1766.96**	175.09**	5455.04**
<i>Bradyrhizobium</i> Inoculation (BRh) conditions	2	3367.97**	1077.22*	2.02**	1.23ns	128.2**	61.57**	108.24**	928.22**	433.93**	703.9**
IR × NF	4	76.31ns	15759.76**	0.28ns	13.36**	6.1ns	1.21ns	4.66 ns	27.45 ns	16.4 ns	52.36 ns
IR × BRh	4	44.08ns	1387.28**	0.33ns	3.06**	5.89ns	0.89ns	13.6 ns	27.13 ns	3.99 ns	23.98 ns
NF × BRh	4	832.44**	778.02ns	0.04ns	0.51ns	1.51ns	6.34ns	7.86 ns	40.51 ns	19.7 ns	32.15 ns
IR × NF × BRh	8	28.95ns	728.95*	0.014ns	0.09ns	1.61ns	5.19ns	0.84 ns	19.93 ns	0.29 ns	28.92 ns
Eb	48	141.08	334.61	0.25	0.42	8.29	4.36	11.14	41.54	20.66	59.21
CV (%)		6.32	6.37	9.96	7.25	10.83	7.75	11.00	15.16	8.49	10.48

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

Ea: Error a; Eb: Error b; CV: Coefficient of Variation; SS: Soluble sugars; PC: Proline content; GPr: Grain protein; CAT: Catalase; SOD: Superoxide dismutase.

For SS content, the response of guar plants to different IR during the first year of the experiment (Fig. 7) indicated that deficit irrigation up to 50% podding produced the highest soluble sugar content ($216.72 \mu\text{g g}^{-1} \text{DW}$), representing significant increases of 34.65% and 17.53% compared with full irrigation ($160.95 \mu\text{g g}^{-1} \text{DW}$) and deficit irrigation up to 50% flowering ($184.40 \mu\text{g g}^{-1} \text{DW}$), respectively. Furthermore, the NF × BRh interaction significantly influenced SS accumulation (see Table 7), with the highest SS level ($219.07 \mu\text{g g}^{-1} \text{DW}$) recorded under $100 \text{ kg N ha}^{-1} \times \text{RS-153}$. In contrast, the lowest value ($160.59 \mu\text{g g}^{-1} \text{DW}$) was observed for the control (no NF) × RS-150 interaction. These results highlight that water deficit during the podding stage strongly induces osmolyte accumulation, while the synergistic application of optimal nitrogen levels and *Bradyrhizobium* inoculation further enhances osmotic adjustment and soluble sugar accumulation in guar.




Fig. 7. Effects of IR on the content of SS (adapted from first-year data)
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Based on the second-year data, a significant three-way interaction of IR $\times$ NF $\times$ BRh was observed for SS (Fig. 8), showing that the highest SS content ($490.16 \mu\text{g g}^{-1} \text{DW}$) occurred under deficit irrigation up to 50% podding $\times 100 \text{ kg N ha}^{-1} \times \text{RS-150}$, whereas the lowest value ($112.41 \mu\text{g g}^{-1} \text{DW}$) was recorded under full irrigation $\times 100 \text{ kg N ha}^{-1} \times \text{non-inoculated}$ conditions. These results indicate that the combined application of adequate nitrogen and *Rhizobium* inoculation, particularly under moderate water deficit, effectively enhances SS accumulation, likely contributing to improved osmotic adjustment and stress tolerance in guar plants.





Fig. 9. Effects of the interaction of IR $\times$ NF $\times$ BRh on the content of SS trait
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Mean squares for the PC trait obtained from the first-year data (Table 10) showed that the highest value ($5.66 \mu\text{mol g}^{-1} \text{FW}$) was recorded in plants subjected to deficit irrigation up

to 50% podding. Regarding nitrogen fertilization, both 50 and 100 kg N ha⁻¹ treatments (averaging 5.77 $\mu\text{mol g}^{-1}$ FW) led to a 65.33% increase in leaf proline content compared with the control (3.49 $\mu\text{mol g}^{-1}$ FW). Moreover, plants under non-inoculation conditions (5.31 $\mu\text{mol g}^{-1}$ FW) exhibited significantly higher PC values compared to those inoculated with RS-150 and RS-153 strains (4.94 and 4.77 $\mu\text{mol g}^{-1}$ FW, respectively), with no significant difference between the two inoculated treatments.

In the second year of the study, PC was significantly affected by the IR $\times$ NF and IR $\times$ BRh interactions. Results of the IR $\times$ NF interaction (Fig. 9a) showed that the highest proline contents (13.67 and 13.44 $\mu\text{mol g}^{-1}$ FW) were recorded under deficit irrigation up to 50% podding combined with 100 and 50 kg N ha⁻¹, respectively. In contrast, the lowest PC (6.18 $\mu\text{mol g}^{-1}$ FW) was observed under full irrigation $\times$ non-inoculation treatment (Fig. 9a). Similarly, the IR $\times$ BRh interaction (Fig. 9b) revealed that the highest proline accumulation occurred under deficit irrigation up to 50% podding $\times$ non-inoculation, which was not significantly different from the deficit irrigation up to 50% podding $\times$ RS-150 inoculation treatment (11.74 $\mu\text{mol g}^{-1}$ FW). Conversely, the lowest significant PC (5.97 $\mu\text{mol g}^{-1}$ FW) was recorded under full irrigation $\times$ non-inoculation conditions (Fig. 9b).






Fig. 9. Effects of interactions of (a) IR × NF and (b) IR × BRh on the PC trait
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

According to ANOVA results, the main effects of IR, NF levels, and BRh inoculation significantly influenced GPr in both years of the experiment. In the first year (Table 8), full irrigation produced the highest GPr (29.27%), representing increases of 11.89% and 22.19% compared to deficit irrigation up to 50% flowering and 50% podding, respectively. Regarding NF application, the maximum and minimum GPr values (29.34% and 24.19%) were recorded under 100 kg N ha⁻¹ and the control (no NF), respectively. Seed inoculation with RS-153 also resulted in the highest GPr (28.15%), while non-inoculated plants showed the lowest value (24.01%). In the second year, full irrigation again resulted in the highest GPr (26.92%), which was not significantly different from deficit irrigation up to 50% flowering (26.60%), whereas deficit irrigation up to 50% podding produced the lowest GPr (24.27%). Nitrogen application exhibited a concentration-dependent effect, with 100 kg N ha⁻¹ increasing GPr by 23.06% and 12.33% compared with the control and 50 kg N ha⁻¹ treatments, respectively. Additionally, *Bradyrhizobium* inoculation significantly enhanced GPr relative to non-inoculated plants, with RS-153 (28.06%) outperforming both RS-150 (27.52%) and the non-inoculated control (25.21%).

4.2.3. Enzymatic activities of catalase (CAT) and superoxide dismutase (SOD)

The ANOVA results (Table 9) revealed that the experimental treatments had significant effects on the enzymatic activities examined in this study. Specifically, IR, NF levels, and BRh inoculation significantly influenced CAT and SOD in both years of the experiment at $p < 0.01$.

According to the first-year data (Table 8), the highest and lowest significant CAT activities (70.35 and 23.98 μmol H₂O₂.mg⁻¹ protein min⁻¹) and SOD activities (11.61 and 27.45 U mg⁻¹ protein min⁻¹) were observed under deficit irrigation up to 50% podding and full irrigation treatments, respectively. Regarding NF effects, increasing NF levels significantly reduced CAT and SOD activities. Under control (no NF application), CAT and SOD activities were 81 μmol H₂O₂.mg⁻¹ protein min⁻¹ and 37.56 U mg⁻¹ protein min⁻¹, respectively. Additionally, seed inoculation with RS-153 decreased CAT activity by 12.32% and 18.8%, and SOD activity by 13.63% and 5.10% compared with the control and RS-150 inoculation, respectively. In the second year of the experiment, mean comparisons (Table 8) showed that the highest and

lowest values for CAT (20.52 and 31.52 $\mu\text{mol H}_2\text{O}_2\cdot\text{mg}^{-1}\text{ protein min}^{-1}$) and SOD (21.90 and 17.56 $\text{U}\cdot\text{mg}^{-1}\text{ protein min}^{-1}$) were observed under deficit irrigation up to 50% podding and full irrigation, respectively. Increasing NF levels significantly decreased CAT and SOD activities, with minimum and maximum activities observed under control (no NF application) and 100 kg N ha^{-1} , respectively. Furthermore, inoculation with *Rhizobium* strains (RS-150 and RS-153) led to a significant reduction in enzymatic activities compared to non-inoculated plants (Table 8).

4.3. Quantitative traits of grain number (GN), 1000-grain weight (TGW), and grain yield (GY)

In general, ANOVA results (Table 10) indicated that all main factors, including IR, NF, and inoculation/non-inoculation conditions, significantly influenced TGW, GN, and GY during the first year of the study. In the second year, interactions between treatments also showed notable effects: GN was significantly affected by IR $\times$ NF ($p < 0.01$) and IR $\times$ BRh ($p < 0.05$), while GY was significantly influenced by the IR $\times$ NF interaction ($p < 0.01$).

Table 10. Effects of IR, NF levels, and inoculation/non-inoculation of *Rhizobium* bacteria on yield and yield components

Sources of variations	d f	GN		TGW		GY	
		First year	Second year	First year	Second year	First year	Second year
Block	2	426847.49 ns	558479 ns	1.08 ns	34.53 ns	59422.08 ns	269631.01 ns
Irrigation regimes (IR)	2	4133915.0 1**	138803101 .2**	159.12 **	220.22**	1370373.4 6**	20548539. 5**
Ea	4	37597.11	85619.8	6.74	17.78	16047.06	109666.86
Nitrogen Fertilizer (NF) levels	2	2414560.6 4**	18270242* *	128.03 **	185.46**	740600.74 **	4628131.8 **
<i>Bradyrhizobium</i> Inoculation (BRh) conditions	2	2197763.1 2**	3754212.3* *	113.07 **	80.79**	691384.45 **	1218157.8 **
IR $\times$ NF	4	57477.42 ns	3953471.6* *	4.86 ns	2.12 ns	15520.94 ns	603128.49 **
IR $\times$ BRh	4	44569.29 ns	969797.5 * *	0.15 ns	1.66 ns	18096.77 ns	82825.6 ns
NF $\times$ BRh	4	30671.75 ns	228938.3 ns	3.18 ns	3.68 ns	5709.68 ns	36561.09 ns
IR $\times$ NF $\times$ BRh	8	39297.75 ns	276656.8 ns	3.4 ns	2.99 ns	26836.92 ns	21894.72
Eb	4 8	147123.3	340506.2	11.23	12.27	28541.09	97098.02
CV (%)		9.1	9.22	11.53	12.64	13.71	16.41

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

Based on the first-year data (Table 11), full irrigation (4662.52 grains m^{-2}) significantly increased GN m^{-2} by 15.70% and 18.12% compared with deficit irrigation up to 50% flowering (4029.78 grains m^{-2}) and 50% podding (3947.33 grains m^{-2}), respectively. Similarly, the application of 100 kg N ha^{-1} resulted in 4544.8 grains m^{-2} , representing significant increases of 14.65% and 10.02% relative to the control (3964 grains m^{-2}) and 50 kg N ha^{-1} (4130.8 grains m^{-2}), respectively. Seed inoculation with RS-150 and RS-153 strains also improved GN m^{-2} compared with the non-inoculated control, with the highest (4467.1 grains m^{-2}) and lowest (3904.4 grains m^{-2}) values recorded for RS-153 and non-inoculated treatments, respectively. Overall, RS-153 inoculation increased GN m^{-2} by 14.41% and 4.66% compared with non-inoculated and RS-150-inoculated plants, respectively. Regarding TGW, the highest value in the first year (31.6 g) was obtained under full irrigation, corresponding to significant increases of 9.57% and 18.07% relative to deficit irrigation up to 50% flowering and 50% podding, respectively. Application of 100 kg N ha^{-1} increased TGW by 15.90% and 4.25% compared with the control and 50 kg N ha^{-1} treatments, respectively. Among inoculation treatments, TGW was highest in RS-153-inoculated plants (30.64 g) and lowest in the non-inoculated control (26.75 g). In the first year of the experiment, GY of guar plants

ranged from 1042.1 to 1481.3 kg ha⁻¹, with the highest value recorded under full irrigation and the lowest under deficit irrigation up to 50% podding. Nitrogen fertilization also significantly influenced GY, where 100 kg N ha⁻¹ resulted in the maximum yield (1481.3 kg ha⁻¹). Similarly, inoculation treatments significantly affected GY, with the highest and lowest values observed in RS-153-inoculated (1363.60 kg ha⁻¹) and non-inoculated control plants (1054.27 kg ha⁻¹), respectively (Table 11).

Table 11. Effects of IR, NF levels, and inoculation conditions on some traits of guar

Sources of variations	GN (number)	TGW (g)		GY (kg ha ⁻¹)
	First year	First year	Second year	First year
Irrigation regimes (IR)				
Full irrigation	4662.52 a	31.6 a	32.1 a	1481.3 a
Deficit irrigation up to 50% flowering	4029.78 b	28.84 b	28.37 b	1174.6 b
Deficit irrigation up to 50% podding	3947.33 b	26.76 c	26.49 b	1042.1 c
LSD	146.52	1.96	3.19	95.7
NF levels (kg ha⁻¹)				
Control	3964 b	26.66 b	26.4 c	1068.6 c
50	4130.8 b	29.64 b	28.93 b	1299.51 b
100	4544.8 a	30.9 a	31.64 a	1399.79 a
BRh Inoculation conditions				
Control (non-inoculation)	3904.4 b	26.75 b	27 b	1054.27 b
RS 150	4268.1 a	29.8 a	29.76 a	1280.03 a
RS 153	4467.1 a	30.64 a	30.19 a	1363.60 a
LSD	209.9	1.83	1.95	92.45

Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Based on the second-year data (Fig. 10a), GN per m² was significantly affected by the interaction of IR × NF. The highest GN (5066.33 grains) was recorded under full irrigation × 100 kg N ha⁻¹, which showed a significant increase compared to other treatment combinations. In contrast, the lowest GN (3675 grains) occurred under deficit irrigation up to 50% podding × no N application, although it was not significantly different from the combinations of deficit irrigation up to 50% flowering × no N fertilizer (3866 grains), deficit irrigation up to 50% flowering × 50 kg N ha⁻¹ (3875 grains), and deficit irrigation up to 50% podding × 50 kg N ha⁻¹ (3870.33 grains). Furthermore, according to the results of the GN trait in the second year (Fig. 10b), the highest GN (4943 grains) was obtained under full irrigation × seed inoculation with the RS-153 strain, which was statistically similar to the combination of full irrigation × RS-150 strain (4700 grains). The lowest GN (3648 grains) was recorded under deficit irrigation up to 50% podding × non-inoculation, although it did not differ significantly from the combination of deficit irrigation up to 50% flowering × non-inoculation (3721 grains).




Fig. 10. Interaction effects of (a) IR × NF levels and (b) IR × Rh inoculation conditions on GN per square meter (adapted from the second-year data)

Means followed by the same letter within each column do not differ significantly at $p < 0.05$

According to the second-year data (Table 11), TGW was significantly influenced by irrigation regimes, showing a direct positive relationship with IR levels. The highest TGW (32.1 g) was obtained under full irrigation, which showed significant increases of 13.15% and 21.18% compared with deficit irrigation up to 50% flowering and up to 50% podding, respectively. Consistent with the first-year results, applying 100 kg N ha⁻¹ produced the highest TGW (31.64 g), which was significantly higher than those of the control (26.40 g) and 50 kg N ha⁻¹ (28.93 g) treatments, corresponding to increases of 19.85% and 9.37%, respectively. Furthermore, comparison of means related to the inoculation/non-inoculation factor indicated that the highest TGW (30.19 g) was recorded for seeds inoculated with the RS-153 strain, which was not significantly different from that obtained with RS-150 (29.76 g).

Based on the second-year data (Fig. 11), comparing means related to the interaction effects of IR × NF on GY revealed that drought stress significantly reduced GY, although this reduction was partially compensated by N fertilizer application. Overall, the highest GY (3658.2 kg ha⁻¹) was obtained from the interaction of full irrigation × 100 kg N ha⁻¹, which showed a significant difference compared with other interaction treatments. In contrast, the lowest GY value (1017.8 kg ha⁻¹) was recorded for the interaction of deficit irrigation up to

50% podding $\times$ non-application of NF. Overall, these results demonstrate that optimized water and nitrogen management, along with seed inoculation, can effectively moderate yield losses under drought stress, an aspect discussed in detail in the following section.




Fig. 11. Interaction effects of IR $\times$ NF levels on GN per square meter (adapted from the second-year data)
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

1.1. Forage quality parameters

This section presents the effects of IR, NF levels, and BRh inoculation on key forage quality traits over two consecutive years (2021–2022 and 2022–2023). For clarity, comparisons focused on the best- and worst-performing treatments at each stage. The ANOVA results (Table 12) indicated that both IR and NF exerted significant effects ($p < 0.01$) on CrPro, ash%, ADF, DMD, and ME across both years, highlighting the critical role of water and nitrogen management in determining forage nutritive value. In contrast, BRh inoculation significantly influenced CrPro in the second year ($p < 0.01$), DMD in the first year ($p < 0.01$), ME in the first year ($p < 0.05$), and ash% in both years ($p < 0.01$). Notably, NDF was significantly affected by IR and the IR $\times$ NF interaction in the first year ($p < 0.01$) and by IR alone in the second year ($p < 0.01$). These findings suggest that, while water and nitrogen supply predominantly govern overall forage quality, microbial inoculation can further enhance specific traits, underscoring the potential of integrated management practices to optimize both yield and nutritive value of guar under varying environmental conditions.

Table 12. ANOVA (mean squares) for selected forage quality traits of guar under IR, NF levels, and *Bradyrhizobium japonicum* inoculation

Source of variation	df	CrPr		NDF		ADF		Ash		DMD		ME	
		First year	Second year	First year	Second year	First year	Second year	First year	Second year	First year	Second year	First year	Second year
Block	2	1.05 ns	3.88ns	10.3 ns	12.1 ns	18.21n s	0.43 ns	1.33 ns	1.09ns	10.96n s	0.99 ns	0.32 ns	0.03 ns
Irrigation regimes (IR)	1	167.11**	233.48**	4266.5**	1025.3**	2158.22**	673.92**	59.2* *	121.37**	1845.6**	724.69**	53.3* *	20.98**
Ea	2	12.3	1.27	14.34	0.27	14.8	0.41	2.8	2.11	19.11	0.03	0.56	0.0009
Nitrogen fertilizer levels (NF)	1	48.85**	34.61* *	0.99 ns	13.2 ns	113.92**	42.29**	65.34**	55.16* *	128.2* *	56.1* *	3.69* *	1.63* *
<i>Bradyrhizobium</i> (BRh) inoculum	2	4.99 ns	11.48* *	0.67 ns	0.14 ns	38.54n s	1.03 ns	18.01**	17.99* *	34.2**	3.93 ns	0.99* *	0.11 ns

IR × NF	1	19.8 ns	0.0001 ns	73.2* *	5.33 ns	0.29ns	0.19 ns	0.47 ns	1.06ns	1.38ns	0.13 ns	0.04 ns	0.00 4ns
IR × BRh	2	0.34 ns	1.18ns	9.83 ns	0.71 ns	5.27ns	0.25 ns	3.1ns	0.28ns	3.74ns	0.01 ns	0.11 ns	0.00 04ns
NF × BRh	2	0.21 ns	0.37ns	0.77 ns	0.82 ns	0.7ns	0.02 ns	0.34 ns	1.02ns	0.62ns	0.01 ns	0.03 ns	0.00 03ns
IR × NF × BRh	2	1.65 ns	27ns	5.35 ns	0.08 ns	0.89ns	0.11 ns	1.11 ns	0.37ns	0.13ns	0.09 ns	0.00 4ns	0.00 2ns
Eb	2 5	5.06	1.64	14.6 2	8.99	13.97	3.17	2.81	2.92	9.59	2.06	0.28	0.06
CV		18.4 8	9.22	11.2 9	10.3 2	14.3	12.2 6	15.8 6	13.15	4.66	1.87	5.66	2.22

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

According to Table 13, full irrigation and 100 kg N ha⁻¹ led to significant improvements in all forage quality parameters studied in both experimental years (except for ADF). Accordingly, full irrigation in both trial years resulted in notable increases of 43.01 and 44.89% for CrPro, 21.02 and 15.53% for NA, 27.69 and 32.74% for ash, 24.16 and 12.45% for DMD, and 29.91 and 14.91% for ME, and significant decreases of 45.71 and 45.95% for ADF, respectively. Also, the application of 100 kg N ha⁻¹ in comparison to the control treatment significantly increased CrPro (18.44 and 15.27%), ash (29.28 and 21%), DMD (5.84 and 3.32%), and ME (7.13 and 3.98%), and decreased ADF (12.75 and 13.89%), respectively (Table 13).

Table 13. Mean comparisons (mean squares) of the effects of IR and NF levels on selected forage quality traits of guar

Source of variation	CrPro (%)		ADF (%)		Ash (%)		DMD (%)		ME (MJ.Kg ⁻¹ DM)	
	First year	Second year	First year	Second year	First year	Second year	First year	Second year	First year	Second year
IR										
Full irrigation	14.33a	16.43a	18.4b	10.2b	11.85a	14.88a	73.6a	81.12a	10.51a	11.79a
Deficit irrigation up to 50% podding	10.02b	11.34b	33.89a	18.67a	9.28b	11.21b	59.28b	72.14b	8.09b	10.26b
LSD	4.03	1.62	5.52	0.92	2.4	2.09	6.27	0.25	1.07	0.04
NF levels (Kg ha⁻¹)										
Control	11.01b	12.9b	27.92a	15.62a	9.22b	11.81b	64.56b	75.38b	8.98b	10.81b
100	13.34a	14.87a	24.36b	13.45b	11.92a	14.29a	68.33a	77.88a	9.62a	11.24a
LSD	1.56	0.89	2.6	1.24	1.17	1.19	2.15	0.99	0.37	0.17

Means followed by the same letter within each column do not differ significantly at $p < 0.05$

In addition, seed inoculation with *Bradyrhizobium japonicum* strains had a significant effect on CrPro values in the second year of the experiment. Under non-inoculated conditions, the CrPro value (12.77%) showed significant reductions of 12.41% and 10.70% compared to the values obtained from seeds inoculated with RS-150 and RS-153 strains, respectively (Fig. 12).





Fig. 12. Effects of inoculation conditions on the CPro trait (Based on the first-year data)
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Based on the significant effects of the inoculation/non-inoculation factor on the ash percentage of guar plants in both years of the experiment, it was observed (Fig. 13a) that the highest ash percentages (11.85% in the first year and 13.95% in the second year) were obtained under the application of the RS-153 strain, which showed a significant difference compared to the control (non-inoculated) treatment. The lowest significant values of ash percentage were recorded in the control treatment (9.21% in the first year and 11.65% in the second year). In evaluating the effects of inoculation/non-inoculation treatments with different strains of *Bradyrhizobium* on DMD percentage in the first year of the experiment (Fig. 13b), the highest value of this trait (67.49%) was obtained from the application of the RS-150 strain, which showed no significant difference compared to the DMD value recorded under the RS-153 strain (67.35%). The lowest significant DMD value (64.49%) was observed in the non-inoculated (control) treatment. Finally, comparison of mean effects of inoculation conditions on ME percentage (adopted from first-year data; Fig. 11c) showed that non-inoculation of guar seeds caused significant decreases in ME. The ME values for the non-inoculated control, RS-150, and RS-153 inoculated treatments were estimated at 8.96, 9.40, and 9.47 MJ kg⁻¹, respectively (Fig. 13c).






Fig. 13. Effects of inoculation conditions on (a) Ash, (b) DMD, and (c) ME attributes
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

Based on the first-year mean comparisons for the IR $\times$ NF interaction on NDF (Fig. 14a), the highest NDF value (46%) was observed under deficit irrigation up to 50% podding $\times$ no NF. In contrast, the lowest NDF values (37.21% and 36.24%) were recorded for full irrigation $\times$ no NF and full irrigation $\times$ 100 kg N ha⁻¹, respectively. In the second year of the experiment (Fig. 14b), plants grown under full irrigation exhibited 23.73% NDF, which was 31.02% lower than those subjected to deficit irrigation up to 50% podding (34.4% NDF). These results highlight that both water availability and nitrogen supply strongly influence NDF accumulation, with water deficit substantially increasing fiber content, whereas full irrigation and adequate nitrogen reduce NDF and potentially improve forage digestibility.




Fig. 14. Effects (a) IR $\times$ NF interaction and (b) IR on NDF
Means followed by the same letter within each column do not differ significantly at $p < 0.05$

2. Discussion

As stated earlier, drought stress markedly altered phenological progression in guar, primarily through osmotic imbalance and reduced cellular turgor that constrain cell expansion and carbon assimilation [84,85]. This reduction in soil moisture not only shortens critical growth phases such as flowering (D50F), pod formation (D50P), and physiological maturity (DH), but also reflects a strategic shift in life-history traits, where the plant prioritizes early reproductive completion to escape severe stress. Rather than merely reflecting developmental acceleration, the shortened phenology observed under deficit

irrigation suggests a strategic shift toward drought escape, where life-cycle completion precedes severe stress but at the expense of assimilate accumulation and sink development [86,87,88,89,90]. This pattern highlights a key adaptive mechanism in guar, emphasizing survival over maximal reproductive output, which has implications for timing irrigation and nutrient management to support yield stability. Similar stage-specific sensitivity has been reported across legumes, with reproductive stages exhibiting higher vulnerability due to their dependence on sustained carbon supply and translocation [91,92,93]. The observed reduction in phenological duration in the present study reflects a source/sink imbalance under water limitation, consistent with recent syntheses on drought-induced carbon partitioning trade-offs [94]). Although guar is recognized as a drought-adapted legume, its adaptive responses, such as stomatal regulation, reduced leaf expansion, and altered carbon allocation, primarily preserve survival rather than maximizing reproductive output [26,95,96,97]. Such mechanistic insights indicate that deficit irrigation management must consider phenological shifts to prevent critical yield losses.

Nitrogen supply significantly modulated plant responses to deficit irrigation. Adequate N availability sustains chlorophyll biosynthesis, Rubisco activity, and antioxidant metabolism, thereby maintaining photosynthetic efficiency under stress [98]. In this study, the application of 50-100 kg N ha⁻¹ not only prolonged flowering and podding phases but likely enhanced assimilate remobilization and reinforced reproductive sink strength, mitigating the source/sink imbalance induced by water limitation. This response aligns with findings that nitrogen enhances root proliferation, hydraulic conductance, and water uptake capacity, partially compensating for drought-induced reductions in turgor [99,100,101]. These interactions underline the synergistic potential of water and nitrogen management, suggesting that optimized N supply can maintain photosynthetic and reproductive function even under deficit irrigation. Importantly, water-nitrogen interactions exert a significant control on nitrogen use efficiency (NUE) under deficit irrigation; optimized nitrogen supply enhances nitrogen uptake and translocation, thereby preventing both dilution effects and metabolic limitation commonly observed under water stress. Such integrative management strategies may therefore improve both carbon and nitrogen use efficiency in water-limited systems, reinforcing yield stability. For instance, combined increases in irrigation and nitrogen markedly improved NUE components such as pre- and post-anthesis N translocation in wheat, indicating a synergistic effect of water and nitrogen on crop nitrogen dynamics [102,103]. Collectively, these findings emphasize the importance of synchronizing irrigation and nitrogen application to sustain both vegetative and reproductive functions under drought.

Rhizobium inoculation further reinforced drought resilience through multiple complementary mechanisms. Beyond biological nitrogen fixation, symbiotic bacteria enhance root architecture, exopolysaccharide production, and soil aggregation, improving rhizosphere water retention and nutrient availability [104,105,106,107]. PGPR-mediated hormonal modulation, particularly through the production and regulation of auxin (IAA) and cytokinins, stimulates root system growth and mitigates drought-induced senescence by modulating endogenous hormonal pathways controlling cell division, lateral root development, and stress signaling [84,108]. This integrated microbial effect likely underpins the observed improvements in chlorophyll, carotenoid content, and grain protein under combined N and *Rhizobium* treatments. These results suggest that integrating biological and mineral N sources enhances carbon allocation efficiency and strengthens reproductive sink establishment, providing a mechanistic explanation for improved yield components under water-limited conditions.

Biochemically, drought stress elevated soluble sugars and antioxidant enzyme activities (CAT, SOD), reflecting activation of osmotic adjustment and ROS detoxification pathways [109,110,111]. The accumulation of osmolytes and enhancement of enzymatic antioxidants stabilizes membranes, preserves photosystem II efficiency, and maintains cellular redox

balance [112,113,114,115,116]. Under nitrogen fertilization and *Rhizobium* inoculation, these protective mechanisms were further strengthened, indicating that nutrient sufficiency supports metabolic resilience by sustaining both osmotic regulation and antioxidant capacity [117,118,119]. It should be noted that the observed decrease in CAT and SOD activities under higher nitrogen supply and *Rhizobium* inoculation likely reflects reduced oxidative stress resulting from improved water and nutrient status. In other words, the plants experience lower ROS accumulation and therefore require less enzymatic scavenging. This interpretation is consistent with the understanding that antioxidant enzyme activity is dynamic, responding to the intensity of oxidative challenge rather than serving as a static indicator of defense capacity. This interpretation is supported by evidence that antioxidant enzyme activities are dynamic and closely reflect the level of oxidative challenge; when ROS accumulation is reduced under improved water and nutrient status, the requirement for high CAT and SOD activity diminishes accordingly, indicating lower oxidative stress rather than impaired defense capacity, e.g., ROS have dual roles in plant stress signalling and antioxidant responses; both ROS production and antioxidant activity adjust dynamically to stress intensity [120,121]. From a functional perspective, these biochemical adjustments help maintain photosynthetic performance and partially protect yield formation under deficit irrigation.

Forage quality responses were closely aligned with physiological adjustments. Deficit irrigation increased fiber fractions (ADF, NDF) while reducing crude protein and digestible dry matter, consistent with stress-induced shifts toward structural carbohydrate accumulation [122,123,124]. This indicates that water limitation triggers a trade-off between structural integrity and forage nutritive value, which can influence livestock performance. In contrast, full irrigation combined with 100 kg N ha⁻¹ improved protein concentration, metabolizable energy, and digestibility, reflecting enhanced nitrogen assimilation and sustained carbon metabolism. Comparable irrigation-nitrogen interactions affecting forage nutritive value have been documented in annual ryegrass [125]. *Bradyrhizobium* inoculation further improved protein and digestibility parameters in a strain-dependent manner, consistent with enhanced N fixation and nutrient uptake efficiency [126,127,128,129]. These results collectively highlight that reproductive-stage water availability is critical for nutrient remobilization and forage quality, and that combined nutrient-microbial management can buffer quality losses under water stress.

A comparison with previous studies further strengthens the interpretation of the present findings. Consistent with our results, drought stress has been widely reported to reduce growth, seed yield, and forage quality in legumes, primarily through limitations in photosynthetic carbon assimilation and assimilate partitioning [130,131], and in the present study, the observed reductions in phenological duration and forage quality under deficit irrigation are in line with these established responses. However, inoculation with *Bradyrhizobium* partially alleviated these constraints, which agrees with earlier reports demonstrating that rhizobial symbiosis enhances biological nitrogen fixation, improves root system architecture, and maintains plant water relations under drought conditions [84,132]. Similar enhancements in protein content and forage nutritive value following inoculation have been reported in legumes such as soybean and cowpea, where improved nitrogen availability and metabolic activity contributed to better yield quality under water-limited environments [105,133,134]. Furthermore, previous studies have shown that the combined application of deficit irrigation and plant growth-promoting rhizobacteria (PGPRs) leads to greater yield stability compared with non-inoculated treatments, highlighting the buffering role of plant-microbe interactions under stress conditions [135–137]. Nevertheless, some variability among studies has been noted, which may be attributed to differences in environmental conditions, soil fertility status, and rhizobial strain specificity [138,139]. In summary, the present findings corroborate existing evidence and further demonstrate that integrating microbial inoculation with optimized water management can effectively mitigate

drought-induced limitations while sustaining both yield and forage quality in legume-based systems.

From a management perspective, integrated nitrogen and PGPRs are more effective to cope with drought conditions than single-factor interventions. Coordinated deficit irrigation with optimized nitrogen supply and symbiotic inoculation enhances physiological resilience, stabilizes sink development, and improves forage quality under semi-arid conditions. This integrative framework aligns with sustainable intensification principles highlighted by some researchers, where resource-use efficiency and stress buffering through coordinated water-nutrient management are central to productivity in water-limited agroecosystems [130,131]. Thus, these findings provide actionable guidance for optimizing irrigation scheduling, fertilization, and microbial inoculation to sustain both yield and forage quality in semi-arid legume systems.

Nevertheless, certain limitations should be acknowledged for this study. The study was conducted under specific environmental conditions and may not fully capture interannual climatic variability. Direct measurements of leaf water potential, gas exchange parameters, and nitrogen use efficiency indices were not included, limiting mechanistic quantification. Future research integrating physiological, isotopic, and molecular analyses would allow more precise causal inference regarding water, nitrogen, and PGPRs interactions and improve the transferability of results across diverse agroecosystems. Overall, the results demonstrate that deficit irrigation primarily constrains guar productivity through phenological shortening, source/sink imbalance, and oxidative stress, whereas optimized nitrogen fertilization and *Rhizobium* inoculation mitigate these constraints by sustaining photosynthetic capacity, antioxidant defense, and reproductive allocation. The synergistic integration of irrigation scheduling, nitrogen management, and microbial inoculation therefore represents a robust strategy for stabilizing guar yield and forage quality under arid and semi-arid conditions, providing both mechanistic insight and practical guidance for water-limited agroecosystems.

Conclusion

This study demonstrated that drought stress significantly reduced phenological duration, disrupted assimilate partitioning, and impaired grain development in guar, leading to decreased yield and forage quality. Shortened flowering and pod-filling phases under water deficit limited the accumulation of storage compounds, while osmotic imbalance and oxidative stress further constrained plant growth. Conversely, nitrogen fertilization enhanced chlorophyll content, photosynthetic efficiency, and protein synthesis, providing additional resources for seed filling and forage quality. Inoculation with *Bradyrhizobium* strains improved symbiotic nitrogen fixation and nutrient assimilation, contributing to higher crude protein content and improved forage digestibility. The combined application of nitrogen and *Bradyrhizobium* effectively mitigated drought effects by promoting osmotic adjustment, activating antioxidant defenses, and facilitating translocation of photosynthates to developing seeds. These findings highlight that integrating optimized irrigation scheduling with balanced nitrogen fertilization and effective rhizobial inoculation is a practical strategy to maintain yield and forage nutritive value under limited water availability. Future research should validate these findings over multiple growing seasons and investigate genotype-specific responses to refine sustainable management strategies for guar cultivation.

Data availability

The datasets generated, used, deployed, and analyzed during the present study are not accessible to the general public. However, they are available from the corresponding author upon adequate request.

References

1. Ogunleye, O. O., Owolabi, O. A. & Mubarak, M. Population growth and economic growth in Nigeria: an appraisal. *Int. J. Manag. Account. Econ.* **5**, 282–299 (2018).
2. Norouzi, S. & Akbari, G. Differential sensitivity and phytoremediation potential of faba bean (*Vicia faba* L.) cultivars irrigated with urban wastewater effluent. *Plant Soil* **520**, 1605–1630; <https://doi.org/10.1007/s11104-026-08346-5> (2026).
3. Miner, G. L., Delgado, J. A., Ippolito, J. A. & Stewart, C. E. Soil health management practices and crop productivity. *Agric. Environ. Lett.* **5**, e20023; <https://doi.org/10.1002/ael2.20023> (2020).
4. Țopa, D. C., Căpșună, S., Calistru, A. E. & Ailincăi, C. Sustainable practices for enhancing soil health and crop quality in modern agriculture: a review. *Agriculture (Basel)* **15**, 998; <https://doi.org/10.3390/agriculture15090998> (2025).
5. Jiang, N. H. Impacts of chemical fertilizer reduction and organic amendments supplementation on soil nutrient, enzyme activity and heavy metal content. *J. Integr. Agric.* **16**, 1819–1831; [https://doi.org/10.1016/S2095-3119\(16\)61476-4](https://doi.org/10.1016/S2095-3119(16)61476-4) (2017).
6. Azam, M. F. et al. Planting pattern and nitrogen management strategies: positive effect on yield and quality attributes of *Triticum aestivum* L. crop. *BMC Plant Biol.* **24**, 845; <https://doi.org/10.1186/s12870-024-05537-z> (2024).
7. Shekara, B. G. et al. Optimizing yield, quality, and nitrogen use efficiency in fodder maize through precision N-management. *Discov. Appl. Sci.* **7**, 812; <https://doi.org/10.1007/s42452-025-07232-6> (2025).
8. Vaghefi, S. A. et al. The future of extreme climate in Iran. *Sci. Rep.* **9**, 1464; <https://doi.org/10.1038/s41598-018-38071-8> (2019).
9. Farahani, E., Emami, H. & Forouhar, M. Effects of tillage systems on soil organic carbon and some soil physical properties. *Land Degrad. Dev.* **33**, 1307–1320; <https://doi.org/10.1002/ldr.4221> (2022).
10. Hua, W. et al. Manure application increased crop yields by promoting nitrogen use efficiency in the soils of 40-year soybean-maize rotation. *Sci. Rep.* **10**, 14882; <https://doi.org/10.1038/s41598-020-71932-9> (2020).
11. Barłóg, P., Grzebisz, W. & Łukowiak, R. Fertilizers and fertilization strategies mitigating soil factors constraining efficiency of nitrogen in plant production. *Plants* **11**, 1855; <https://doi.org/10.3390/plants11141855> (2022).
12. Tyagi, J., Ahmad, S. & Malik, M. Nitrogenous fertilizers: impact on environment sustainability, mitigation strategies, and challenges. *Int. J. Environ. Sci. Technol.* **19**, 11649–11672; <https://doi.org/10.1007/s13762-022-04027-9> (2022).
13. Anas, M., Liao, F., Verma, K.K. et al. Fate of nitrogen in agriculture and environment: agronomic, eco-physiological and molecular approaches to improve nitrogen use efficiency. *Biol Res* **53**, 47; <https://doi.org/10.1186/s40659-020-00312-4> (2020).
14. Gai, X. et al. Long-term benefits of combining chemical fertilizer and manure applications on crop yields and soil carbon and nitrogen stocks in North China Plain. *Agric. Water Manag.* **208**, 384–392; <https://doi.org/10.1016/j.agwat.2018.07.002> (2018).
15. Kebede, E. Contribution, utilization, and improvement of legumes-driven biological nitrogen fixation in agricultural systems. *Front. Sustain. Food Syst.* **5**, 767998; <https://doi.org/10.3389/fsufs.2021.767998> (2021).
16. Huang, Q. Enhancing soil health and biodiversity through nitrogen fixation symbiosis in leguminous plants. *Mol. Microbiol. Res.* **14**, 1–12; <https://doi.org/10.5376/mmr.2024.14.0006> (2024).
17. Gou, X. et al. Leguminous plants significantly increase soil nitrogen cycling across global climates and ecosystem types. *Glob Chang Biol.* **29**, 4028–4043; <https://doi.org/10.1111/gcb.16742> (2023).
18. Ma, J. et al. Legume-non-legume cover crop mixtures enhance soil nutrient availability and physical properties: a meta-analysis across Chinese agroecosystems. *Agron.* **15**, 1756; <https://doi.org/10.3390/agronomy15081756> (2025).

19. Suthar, J. D. et al. Evaluation of guar (*Cyamopsis tetragonoloba* L.) genotypes performance under different irrigation water salinity levels: growth parameters and seed yield. *Ind. Crop. Prod.* **123**, 247-253; <https://doi.org/10.1016/j.indcrop.2018.06.084> (2018).
20. Avola, G. et al. Deficit irrigation on guar genotypes (*Cyamopsis tetragonoloba* (L.) Taub.): effects on seed yield and water use efficiency. *Agron.* **10**, 789; <https://doi.org/10.3390/agronomy10060789> (2020).
21. Vishnyakova, M. A. et al. Drought stress response in guar (*Cyamopsis tetragonoloba* (L.) Taub.): physiological and molecular genetic aspects. *Plants* **12**, 3955; <https://doi.org/10.3390/plants12233955> (2023).
22. de Souza, M. et al. Guar gum as galactomannan source induces dysbiosis and reduces performance in broiler chickens and dietary β -mannanase restores the gut homeostasis. *Poult. Sci.* **102**, 102810; <https://doi.org/10.1016/j.psj.2023.102810> (2023).
23. Dodi, G., Sabau, R. E., Crețu, B. E. B. & Gardikiotis, I. Exploring the antioxidant potential of gellan and guar gums in wound healing. *Pharmaceutics* **15**, 2152; <https://doi.org/10.3390/pharmaceutics15082152> (2023).
24. Tahmouzi, S. et al. Application of guar (*Cyamopsis tetragonoloba* L.) gum in food technologies: a review of properties and mechanisms of action. *Food Sci. Nutr.* **11**, 4869-4897; <https://doi.org/10.1002/fsn3.3383> (2023).
25. Awan, S. J. et al. Guar gum in therapeutics: a succinct exploration. *Bull. Biol. Allied Sci. Res.* **2024**, 60; <https://doi.org/10.54112/bbasr.v2024i1.60> (2024).
26. Soltani-Gerdefaramarzi, S. et al. Differential responses of two local and commercial guar cultivars for nutrient uptake and yield components under drought and biochar application. *Sci. Rep.* **14**, 23665; <https://doi.org/10.1038/s41598-024-74849-9> (2024).
27. Basu, A. et al. Plant growth promoting rhizobacteria (PGPR) as green bioinoculants: recent developments, constraints, and prospects. *Sustainability* **13**, 1140; <https://doi.org/10.3390/su13031140> (2021).
28. Kebede, M. T. & Mengstie, G. Y. Optimization the efficacy of plant growth-promoting rhizobacteria via genetic engineering. *Discov. Plants* **2**, 154; <https://doi.org/10.1007/s44372-025-00240-1> (2025).
29. Bhardwaj, D. et al. Biofertilizers function as key player in sustainable agriculture by improving soil fertility, plant tolerance and crop productivity. *Microb. Cell Fact.* **13**, 66; <https://doi.org/10.1186/1475-2859-13-66> (2014).
30. Ajeng, A. A. et al. The effects of biofertilizers on growth, soil fertility, and nutrients uptake of oil palm (*Elaeis guineensis*) under greenhouse conditions. *Process.* **8**, 1681; <https://doi.org/10.3390/pr8121681> (2020).
31. Ahmad, H. M., Fiaz, S., Hafeez, S. et al. Plant growth-promoting rhizobacteria eliminate the effect of drought stress in plants: a review. *Front. Plant Sci.* **13**, 875774; <https://doi.org/10.3389/fpls.2022.875774> (2022).
32. Chaudhary, P., Singh, S., Chaudhary, A. et al. Overview of biofertilizers in crop production and stress management for sustainable agriculture. *Front. Plant Sci.* **13**, 930340; <https://doi.org/10.3389/fpls.2022.930340> (2022).
33. Shahwar, D., Mushtaq, Z., Mushtaq, H. et al. Role of microbial inoculants as bio fertilizers for improving crop productivity: a review. *Heliyon* **9**, e16134; <https://doi.org/10.1016/j.heliyon.2023.e16134> (2023).
34. He, S. et al. PGPR: key to enhancing crop productivity and achieving sustainable agriculture. *Curr. Microbiol.* **81**, 377; <https://doi.org/10.1007/s00284-024-03893-5> (2024).
35. Khan, A. et al. Applications of beneficial plant growth promoting rhizobacteria and mycorrhizae in rhizosphere and plant growth: a review. *Int. J. Agric. Biol. Eng.* **13**, 199-208; <https://doi.org/10.25165/j.ijabe.20201305.5762> (2020).
36. Fasusi, O. A., Babalola, O. O. & Adejumo, T. O. Harnessing of plant growth-promoting rhizobacteria and arbuscular mycorrhizal fungi in agroecosystem sustainability. *CABI Agric. Biosci.* **4**, 26; <https://doi.org/10.1186/s43170-023-00168-0> (2023).

37. Nagrale, D. T. et al. PGPR: the treasure of multifarious beneficial microorganisms for nutrient mobilization, pest biocontrol and plant growth promotion in field crops. *World J. Microbiol. Biotechnol.* **39**, 100; <https://doi.org/10.1007/s11274-023-03536-0> (2023).
38. Ibrahim, K. A. et al. Response of guar (*Cyamopsis teteragonolopa* L.) to *Bradyrhizobium* inoculations in semi-arid environment. *IJAF* **6**, 137-141; <https://doi.org/10.5923/j.ijaf.20160604.01> (2016).
39. Gopalakrishnan, S. et al. Plant growth promoting rhizobia: challenges and opportunities. *3 Biotech* **5**, 355-377; <https://doi.org/10.1007/s13205-014-0241-x> (2015).
40. Benmoussa, S. et al. Growth performance and nitrogen fixing efficiency of faba bean (*Vicia faba* L.) genotypes in symbiosis with rhizobia under combined salinity and hypoxia stresses. *Agron.* **12**, 606; <https://doi.org/10.3390/agronomy12030606> (2022).
41. Goyal, R. K. & Habtewold, J. Z. Evaluation of legume-rhizobial symbiotic interactions beyond nitrogen fixation that help the host survival and diversification in hostile environments. *Microorganisms* **11**, 1454; <https://doi.org/10.3390/microorganisms11061454> (2023).
42. Yeremko, L. et al. Role of environmental factors in legume-rhizobium symbiosis: a review. *Biomolecules* **15**, 118; <https://doi.org/10.3390/biom15010118> (2025).
43. Ochieno, D. M. et al. Rhizobium-linked nutritional and phytochemical changes under multitrophic functional contexts in sustainable food systems. *Front. Sustain. Food Syst.* **4**, 604396; <https://doi.org/10.3389/fsufs.2020.604396> (2021).
44. Vitale, L., Vitale, E., Guercia, G. et al. Effects of different light quality and biofertilizers on structural and physiological traits of spinach plants. *Photosynthetica* **58**, 932-943 (2020).
45. Ammar, E. E. et al. A comprehensive overview of eco-friendly bio-fertilizers extracted from living organisms. *Environ. Sci. Pollut. Res.* **30**, 113119-113137; <https://doi.org/10.1007/s11356-023-30260-x> (2023).
46. Nabati, J. et al. Biofertilizers containing plant growth promoting rhizobacteria enhance nutrient uptake and improve the growth and yield of chickpea plants in an arid environment. *Sci. Rep.* **15**, 8331; <https://doi.org/10.1038/s41598-025-93070-w> (2025).
47. Norouzi, S. & Akbari, G. Changes in biochemical compositions and salinity tolerance responses of different bread wheat varieties cultivated in an arid and semi-arid climate. *Sci. Rep.* **15**, 44914; <https://doi.org/10.1038/s41598-025-28944-0> (2025).
48. Dubois, M., Inzé, D. Plant growth under suboptimal water conditions: early responses and methods to study them. *J. Exp. Bot.* **71**, 1706-1722; <https://doi.org/10.1093/jxb/eraa037> (2020).
49. Flynn, N. E. et al. High N availability decreases N uptake and yield under limited water availability in maize. *Sci. Rep.* **13**, 14269; <https://doi.org/10.1038/s41598-023-40459-0> (2023).
50. Kapoor, D. et al. The impact of drought in plant metabolism: how to exploit tolerance mechanisms to increase crop production. *Appl. Sci.* **10**, 5692; <https://doi.org/10.3390/app10165692> (2020).
51. Torkaman Pary, A. et al. Impacts of drought severity and frequency on natural vegetation across Iran. *Water* **16**, 3334; <https://doi.org/10.3390/w16223334> (2024).
52. Sohrabi, R. The politics of drought in the Middle East, case study from Iran. *Discov. Environ.* **3**, 122; <https://doi.org/10.1007/s44274-025-00334-3> (2025).
53. Lakhiar, I. A. et al. A review of precision irrigation water-saving technology under changing climate for enhancing water use efficiency, crop yield, and environmental footprints. *Agriculture* **14**, 1141; <https://doi.org/10.3390/agriculture14071141> (2024).
54. Li, M. et al. Climate-smart irrigation strategy can mitigate agricultural water consumption while ensuring food security under a changing climate. *Agric. Water Manag.* **292**, 108663; <https://doi.org/10.1016/j.agwat.2023.108663> (2024).

55. Xue, R. et al. Review of advancements and challenges in delayed irrigation: enhancing crop water productivity and sustainable crop production. *Agric. Water Manag.* **318**, 109739; <https://doi.org/10.1016/j.agwat.2025.109739> (2025).
56. Lesk, C. et al. Compound heat and moisture extreme impacts on global crop yields under climate change. *Nat. Rev. Earth Environ.* **3**, 872–889; <https://doi.org/10.1038/s43017-022-00368-8> (2022).
57. Bistgani, Z. E., Barker, A. V. & Hashemi, M. Physiology of medicinal and aromatic plants under drought stress. *Crop J.* **12**, 330–339; <https://doi.org/10.1016/j.cj.2023.12.003> (2024).
58. Jiang, X., Li, G. & Luo, S. The research progress on the mechanisms of drought stress at different growth stages affecting rice yield formation. *Resour. Data J.* **4**, 302–321; https://doi.org/10.50908/rdj.4.0_302 (2025).
59. Sharma, S. et al. The effects of water-deficit stress on *Cannabis sativa* L. development and production of secondary metabolites: a review. *Horticulturae* **11**, 646; <https://doi.org/10.3390/horticulturae11060646> (2025).
60. Gill, S. S. & Tuteja, N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* **48**, 909–930 (2010).
61. Deeba, F. et al. Physiological and proteomic responses of cotton (*Gossypium herbaceum* L.) to drought stress. *Plant Physiol. Biochem.* **53**, 6–18; <https://doi.org/10.1016/j.plaphy.2012.01.002> (2012).
62. Wallace, J. G. et al. Genome-wide association for plant height and flowering time across 15 tropical maize populations under managed drought stress and well-watered conditions in Sub-Saharan Africa. *Crop Sci.* **56**, 2365–2378; <https://doi.org/10.2135/cropsci2015.10.0632> (2016).
63. Gao, S. et al. Effects of drought stress on growth, physiology and secondary metabolites of two *Adonis* species in Northeast China. *Sci. Hortic.* **259**, 108795; <https://doi.org/10.1016/j.scienta.2019.108795> (2020).
64. Meftahizade, H. et al. Effect of sowing date and irrigation regimes on yield components, protein and galactomannan content of guar (*Cyamopsis tetragonoloba* L.) in Iran climate. *Aust. J. Crop Sci.* **11**, 1481–1487 (2017).
65. Hussain, T. et al. Impacts of drought stress on water use efficiency and grain productivity of rice and utilization of genotypic variability to combat climate change. *Agronomy* **12**, 2518; <https://doi.org/10.3390/agronomy12102518> (2022).
66. Quiloango-Chimarro, C. A. et al. Physiology, yield, and water use efficiency of drip-irrigated upland rice cultivars subjected to water stress at and after flowering. *Exp. Agric.* **58**, e19; <https://doi.org/10.1017/S0014479722000205> (2022).
67. Tippayawat, A. et al. Timing and duration of drought differentially affect growth and yield components among sugarcane genotypes. *Plants* **14**, 796; <https://doi.org/10.3390/plants14050796> (2025).
68. Drobnitch, S. T. et al. Can nitrogen availability impact plant performance under water stress? A review of traits, mechanisms, and whole plant effects. *Plant Soil* **511**, 45–67; <https://doi.org/10.1002/ldr.5667> (2025).
69. Bitire, T. D. et al. Effect of *Bradyrhizobium japonicum* strains and inorganic nitrogen fertilizer on the growth and yield of bambara groundnut (*Vigna subterranea* (L.) Verdc) accessions. *Front. Sustain. Food Syst.* **6**, 913239; <https://doi.org/10.3389/fsufs.2022.913239> (2022).
70. Galindo, F. S. et al. Co-inoculation with *Azospirillum brasilense* and *Bradyrhizobium* sp. enhances nitrogen uptake and yield in field-grown cowpea and did not change N-fertilizer recovery. *Plants* **11**, 1847; <https://doi.org/10.3390/plants11141847> (2022).
71. Lancashire, P. D. et al. A uniform decimal code for growth stages of crops and weeds. *Ann. Appl. Biol.* **119**, 561–601; <https://doi.org/10.1111/j.1744-7348.1991.tb04895.x> (1991).
72. Arnon, D. I. Copper enzymes in isolated chloroplasts. Polyphenol-oxidase in *Beta vulgaris*. *Plant Physiol.* **24**, 1–15; <https://doi.org/10.1104/pp.24.1.1> (1949).

73. Lichtenthaler, H. K. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* **148**, 350–382; [https://doi.org/10.1016/0076-6879\(87\)48036-1](https://doi.org/10.1016/0076-6879(87)48036-1) (1987).
74. Bates, L., Waldren, R. P. & Teare, I. D. Rapid determination of free proline for water-stress studies. *Plant Soil* **39**, 205–207; <https://doi.org/10.1007/BF00018060> (1973).
75. Yemm, E. W. & Willis, A. The estimation of carbohydrates in plant extracts by anthrone. *Biochem. J.* **57**, 508–514; <https://doi.org/10.1042/bj0570508> (1954).
76. Bailly, C. et al. Catalase activity and expression in developing sunflower seeds as related to drying. *J. Exp. Bot.* **55**, 475–483; <https://doi.org/10.1093/jxb/erh050> (2004).
77. Giannopolitis, C. N. & Ries, S. K. Superoxide dismutases: I. Occurrence in higher plants. *Plant Physiol.* **59**, 309–314 (1977).
78. Jones, D. B. Factors for converting percentages of nitrogen in foods and feeds into percentages of proteins. United States Department of Agriculture, Washington (1931).
79. Thiex, N., Novotny, L. & Crawford, A. Determination of ash in animal feed: AOAC official method 942.05 revisited. *J. AOAC Int.* **95**, 1392–1397; <https://doi.org/10.5740/jaoacint.12-129> (2012).
80. Van Soest, P. V., Robertson, J. B. & Lewis, B. A. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **74**, 3583–3597; [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2) (1991).
81. Oddy, V. H. In Robards, G. E. & Packham, R. G. (eds.) Feed Information and Animal Production 395–398 (Commonwealth Agricultural Bureaux, Farnham Royal, UK, 1983).
82. Standing Committee on Agriculture. Feeding standards for Australian livestock (Commonwealth Scientific and Industrial Research Organisation (CSIRO), East Melbourne, Victoria, Australia, 1990).
83. Williams, Y. J. et al. Variation in residual feed intake in Holstein-Friesian dairy heifers in southern Australia. *J. Dairy Sci.* **94**, 4715–4725; <https://doi.org/10.3168/jds.2010-4015> (2011).
84. Chieb, M. & Gachomo, E. W. The role of plant growth promoting rhizobacteria in plant drought stress responses. *BMC Plant Biol.* **23**, 407; <https://doi.org/10.1186/s12870-023-04403-8> (2023).
85. Kopecká, R. et al. Abiotic stress in crop production. *Int. J. Mol. Sci.* **24**, 6603; <https://doi.org/10.3390/ijms24076603> (2023).
86. Kazan, K. & Lyons, R. The link between flowering time and stress tolerance. *J. Exp. Bot.* **67**, 47–60; <https://doi.org/10.1093/jxb/erv441> (2016).
87. Seleiman, M. F. et al. Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants* **10**, 259; <https://doi.org/10.3390/plants10020259> (2021).
88. Shavrukov, Y. et al. Early flowering as a drought escape mechanism in plants: how can it aid wheat production? *Front. Plant Sci.* **8**, 1950; <https://doi.org/10.3389/fpls.2017.01950> (2017).
89. Pour-Aboughadareh, A. et al. Effects of drought stress on some agronomic and morpho-physiological traits in durum wheat genotypes. *Sustainability* **12**, 5610; <https://doi.org/10.3390/su12145610> (2020).
90. Wach, D. & Skowron, P. An overview of plant responses to the drought stress at morphological, physiological and biochemical levels. *Pol. J. Agron.* **50**, 25–34; <https://doi.org/10.26114/pja.iung.435.2022.04> (2022).
91. He, M. & Ding, N. Z. Plant unsaturated fatty acids: multiple roles in stress response. *Front. Plant Sci.* **11**, 562785; <https://doi.org/10.3389/fpls.2020.562785> (2020).
92. Wang, X. et al. Response of rice with overlapping growth stages to water stress by assimilates accumulation and transport and starch synthesis of superior and inferior grains. *Int. J. Mol. Sci.* **23**, 11157; <https://doi.org/10.3390/ijms231911157> (2022).

93. Li, Y. et al. Securing maize reproductive success under drought stress by harnessing CO₂ fertilization for greater productivity. *Front. Plant Sci.* **14**, 1221095; <https://doi.org/10.3389/fpls.2023.1221095> (2023).
94. Stefaniak, E. Z., Tissue, D. T., Dewar, R. C. & Medlyn, B. E. Optimal carbon storage during drought. *Tree Physiol.* **44**, 34-45; <https://doi.org/10.1093/treephys/tpae032> (2024).
95. Upreti, P. et al. Physiological attributes associated with leaf spectral alterations in guar [*Cyamopsis tetragonoloba* (L.) Taub.] under drought. *3 Biotech* **11**, 125; <https://doi.org/10.1007/s13205-021-02674-2> (2021).
96. Reddy, B. M. et al. De novo transcriptome analysis of drought-adapted cluster bean (Cultivar RGC-1025) reveals the wax regulatory genes involved in drought resistance. *Front. Plant Sci.* **13**, 868142; <https://doi.org/10.3389/fpls.2022.868142> (2022).
97. Khurizadeh, S. et al. Foliar application of methyl jasmonate and chitosan improve growth, yield, and quality of guar (*Cyamopsis tetragonoloba* L.) under water-deficit stress. *Plants* **13**, 3099; <https://doi.org/10.3390/plants13213099> (2024).
98. Sulaman, S. et al. Exogenous application of nitrogen (N) and potassium (K) improves drought tolerance in plants: a review. *J. Soil Sci. Plant Nutr.* 1-16; <https://doi.org/10.1007/s42729-025-02433-9> (2025).
99. Zayed, O. et al. Nitrogen journey in plants: from uptake to metabolism, stress response, and microbe interaction. *Biomolecules* **13**, 1443; <https://doi.org/10.3390/biom13101443> (2023).
100. Guo, P. et al. Optimized nitrogen application ameliorates the photosynthetic performance and yield potential in peanuts as revealed by OJIP chlorophyll fluorescence kinetics. *BMC Plant Biol.* **24**, 774; <https://doi.org/10.1186/s12870-024-05482-x> (2024).
101. Liu, J. et al. Effect of nitrogen, phosphorus and potassium fertilization management on soil properties and leaf traits and yield of *Sapindus mukorossi*. *Front. Plant Sci.* **15**, 1300683; <https://doi.org/10.3389/fpls.2024.1300683> (2024).
102. Ye, T. et al. Interaction effects of irrigation and nitrogen on the coordination between crop water productivity and nitrogen use efficiency in wheat production on the North China Plain. *Agric. Water Manage.* **271**, 107787; <https://doi.org/10.1016/j.agwat.2022.107787> (2022).
103. Abdelrhman, A. A. et al. Optimizing drip irrigation and nitrogen fertilization for sustainable wheat production in arid soils: Water-nitrogen use efficiency. *Water* **17**, 2708; <https://doi.org/10.3390/w17182708> (2025).
104. Egamberdieva, D. et al. Interactive effects of nutrients and Bradyrhizobium japonicum on the growth and root architecture of soybean (*Glycine max* L.). *Front. Microbiol.* **9**, 1000; <https://doi.org/10.3389/fmicb.2018.01000> (2018).
105. Jabborova, D. et al. Co-inoculation of rhizobacteria promotes growth, yield, and nutrient contents in soybean and improves soil enzymes and nutrients under drought conditions. *Sci. Rep.* **11**, 22081; <https://doi.org/10.1038/s41598-021-01337-9> (2021).
106. Meel, S. & Saharan, B. S. Enhancing crop resilience towards drought by integrating nanotechnology, microbiomes, and growth-promoting rhizobacteria. *Discov. Agric.* **2**, 112; <https://doi.org/10.1007/s44279-024-00131-1> (2024).
107. Rani, S. et al. Harnessing plant-bacterial interactions to enhance abiotic stress tolerance in plants: a review. *Discov. Plants* **2**, 250; <https://doi.org/10.1007/s44372-025-00330-0> (2025).
108. Timofeeva, A. M., Galyamova, M. R. & Sedykh, S. E. How do plant growth-promoting bacteria use plant hormones to regulate stress reactions? *Plants* **13**, 2371; <https://doi.org/10.3390/plants13172371> (2024).
109. Gill, S. S. & Tuteja, N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* **48**, 909-930 (2010).
110. Zhou, Y. et al. Edible plant oil: global status, health issues, and perspectives. *Front. Plant Sci.* **11**, 1315; <https://doi.org/10.3389/fpls.2020.01315> (2020).

111. Patel, T. B. B. et al. Myo-inositol reduces drought-induced oxidative damage and improves mungbean photosynthetic efficiency and antioxidant machinery. *J. Plant Growth Regul.*; <https://doi.org/10.1007/s00344-025-11819-z> (2025).
112. Hussain, T. et al. Impacts of drought stress on water use efficiency and grain productivity of rice and utilization of genotypic variability to combat climate change. *Agronomy* **12**, 2518; <https://doi.org/10.3390/agronomy12102518> (2022).
113. Du, Y. et al. Effect of drought stress during soybean R2-R6 growth stages on sucrose metabolism in leaf and seed. *Int. J. Mol. Sci.* **21**, 618; <https://doi.org/10.3390/ijms21020618> (2020).
114. Kaur, G. et al. Role of osmolytes dynamics in plant metabolism to cope with salinity induced osmotic stress. *Discov. Agric.* **2**, 59; <https://doi.org/10.1007/s44279-024-00070-x> (2024).
115. Sheoran, A. R. et al. PGPR-mediated drought tolerance in Brassica juncea by modulating morpho-physiological, biochemical, and antioxidant responses. *J. Plant Growth Regul.*; <https://doi.org/10.1007/s00344-025-11887-1> (2025).
116. Fujita, M. & Hasanuzzaman, M. Approaches to enhancing antioxidant defense in plants. *Antioxidants* **11**, 925; <https://doi.org/10.3390/antiox11050925> (2022).
117. Atieno, M. & Lesueur, D. Opportunities for improved legume inoculants: Enhanced stress tolerance of rhizobia and benefits to agroecosystems. *Symbiosis* **77**, 191-205; <https://doi.org/10.1007/s13199-018-0585-9> (2019).
118. Mushtaq, Z. et al. Inoculation of Rhizobium alleviates salinity stress through modulation of growth characteristics, physiological and biochemical attributes, stomatal activities and antioxidant defence in Cicer arietinum L. *J. Plant Growth Regul.* **40**, 2148-2163; <https://doi.org/10.1007/s00344-020-10267-1> (2021).
119. Sun, X. et al. Research on the physiological mechanisms of nitrogen in alleviating plant drought tolerance. *Plants* **14**, 2928; <https://doi.org/10.3390/plants14182928> (2025).
120. Gulcin, İ. Antioxidants: a comprehensive review. *Arch Toxicol* **99**, 1893-1997. <https://doi.org/10.1007/s00204-025-03997-2> (2025).
121. Sood, M. Reactive oxygen species (ROS): plant perspectives on oxidative signalling and biotic stress response. *Discov. Plants* **2**, 187; <https://doi.org/10.1007/s44372-025-00275-4> (2025).
122. Jahanzad, E. et al. Response of a new and a commonly grown forage sorghum cultivar to limited irrigation and planting density. *Agric. Water Manag.* **117**, 62-69; <https://doi.org/10.1016/j.agwat.2012.11.001> (2013).
123. Bista, D. R. et al. Effects of drought on nutrient uptake and the levels of nutrient-uptake proteins in roots of drought-sensitive and tolerant grasses. *Plants* **7**, 28; <https://doi.org/10.3390/plants7020028> (2018).
124. Smit, M. et al. Drought impact on the nutrients of forage plants in a semi-arid rangeland and its potential implications for sustaining herbivores. *J. Arid Environ.* **227**, 105299; <https://doi.org/10.1016/j.jaridenv.2024.105299> (2025).
125. Varol, I. S. et al. Water and nitrogen impacts on water use, forage yield and quality of annual ryegrass. *Sci. Rep.* **15**, 34143; <https://doi.org/10.1038/s41598-025-08717-5> (2025).
126. Jovino, R. S. et al. Elite Bradyrhizobium strains boost biological nitrogen fixation and peanut yield in tropical drylands. *Braz. J. Microbiol.* **53**, 1623-1632; <https://doi.org/10.1007/s42770-022-00792-4> (2022).
127. Ramya, S. et al. Improvement in the quality attributes of forage cowpea by use of liquid microbial inoculants. *Indian J. Anim. Res.* **56**, 959-965; <https://doi.org/10.18805/IJAR.B-4503> (2022).
128. Kyei-Boahen, S. et al. Symbiotic effectiveness of Bradyrhizobium strains on soybean growth and productivity in Northern Mozambique. *Front. Sustain. Food Syst.* **6**, 1084745; <https://doi.org/10.3389/fsufs.2022.1084745> (2023).

129. Ajayi, O. O. et al. Native symbiotic rhizobia strains improve yield and nitrogen fixation in underutilized Bambara groundnut (*Vigna subterranea* L. Verdc.). *Agrosyst. Geosci. Environ.* **8**, e70211; <https://doi.org/10.1002/agg2.70211> (2025).
130. Farooq, M., Hussain, M., Wahid, A. & Siddique, K. H. M. Drought stress in plants: an overview. *Agron. Sustain. Dev.* **32**, 1–33 (2012).
131. Ullah, A. & Farooq, M. The challenge of drought stress for grain legumes and options for improvement. *Arch. Agron. Soil Sci.* **68**, 1601–1618; <https://doi.org/10.1080/03650340.2021.1906413> (2022).
132. Wang, Z. et al. From phenotyping to genetic mapping: identifying water-stress adaptations in legume root traits. *BMC Plant Biol.* **24**, 749; <https://doi.org/10.1186/s12870-024-05477-8> (2024).
133. Becker, J. N. et al. Effects of cowpea (*Vigna unguiculata*) inoculation on nodule development and rhizosphere carbon and nitrogen content under simulated drought. *Plant Soil* **500**, 33–51; <https://doi.org/10.1007/s11104-023-06051-1> (2024).
134. Pathak, A. et al. Mechanistic insights into plant growth promoting rhizobacteria with focus on root soil interactions, functional attributes and agricultural sustainability. *Discov. Biotechnol.* **3**, 1; <https://doi.org/10.1007/s44340-025-00046-7> (2026).
135. Abd El-Mageed, T. A. et al. Plant growth-promoting rhizobacteria improve growth, morph-physiological responses, water productivity, and yield of rice plants under full and deficit drip irrigation. *Rice* **15**, 16; <https://doi.org/10.1186/s12284-022-00564-6> (2022).
136. Demir, H. et al. Ameliorative effects of biofertilizers on yield, water use efficiency and quality of bean under drought stress conditions. *BMC Plant Biol.* **25**, 1512; <https://doi.org/10.1186/s12870-025-07339-3> (2025).
137. Liu, K. et al. Plant growth-promoting rhizobacteria improve drought tolerance of crops: a review. *Plant Growth Regul.* **105**, 567–581; <https://doi.org/10.1007/s10725-025-01300-y> (2025).
138. Allito, B. B. et al. Legume-rhizobium specificity effect on nodulation, biomass production and partitioning of faba bean (*Vicia faba* L.). *Sci. Rep.* **11**, 3678; <https://doi.org/10.1038/s41598-021-83235-8> (2021).
139. Iqbal, N. et al. Mesorhizobia strain and chickpea variety drive phenotypic plasticity of plant growth and nodulation. *Plant Growth Regul.* **104**, 475–485; <https://doi.org/10.1007/s10725-024-01177-3> (2024).
140. Rockström, J. et al. Sustainable intensification of agriculture for human prosperity and global sustainability. *Ambio* **46**, 4–17; <https://doi.org/10.1007/s13280-016-0793-6> (2017).
141. Gao, Y. et al. Regulation effects of water and nitrogen on yield, water, and nitrogen use efficiency of wolfberry. *J. Arid Land* **16**, 29–45; <https://doi.org/10.1007/s40333-024-0003-5> (2024).

Acknowledgments

The authors thank Dr. Hamidreza Ramazani (University of Birjand, Iran) for providing guar seeds and confirming the taxonomic identification of plant material. The authors also acknowledge Green Biotech Company (Qom Province, Iran) for supplying *Bradyrhizobium* strains RS-150 and RS-153 used in this study.

Author contributions

G.A.A. and S.N., as corresponding authors, jointly designed the experimental treatments, conducted the experiments, wrote the main manuscript, and prepared all figures and tables. I.A., E.S., and M.A.N., as co-authors, also analyzed the data and contributed to the implementation of the experiments, development of the experimental design, and revised the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Declaration of compliance

We confirm and ensure that the study follows all relevant guidelines and regulations and complies with the IUCN Policy Statement on Research Involving Species at Risk of Extinction and the Convention on the Trade in Endangered Species of Wild Fauna and Flora

Additional information

Correspondence and requests for materials should be addressed to G.A.A.

Funding Declaration

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

ARTICLE IN PRESS