


RESEARCH ARTICLE

The influence of vermicomposting on photosynthetic activity and productivity of maize (*Zea mays* L.) crop under semi-arid climate

Muhammad Younas^{1,2}, Huasong Zou³, Tasmia Laraib⁴, Waseem Abbas⁵, Muhammad Waqar Akhtar⁶, Muhammad Naveed Aslam⁷, Luqman Amrao², Shoukat Hayat⁸, Tariq Abdul Hamid⁹, Akhtar Hameed², Ghalib Ayaz Kachelo², Mona M. Elseehy¹⁰, Ahmed M. El-Shehawi¹¹, Ali Tan Kee Zuan^{12*}, Yunzhou Li^{1*}, Muhammad Arif^{13*}

1 Department of Plant Pathology, College of Agriculture, Guizhou University, Guiyang, Guizhou, China, **2** Department of Plant Pathology, University of Agriculture Faisalabad, Punjab, Pakistan, **3** College of Plant Protection, Fujian Agriculture and Forestry University, Fujian, Fuzhou, China, **4** Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Punjab, Pakistan, **5** Ayub Agricultural Research Institute, Faisalabad, Punjab, Pakistan, **6** Department of Soil Science, The Islamia University of Bahawalpur, Bahawalpur, Punjab, Pakistan, **7** Department of Plant Pathology, The Islamia University of Bahawalpur, Bahawalpur, Punjab, Pakistan, **8** Department of Forestry, Department of Forestry, The Islamia University of Bahawalpur, Bahawalpur, Punjab, Pakistan, **9** Assistant Director, office of Deputy Director of Agriculture, Pest Warning & Quality Control of Pesticides (PW&QCP) Lahore, Lahore, Pakistan, **10** Department of Genetics, Faculty of Agriculture, University of Alexandria, Alexandria, Egypt, **11** Department of Biotechnology, College of Science, Taif University, Taif, Saudi Arabia, **12** Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Selangor, Malaysia, **13** Department of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

* tkz@upm.edu.my (ATKZ); liyunzhou2007@126.com (YL); arif_1821uaf@yahoo.com (MA)



OPEN ACCESS

Citation: Younas M, Zou H, Laraib T, Abbas W, Akhtar MW, Aslam MN, et al. (2021) The influence of vermicomposting on photosynthetic activity and productivity of maize (*Zea mays* L.) crop under semi-arid climate. PLoS ONE 16(8): e0256450. <https://doi.org/10.1371/journal.pone.0256450>

Editor: Sagheer Atta, Ghazi University Dera Ghazi Khan, PAKISTAN

Received: June 8, 2021

Accepted: August 9, 2021

Published: August 25, 2021

Copyright: © 2021 Younas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: This work was supported by projects of the national Nature Science Foundation (No. 32060679) and projects of Guizhou University (No. GuidapeiYU[2019]52 and No. [2017]50). The current work was funded by Taif University Researchers Supporting Project number (TURSP-2020/75), Taif University, Taif, Saudi Arabia. There were no additional external funding involved in the study. The funders had no role in study design,

Abstract

Food production and waste recycling are the two major issues faced globally with rapidly increasing population. Recycling organic wastes to crop amendments could be a possible solution to these issues. Earthworms transfer organic waste to compost, which is used to grow crops and increase crop productivity. This study assessed the impact of vermicompost produced from the residues of six desert plant species, i.e., (*Ziziphus mauritiana*, *Aerva javanica*, *Calligonum comosum*, *Sacchrum benghalens*, *Calligonum polygonoides* and *Prosopis cineraria*) combined with farmyard manure (5 t ha⁻¹) on growth, yield and photosynthetic activity of maize crop. Earthworm species *Eisenia fetida* (Savigny, 1826) was used to prepare vermicomposting of all tested plant species. The desert species were collected from natural habitats, chopped, dried, mixed with FYM and then earthworms were released to prepare the vermicompost. The earthworms were excluded twenty days after release and resultant was considered as compost and used in the experiment. Results revealed that application of *P. cineraria* vermicompost resulted in the highest plant height (75.33 cm), stem diameter (22.66 mm), cob length (17.66 cm), number of grains/cob (374.67), 1000-grain weight (260.41 g) and grains yield (3.20 t/ha). Application of *P. cineraria* vermicompost resulted in the highest uptake of macronutrients, i.e., N (91.01%), P (22.07%), K (80.41%), micronutrients, i.e., Fe (19.07 ppm), Zn (40.05 ppm), and phenolic contents (150). Application of *P. cineraria* vermicompost also resulted in the highest quantum photosynthetic yield

data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

(0.42 mole C/mole of photon), chlorophyll fluorescence (355.18 moles of photon $\text{m}^{-2}\text{s}^{-1}$) and electron transport rate (310.18 micro mole $\text{m}^{-2}\text{s}^{-1}$). It is concluded that vermicomposting has the potential to improve growth and yield of maize crop. Particularly, application of vermicompost obtained from *P. cineraria* can be used to improve the growth and yield of maize crop. Nonetheless, field trials are necessary for a wide scale recommendation.

Introduction

Conventional agriculture is reliant on excessive application of pesticides, herbicides, chemical fertilizers and low use of organic fertilizers [1, 2]. Frequent and non-judicious application of chemicals exerts various negative effects on environment [3], causes food pollution [4, 5], decreases soil quality and results in the loss of agricultural biodiversity [6–8]. Sustainable agricultural practices relying on natural or organic sources are needed to overcome these hazards.

Organic farming is the widely accepted option for maintaining the environmental sustainability and biodiversity [9, 10]. Organic farming is being adopted by the environment-sensitive populations in developed countries due to its enormous ecological and economic benefits [11, 12]. Previous studies revealed that vermicomposting improves nutrient status as well as biological characters of soil [13]. It is highly beneficial as it allows good drainage, improves water storage capacity and soil ventilation [14].

The production of livestock manure has been considerably increased due to rapidly increasing industrialization and human population [15, 16]. However, it caused numerous environmental drawbacks including contaminated ground water and offensive odors [17, 18]. The conversion of waste to valuable materials may could play a significant role in environmental cleaning and resource recycling [19]. Various microbiological, physical and chemical techniques are used for the disposal of livestock manure. Unfortunately, all these strategies are labor intensive and time-consuming [20–23]. Traditional composting is cheap and unadorned; however, it causes nutrient losses [24, 25].

Vermicomposting is a viable and cost effective technique for the management of livestock manure. Earthworms' activity initiates the decomposition of organic matter and prepares nutrients for plant growth [26, 27]. Thus, composting of livestock manure is helpful in reducing environmental pollution and also provides a healthier substitute of chemical fertilizers [28]. Studies relating to photosynthesis in maize crop are important for the selection of superior genotypes and improve yield under abiotic stresses [29, 30]. Chlorophyll fluorescence and electron transport rate directly influence plant vegetative growth, which ultimately leads to higher yield [31].

Although several studies have reported the benefits of vermicomposting, rare studies have combined plant residues and FYM to produce a nutrient rich compost. Therefore, this study produced compost of FYM combined with plant residues and tested their impacts on growth, photosynthesis and yield of maize crop. The major aim of the study was to select the most suitable desert species for vermicomposting. Investigate the impacts of applied vermicompost on the yield of maize crop was the second objective of the study.

Materials and methods

Experimental site

The experimental site is located at Students Research Farm, University of Agriculture Faisalabad, Pakistan (31.4278° N, 73.0758° E). The climate of the study area is semi-arid. The average

temperature is 24.8°C, while annual rainfall is 526 mm. Soil type is silt-loam to sandy-loam. Physio-chemical properties of soil were determined before the initiation of experiment and are given in Table 1.

Experimental design

Six desert plant species, i.e., *Ziziphus mauritiana*, *Aerva javanica*, *Calligonum comosum*, *Saccharum benghalens*, *Calligonum polygonoides* and *Prosopis cineraria* were collected from the surroundings of Faisalabad and Bhakkar, Pakistan. Similarly, farmyard manure (FYM) was collected from Student Research Farm, University of Agriculture, Faisalabad, Pakistan. Collected materials were dried and crushed at biogas plant located at Research farm, University of Agriculture, Faisalabad, Pakistan. After processing, the material was mixed with FYM and soil. Twenty individuals of earthworm [*Eisenia fetida* (Savigny, 1826)] were added into each pot and kept wet for twenty days [32]. The earthworms were discarded by following the sieve method and vermicompost was transferred to pots (Table 2). Maize plants were grown in the pots. Experiment was laid out by following the complete randomized design. The treatments used in the experiment are given in Table 2.

Photosynthetic yield analyzer

Photosynthetic yield analyzer was used to measure the photosynthetic efficiency of photosystem II and physiological parameters of photosystem II, including, quantum photosynthetic yield (mol C/mole of photon), chlorophyll fluorescence (moles of photons m^{-2}/s) and electron transport rate ($\mu mol m^{-2} s^{-1}$). The Kjeldahl and Champ and Marker method [33] was used for determining the nitrogen, phosphorus and potassium. This Kjeldahl method is mainly divided in three steps, i.e., digestion, distillation and titration. Homogenous samples were boiled in concentrated sulfuric acid, which lead to the formation of ammonium sulfate solution. Excess base was added for converting NH_4 to NH_3 and then amount of nitrogen was calculated from the ammonia [34].

Determination of micronutrients

Atomic absorption spectrophotometer (Hitachi Polarized Zeeman AAS, Z-8200, Japan) was used for determining the minerals, i.e., iron (Fe) and zinc (Zn) following the conditions described in AOAC. The instrumental operating conditions are summarized in Table 3.

Standards' preparation

Calibrated standards were prepared from the commercially available stock solution (Appli-Chem[®]) in the form of aqueous solution (1000 ppm). Highly purified de-ionized water was

Table 1. Physio-chemical parameters of experimental soil.

Soil properties	Values
Total nitrogen (%)	0.038
Organic matter (%)	0.77
pH	8.4
Electric conductivity ($mS\ cm^{-1}$)	1.35
Available potassium (%)	160
Available phosphorus (%)	7.2
Textural class	Loam

<https://doi.org/10.1371/journal.pone.0256450.t001>

Table 2. Experimental treatments used in the study, their abbreviations and explanations.

Abbreviation	Treatments
T ₀	Absolute control*
T ₁	Vermicompost of <i>Ziziphus mauritiana</i> + FYM* *
T ₂	Vermicompost of <i>Aerva javanica</i> + FYM
T ₃	Vermicompost of <i>Calligonum comosum</i> + FYM
T ₄	Vermicompost of <i>Sacchrum benghalens</i> + FYM
T ₅	Vermicompost of <i>Calligonum polygonoides</i> + FYM
T ₆	Vermicompost of <i>Prosopis cineraria</i> + FYM

*The plants were grown on simple soil,

**Five tons per hectare farmyard manure (FYM) was added

<https://doi.org/10.1371/journal.pone.0256450.t002>

used for the preparation of working standards. All the glass apparatus used throughout the process of analytical work were immersed in 8N HNO₃ overnight and washed with several changes of de-ionized water prior to use.

Statistical analysis

The data were recorded using the standard procedures and statistically analyzed at 5% level of significance using Fisher's analysis of variance (ANOVA) technique. One-way ANOVA was used to test the significance among treatments and data were normalized prior to execute ANOVA. The treatment means were compared using least significant different test as a post-hoc test where ANOVA indicated significant differences [35].

Results and discussion

Growth and yield attributes

Growth parameters such as stem diameter and plant height were significantly influenced by various vermicompost treatments. The vermicompost of *P. cineraria* proved the most effective improving growth and yield attributes maize crop. The highest values of stem diameter, plant height, cob length, number of grains per cob, 1000-grain weight and grain yield were recorded with the application of *P. cineraria* vermicompost, whereas control treatment recorded the lowest values of these traits (Table 4).

Application vermicomposting enhances nutrient mobility; thus, plant uptakes more nutrient, which improves growth and yield attributes. Control plants resulted in the lowest stem

Table 3. Operational conditions used in atomic absorption spectrophotometer for the determination of iron (Fe) and zinc (Zn).

Parameters	Set value	
	Fe	Zn
Wavelength (nm)	248.3	213.9
Slit Width (nm)	0.2	1.3
Lamp Current (mA)	10.0	10.0
Burner Head	Standard type	Standard type
Flame	Air-C ₂ H ₂	Air-C ₂ H ₂
Burner Height (mm)	7.5	7.5
Oxidant gas pressure (Flow rate) (kpa)	160	160
Fuel gas pressure (Flow rate) (kpa)	6	6

<https://doi.org/10.1371/journal.pone.0256450.t003>

Table 4. The impact of different vermicompost treatments on growth and yield attributes of maize crop.

Vermicompost source	Stem diameter (mm)	Plant height (cm)	Cob length (cm)	Number of grains/cob	1000-grain weight (g)	Grain yield t/ha
Absolute control	14.83 e	49.55 g	7.08	129.67 g	180.34 g	1.60 e
<i>Ziziphus mauritiana</i>	16.16 d	54.00 f	11.00 e	289.67 f	225.43 f	2.74 d
<i>Aerva javanica</i>	18.00 c	59.66 e	12.91 d	328.67 e	230.36 e	3.10 c
<i>Calligonum comosum</i>	20.08 b	65.00 d	15.16 c	340.00 d	241.30 d	3.09 c
<i>Sacchrum benghalens</i>	20.83 b	70.0 c	15.75 b	355.00 c	250.42 c	3.10 c
<i>Calligonum polygonoides</i>	21.03 b	72.57 b	16.08 b	360.00 b	255.48 b	3.15 b
<i>Prosopis cineraria</i>	22.66 a	75.33 a	17.66 a	374.67 a	260.41 a	3.20 a
LSD (0.05)	1.08	1.4	0.84	4.39	3.66	0.04

Means sharing the same letters within a column are statistically non-significant

<https://doi.org/10.1371/journal.pone.0256450.t004>

diameter and plant height, which can be explained as a direct effect of nutrient availability. Results of this study are in line with several earlier work of [36–38].

The yield attributes, mainly yield and is the result of vegetative development of plant. All yield attributes were significantly influenced by vermicomposting of different desert species. These results have significantly attributed the importance of vermicomposting with different earthworm activities to increase the yield of several crops and improve defense mechanism against several pathogens and insects. Higher growth characters quantify higher yield attributes, particularly stem diameter and plant height, which ultimately lead to higher yield. Results of present study are supported by earlier studies [39, 40]. Improvement in cob length and number of grains/cob significantly improved yield in earlier studies [41–43].

Nutrient uptake

Average yield potential is the result of co-ordinate interplay of different growth characters and nutrient uptake. Grain and stover yield is increased by increased nutrient uptake. The highest nutrient (N, P, K) uptake was observed with the application of *P. cineraria* vermicompost, whereas control treatment resulted in the lowest nutrient uptake (Table 5). Micronutrients and phenolic contents also increased with the application of vermicompost obtained from different plant species and *P. cineraria* (resulted in the highest values of micronutrients and phenolic contents (Table 5). Results of our study are in accordance with the findings of earlier studies [44–47]. The organic matter, i.e., FYM mainly supplies beneficial nutrients to crop and enhances its productivity significantly [48].

Table 5. The impact of different vermicompost treatments on the uptake of macro and micronutrients.

Treatment combinations	N (%)	P (%)	K (%)	Micro-nutrients		
				Iron (ppm)	Zinc (ppm)	Phenolic contents
Absolute control	29.71 g	8.85 g	24.79 g	0.00 f	0.00 g	0.00 g
<i>Ziziphus mauritiana</i>	70.05 f	13.16 f	60.25 f	10.19 e	23.16 f	109.93 f
<i>Aerva javanica</i>	78.05 e	16.18 e	63.79 e	11.80 d	25.05 e	114.77 e
<i>Calligonum comosum</i>	80.71 d	17.70 d	65.06 d	15.04 c	30.05 d	120.05 d
<i>Sacchrum benghalens</i>	83.00 c	19.03 c	70.25 c	16.30 b	31.07 c	125.29 c
<i>Calligonum polygonoides</i>	86.96 b	20.31 b	75.05 b	16.65 b	35.05 b	140.08 b
<i>Prosopis cineraria</i>	91.01 a	22.07 a	80.41 a	19.07 a	40.05 a	150.00 a
LSD (0.05)	1.16	0.89	1.21	0.90	0.82	1.12

Means sharing the same letters within a column are statistically non-significant

<https://doi.org/10.1371/journal.pone.0256450.t005>

Table 6. The impact of different vermicompost treatments on quantum photosynthetic yield, chlorophyll florescence and electron transport rate.

Treatments	Quantum Photosynthetic Yield (mole C/mole of photon)	Chlorophyll Florescence (moles of photon $\text{m}^{-2}\text{s}^{-1}$)	Electron Transport Rate.(micro mole $\text{m}^{-2}\text{s}^{-1}$)
Absolute control	0.28 g	290.00 g	185.13 g
<i>Ziziphus mauritiana</i>	0.31 f	315.17 f	220.80 f
<i>Aerva javanica</i>	0.32 e	330.45 e	231.44 e
<i>Calligonum comosum</i>	0.34 d	332.97 d	234.56 d
<i>Sacchrum benghalens</i>	0.36 c	343.75 c	270.61 c
<i>Calligonum polygonoides</i>	0.40 b	344.34 b	272.42 b
<i>Prosopis cineraria</i>	0.42 a	355.18 a	310.18 a
LSD _(0.05)	0.01	1.27	1.49

Means sharing the same letters within a column are statistically non-significant

<https://doi.org/10.1371/journal.pone.0256450.t006>

Physiological parameters of photosystem II

Physiological parameters of photosystem II, i.e., quantum photosynthetic yield (mol C/mole of photon) chlorophyll florescence (moles of photons m^{-2}/S) and electron transport rate ($\mu\text{molm}^{-2}\text{S}^{-1}$) were significantly altered by different vermicomposting treatments. The vermicompost of *P. cineraria* recorded the highest values of these traits, whereas the lowest values were recorded for control treatment (Table 6). Inappropriate function of photosynthetic apparatus may lead to less chlorophyll contents [49–51], and the application of vermicompost of *P. cineraria* and FYM improved photosynthetic parameters in the current study.

Closure of stomata reduces internal concentration of CO_2 and inhibits the functioning of Rubisco enzymes [52, 53], which mainly decrease the rate of photosynthesis. Numerous earth-worm species utilize decaying organic materials and convert material to nutrient-rich elements, which further enhances vegetative and reproductive growth of plants [54]. However, very few studies have reported the role of vermicomposting on physiological parameters of photosystem II. Results of present study are also in line with the findings of [53, 55] that vermicomposting significantly increased chlorophyll contents, net photosynthetic rate and electron transport rate.

Conclusion

It can be concluded that higher growth, nutrient uptake, quality and maximum yield potential can be achieved through combined application of vermicompost of *Prosopis cineraria* and FYM in maize. Results also proved that integrated nutrient management, i.e., application of vermicomposting strategies have multipurpose on the improvement of crop productivity in sustainable manners.

Supporting information

S1 Dataset.
(XLSX)

Author Contributions

Conceptualization: Tasmia Laraib, Muhammad Waqar Akhtar, Akhtar Hameed, Mona M. Elseehy, Ahmed M. El-Shehawi, Ali Tan Kee Zuan, Yunzhou Li, Muhammad Arif.

Formal analysis: Huasong Zou, Waseem Abbas, Muhammad Naveed Aslam.

Funding acquisition: Mona M. Elseehy, Ali Tan Kee Zuan.

Investigation: Shoukat Hayat, Yunzhou Li.

Methodology: Waseem Abbas, Shoukat Hayat, Ghalib Ayaz Kachelo.

Project administration: Luqman Amrao.

Resources: Muhammad Younas, Muhammad Waqar Akhtar, Muhammad Naveed Aslam, Ghalib Ayaz Kachelo.

Software: Muhammad Waqar Akhtar, Tariq Abdul Hamid.

Supervision: Muhammad Arif.

Validation: Huasong Zou, Muhammad Naveed Aslam.

Visualization: Luqman Amrao, Tariq Abdul Hamid.

Writing – original draft: Muhammad Younas.

Writing – review & editing: Tasmia Laraib, Akhtar Hameed, Mona M. Elseehy, Ahmed M. El-Shehawi, Ali Tan Kee Zuan, Yunzhou Li.

References

1. Durán-Lara EF, Valderrama A, Marican A (2020) Natural organic compounds for application in organic farming. *Agriculture* 10: 41.
2. Nawaz A, Sufyan M, Gogi MD, Javed MW (2019) Sustainable Management of Insect-Pests. *Innovations in Sustainable Agriculture*: Springer. pp. 287–335.
3. Gu B, Ju X, Chang SX, Ge Y, Chang J (2017) Nitrogen use efficiencies in Chinese agricultural systems and implications for food security and environmental protection. *Regional Environmental Change* 17: 1217–1227.
4. Wang F, Chen S, Wang Y, Zhang Y, Hu C, et al. (2018) Long-term nitrogen fertilization elevates the activity and abundance of nitrifying and denitrifying microbial communities in an upland soil: implications for nitrogen loss from intensive agricultural systems. *Frontiers in microbiology* 9: 2424. <https://doi.org/10.3389/fmicb.2018.02424> PMID: 30405543
5. Hou Y, Wei S, Ma W, Roelcke M, Nieder R, et al. (2018) Changes in nitrogen and phosphorus flows and losses in agricultural systems of three megacities of China, 1990–2014. *Resources, Conservation and Recycling* 139: 64–75.
6. Li W, Guo S, Liu H, Zhai L, Wang H, et al. (2018) Comprehensive environmental impacts of fertilizer application vary among different crops: Implications for the adjustment of agricultural structure aimed to reduce fertilizer use. *Agricultural Water Management* 210: 1–10.
7. Hou P, Liu Y, Liu W, Liu G, Xie R, et al. (2020) How to increase maize production without extra nitrogen input. *Resources, Conservation and Recycling* 160: 104913.
8. Gill HK, Garg H (2014) Pesticide: environmental impacts and management strategies. *Pesticides-toxic aspects* 8: 187.
9. Leite LF, Oliveira FC, Araújo AS, Galvão SR, Lemos JO, et al. (2010) Soil organic carbon and biological indicators in an Acrisol under tillage systems and organic management in north-eastern Brazil. *Soil Research* 48: 258–265.
10. Santos VB, Araújo AS, Leite LF, Nunes LA, Melo WJ (2012) Soil microbial biomass and organic matter fractions during transition from conventional to organic farming systems. *Geoderma* 170: 227–231.
11. Lobley M, Butler A, Reed M (2009) The contribution of organic farming to rural development: An exploration of the socio-economic linkages of organic and non-organic farms in England. *Land use policy* 26: 723–735.
12. Lobley M, Butler A, Winter M (2013) Local organic food for local people? Organic marketing strategies in England and Wales. *Regional Studies* 47: 216–228.
13. Pant A, Radovich T, Hue N, Arancon N (2011) Effects of vermicompost tea (aqueous extract) on pak choi yield, quality, and on soil biological properties. *Compost Science & Utilization* 19: 279–292.

14. Kim MJ, Shim CK, Kim YK, Hong SJ, Park JH, et al. (2015) Effect of aerated compost tea on the growth promotion of lettuce, soybean, and sweet corn in organic cultivation. *The plant pathology journal* 31: 259. <https://doi.org/10.5423/PPJ.OA.02.2015.0024> PMID: 26361474
15. Yadav P, Jaiswal DK, Sinha RK (2021) Climate change: Impact on agricultural production and sustainable mitigation. *Global Climate Change: Elsevier*. pp. 151–174.
16. Kour D, Kaur T, Devi R, Yadav A, Singh M, et al. (2021) Beneficial microbiomes for bioremediation of diverse contaminated environments for environmental sustainability: Present status and future challenges. *Environmental Science and Pollution Research*: 1–23. <https://doi.org/10.1007/s11356-020-11060-z> PMID: 33009614
17. Yadav A, Garg V (2011) Recycling of organic wastes by employing *Eisenia fetida*. *Bioresource technology* 102: 2874–2880. <https://doi.org/10.1016/j.biortech.2010.10.083> PMID: 21078553
18. Suthar S (2010) Recycling of agro-industrial sludge through vermitechnology. *Ecological Engineering* 36: 1028–1036.
19. Singh J, Kaur A, Vig A, Rup P (2010) Role of *Eisenia fetida* in rapid recycling of nutrients from bio sludge of beverage industry. *Ecotoxicology and Environmental Safety* 73: 430–435. <https://doi.org/10.1016/j.ecoenv.2009.08.019> PMID: 19945748
20. Garg V, Gupta R (2011) Optimization of cow dung spiked pre-consumer processing vegetable waste for vermicomposting using *Eisenia fetida*. *Ecotoxicology and environmental safety* 74: 19–24. <https://doi.org/10.1016/j.ecoenv.2010.09.015> PMID: 20951432
21. Vig AP, Singh J, Wani SH, Dhaliwal SS (2011) Vermicomposting of tannery sludge mixed with cattle dung into valuable manure using earthworm *Eisenia fetida* (Savigny). *Bioresource technology* 102: 7941–7945. <https://doi.org/10.1016/j.biortech.2011.05.056> PMID: 21676611
22. Garg P, Gupta A, Satya S (2006) Vermicomposting of different types of waste using *Eisenia foetida*: A comparative study. *Bioresource technology* 97: 391–395. <https://doi.org/10.1016/j.biortech.2005.03.009> PMID: 16168639
23. Wani K, Rao R (2013) Bioconversion of garden waste, kitchen waste and cow dung into value-added products using earthworm *Eisenia fetida*. *Saudi journal of biological sciences* 20: 149–154. <https://doi.org/10.1016/j.sjbs.2013.01.001> PMID: 23961230
24. Hagemann N, Kammann CI, Schmidt H-P, Kappler A, Behrens S (2017) Nitrate capture and slow release in biochar amended compost and soil. *PloS one* 12: e0171214. <https://doi.org/10.1371/journal.pone.0171214> PMID: 28199354
25. Qasim W, Moon BE, Okyere FG, Khan F, Nafees M, et al. (2019) Influence of aeration rate and reactor shape on the composting of poultry manure and sawdust. *Journal of the Air & Waste Management Association* 69: 633–645. <https://doi.org/10.1080/10962247.2019.1569570> PMID: 30640581
26. Obalum S, Chibuike G, Peth S, Ouyang Y (2017) Soil organic matter as sole indicator of soil degradation. *Environmental monitoring and assessment* 189: 176. <https://doi.org/10.1007/s10661-017-5881-y> PMID: 28324279
27. Deepthi MP, Nivethitha S, Saminathan K, Narendhirakannan RT, Karmegam N, et al. (2021) Effect of vermiwash prepared from livestock biowaste as vermiponics medium on the growth and biochemical indices of *Amaranthus viridis* L. *Environmental Technology & Innovation* 21: 101300.
28. Fatima S, Riaz M, Al-Wabel MI, Arif MS, Yasmeen T, et al. (2021) Higher biochar rate strongly reduced decomposition of soil organic matter to enhance C and N sequestration in nutrient-poor alkaline calcareous soil. *Journal of Soils and Sediments* 21: 148–162.
29. Prasanna BM, Cairns JE, Zaidi P, Beyene Y, Makumbi D, et al. (2021) Beat the stress: breeding for climate resilience in maize for the tropical rainfed environments. *Theoretical and Applied Genetics*: 1–24. <https://doi.org/10.1007/s00122-020-03709-7> PMID: 33136168
30. Wijewardana C, Henry WB, Reddy KR (2017) Evaluation of drought tolerant maize germplasm to induced drought stress. *Miss Acad Sci* 62: 316–329.
31. Meena RK, Reddy KS, Gautam R, Maddela S, Reddy AR, et al. (2021) Improved photosynthetic characteristics correlated with enhanced biomass in a heterotic F1 hybrid of maize (*Zea mays* L.). *Photosynthesis Research* 147: 253–267. <https://doi.org/10.1007/s11120-021-00822-6> PMID: 33555518
32. Yuvaraj A, Thangaraj R, Maheswaran R (2019) Decomposition of poultry litter through vermicomposting using earthworm *Drawida sulcata* and its effect on plant growth. *International Journal of Environmental Science and Technology* 16: 7241–7254.
33. Riboulet C, Fabre F, Dénoue D, Martinant J, Lefevre B, et al. (2008) QTL mapping and candidate gene research for lignin content and cell wall digestibility in a top-cross of a flint maize recombinant inbred line progeny harvested at silage stage. *Maydica* 53: 1–9.
34. Dalton RL (2014) Effects of agrochemicals on riparian and aquatic primary producers in an agricultural watershed: Université d'Ottawa/University of Ottawa.

35. Steel RG, Torrie JH, Dickey DA (1997) Principles and procedures of statistics: A biological approach: McGraw-Hill.
36. Ravi N, Basavarajappa R, Chandrashekar C, Harlapur S, Hosamani M, et al. (2012) Effect of integrated nutrient management on growth and yield of quality protein maize. *Karnataka Journal of Agricultural Sciences* 25: 395–396.
37. Kannan RL, Dhivya M, Abinaya D, Krishna RL, Krishnakumar S (2013) Effect of integrated nutrient management on soil fertility and productivity in maize. *Bulletin of Environment, Pharmacology and Life Sciences* 2: 61–67.
38. Lone AA, Allai B, Nehvi F (2013) Growth, yield and economics of baby corn (*Zea mays* L.) as influenced by Integrated Nutrient Management (INM) practices. *African journal of agricultural research* 8: 4537–4540.
39. Haque S, Patra PS, Saha A (2012) Growth attributes, cob yield and economics of maize (*Zea mays* L.) cultivars as affected by integrated nutrient management under Terai region of West Bengal. *Indian Journal of Agricultural Research* 46.
40. Gundlur S, Patil P, Rajkumara S, Ashoka P, Neelakantha J (2015) Influence of integrated nutrient management on yield and uptake of nutrients by maize and soil fertility under irrigated conditions in Vertisol. *Karnataka Journal of Agricultural Sciences* 28: 172–175.
41. Mahala H, Shaktawat M, Shivran R (2006) Direct and residual effects of sources and levels of phosphorus and farmyard manure in maize (*Zea mays*)-mustard (*Brassica juncea*) cropping sequence. *Indian Journal of Agronomy* 51: 10–13.
42. Reza S, Pal S, Singh S (2012) Rock Phosphate-enriched Pressmud Compost: Direct Effect in Pearl Millet (*Pennisetum glaucum* L.) and Residual Effect in Mustard (*Brassica juncea*) in a Typic Haplustept. *Journal of the Indian Society of Soil Science* 60: 138–144.
43. Zaremanesh H, Nasiri B, Amiri A (2017) The effect of vermicompost biological fertilizer on corn yield. *J Mater Environ Sci* 8: 154.
44. Sharma K, Sharma D, Shukla K, Upadhyay B (2008) Growth, partitioning and productivity of wheat as influenced by fertilization and foliar application of bioregulators. *Indian J Plant Physiol* 13: 387–393.
45. Baraich A, Baraich A, Chattha S, Salarzi A (2012) Impact of phosphorus and potassium on the growth and yield of wheat cv. Mehran-89. *Pakistan Journal of Agriculture: Agricultural Engineering Veterinary Sciences (Pakistan)*.
46. Arif M, Atta S, Bashir MA, Khan MI, Hussain A, et al. (2021) The impact of Fosetyl-Aluminium application timing on Karnal bunt suppression and economic returns of bread wheat (*Triticum aestivum* L.). *Plos one* 16: e0244931. <https://doi.org/10.1371/journal.pone.0244931> PMID: 33428646
47. Khalofah A, Khan MI, Arif M, Hussain A, Ullah R, et al. (2021) Deep placement of nitrogen fertilizer improves yield, nitrogen use efficiency and economic returns of transplanted fine rice. *Plos one* 16: e0247529. <https://doi.org/10.1371/journal.pone.0247529> PMID: 33630922
48. Mamuye M, Nebiyu A, Elias E, Berecha G (2021) Combined Use of Organic and Inorganic Nutrient Sources Improved Maize Productivity and Soil Fertility in Southwestern Ethiopia. *International Journal of Plant Production*: 1–12.
49. Mateos-Naranjo E, Andrades-Moreno L, Davy AJ (2013) Silicon alleviates deleterious effects of high salinity on the halophytic grass *Spartina densiflora*. *Plant Physiology and Biochemistry* 63: 115–121. <https://doi.org/10.1016/j.plaphy.2012.11.015> PMID: 23257076
50. Khoshgoftarmansh AH, Khodarahmi S, Haghighi M (2014) Effect of silicon nutrition on lipid peroxidation and antioxidant response of cucumber plants exposed to salinity stress. *Archives of Agronomy and Soil Science* 60: 639–653.
51. Abbas T, Balal RM, Shahid MA, Pervez MA, Ayyub CM, et al. (2015) Silicon-induced alleviation of NaCl toxicity in okra (*Abelmoschus esculentus*) is associated with enhanced photosynthesis, osmoprotectants and antioxidant metabolism. *Acta Physiologiae Plantarum* 37: 6.
52. Rahbarian R, Khavari-Nejad R, Ganjeali A, Bagheri A, Najafi F (2011) Drought stress effects on photosynthesis, chlorophyll fluorescence and water relations in tolerant and susceptible chickpea (*Cicer arietinum* L.) genotypes. *Acta Biologica Cracoviensia s Botanica*.
53. Hosseinzadeh S, Amiri H, Ismaili A (2016) Effect of vermicompost fertilizer on photosynthetic characteristics of chickpea (*Cicer arietinum* L.) under drought stress. *Photosynthetica* 54: 87–92.
54. Atik A (2013) Effects of planting density and treatment with vermicompost on the morphological characteristics of oriental beech (*Fagus orientalis* Lipsky.). *Compost Science & Utilization* 21: 87–98.
55. Amiri H, Ismaili A, Hosseinzadeh SR (2017) Influence of vermicompost fertilizer and water deficit stress on morpho-physiological features of chickpea (*Cicer arietinum* L. cv. karaj). *Compost Science & Utilization* 25: 152–165.