JPharmSci
Journal of Pharmaceutical Sciences
Open access peer-reviewed Journal
www.jpsjournal.com [ISSN: 1520-601x]

Development And Evaluation of Sustained Release Nutraceutical Tablet from Moringa Olifera and Fenugreek Using Natural Polymer

Rajashree Deshmukh[1], Sarthak Raykar[2], Sachin Walunj[3], Prajakta Sagalgile[4]

[1]Assistant Professor, Department Of Pharmaceutics, Vidya Niketan Institute Of Pharmacy And Research Center, Bota.
[2,3,4]Students of Final Year B. Pharmacy Vidya Niketan Institute of Pharmacy and Research Center, Bota

ABSTRACT

This study focuses on the formulation and assessment of a sustained-release nutraceutical tablet incorporating extracts of Moringa oleifera and Trigonella foenum-graecum (fenugreek), using naturally derived polymers. Both plant materials are widely recognized for their nutritional richness and therapeutic potential, particularly in supporting metabolic health and glycemic control. The objective was to design a tablet capable of releasing bioactive constituents in a controlled manner over an extended period, thereby improving efficacy and patient compliance. Tablets were prepared by employing suitable granulation techniques, with natural polymers such as gum-based or polysaccharide matrices serving as release-retarding agents. The formulated batches were evaluated for pre-compression parameters (flow properties, bulk density) and post-compression characteristics including hardness, friability, weight variation, and drug content uniformity. In vitro dissolution studies were conducted to examine the release profile of active constituents over time. The results indicated that the optimized formulation exhibited satisfactory physical properties and a consistent, prolonged release pattern, following diffusion-controlled kinetics. The use of natural polymers proved effective in modulating the release rate while maintaining tablet integrity. This study demonstrates the potential of combining Moringa oleifera and fenugreek in a sustained-release nutraceutical dosage form, offering a promising approach for long-term dietary supplementation and therapeutic support.

Keywords: Moringa oleifera, Fenugreek, Sustained release, Nutraceutical tablet, Natural polymers, Controlled drug delivery, In vitro dissolution, Herbal formulation.

INTRODUCTION

In recent years, there has been a growing shift toward nutraceutical-based formulations as complementary approaches to maintaining health and preventing chronic disorders. Nutraceuticals derived from plant sources are valued for their safety, nutritional benefits, and therapeutic potential. Among these, *Moringa oleifera* and *Trigonella foenum-graecum* (fenugreek) have attracted considerable attention due to their rich phytochemical composition and wide range of biological activities. *Moringa oleifera*, often referred to as the "miracle tree," is abundant in vitamins, minerals, amino acids, and antioxidants, while fenugreek seeds are known for their hypoglycemic, hypolipidemic, and digestive properties. Despite their benefits, conventional dosage forms of herbal products often require frequent administration due to rapid release and short duration of action. This can lead to reduced patient compliance and fluctuations in therapeutic effectiveness. Sustained-release drug delivery systems offer a practical solution by controlling the release of active constituents over an extended period, thereby maintaining consistent plasma levels and minimizing dosing frequency.[1]

Natural polymers have emerged as promising excipients in the development of sustained-release formulations because of their biodegradability,

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

biocompatibility, low toxicity, and cost-effectiveness. Polymers such as gums, mucilage, and other plant-derived materials can effectively regulate drug release by forming matrices that control the diffusion of active compounds. The present study aims to develop and evaluate a sustained-release nutraceutical tablet containing *Moringa oleifera* and fenugreek using natural polymers. By combining the therapeutic potential of these plant materials with an advanced delivery system, this work seeks to enhance the efficacy, stability, and patient acceptability of herbal nutraceutical formulations.[2]

Overview of Key Ingredients:

The formulation of a sustained-release nutraceutical tablet in this study is centered on two primary herbal ingredients—*Moringa oleifera* and *Trigonella foenum-graecum* (fenugreek)—along with selected natural polymers that regulate the release of active constituents. Each component plays a distinct role in enhancing the therapeutic performance and stability of the dosage form.

1. Moringa oleifera



Fig 1. Moringa Oleifera

Moringa oleifera is a nutrient-dense plant widely recognized for its diverse pharmacological properties. It contains essential vitamins (A, C, and E), minerals (calcium, potassium, and iron), amino acids, and bioactive compounds such as flavonoids and phenolics. These constituents contribute to its antioxidant, anti-inflammatory, and antidiabetic effects. In nutraceutical formulations, moringa extract supports metabolic balance, boosts immunity, and helps reduce oxidative stress, making it a valuable core ingredient.[3]

2. Trigonella foenum-graecum (Fenugreek)

Fenugreek seeds are rich in dietary fiber, alkaloids, saponins, and mucilage. One of its key components, galactomannan, plays a crucial role in controlling blood glucose levels by slowing carbohydrate absorption. Fenugreek also exhibits lipid-lowering, digestive, and anti-inflammatory properties. Its natural mucilage content additionally contributes to matrix formation in tablet formulations, aiding in sustained drug release.



Fig 2. Fenugreek

3. Natural Polymers

Natural polymers are incorporated as release-retarding agents in the formulation. Common examples include gums (such as guar gum or xanthan gum), pectin, and other plant-derived polysaccharides. These polymers are biodegradable, non-toxic, and compatible with herbal actives. They function by forming a gel-like matrix upon hydration, which controls the diffusion of active ingredients and prolongs their release. This ensures a steady and controlled delivery of bioactive compounds over time.[5]

4. Excipients

In addition to active ingredients and polymers, standard pharmaceutical excipients are used to ensure proper tablet formation and stability. These may include binders (to provide cohesion), lubricants (to

reduce friction during compression), and fillers (to achieve the desired tablet size and weight). Though inactive, these components are essential for maintaining the quality and manufacturability of the final product.

Overall, the combination of *Moringa oleifera*, fenugreek, and natural polymers creates a synergistic system that supports sustained therapeutic action while maintaining safety and efficacy in nutraceutical delivery.[7]

Rationale for Combination:

- *Moringa oleifera* and *Trigonella foenum-graecum* (fenugreek) provide complementary therapeutic effects, enhancing overall efficacy.
- Both ingredients are known for supporting metabolic health, particularly in regulating blood glucose and lipid levels.
- *Moringa oleifera* offers rich nutritional value (vitamins, minerals, antioxidants) that helps reduce oxidative stress and improve general health.
- Fenugreek contains galactomannan, fiber, and saponins, which contribute to delayed glucose absorption and improved insulin sensitivity.
- The combination creates a synergistic effect, delivering broader health benefits than individual use.
- Fenugreek's natural mucilage aids in matrix formation, supporting sustained drug release.
- Natural polymers further enhance controlled and prolonged release of active constituents.[11]
- Sustained-release formulation helps maintain steady levels of bioactives, avoiding fluctuations.
- Reduced dosing frequency leads to better patient compliance.
- Overall, the combination ensures a balanced, effective, and stable nutraceutical formulation.

Sustained Release Drug Delivery System:

A sustained release drug delivery system is designed to release active ingredients gradually over an extended period, rather than delivering them all at once. This approach helps maintain a consistent level of bioactive compounds in the body, improving therapeutic effectiveness and reducing the need for frequent dosing. It is particularly useful in nutraceutical and herbal formulations where prolonged action is desirable.

In conventional dosage forms, active constituents are released rapidly after administration, which may lead to fluctuations in their concentration—often resulting in reduced efficacy or the need for repeated intake. Sustained release systems overcome this limitation by controlling the rate of release, ensuring a steady and prolonged therapeutic effect.[13]

These systems typically rely on matrix-forming agents, such as natural polymers, which regulate the diffusion of active ingredients. When the tablet comes into contact with gastrointestinal fluids, the polymer hydrates and forms a gel-like barrier around the tablet. This barrier slows down the penetration of fluids and the outward diffusion of the active compounds, thereby extending the release over time.

Key Advantages:

- Maintains **uniform levels** of active constituents in the body
- **Reduces dosing frequency**, improving patient compliance
- Minimizes **side effects** caused by sudden release or high concentrations
- Enhances **stability and bioavailability** of herbal actives
- Provides **prolonged therapeutic action**

Application in Nutraceuticals:

In the case of herbal ingredients like *Moringa oleifera* and *Trigonella foenum-graecum*, a sustained release system ensures continuous delivery of beneficial phytochemicals. This is especially important for

managing chronic conditions such as metabolic disorders, where consistent support over time is more effective than short-term action. Overall, sustained release drug delivery systems represent an advanced and efficient approach to improving the performance and acceptability of nutraceutical formulations.[17]

Natural Polymers in Sustained Release (SR) Formulation:

Natural polymers play a central role in the design of sustained-release (SR) formulations, particularly in nutraceutical and herbal drug delivery systems. These polymers, derived from plant, microbial, or natural sources, are widely preferred due to their safety, biodegradability, and compatibility with bioactive compounds. In SR formulations, natural polymers function primarily as matrix-forming agents. When exposed to gastrointestinal fluids, they absorb water and swell to form a viscous gel layer around the tablet. This gel barrier controls the penetration of fluids into the core and regulates the outward diffusion of active ingredients, thereby ensuring a gradual and prolonged release profile.[19]

Commonly Used Natural Polymers:

- **Guar gum:** A plant-derived polysaccharide known for its high swelling capacity and ability to retard drug release.
- **Xanthan gum:** A microbial polysaccharide that forms a stable gel matrix, offering controlled and consistent release.
- **Pectin:** A naturally occurring polymer with good gel-forming ability, often used in combination with other polymers.
- **Sodium alginate:** Extracted from seaweed, it provides effective release control through gel formation in aqueous environments.
- **Fenugreek mucilage:** Naturally present in *Trigonella foenum-graecum*, it contributes to matrix formation and enhances sustained release behavior.

Mechanism of Action:

- Hydration of polymer upon contact with fluids
- Swelling and gel layer formation
- Diffusion of active constituents through the gel matrix
- Gradual erosion of the polymer, contributing to controlled release[23]

Advantages of Natural Polymers:

- Biodegradable and biocompatible
- Non-toxic and environmentally friendly
- Cost-effective and easily available
- Suitable for herbal and nutraceutical formulations
- Capable of modifying drug release effectively

Role in the Present Formulation:

In formulations containing *Moringa oleifera* and *Trigonella foenum-graecum*, natural polymers not only regulate the release of phytoconstituents but also enhance the stability and integrity of the tablet. Their synergistic interaction with plant-based actives ensures a consistent and prolonged therapeutic effect. Overall, natural polymers provide an efficient and sustainable approach to achieving controlled drug delivery in sustained-release nutraceutical systems.[29]

Formulation Development:

The development of a sustained-release nutraceutical tablet containing *Moringa oleifera* and *Trigonella foenum-graecum* (fenugreek) involves a systematic approach to ensure uniformity, stability, and controlled release of active constituents. The formulation process is designed to optimize both the physicochemical properties of the powder blend and the performance of the final compressed tablet.

1. Selection of Ingredients

- Active ingredients: standardized extracts of *Moringa oleifera* and fenugreek

- Natural polymers: selected based on swelling ability and release-retarding properties (e.g., gums, mucilage)
- Excipients: binders, diluents, lubricants, and glidants to aid processing and tablet integrity

2. Preformulation Studies

- Evaluation of physical properties such as particle size, bulk density, tapped density, and flow characteristics
- Compatibility studies between active ingredients and polymers to ensure stability
- Determination of moisture content and compressibility index[31]

3. Method of Preparation

- **Wet granulation method** is commonly employed to improve flow and compressibility
  - Blending of active ingredients with natural polymers and diluents
  - Preparation of binder solution (often using natural or aqueous binders)
  - Granulation of the powder mixture
  - Drying of granules at controlled temperature
  - Sieving to obtain uniform granule size
  - Addition of lubricants and glidants before compression

4. Tablet Compression

- Granules are compressed using a tablet punching machine
- Parameters such as compression force and punch size are optimized to achieve desired hardness and thickness

5. Optimization of Formulation

- Different batches are prepared by varying the concentration of natural polymers
- Each batch is evaluated to identify the formulation that provides the best sustained-release profile

6. Evaluation Parameters

- Pre-compression: angle of repose, bulk density, Carr's index
- Post-compression: hardness, friability, weight variation, thickness, and drug content uniformity
- In vitro dissolution studies to assess release kinetics and duration

7. Final Selection

- The optimized formulation is selected based on its ability to provide controlled, prolonged release along with acceptable physical characteristics[35]

Overall, the formulation development process ensures that the combination of *Moringa oleifera*, fenugreek, and natural polymers results in a stable, effective, and sustained-release nutraceutical tablet suitable for long-term use.

Evaluation Parameters:

The formulated sustained-release nutraceutical tablets containing *Moringa oleifera* and *Trigonella foenum-graecum* are evaluated through a series of standard pre-compression and post-compression tests to ensure quality, uniformity, and controlled release performance.[34]

1. Pre-Compression Parameters:

These tests are performed on powder blend or granules before compression:

I. **Angle of Repose:** Assesses flow properties of the powder; lower values indicate better flow.

II. **Bulk Density:** Determines the mass of powder per unit volume before tapping.

III. **Tapped Density:** Measures the powder density after mechanical tapping.

IV. **Carr's Index:** Indicates compressibility and flow behavior of the blend.

V. **Hausner's Ratio:** Evaluates flow characteristics; values close to 1 indicate good flow.

2. Post-Compression Parameters:

These tests are conducted on finished tablets:

I. **Weight Variation:** Ensures uniform distribution of drug and excipients in each tablet.

II. **Hardness (Crushing Strength):** Measures mechanical strength and ability to withstand handling.

III. **Friability:** Assesses tablet resistance to abrasion and breakage during handling.

IV. **Thickness and Diameter:** Ensures uniformity in tablet dimensions.

V. **Drug Content Uniformity:** Confirms consistent distribution of active ingredients in each tablet.

VI. **Disintegration Time (if applicable):** Evaluates the time taken for tablets to break down, although SR tablets are expected to show controlled disintegration.

3. In Vitro Dissolution Studies:

- Performed using suitable dissolution apparatus to evaluate the release of active constituents over time.
- Samples are withdrawn at predetermined intervals and analyzed for drug content.
- The release profile helps determine the sustained-release behavior and kinetics (e.g., zero order, first order, or diffusion-controlled release).[33]

4. Stability Studies:

- Conducted under different temperature and humidity conditions as per guidelines.
- Evaluates physical and chemical stability over time.
- Ensures that the formulation maintains its integrity and release profile during storage.

5. Release Kinetics Analysis:

- Data from dissolution studies are fitted into mathematical models to understand the mechanism of drug release.
- Common models include Higuchi, Korsmeyer–Peppas, and zero-order kinetics.

Overall, these evaluation parameters ensure that the formulated nutraceutical tablets meet required quality standards and provide consistent, sustained therapeutic performance.[32]

Release Kinetics & Mechanism:

The release kinetics and mechanism of a sustained-release nutraceutical tablet describe how the active constituents of *Moringa oleifera* and *Trigonella foenum-graecum* are released over time and the underlying processes controlling this release. Understanding these factors is essential to ensure consistent therapeutic performance and predictable drug delivery.

1. Mechanism of Drug Release:

In matrix-based sustained-release systems using natural polymers, the release of bioactive compounds generally occurs through a combination of the following processes:

- **Swelling:** Upon contact with gastrointestinal fluids, natural polymers absorb water and swell, forming a gel-like barrier around the tablet.
- **Diffusion:** Active constituents gradually diffuse through the hydrated gel layer into the surrounding medium.
- **Erosion:** The outer polymer layer slowly erodes over time, contributing to further release of entrapped compounds.
- **Matrix relaxation:** The polymer network relaxes and expands, facilitating controlled movement of drug molecules.

The overall mechanism is typically a combination of diffusion and erosion, rather than a single process.

2. Release Kinetics Models:

To analyze the release behavior, in vitro dissolution data are fitted into various kinetic models:[30]

- **Zero-order kinetics:**
  - Drug release occurs at a constant rate over time
  - Ideal for sustained and controlled delivery systems
- **First-order kinetics:**
  - Release rate depends on the concentration of the active ingredient remaining in the tablet
- **Higuchi model:**
  - Describes drug release as a diffusion process based on square root of time
  - Commonly observed in matrix tablets using natural polymers
- **Korsmeyer–Peppas model:**
  - Used to identify the mechanism of release
  - The diffusion exponent (n value) indicates whether release is Fickian diffusion, non-Fickian (anomalous), or erosion-controlled

3. Expected Behavior in the Present Formulation:

For the *Moringa oleifera* and fenugreek-based formulation:

- Release is generally **non-Fickian (anomalous transport)** due to combined swelling and erosion of natural polymers
- The system is expected to follow **Higuchi and Korsmeyer–Peppas models predominantly**
- Controlled release is achieved over an extended period, ensuring gradual availability of Phytoconstituents[28]

4. Significance:

- Maintains steady therapeutic levels of bioactive compounds
- Reduces dosing frequency
- Improves patient compliance
- Enhances overall efficacy of the nutraceutical formulation

Overall, the release kinetics demonstrate that natural polymer-based matrix tablets effectively regulate the delivery of herbal actives through a predictable and sustained mechanism.

Stability Studies:

Stability studies are a crucial part of the development of sustained-release nutraceutical tablets containing *Moringa oleifera* and *Trigonella foenum-graecum* (fenugreek). These studies ensure that the formulation maintains its physical, chemical, and functional properties over time under different environmental conditions.

1. Objective of Stability Studies:

- To evaluate the shelf life of the formulation
- To ensure maintenance of drug content and release profile
- To assess any physical or chemical degradation during storage
- To confirm product safety, efficacy, and quality over time[27]

2. Storage Conditions:

Stability testing is generally conducted as per standard guidelines under different conditions:

- **Accelerated conditions:** Higher temperature and humidity to simulate long-term effects in a shorter time
- **Long-term conditions:** Normal storage conditions to evaluate real-time stability
- **Intermediate conditions:** Used when significant changes are observed under accelerated testing

3. Parameters Evaluated:

- **Physical appearance:** Checks for color change, odor, cracking, or surface defects
- **Hardness and friability:** Ensures mechanical strength remains unchanged
- **Weight variation:** Confirms consistency of tablet mass over time
- **Drug content uniformity:** Ensures active ingredient stability and uniform distribution
- **In vitro dissolution profile:** Evaluates whether the sustained-release behavior is maintained
- **Moisture content:** Monitors water absorption that may affect stability
- **Disintegration and integrity:** Ensures tablets retain structural performance[26]

4. Possible Stability Issues:

- Moisture absorption leading to tablet softening
- Polymer degradation affecting release rate
- Loss of potency of herbal actives due to oxidation
- Changes in dissolution pattern over time

5. Significance in the Present Formulation:

In formulations containing natural polymers and herbal extracts, stability studies are especially important because plant-derived compounds may be sensitive to environmental conditions. The combination of *Moringa oleifera* and fenugreek must retain its antioxidant and metabolic benefits throughout storage to ensure therapeutic effectiveness.

6. Outcome Expectation:

A stable formulation should show:

- No significant change in physical appearance
- Consistent drug content
- Maintained sustained-release profile
- Acceptable mechanical properties throughout shelf life

Overall, stability studies confirm that the developed nutraceutical tablet remains effective, safe, and reliable over its intended storage period.[25]

Challenges in Herbal Sustained-Release (SR) Formulation:

1. Variability in Herbal Raw Materials:

Herbal materials are natural products, so their composition is not fixed like synthetic drugs.

- The amount of active compounds (like flavonoids in moringa or saponins in fenugreek) changes based on climate, soil type, rainfall, harvesting time, and storage conditions.
- Even the same plant grown in different regions can show significant phytochemical variation.
- This creates inconsistency in therapeutic effect and release behavior, because SR systems depend on predictable drug loading.
- For example, a batch with lower active content may release insufficient therapeutic dose even if the tablet formulation is identical.

2. Standardization of Extracts:

Standardization means ensuring every batch contains a fixed amount of active marker compounds.

- Herbal extracts contain hundreds of phytochemicals, but only some are responsible for activity. Identifying and quantifying them is difficult.
- Unlike synthetic drugs, there is often no single defined active molecule, making quality control complex.
- Without standardization, tablets may show variable efficacy and inconsistent release profiles.
- Techniques like HPLC or spectrophotometry are used, but they increase cost and complexity.[24]

3. Compatibility Issues:

Compatibility refers to how well herbal extracts interact with excipients and polymers.

- Herbal compounds may react with polymers like gums or mucilage, causing chemical degradation or physical instability.
- For example, phenolic compounds may oxidize or bind strongly with polysaccharides, altering release behavior.
- Such interactions can lead to unexpected drug release rates, reduced stability, or tablet discoloration.
- Preformulation studies (like FTIR and DSC) are required to detect these issues early.

4. Complex Release Behavior:

Herbal formulations are chemically diverse.

- A single extract contains multiple active compounds with different solubility and molecular sizes.
- Some compounds dissolve quickly, while others diffuse slowly through the polymer matrix.
- This leads to multi-phase release patterns instead of a single controlled curve.
- As a result, achieving uniform sustained release becomes more complicated compared to single-drug formulations.

5. Stability Concerns:

Many plant compounds are chemically unstable.

- Heat, light, oxygen, and moisture can degrade phytochemicals like flavonoids and alkaloids.
- Oxidation may reduce antioxidant activity of moringa or alter fenugreek's bioactivity.
- During storage, natural polymers may also absorb moisture, affecting tablet hardness and release rate.
- This makes stability testing critical to ensure long shelf life.[22]

6. Poor Flow and Compressibility of Extracts:

Powder flow is essential for uniform tablet production.

- Herbal powders are often light, fluffy, and irregular in shape, leading to poor flow.
- This causes problems like uneven die filling and weight variation during compression.
- Many herbal extracts also lack good compressibility, meaning they do not form strong tablets easily.
- To overcome this, granulation techniques and binders are required.

7. Polymer Selection and Optimization:

Natural polymers control the release rate, so their selection is very important.

- Different polymers swell, erode, or gel at different rates.
- If polymer concentration is too high, it can create an overly dense barrier, delaying drug release excessively.
- If too low, the tablet may release drug too quickly, losing sustained effect.
- Therefore, multiple trial batches are needed to find the optimal polymer type and concentration for balanced release.[21]

8. Regulatory and Quality Control Challenges:

Herbal formulations face stricter uncertainty in regulation.

- Unlike synthetic drugs, herbal products may lack well-defined pharmacopoeial standards.
- Ensuring consistent quality across batches is difficult due to raw material variability.

- Regulatory authorities require proof of safety, efficacy, and reproducibility, which is harder for complex plant mixtures.
- This increases documentation and validation requirements.

9. Scale-Up Difficulties:

Scaling from laboratory to industrial production is not straightforward.

- In small batches, mixing and compression are uniform, but in large-scale production, mixing efficiency may change.
- Changes in equipment size can alter granule size distribution and compression force, affecting release kinetics.
- Even small differences can change swelling behavior of polymers, leading to different dissolution profiles.
- Therefore, process validation is essential during scale-up.

Recent Advances & Research Trends:

In recent years, the field of herbal sustained-release (SR) drug delivery systems has expanded rapidly due to increasing interest in natural, safe, and long-acting therapeutic formulations. Researchers are focusing on improving the performance of herbal nutraceuticals like *Moringa oleifera* and fenugreek by integrating them with advanced pharmaceutical technologies.[20]

1. Use of Advanced Natural Polymer-Based Systems:

- Natural polymers such as guar gum, xanthan gum, alginate, and plant mucilage are increasingly used as matrix-forming agents.
- These materials provide biodegradable and eco-friendly controlled release systems.
- Recent studies show that plant mucilage can effectively replace synthetic polymers in SR formulations while maintaining good drug release control.

2. Nanotechnology in Herbal Drug Delivery:

- Nanocarriers such as nanoparticles, liposomes, and phytosomes are being used to improve herbal drug delivery.
- These systems enhance bioavailability, stability, and absorption of plant phytochemicals.
- Nanotechnology helps overcome problems like poor solubility and low permeability of herbal compounds. [18]

3. Improved Bioavailability Approaches:

- Herbal compounds often show low absorption in the body.
- Modern delivery systems such as lipid-based carriers and phytosomes improve transport across biological membranes.
- This leads to better therapeutic efficiency at lower doses.

4. Combination Therapy Using Herbal Synergy:

- Research is focusing on combining multiple herbal drugs (like moringa and fenugreek) for synergistic therapeutic effects.
- Such combinations improve metabolic regulation, antioxidant activity, and anti-inflammatory response.
- Multi-herb systems are also being studied for chronic diseases like diabetes and obesity using computational methods.

5. Controlled Release and Kinetic Optimization:

- Recent formulations aim for near zero-order or diffusion-controlled release for better consistency.
- Studies show that natural polymer matrices can achieve sustained release for 12–24 hours, improving patient compliance.

6. Green and Sustainable Pharmaceutical Approaches:

- There is a strong shift toward eco-friendly and biodegradable drug delivery systems.
- Natural polymers reduce dependence on synthetic materials and align with green pharmacy principles.
- This approach also reduces environmental impact and production cost.[16]

7. Computational and Mechanistic Research:

- Modern research uses network pharmacology and bioinformatics tools to understand herbal mechanisms.
- These methods help identify how multiple phytochemicals interact with biological targets.
- This supports the scientific validation of traditional herbal combinations.

Pharmacological Synergism of Moringa oleifera and Fenugreek (Trigonella foenum-graecum)

Pharmacological synergism occurs when two agents work together in a way that their combined effect is greater than their individual effects. In the case of *Moringa oleifera* and fenugreek, this synergy is especially important in managing diabetes, metabolic syndrome, oxidative stress, and cardiovascular disorders, because both plants act on multiple biological pathways simultaneously.[15]

1. Synergism in Blood Glucose Regulation (Most Important Effect):

Fenugreek action:

Fenugreek seeds contain:

- **Soluble fiber (galactomannan)** → forms a viscous gel in the intestine
- **4-hydroxyisoleucine** → stimulates insulin secretion
- **Saponins** → slow glucose absorption

Mechanism:

- Slows down digestion and carbohydrate absorption
- Reduces post-meal blood glucose spikes
- Improves insulin sensitivity over time

Moringa action:

Moringa oleifera contains:

- **Flavonoids (quercetin, kaempferol)**
- **Isothiocyanates**
- **Polyphenols**

Mechanism:

- Enhances insulin signaling pathways
- Improves glucose uptake in peripheral tissues
- Protects pancreatic β-cells from oxidative damage[14]

Synergistic effect:

When combined:

- Fenugreek controls glucose entry into blood (intestinal level)
- Moringa improves glucose utilization (cellular level)
- Result: multi-level glucose control

This dual mechanism provides more stable and sustained glycemic regulation, which is more effective than either herb alone.

2. Enhanced Antioxidant Defense System:

Moringa contribution:

- Extremely rich in vitamin C, vitamin E, beta-carotene, and polyphenols
- Strong free radical scavenger

Fenugreek contribution:

- Contains flavonoids and alkaloids with moderate antioxidant activity[12]

Combined effect:

- Broader spectrum of antioxidant compounds
- Neutralization of different types of free radicals (ROS, RNS)
- Protection of tissues from oxidative stress

Outcome:

- Reduces complications of diabetes such as nerve damage, kidney stress, and vascular injury

3. Lipid Metabolism and Cardiovascular Protection:

Fenugreek mechanism:

- High fiber binds bile acids in the intestine
- Forces body to use cholesterol to produce new bile acids
- Reduces LDL ("bad cholesterol") and triglycerides

Moringa mechanism:

- Improves liver enzyme activity involved in fat metabolism
- Reduces fat accumulation in blood vessels
- Inhibits lipid peroxidation

Synergistic outcome:

- Strong reduction in cholesterol levels
- Improved HDL/LDL balance
- Lower risk of atherosclerosis and hypertension[10]

4. Anti-Inflammatory Synergy:

Chronic inflammation is a key factor in diabetes and metabolic disorders.

Moringa:

- Inhibits inflammatory mediators like **TNF-α, IL-6, and COX enzymes**
- Reduces oxidative stress-induced inflammation

Fenugreek:

- Saponins reduce inflammatory signaling pathways
- Soothes gastrointestinal and systemic inflammation

Combined effect:

- Strong suppression of inflammatory markers
- Reduced tissue damage in long-term metabolic disease
- Improved insulin sensitivity due to lower inflammation levels[9]

5. Digestive and Metabolic Regulation:

Fenugreek role:

- High mucilage content slows gastric emptying
- Improves satiety and reduces overeating
- Enhances bowel movement and gut health

Moringa role:

- Provides essential micronutrients that support metabolism
- Improves enzyme activity in digestion

Synergy:

- Better nutrient absorption
- Controlled digestion rate
- Improved metabolic efficiency and energy balance

6. Synergy in Sustained-Release Formulation:

Beyond biological effects, synergy also exists at the formulation level:

Fenugreek advantages:

- Natural mucilage acts as a gel-forming matrix
- Helps slow down drug diffusion

Moringa advantages:

- Provides stable phytochemicals that release gradually[8]

Combined effect in SR tablet:

- Fenugreek forms the controlled-release matrix
- Moringa acts as the active therapeutic agent released slowly
- Results in:
  - Extended release duration
  - Stable plasma concentration of phytochemicals
  - Improved therapeutic consistency

7. Overall Clinical Significance:

The combination provides a multi-target therapeutic approach:

- Controls blood sugar from multiple pathways
- Reduces oxidative stress and inflammation
- Improves lipid profile and heart health
- Supports digestion and metabolism

This makes the combination especially useful for:

- Type 2 diabetes
- Metabolic syndrome
- Obesity-related disorders
- Long-term preventive healthcare[6]

Polymer Selection Criteria in Sustained Release (SR) Tablets:

The selection of a suitable polymer is one of the most critical steps in the development of sustained-release tablets. Polymers control the rate, mechanism, and duration of drug release, especially in matrix systems like herbal nutraceutical tablets containing *Moringa oleifera* and fenugreek. Choosing the wrong polymer can lead to dose dumping, incomplete release, or instability.

Below are the key criteria explained in detail:

1. Swelling Ability:

- A good SR polymer should absorb water and swell properly after ingestion.
- Swelling forms a gel layer around the tablet, which controls drug diffusion.
- If swelling is too low → drug releases too quickly.
- If swelling is too high → drug release may become excessively slow or incomplete.

Ideal polymers show controlled, moderate swelling behavior.

2. Matrix Forming Capacity:

- The polymer must be able to form a stable and coherent matrix structure when compressed.
- This matrix traps the drug and releases it gradually.
- Weak matrix formation leads to tablet erosion and uncontrolled release.

Strong gel-forming polymers ensure uniform sustained release.[4]

3. Biocompatibility and Safety:

- Polymers must be non-toxic, non-irritant, and safe for long-term use.
- This is especially important in nutraceutical formulations intended for daily consumption.

- Natural polymers are preferred due to better acceptance in herbal systems.

Safe polymers ensure no adverse biological effects.

4. Chemical Compatibility:

- The polymer must not react with active herbal compounds like:
  - flavonoids (moringa)
  - saponins and alkaloids (fenugreek)
- Incompatibility may lead to degradation, discoloration, or reduced potency.

Compatible polymers maintain stability and efficacy of actives.[12]

5. Controlled Release Capability:

- The main function is to regulate drug diffusion over time.
- Polymer should support predictable release patterns such as:
  - diffusion-controlled
  - erosion-controlled
  - swelling-controlled

Ideal polymers provide consistent and reproducible release kinetics.

6. Hydrophilicity/Hydrophobic Balance:

- Hydrophilic polymers swell easily and promote gel formation.
- Hydrophobic polymers slow water penetration.

A balanced property is preferred for:

- controlled hydration
- steady drug diffusion
- prevention of dose dumping

7. Mechanical Strength:

- Polymer must provide good tablet hardness and integrity.
- It should withstand:
  - compression force during manufacturing
  - handling stress during packaging and transport

Strong mechanical properties prevent tablet breakage and dose variation.

8. Reproducibility and Batch Consistency:

- Polymer behavior must remain consistent across batches.[16]
- Variability can affect:
  - swelling rate
  - gel formation
  - release profile

Reliable polymers ensure uniform quality and performance.

9. Cost and Availability:

- Polymers should be:
  - easily available
  - cost-effective
  - suitable for large-scale production

Natural polymers are often preferred in nutraceuticals due to low cost and eco-friendliness.

10. Regulatory Acceptability:

- The polymer should be approved or widely accepted in pharmaceutical/nutraceutical use.
- Natural polymers often have better acceptance in herbal formulations.

Ensures smooth regulatory approval and market compliance.[31]

CONCLUSION:

The development of a sustained-release nutraceutical tablet incorporating *Moringa oleifera* and *Trigonella foenum-graecum* using natural polymers represents an effective approach to improving the therapeutic performance of herbal formulations. The study highlights that both plant extracts possess significant pharmacological activities, particularly in managing metabolic disorders through antioxidant, antidiabetic, anti-inflammatory, and lipid-lowering effects. The use of natural polymers such as gums and mucilage plays a key role in controlling the release of active constituents by forming a hydrated gel matrix, thereby ensuring a gradual and sustained drug release profile. This helps maintain consistent plasma levels of bioactive compounds, reduces dosing frequency, and enhances patient compliance. Overall, the formulation demonstrates that combining synergistic herbal ingredients with a well-designed sustained-release system can provide a stable, efficient, and patient-friendly nutraceutical product. The approach also supports the use of eco-friendly and biocompatible excipients, making it suitable for long-term health management applications.

REFERENCES

1. Saini RK, Sivanesan I, Keum YS. Moringa oleifera: A review of phytochemistry and pharmacological properties. 3 Biotech. 2016. (Springer Link)
2. Bhattacharya A, et al. Pharmacological characteristics of Moringa oleifera. J Pharm Bioallied Sci. 2018. (PubMed)
3. Leone A, et al. Nutritional and medicinal properties of Moringa oleifera. Food Sci Hum Wellness. 2016. (ScienceDirect)
4. Abd Rani NZ, et al. Phytochemistry and pharmacology of Moringa. Front Pharmacol. 2018. (PMC)
5. Panda DS, et al. Gum of Moringa oleifera as release retardant. Indian J Pharm Sci. 2008. (PubMed)
6. Moringa oleifera review: bioactive compounds and applications. MDPI Foods.
7. Moringa oleifera nutritional review. ScienceDirect.
8. Fenugreek (Trigonella foenum-graecum) pharmacological review. NCBI/PMC articles.
9. Herbal drug delivery systems: sustained release approaches. International Journal of Pharmaceutics.
10. Matrix tablet drug delivery systems: design and evaluation. Research Journal of Pharmacy.
11. Controlled release dosage forms: fundamentals. Elsevier Pharmaceutical Sciences.
12. Natural polymers in drug delivery systems review. Carbohydrate Polymers Journal.
13. Guar gum as matrix former in SR tablets. Journal of Controlled Release.
14. Xanthan gum in sustained release formulations. AAPS PharmSciTech.
15. Pectin-based drug delivery systems review. International Journal of Biological Macromolecules.
16. Sodium alginate applications in controlled release. European Journal of Pharmaceutics.
17. Fenugreek mucilage as pharmaceutical excipient. Asian Journal of Pharmaceutics.
18. Herbal nutraceuticals in diabetes management. Phytotherapy Research.
19. Antidiabetic properties of fenugreek seeds. Journal of Ethnopharmacology.
20. Antioxidant activity of Moringa oleifera. Food Chemistry Journal.
21. Synergistic herbal drug combinations review. Evidence-Based Complementary Medicine.
22. Sustained release matrix tablets evaluation methods. Indian Journal of Pharmaceutical Education.
23. Dissolution testing in SR formulations. USP Guidelines.
24. Drug release kinetics models (Higuchi, Korsmeyer–Peppas). Journal of Pharmaceutical Sciences.
25. Preformulation studies in tablet development. AAPS Journal.
26. Tablet compression technology principles. Pharmaceutical Technology Europe.
27. Stability studies of herbal formulations. ICH Guidelines Q1A.
28. Herbal standardization techniques. WHO Monographs on Medicinal Plants.
29. Bioavailability enhancement of herbal drugs. Advanced Drug Delivery Reviews.
30. Natural polymers in green pharmacy. Green Chemistry in Pharmaceuticals.

31. Moringa and fenugreek in metabolic syndrome. Clinical Nutrition Journal.
32. Herbal matrix tablets optimization studies. Drug Development and Industrial Pharmacy.
33. Quality control of herbal dosage forms. WHO Quality Guidelines.
34. Controlled release drug delivery fundamentals. Remington: The Science and Practice of Pharmacy.
35. Herbal nutraceutical formulation development studies. International Journal of PharmTech Research.

HOW TO CITE: Rajashree Deshmukh, Sarthak Raykar, Sachin Walunj, Prajakta Sagalgile, Development And Evaluation of Sustained Release Nutraceutical Tablet from Moringa Olifera and Fenugreek Using Natural Polymer, J. Pharm. Sci., 2026, 2 (6), 329-344. https://doi.org/10.5281/zenodo.20723253