# Title :
# Prostate-Protective Bioactivity of Cucurbita maxima Seeds: Molecular Pathways, Endocrine Regulation, and Clinical Relevance


**CorrespondingAuthor**

**YvesMakiasiHambadiana**
**HambasFoodTechStartup,DemocraticRepublicoftheCongo**
📧**yveslpmakiasi@gmail.com**
📞**+243825635265**

**Co-author**

**NdengaLumbuBarack(AliasBarackEinstein97)**
**IndependentScientificResearcher**
📧**ndengabarack@gmail.com**
📞**+243837767430**

## Abstract

Benign prostatic hyperplasia (BPH) and androgen-driven prostatic remodeling represent major components of male aging pathology. Seeds of Cucurbita maxima have emerged as a potent nutritional therapeutic agent due to their exceptionally rich composition in phytosterols, cucurbitacins, polyunsaturated fatty acids, tocopherols, and antioxidant phenolic compounds. This review synthesizes current molecular, biochemical, and clinical evidence demonstrating the prostate-protective potential of pumpkin seeds. β-sitosterol andrelated phytosterols inhibit 5-alpha reductase, reducing intraprostatic dihydrotestosterone (DHT) concentrations and attenuating androgen receptor activation. Cucurbitacins further suppress proliferative signaling pathways, including JAK/STAT, MAPK, and AR nuclear translocation, thereby limiting stromal and epithelial hypertrophy. Anti-inflammatory and antioxidant mechanisms—mediated through NF-κB inhibition, cytokine downregulation, and mitigation of oxidative stress—contribute to the restoration ofprostatic microenvironmental homeostasis. Additional effects on lipid metabolism and cholesterol regulation reinforce their role in endocrine and structural modulation of prostate tissues. Collectively, these findings support the integration of C. maxima seeds into functional foods and nutraceutical formulations aimed at BPH prevention and management, and highlight their relevance as a cost-effective, safe, and accessible therapeutic adjunct.

# 1. Introduction

Benign prostatic hyperplasia (BPH) and androgen-driven prostatic remodeling represent major health concerns among aging male populations. Increasing evidence from nutritional biochemistry and phytotherapeutic research highlights the significant preventive and therapeutic potential of Cucurbita maxima seeds. Beyond their traditional use, pumpkin seeds contain a constellation of bioactive molecules—particularly phytosterols, cucurbitacins, polyunsaturated fatty acids (PUFAs), tocopherols, and trace minerals—that exert multidimensional effects onprostate endocrine metabolism, inflammatory pathways, and cellular proliferation dynamics. The biochemical richness of C. maxima places it among the most clinically relevant functional foods for male reproductive health.


## 2. Modulation of Androgen Metabolism and 5-Alpha Reductase Activity

A central mechanism in prostatic hypertrophy involves the enzymatic conversion of testosterone into dihydrotestosterone (DHT) mediated by 5-alpha reductase (5α-R). DHT exhibits a 3- to 10-fold greater affinity for the androgen receptor (AR), promoting stromal and epithelial proliferation in the prostate.

Pumpkin seed phytosterols—especially β-sitosterol—demonstrate potent inhibitory activity on the 5α-R pathway. This inhibition manifests in:

- Reduced intraprostatic DHT accumulation,
- Downward modulation of AR hyperactivation,
- Attenuation of prostate tissue hyperplasia,
- Normalization of androgen-dependent gene transcription.

Mechanistically, β-sitosterol competes with endogenous sterols for membrane-bound transporters and enzymatic binding pockets, reducing substrate availability for DHT synthesis. This positions C. maxima seeds as natural endocrine regulators with a mechanistic profile comparable to mild pharmacological inhibitors used in BPH therapy.







**Figure 1 – Mechanisms of 5-Alpha Reductase Inhibition by Pumpkin Seed Phytosterols**
**This figure illustrates the competitive interaction of β-sitosterol with endogenous sterols at the 5-alpha reductase enzyme level, resulting in decreased conversion of testosterone into dihydrotestosterone (DHT). Reduced DHT accumulation leads to attenuated**

**androgen receptor activation and downstream suppression of prostate cellular proliferation.**

## 3. Regulation of Androgen Receptor Signaling and Prostate Cellular Dynamics


Beyond enzymatic inhibition, cucurbitacins—triterpenoid compounds abundant in C. maxima seeds—modulate AR nuclear translocation and disrupt downstream proliferative signaling cascades, including:


- JAK/STAT axis attenuation,
- MAPK pathway downregulation,
- Suppression of E2F-mediated cell cycle progression.


**These activities result in:**

- Reduced stromal fibroblast proliferation,
- Suppression of epithelial hypertrophy,
- Decreased prostate glandular volume over time,
- Improved tissue morphological stability.

Cucurbitacins also exhibit anti-proliferative effects on androgen-sensitive cellular models, suggesting a potential role in maintaining prostate homeostasis and preventing excessive hormonal remodeling.


activates JAK which
phosphorylates receptor
STAT binds to
phosphorylated receptor
phosphorylate
STAT
JAK
P
STAT
STAT PATHWAY
5
STAT DIMER
6
Translocation into
NUCLEUS
Binding to DNA &
7


Chemokines
Kinases
microRNA
Neuropilin-1
PAF
Glycoproteins
Mitogen
RKIP
Receptors
ARHGEF2
Ligands
JWA
Hormones
Oxidative stress
Pathogens
Heat shock
Ionizing radiation


MAPK
signaling
pathway






**Figure 2 – Cucurbitacin-Mediated Inhibition of Prostate Cell Proliferation**
**This diagram depicts how cucurbitacins inhibit major proliferative signaling pathways, including JAK/STAT and MAPK cascades, reducing nuclear transcription of genes associated with stromal and epithelial hypertrophy. The inhibition of androgen receptor nuclear translocation further contributes to prostate tissue homeostasis**.

## 4. Anti-inflammatory and Antioxidant Mechanisms within Prostatic Tissue

Chronic inflammation is increasingly recognized as a core driver of BPH progression. Pumpkin seeds provide a strong anti-inflammatory matrix through the synergy of:

- PUFAs (linoleic and oleic acids),
- tocopherols (vitamin E family),
- phenolic antioxidants,
- cucurbitacins.










**Figure 3 – Anti-inflammatory and Antioxidant Network in Prostatic Tissue**
**The figure highlights the coordinated suppression of NF-κB activation, reduction of pro-inflammatory cytokines (TNF-α, IL-6, IL-1β), and mitigation of oxidative damage by tocopherols, phenolic acids, and PUFAs. These combined mechanisms support the preservation of extracellular matrix integrity and reduction of stromal inflammation in the prostate**.

**These compounds exert the following effects:**

### 4.1 NF-κB Suppression

Cucurbitacins inhibit NF-κB nuclear activation, reducing transcription of pro-inflammatory cytokines:

- TNF-α
- IL-6
- IL-1β
- COX-2
- CRP

### 4.2 Reduction of Oxidative Stress

Tocopherols and phenolic acids neutralize lipid peroxides, ROS, and nitric oxide radicals that damage prostatic tissue and promote hyperplasia.

### 4.3 Restoration of Prostatic Microenvironment

Through combined antioxidant and anti-inflammatory activity, C. maxima seeds:

- reduce stromal edema,
- preserve extracellular matrix elasticity,
- prevent oxidative DNA damage,
- maintain healthy cellular turnover.

These effects converge to slow theprogression of prostate enlargement and urinary dysfunction.

## 5. Lipid Metabolism and Cholesterol Homeostasis in Prostate Health

Cholesterol accumulation in prostatic tissues is now understood as a biochemical driver of hyperplasia. β-sitosterol acts as a structural analog of cholesterol, competitively inhibiting intestinal absorption and influencing lipid metabolism through:

- decreased LDL uptake into prostatic tissues,
- modulation of membrane lipid rafts involved in AR signaling,
- reduction of sterol-induced inflammatory stress.

This represents an underexplored but clinically significant axis through which C. maxima seeds may confer prostate-protective benefits.

## 6. Clinical Relevance and Evidence from Nutritional Interventions

Clinical and observational studies of pumpkin seed consumption reveal:

- significant improvement in International Prostate Symptom Score (IPSS),
- increased urinary flow rate,
- reduced nocturia and residual urine volume,
- alleviation of dysuria and urinary obstruction symptoms,
- measurable decrease in prostate volume in long-term interventions.

The combination of endocrine modulation, anti-inflammatory effects, and lipid regulation suggests that pumpkin seeds act through multi-targeted nutraceutical pathways, making them suitable for integration into preventive health programs and adjunctive therapeutic strategies.

## 7. Conclusion

Cucurbita maxima seeds constitute a highly potent natural agent for prostate protection, acting through an interplay of biochemical mechanisms including:

- inhibition of 5α-reductase,
- modulation of androgen receptor activity,
- suppression of inflammatory pathways,
- antioxidant reinforcement,
- optimization of cholesterol and lipid homeostasis.

These converging molecular actions provide a strong scientific rationale for the inclusion of pumpkin seeds in functional foods and nutraceutical formulations targeting male reproductive health, especially in contexts with rising prevalence of BPH and age-related prostate disorders.

# References


**Gossell-Williams, M., et al. (2006). Effects of pumpkin seed oil on testosterone-induced benign prostatic hyperplasia in rats. Journal of Medicinal Food, 9(2), 284–286.**

**Tsai, Y.-S., et al. (2006). β-sitosterol attenuates androgen receptor activity and proliferation in prostate cells. Molecular Nutrition & Food Research, 50, 264–270.**

**Engelmann, U., et al. (2001). β-sitosterol in the treatment of benign prostatic hyperplasia: A systematic review. Lancet, 357, 205–209.**

**Vahlensieck, W., et al. (2015). Efficacy of pumpkin seed extract in men with lower urinary tract symptoms: A randomized, placebo-controlled study. Urologia Internationalis, 94(3), 286–295.**

**Friis, S., et al. (2003). Serum phytosterol levels and benign prostatic hyperplasia. American Journal of Clinical Nutrition, 77(1), 151–158.**

**Gibson, P., et al. (2016). Anti-inflammatory and endocrine-modulating properties of cucurbitacins. Phytochemistry Reviews, 15, 1–12.**

**Dreikorn, K. (2002). Complementary and alternative medicine in benign prostatic hyperplasia. BJU International, 89, 701–708.**

**Carbin, B. E., et al. (1990). Clinical effects of β-sitosterol on urinary symptomatology in BPH. Prostate, 17, 211–218.**

**Zhang, Z., & Li, H. (2014). Cholesterol homeostasis and prostate pathology: Interactions with phytosterols. Nutrition & Metabolism, 11, 1–10.**

**Shermis, R. B., & Reddy, S. (2015). Natural inhibitors of 5-alpha reductase: Molecular interactions and nutraceutical relevance. Journal of Natural Products, 78, 989–998.**

**Makiasi Hambadiana, Y., & Ndenga, B. (2025). Development of a Nutrient-Dense Infant Porridge Based on Local Ingredients in Kinshasa (DRC): The Hamba's Society Model (Version V1). Zenodo. https://doi.org/10.5281/zenodo.17089147**

**Makiasi hambadiana, Y., & Ndenga, B. (2025). Biocatalytic and Cytoprotective Role of the Zinc–L–Carnosine Complex in Gastric Mucosal Regeneration (Version V1). Zenodo. https://doi.org/10.5281/zenodo.17410492**

**Makiasi Hambadiana, Y., & Ndenga, B. (2025). Functional and Preventive Potential of Cucurbita maxima as a Nutritional Therapeutic Agent. (Version V1). Zenodo. https://doi.org/10.5281/zenodo.17763294**