

A de novo assembled high-quality chromosome-scale *Trifolium pratense* genome and fine-scale phylogenetic analysis

Zhenfei Yan

College of Grassland Science, Qingdao Agricultural University

Lijun Sang

College of Grassland Science, Qingdao Agricultural University

Yue Ma

College of Grassland Science, Qingdao Agricultural University

Yong He

College of Grassland Science, Qingdao Agricultural University

Juan Sun

College of Grassland Science, Qingdao Agricultural University

Lichao Ma

College of Grassland Science, Qingdao Agricultural University

Shuo Li

College of Grassland Science, Qingdao Agricultural University

Fuhong Miao

College of Grassland Science, Qingdao Agricultural University

Zixin Zhang

College of Grassland Science, Qingdao Agricultural University

Zengyu Wang

College of Grassland Science, Qingdao Agricultural University

Guofeng Yang (✉ yanggf@qau.edu.cn)

College of Grassland Science, Qingdao Agricultural University

Research Article

Keywords: *Trifolium pratense*, Genome, De novo assembly, PacBio HiFi, Genome annotation

Posted Date: April 29th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1582804/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

A de novo assembled high-quality chromosome-scale *Trifolium*

***pratense* genome and fine-scale phylogenetic analysis**

Zhenfei Yan^{1,2#}, Lijun Sang^{1,2#}, Yue Ma^{1,2}, Yong He^{1,2}, Juan Sun^{1,2}, Lichao Ma^{1,2}, Shuo Li^{1,2}, Fuhong Miao^{1,2}, Zixin Zhang¹, Zengyu Wang^{1,2*} and Guofeng Yang^{1,2*}

(¹College of Grassland Science, Qingdao Agricultural University, Qingdao 266109, China. ²Key Laboratory of National Forestry and Grassland Administration on Grassland Resources and Ecology in the Yellow River Delta, Qingdao 266109, China.)

Abstract

Background: Red clover (*Trifolium pratense* L.) is a diploid perennial temperate legume with 14 chromosomes (2n=14) native to Europe and West Asia, with high nutritional and economic value. It is a very important forage grass and is widely grown in marine climates, such as the United States and New Zealand. Genetic research and molecular breeding are limited by the lack of high-quality reference genomes. In this study, we used Illumina, PacBio HiFi, and Hi-C to obtain a high-quality chromosome-scale red clover genome and used genome annotation results to analyze evolutionary relationships among related species.

Results: The red clover genome obtained by PacBio HiFi assembly sequencing was 423 M. The assembly quality was the highest among legume genome assemblies published to date. The contig N50 was 13 Mb, scaffold N50 was 55 Mb, and BUSCO completeness was 97.9%, accounting for 92.8% of the predicted genome. Genome annotation revealed 44,588 gene models with high confidence and 52.81% repetitive elements in red clover genome. Based on a comparison of genome annotation results,

red clover was closely related to *Trifolium medium* and distantly related to *Glycine max*, *Vigna radiata*, *Medicago truncatula*, and *Cicer arietinum* among legumes. Analyses of gene family expansions and contractions and forward gene selection revealed gene families and genes related to environmental stress resistance, protein synthesis, and energy metabolism.

Conclusions: We report a high-quality de novo genome assembly for the red clover at the chromosome level, with a substantial improvement in assembly quality over those of previously published red clover genomes. These annotated gene models can provide an important resource for molecular genetic breeding and legume evolution studies. Furthermore, we analyzed the evolutionary relationships among red clover and closely related species, providing a basis for evolutionary studies of clover leaf and legumes, genomics analyses of forage grass, the improvement of agronomic traits.

Keywords: *Trifolium pratense*; Genome; *De novo* assembly; PacBio HiFi; Genome annotation

Background

Red clover (*Trifolium pratense* L.) (Figure 1a) is a temperate legume (*Leguminosae*) native to Europe and West Asia. It is not only cheap and rich in nutritional value but also contains a crude protein content of about 20%, with dry matter digestibility of about 70%, making it highly palatable and easily eaten by animals[1, 2]. Like alfalfa, red clover is self-incompatible and can be crossed with other subspecies[3, 4]. Its rich genetic diversity enables the production of many excellent agronomic traits. Red clover not only serves as a high-quality livestock feed but also has environmental and

medicinal value[5-8]. As an important feed for dairy and beef cattle, red clover has important implications for the dairy and beef industries[9-11]. Although the world is affected by the new crown rot epidemic, the number of beef and dairy cattle is increasing, with corresponding increases in demand for high-quality forage[4, 12]. Red clover also has implications for environmental protection as well as medicinal value. It can co-produce nitrogen with *Rhizobium*, reducing the need for nitrogen fertilizers[13, 14]. Its developed root system promotes the reproduction of soil microorganisms and the growth of surrounding trees[7, 15]. Red clover contains high amounts of anthocyanins and is a potential candidate for reducing inflammation and oxidative stress[16]. Despite these advantages of red clover, its disadvantages include flatulence in ruminants, cold tolerance, and root sensitivity to soil pH. Accordingly, the development of improved varieties by plant breeding is an important goal.

As a non-model legume, the structure of the red clover genome and important genetic information have not been fully determined, substantially limiting breeding and crop improvement. Relative to closely related species (e.g., white clover, alfalfa, and soybean)[17-19], progress in genomic research on red clover has been slow. The genome structure was initially explored by fluorescence in situ hybridization[20, 21]. Subsequently, simple repeat sequence (SSR) and single nucleotide polymorphism (SNP) markers were used to construct a red clover linkage genetic map[22]; however, few genes were mined, and the impact on variety improvement was limited. Subsequently, the red clover genome was constructed using the Illumina HiSeq 2000 platform; however, the final assembly results were insufficient for data mining in

terms of continuity and completeness[23]. Therefore, it is necessary to construct a high-quality red clover genome to accelerate genetic and genomic research on red clover and future breeding.

In this study, we obtained a high-quality chromosome-level red clover genome by using Illumina, PacBio HiFi, and Hi-C data. The assembly contained 96 scaffolds (~423 Mb), with N50 =55 Mb, accounting for 92.8% of the estimated genome size (456 Mb). Compared with that of the previously reported red clover genome assembly[23], the quality was greatly improved. From the genome, we annotated 44,588 genes. Furthermore, we evaluated gene family expansion and contraction and positive selection and performed phylogenetic analyses. In addition, analyses of the collinear relationship between the chromosomes of red clover and closely related species supported the accuracy and sensitivity of the data for studies of genomic evolution and for predictions of genomic patterns[24, 25]. Complete genomic information for the species and closely related species provides a basis for studying the differentiation and evolution of the species[26, 27]. This study provides a new starting point for evolutionary genomics research and a new research direction for analyzing the evolutionary relationships between red clover and related species.

Results

Genome-survey, sequencing, and assembly

The size, duplication rate, heterozygosity, and other parameters of the red clover genome were assessed[28]. After quality control, Illumina sequencing yielded 26 Gb of data. We randomly selected 10,000 clean reads for comparison with the NT library,

and the rate of successful alignment against plant genomes was 98.63%. A K-mer analysis further predicted a genome size of 456 Mb, with heterozygosity accounting for 1.57% and repeat sequences accounting for 54% of the genome.

We used traditional next-generation sequencing (NGS) data assembly methods to predict the genome size and third-generation HiFi sequencing (TGS) developed by PacBio for assembly of the red clover genome. In addition to making up for shortcomings of NGS in assembly applications, TGS does not require PCR amplification, produces ultra-long reads, and has no GC preference, which simplifies the process and improves the accuracy of assembly results[29-31]. High-quality HiFi reads were obtained after parameter alignment of the output data. In total, 19 Mbp of HiFi reads was generated, and the N50 length was 18 kbp.

Contigs were then generated according to the phased string graph[32]. The assembled genome (423 Mb) contained 194 contigs, with an N50 size of 13 Mbp, and the largest contig was 34 Mbp. The average GC content of the assembled genome was 33.6%, which was slightly lower than those of the previously reported genomes of *White lupin* (33.7%)[33] and *Pisum sativum* (37.6%)[34]. The Illumina reads were then aligned to the DNA library to assess the integrity and quality of the assembly. The concordant paired mapped alignment rate was 91.47%. Furthermore, the single-copy homologous gene pool used to assess genetic spatial integrity showed a BUSCO[35] of 97.9% for the assembled genome, highlighting its good completeness.

Scaffold construction and curation

Hi-C is a high-throughput chromosome conformation capture technology. Taking the

entire nucleus as the research object, this approach fixes and captures the interacting parts of the chromosome, followed by high-throughput sequencing to evaluate the spatial distribution of chromatin DNA throughout the genome and to obtain high-resolution chromosomal regulatory elements from positional relationships[36, 37]. In this study, we used 50 Gb of Hi-C data (100× coverage) to generate chromosome-level super-scaffolds. Subsequent analysis of the Hi-C library results showed that the genome was 423 Mbp with a scaffold N50 of 55 Mb. Compared with those of previous red clover sequence data (scaffold N50 = 223 kb), the quality and integrity were substantially better. After Hi-C-assisted assembly, 412 Mbp of genome sequence was mapped to seven chromosomes, accounting for 97.32% of the total sequence. Inter- and intra-chromosomal linkages were calculated to further verify the accuracy of the assembly. The linkages within chromosomes were much stronger than those between chromosomes, as revealed by the Hi-C heatmap. Furthermore, interactions were stronger for chromosomes at closer physical locations than at more distant physical locations (Figure 1b). These results demonstrate the high accuracy of the assembled genome. Table 1 summarizes the assembly information.

Genome annotation

Gene functions in the genome are inferred by computational homology-based alignments and the prediction of repeat sequences. In this study, we identified miniature inverted-repeat transposable elements (MITEs) and long terminal repeat (LTR) transposable elements. Structural features were used for prediction, and these elements accounted for 52.81% and 30.01% of the total sequences, respectively.

LTR-retrotransposons classified as Copia and Gypsy accounted for 11.26% and 7.70%, respectively, and 7605 simple repeats were found in the assembled genome. In addition, 16,227 ncRNAs were found and included 13 types.

Similarly, 44,588 high-confidence gene models and 46,089 transcripts were obtained using RNA-seq and de novo prediction strategies after removing the gene models containing early stop codons and frameshifts[38]. The gene models were unevenly distributed across the seven chromosomes.

The average lengths of genes and transcripts were 3620 bp and 1730 bp, respectively. Each gene contained an average of 1 transcript, and each transcript contained an average of 5 exons. The average lengths of CDS, exons, and introns were 1585 bp, 348 bp, and 495 bp, respectively. In addition, we compared the genome with those of five closely related species, including *Medicago truncatula* (MtrunA17r5.0-ANR from NCBI), *Trifolium medium* (ASM349008v1 from NCBI), *Vigna radiate* (ver6 from NCBI), *Cicer arietinum* (ASM33114v1 from NCBI), and *Glycine max* (v4.0 from NCBI). Among these, soybean had the largest genome (1012 Mb)[39], which was 2.4 times larger than the red clover genome. The *T. meduium* assembly had the largest number of genes (119102), *C. arietinum* the fewest (28772), and the other three closely related species had similar gene counts and average CDS lengths (Table 2). Using the NR, SwissProt, KEGG, GO, and eggNOG databases, 44018, 30158, 13365, 26349, and 38110 genes were annotated, and gene functions and counts were obtained. A dataset of 10,805 common genes was visualized by a Venn diagram (Figure 2, Table S1).

Gene family and evolution analysis

Comparing the genome of red clover with those of six representative species, a gene family analysis revealed that 44,588 genes clustered in 29,508 gene families. Among species, *T. medium* had the most gene families (5230). As visualized by a Venn diagram, 6602 gene families were shared by all species (Figure 3a). Furthermore, over 11.8 million years of differentiation between red clover and *T. medium*, 619 gene families in red clover expanded and 3 gene families contracted (Figure 3b). A GO enrichment analysis showed that the expanded gene families were related to the generation of energy compounds, defense reactions, protein synthesis, and energy metabolism. Notably, the most highly enriched GO functions were related to energy metabolism, which may be associated with the production of multiple compounds and plant stress resistance (Table S2). A GO enrichment analysis of genes with the signature of positive selection, as detected by the branch-site test, was also performed [40]. Most of the genes under positive selection were related to biosynthetic reactions (Table S3). We speculate that the expanded gene families and genes under positive selection may be involved in plant growth, metabolism, protein synthesis, and environmental stress responses, thus conferring a high protein content, fast growth, and strong stress resistance, which are important agronomic traits.

A phylogenetic tree was constructed based on single-copy homologous genes, with *Arabidopsis* (TAIR10.1 from NCBI) as the outgroup [41]. Red clover clustered with *G. max*, *V. radiata*, *M. truncatula*, *T. medium*, and *C. arietinum* to form a monophyletic group. Red clover was most closely related to *T. medium*, with an estimated

divergence time of about 12 million years ago. The collinearity of several closely related species was then evaluated, revealing that red clover and *M. truncatula* show a certain degree of synteny (Figure S1). The seven chromosomes of red clover and eight chromosomes of *M. truncatula* had good collinearity (Figure 4), indicating high conservation after species divergence.

Whole genome duplication (WGD) events are important indicators of plant evolution and are considered to be the driving force behind plant adaptation to various environments[42]. Using differences in synonymous substitution rates to detect gene duplication and loss in the red clover genome. The results showed that the divergence of red clover occurred prior to the WGD event shared by *M. truncatula* and *V. radiata*. Red clover experienced a common WGD event with *M. truncatula* and *V. radiata* with K_S values of 0.28 and 0.67, respectively (Figure 3c). In addition, a WGD event in red clover also corresponded to a K_S peak of 0.75 (Figure 3d).

Discussion

Genetic and genomic data for leguminous plants with excellent agronomic traits are important resources for the study of genetics, breeding, and functional omics. In this study, we assembled a high-quality diploid genome for a widely important forage crop, the autodiploid red clover. A highly contiguous and complete reference genome is necessary for detailed population genetics analyses and experimental studies. The new genomic data contribute to the legume genome resource bank and provide a good reference for further research on other crops in the clover genus. By combining state-of-the-art technologies (including PacBio HiFi, Hi-C, and high-quality genome

assemblers), we generated a representative red clover diploid genome. Compared with the previously reported red clover reference genome, the continuity and completeness of the newly assembled red clover genome showed a high quality: 13 Mbp contig N50, 55 Mb scaffold N50, and 97.9% BUSCO score. The assembly result was high-quality in comparison with legume genomes published to date. Notably, after combining high-throughput sequencing and Hi-C scaffolding, the total sequence length across all seven chromosomes was 412 Mbp, which was very similar to the size of the assembled genome (423 Mb). Genome annotation revealed 44588 genes with high confidence models, providing an important resource for molecular breeding and evolutionary studies.

Red clover belongs to the legume family, and its evolutionary history is not well-understood. Our analyses of gene family expansion and contraction and positive selection revealed the molecular basis underlying survival under geographical environment and climate change as well as its excellent agronomic traits. The genetic information provides very important resources for future research on legume improvement and stress-resistant species. According to a phylogenetic analysis, *Trifolium* species arose after the differentiation of *G. max*, *V. radiata*, *M. truncatula*, and *C. arietinum*, followed by the divergence of red clover and *T. medium*. Red clover and *Medicago truncatula* showed good genomic collinearity and were closely related based on the phylogeny and sequence similarity. The genome of red clover will help to clarify the evolutionary processes shaping legume species.

Conclusion

We report a high-quality chromosomal-level red clover genome assembled using third-generation HiFi sequencing. The newly generated genome has the highest coverage and integrity among published legume genomes. This study provides an excellent genetic resource for molecular breeding and the improvement of legumes and provides an important basis for research on the evolution of legumes.

Experimental procedures

For genomic DNA sequencing, the leaves of well-grown red clover were collected by a professional graduate student and were collected into vacutainer tubes. Red clover (2n=14) was grown in a light incubator at the Grassland Agri-husbandry Research Center. The study followed ethics norms and was in compliance with Chinese and international regulations.

DNA isolation and sequencing

The red clover Emerson was chosen as the test plant. Plants were grown in an incubator at the Qingdao Agricultural University in Shandong, China. After the leaves were treated with liquid nitrogen, DNA was extracted using the Tiangen DNA Secure Kit for Genome Sequencing (Beijing, China). Genome sequencing was performed by Berry Hekang (Beijing, China) using the third-generation PacBio HiFi assembly sequencing platform. The DNA samples were first subjected to quantity and quality control, followed by library preparation of the processed DNA and paired-end (PE) sequencing of the library using Illumina NovaSeq[30, 43, 44]. Sequence reads containing adapters, repeats, and low-quality reads (quality scores <20) were first filtered out. Then, 10,000 reads were randomly selected for NT library comparison

using the BLAST tool[45]. No apparent external contamination was detected. A K-mer analysis was used to estimate the genome size, duplication rate, and heterozygosity.

Genome assembly and quality evaluation

The quality of genomic DNA was checked using a NanoDrop 2000 spectrophotometer. The purified genome was used to construct the SMRTbell library and sequenced using PacBio SMRT technology[46, 47]. The library size was determined using an Agilent 2100 Bioanalyzer (Santa Clara, CA, USA). The acquired data were filtered and then processed using smrtlink for CCS processing. Reads were assembled using hifiasm, and purge-dups was used to remove hybridized fragments in the contig sequence[48, 49]. A single-copy orthologous gene library using a combination of tblastn, augustus, and hummer was finally used to assess the integrity of the assembled genome[35].

Hi-C data analysis and chromosome construction

The leaf tissue (100 mg) of red clover was soaked in paraformaldehyde, a cell cross-linking agent, for 15 minutes. Glycine was then added to the mixture to stop the chromatin cross-linking reaction, and the treated tissue was collected and frozen in liquid nitrogen. These tissues were then ground into a powder to extract DNA. Biotin-labeled oligonucleotide ends were added during end-repair, and the extracted DNA was subsequently fragmented into 350 bp fragments using a Covaris breaker[36, 50]. The biotin-conjugated DNA was captured and purified using avidin magnetic beads, and the library was constructed and sequenced using the Illumina PE150 platform[51, 52]. The raw reads were filtered, and 10,000 randomly selected

sequencing reads were aligned with the NT library using the BLAST tool. The Hi-C data and draft genome were then compared using JUICER[53]. The Hi-C library results were subsequently analyzed using 3D-DNA alignments to obtain valid Hi-C data and generate the chromosome-level scaffold of the red clover genome[54].

Functional annotation

Repeat sequences were analyzed and predicted using RepeatMasker, MITE Hunter, LTRharvest, LTR Finder, LTR retriever, and RepeatModeler, and MITES and LTR transposable elements were identified using structure-based prediction methods[55-60]. The software parameters for LTRharvest and LTR Finder were -similar 90 -vic 10 -seed 20 -seqids yes -minlenltr 100 -maxlenltr 7000 -mintsd 4 -maxtsd 6 -motif TGCA -motifmis And -D 15000 -d 1000 -L 7000 -l 100 -p 20 -C -M 0.9[61]. The parameters settings for RepeatModeler for the de novo identification of repeat sequences in masked genomes were -engine ncbi -pa 60. Similarly, the parameters for RepeatMasker for masking repeat sequences in the genome were -s -nolow -norna -gff -engine ncbi -parallel 20[57]. tRNAscan-SE was used to predict tRNA ab initio rRNA[62]. The Rfam database was used to search for other types of ncRNAs[63, 64]. Specific information was obtained by a similarity analysis.

Except for tandem repeats, all repeat regions were soft-masked and subsequently used for the annotation of protein-coding genes[65]. The coding sequences of *G. max*, *V. radiata*, *M. truncatula*, *T. medium*, and *C. arietinum* were downloaded. These coding sequences were subjected to Blast (v. 2.2.20) searches against the red clover genome[45]. Homologs containing premature stop codons and frameshifts were

discarded. Red clover -leaf RNA-seq data were aligned to red clover contigs using GeMoMa-1.6.1 and a comprehensive transcriptome database was built using PASA (v. 2.0.1)[66, 67]. Open reading frames were predicted using PASA (v. 2.0.1) and the resulting database was used to train parameters for the following four de novo gene prediction software packages: AUGUSTUS (v. 3.2.2)[68], GeneMarker-ET (v. 4.57)[69], GlimmerHMM (v. 3.0.2)[70], and SNAP[71]. Predictions obtained using these packages were then combined using EVM[72, 73], and 44588 genes were retrieved and functionally annotated by blast searches against databases, including NR, Swiss-Pro, eggNOG, GO, and KEGG[74-76]. A Venn diagram of the genes obtained from the five major databases was then generated to obtain more accurate annotation information.

Comparative analysis

A collinearity analysis of red clover and closely related species was performed using MCscan with parameters -g 1000 -c 90 -l 200[77]. Notably, the OrthoMCL cluster analysis was used to identify seven gene protein families (Red clover, *G. max*, *V. radiata*, *M. truncatula*, *T. medium*, *A. thaliana*, and *C. arietinum*)[78]. First, an all-vs.-all BLAST alignment of the protein-coding sequences of all of the above species was generated, and the similarity between the sequences was calculated[79]. The Markov clustering algorithm (expansion coefficient of 1.5) was used for protein family identification[42, 80, 81]. Single-copy genes in each species were selected as reference markers. Four-fold degenerate sites were selected to construct supergenes[82]. Multiple sequence comparisons of supergenes were subsequently

performed using Mafft[83, 84]. An appropriate base substitution model was selected and a species-based maximum likelihood (ML) phylogenetic tree was constructed using RAxML[85-87]. The mcmctree tool in the PAML software package was used to estimate the differentiation times based on single-copy gene families[88, 89]. The parameters were as follows: burn-in = 5,000,000, sample-number = 1,000,000, sample-frequency = 50. Time calibration points (correction points) were obtained from TimeTree (<http://timetree.org>)[90]. Then, the variation in gene family sizes among species was analyzed using Café, followed by a GO functional enrichment analysis of the expanded gene families[91]. The branch-site model was used to detect positive selection on specific branches and affecting a portion of sites by selecting one-to-one orthologous proteins from red clover and closely related taxa and using PRANK[40]. After aligning homologous protein sequences, Gblocks was used to filter the alignment results[92]. CODEML in PAML was used to evaluate positive selection in specific clades and affecting only certain loci with correction for multiple hypothesis testing using Chi2[89]. The duplicate age distribution was used to detect WGD events, and blastp was used to align the longest protein sequences for genes in the red clover genome[93]. The results were then filtered using MCScanX and synonymous substitution rates were calculated using the Yn00 tool in the PAML software package[94]. A density distribution map based on the K_s values for all paralogous gene pairs and K_s values of ortholog gene pairs between the genomes of red clover, *M. truncatula*, and *V. radiata* were then drawn using Matlab[95].

Abbreviations

NGS: Next-Generation Sequencing; CCS: Circular Consensus Sequencing; BUSCO: Benchmarking Universal Single-Copy Orthologs; Hi-C: High-through chromosome conformation capture; MITEs: miniature inverted repeat transposable elements; LTR: Long terminal repeat; LTR-RT: Long terminal repeat retrotransposons; ncRNA: Non-coding RNA; NR: NCBI nucleotide sequences; GO: Gene Ontology; KEGG: Kyoto Encyclopedia of Genes and Genomes; WGD: Whole genome duplications

Declarations

Acknowledgements

The author would like to thank Professor Guofeng Yang, Professor Zengyu Wang, and Professor Juan Sun (Professor of Grassland Science, Qingdao Agricultural University) for their help in data analysis and article writing. Thank you for the scientific research funding provided by the College of Grassland Science of Qingdao Agricultural University. Thanks for the experimental help provided by Beijing Berry and Kang.

Authors' contributions

ZY, ZW and GY conceived and designed this research. ZY analyzed data and wrote the manuscript. ZY, LS, YM and JS executed the data analyses. JS participated in the discussion of the results. LM, YH, FM and XZ collected samples. GY contributed to the evaluation and discussion of the results and manuscript revisions. All authors have read and approved the final version.

Funding

This study was supported by the National Nature Science Foundation of China (U1906201), Shandong Forage Research System (SDAIT-23-01), China Agriculture

Research System (CARS-34) and the First Class Grassland Science Discipline Program of Shandong Province China (1619002).

Availability of data and materials

All data generated and analyzed during this current study are available in the Grassland Agri-husbandry Research Center, Qingdao Agricultural University with permission from the Competent Authority. All sequencing data were submitted in NCBI Database having BioProject ID PRJNA765108 and details of software used are in Table S4. Biological materials used in this study available from the corresponding author.

Ethics approval and consent to participate

Red clover is not endangered or a protected species in China, and it was purchased from BEST grass industry and planted in a light incubator. The seeds are collected by Professor Guofeng Yang in BEST grass industry. All the study procedures were carried out in accordance with relevant guidelines.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹College of Grassland Science, Qingdao Agricultural University, Qingdao 266109, China. ²Key Laboratory of National Forestry and Grassland Administration on Grassland Resources and Ecology in the Yellow River Delta, Qingdao 266109, China.

1. Niderkorn V, Martin C, Rochette Y, Julien S, Baumont R: **Associative effects between orchardgrass and red clover silages on voluntary intake and digestion in sheep: Evidence of a synergy on digestible dry matter intake.** *Journal of animal science* 2015, **93**(10):4967-4976.
2. Farghaly MM, Youssef IMI, Radwan MA, Hamdon HA: **Effect of feeding Sesbania sesban and reed grass on growth performance, blood parameters, and meat quality of growing lambs.** *Trop Anim Health Pro* 2022, **54**(1).
3. Jones C, De Vega J, Lloyd D, Hegarty M, Ayling S, Powell W, Skot L: **Population structure and genetic diversity in red clover (*Trifolium pratense* L.) germplasm.** *Sci Rep* 2020, **10**(1):8364.
4. Riday H, Krohn AL: **Genetic map-based location of the red clover (*Trifolium pratense* L.) gametophytic self-incompatibility locus.** *TAG Theoretical and applied genetics Theoretische und angewandte Genetik* 2010, **121**(4):761-767.
5. Akbaribazm M, Khazaei F, Naseri L, Pazhouhi M, Zamanian M, Khazaei M: **Pharmacological and therapeutic properties of the Red Clover (*Trifolium pratense* L.): an overview of the new findings.** *J Tradit Chin Med* 2021, **41**(4):642-649.
6. Harlow BE, Flythe MD, Kagan IA, Goodman JP, Klotz JL, Aiken GE: **Isoflavone supplementation, via red clover hay, alters the rumen microbial community and promotes weight gain of steers grazing mixed grass pastures.** *PLoS one* 2020, **15**(3).
7. Nazarova EA, Nazarov AV, Egorova DO, Anan'ina LN: **Influence of destructive bacteria and red clover (*trifolium pratense* L.) on the pesticides degradation in the soil.** *Environmental geochemistry and health* 2022, **44**(2):399-408.
8. Wahdan SFM, Tanunchai B, Wu YT, Sansupa C, Schadler M, Dawoud TM, Buscot F, Purahong W: **Deciphering *Trifolium pratense* L. holobiont reveals a microbiome resilient to future climate changes.** *MicrobiologyOpen* 2021, **10**(4):e1217.
9. Moorby JM, Ellis NM, Davies DR: **Assessment of dietary ratios of red clover and corn silages on milk production and milk quality in dairy cows.** *J Dairy Sci* 2016, **99**(10):7982-7992.
10. Bertilsson J, Aerlind M, Eriksson T: **The effects of high-sugar ryegrass/red clover silage diets on intake, production, digestibility, and N utilization in dairy cows, as measured in vivo and predicted by the NorFor model.** *J Dairy Sci* 2017, **100**(10):7990-8003.
11. Hart EH, Onime LA, Davies TE, Morphew RM, Kingston-Smith AH: **The effects of PPO activity on the proteome of ingested red clover and implications for improving the nutrition of grazing cattle.** *Journal of proteomics* 2016, **141**:67-76.
12. Greenwood PL: **Review: An overview of beef production from pasture and feedlot globally, as demand for beef and the need for sustainable practices increase.** *Animal* 2021, **15** Suppl 1:100295.
13. Stefan A, Van Cauwenberghe J, Rosu CM, Stedel C, Labrou NE, Flemetakis E, Efroze RC: **Genetic diversity and structure of *Rhizobium leguminosarum* populations associated with clover plants are influenced by local environmental variables.** *Systematic and applied microbiology* 2018, **41**(3):251-259.
14. Janczarek M, Urbanik-Sypniewska T: **Expression of the *Rhizobium leguminosarum* bv. *trifolii* *pssA* gene, involved in exopolysaccharide synthesis, is regulated by RosR, phosphate, and the carbon source.** *Journal of bacteriology* 2013, **195**(15):3412-3423.
15. Duodu S, Carlsson G, Huss-Danell K, Svenning MM: **Large genotypic variation but small**

- variation in N₂ fixation among rhizobia nodulating red clover in soils of northern Scandinavia.** *Journal of applied microbiology* 2007, **102**(6):1625-1635.
16. Lee SG, Brownmiller CR, Lee SO, Kang HW: **Anti-Inflammatory and Antioxidant Effects of Anthocyanins of *Trifolium pratense* (Red Clover) in Lipopolysaccharide-Stimulated RAW-267.4 Macrophages.** *Nutrients* 2020, **12**(4).
 17. Griffiths AG, Moraga R, Tausen M, Gupta V, Bilton TP, Campbell MA, Ashby R, Nagy I, Khan A, Larking A *et al*: **Breaking Free: The Genomics of Allopolyploidy-Facilitated Niche Expansion in White Clover.** *The Plant cell* 2019, **31**(7):1466-1487.
 18. Chen H, Zeng Y, Yang Y, Huang L, Tang B, Zhang H, Hao F, Liu W, Li Y, Liu Y *et al*: **Allele-aware chromosome-level genome assembly and efficient transgene-free genome editing for the autotetraploid cultivated alfalfa.** *Nature communications* 2020, **11**(1):2494.
 19. Li YH, Zhou G, Ma J, Jiang W, Jin LG, Zhang Z, Guo Y, Zhang J, Sui Y, Zheng L *et al*: **De novo assembly of soybean wild relatives for pan-genome analysis of diversity and agronomic traits.** *Nature biotechnology* 2014, **32**(10):1045-1052.
 20. Sato S, Isobe S, Asamizu E, Ohmido N, Kataoka R, Nakamura Y, Kaneko T, Sakurai N, Okumura K, Klimenko I *et al*: **Comprehensive structural analysis of the genome of red clover (*Trifolium pratense* L.).** *DNA research : an international journal for rapid publication of reports on genes and genomes* 2005, **12**(5):301-364.
 21. Kataoka R, Hara M, Kato S, Isobe S, Sato S, Tabata S, Ohmido N: **Integration of linkage and chromosome maps of red clover (*Trifolium pratense* L.).** *Cytogenetic and genome research* 2012, **137**(1):60-69.
 22. Lopez-Maestre H, Brinza L, Marchet C, Kielbassa J, Bastien S, Boutigny M, Monnin D, El Filali A, Carareto CM, Vieira C *et al*: **SNP calling from RNA-seq data without a reference genome: identification, quantification, differential analysis and impact on the protein sequence.** *Nucleic acids research* 2016, **44**(19).
 23. De Vega JJ, Ayling S, Hegarty M, Kudrna D, Goicoechea JL, Ergon A, Rognli OA, Jones C, Swain M, Geurts R *et al*: **Red clover (*Trifolium pratense* L.) draft genome provides a platform for trait improvement.** *Sci Rep* 2015, **5**:17394.
 24. Cui F, Taier G, Li M, Dai X, Hang N, Zhang X, Wang X, Wang K: **The genome of the warm-season turfgrass African bermudagrass (*Cynodon transvaalensis*).** *Horticulture research* 2021, **8**(1):93.
 25. International Brachypodium I: **Genome sequencing and analysis of the model grass *Brachypodium distachyon*.** *Nature* 2010, **463**(7282):763-768.
 26. Gao L, Gonda I, Sun H, Ma Q, Bao K, Tieman DM, Burzynski-Chang EA, Fish TL, Stromberg KA, Sacks GL *et al*: **The tomato pan-genome uncovers new genes and a rare allele regulating fruit flavor.** *Nat Genet* 2019, **51**(6):1044-1051.
 27. Hubner S, Bercovich N, Todesco M, Mandel JR, Odenheimer J, Ziegler E, Lee JS, Baute GJ, Owens GL, Grassa CJ *et al*: **Sunflower pan-genome analysis shows that hybridization altered gene content and disease resistance.** *Nature plants* 2019, **5**(1):54-62.
 28. Marçais G, Kingsford C: **A fast, lock-free approach for efficient parallel counting of occurrences of k-mers.** *Bioinformatics* 2011, **27**(6):764-770.
 29. Nurk S, Walenz BP, Rhie A, Vollger MR, Logsdon GA, Grothe R, Miga KH, Eichler EE, Phillippy AM, Koren S: **HiCanu: accurate assembly of segmental duplications, satellites, and allelic variants from high-fidelity long reads.** *Genome research* 2020, **30**(9):1291-1305.

30. Li H, Durbin R: **Fast and accurate short read alignment with Burrows-Wheeler transform.** *Bioinformatics* 2009, **25**(14):1754-1760.
31. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK *et al*: **Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement.** *PloS one* 2014, **9**(11):e112963.
32. Cheng H, Concepcion GT, Feng X, Zhang H, Li H: **Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm.** *Nat Methods* 2021, **18**(2):170-175.
33. Hufnagel B, Marques A, Soriano A, Marques L, Divol F, Doumas P, Sallet E, Mancinotti D, Carrere S, Marande W *et al*: **High-quality genome sequence of white lupin provides insight into soil exploration and seed quality.** *Nature communications* 2020, **11**(1):492.
34. Kreplak J, Madoui MA, Capal P, Novak P, Labadie K, Aubert G, Bayer PE, Gali KK, Syme RA, Main D *et al*: **A reference genome for pea provides insight into legume genome evolution.** *Nat Genet* 2019, **51**(9):1411-1422.
35. Seppey M, Manni M, Zdobnov EM: **BUSCO: Assessing Genome Assembly and Annotation Completeness.** *Methods in molecular biology* 2019, **1962**:227-245.
36. Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I, Lander ES, Aiden AP *et al*: **De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds.** *Science* 2017, **356**(6333):92-95.
37. Mascher M, Gundlach H, Himmelbach A, Beier S, Twardziok SO, Wicker T, Radchuk V, Dockter C, Hedley PE, Russell J *et al*: **A chromosome conformation capture ordered sequence of the barley genome.** *Nature* 2017, **544**(7651):426-+.
38. Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M: **Gene prediction in novel fungal genomes using an ab initio algorithm with unsupervised training.** *Genome research* 2008, **18**(12):1979-1990.
39. Liu YC, Du HL, Li PC, Shen YT, Peng H, Liu SL, Zhou GA, Zhang HK, Liu Z, Shi M *et al*: **Pan-Genome of Wild and Cultivated Soybeans.** *Cell* 2020, **182**(1):162-+.
40. Yang JZNRZ: **Evaluation of an Improved Branch-Site Likelihood Method for Detecting Positive Selection at the Molecular Level.** *Molecular Biology & Evolution* 2005(12):2472-2479.
41. Lynch M, Conery JS: **The evolutionary fate and consequences of duplicate genes.** *Science* 2000, **290**(5494):1151-1155.
42. Berthelot C, Brunet F, Chalopin D, Juanchich A, Bernard M, Noel B, Bento P, Da Silva C, Labadie K, Alberti A *et al*: **The rainbow trout genome provides novel insights into evolution after whole-genome duplication in vertebrates.** *Nature communications* 2014, **5**.
43. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, Schatz MC: **GenomeScope: fast reference-free genome profiling from short reads.** *Bioinformatics* 2017, **33**(14):2202-2204.
44. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools.** *Bioinformatics* 2009, **25**(16):2078-2079.
45. McGinnis S, Madden TL: **BLAST: at the core of a powerful and diverse set of sequence analysis tools.** *Nucleic acids research* 2004, **32**(Web Server issue):W20-25.
46. Koren S, Rhie A, Walenz BP, Dilthey AT, Bickhart DM, Kingan SB, Hiendleder S, Williams JL, Smith TPL, Phillippy AM: **De novo assembly of haplotype-resolved genomes with trio**

- binning. *Nature biotechnology* 2018.
47. Quinlan AR: **BEDTools: The Swiss-Army Tool for Genome Feature Analysis**. *Current protocols in bioinformatics* 2014, **47**:11 12 11-34.
 48. Roach MJ, Schmidt SA, Borneman AR: **Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies**. *BMC bioinformatics* 2018, **19**(1):460.
 49. Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X, Chin CS *et al*: **Improved maize reference genome with single-molecule technologies**. *Nature* 2017, **546**(7659):524-527.
 50. Jarvis DE, Ho YS, Lightfoot DJ, Schmockel SM, Li B, Borm TJA, Ohyanagi H, Mineta K, Michell CT, Saber N *et al*: **Corrigendum: The genome of *Chenopodium quinoa***. *Nature* 2017, **545**(7655):510.
 51. Mascher M, Gundlach H, Himmelbach A, Beier S, Twardziok SO, Wicker T, Radchuk V, Dockter C, Hedley PE, Russell J *et al*: **A chromosome conformation capture ordered sequence of the barley genome**. *Nature* 2017, **544**(7651):427-433.
 52. Teh BT, Lim K, Yong CH, Ng CCY, Rao SR, Rajasegaran V, Lim WK, Ong CK, Chan K, Cheng VKY *et al*: **The draft genome of tropical fruit durian (*Durio zibethinus*)**. *Nat Genet* 2017, **49**(11):1633-1641.
 53. Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL: **Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments**. *Cell Syst* 2016, **3**(1):95-98.
 54. Bickhart DM, Rosen BD, Koren S, Sayre BL, Hastie AR, Chan S, Lee J, Lam ET, Liachko I, Sullivan ST *et al*: **Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome**. *Nat Genet* 2017, **49**(4):643-650.
 55. Gao L, McCarthy EM, Ganko EW, McDonald JF: **Evolutionary history of *Oryza sativa* LTR retrotransposons: a preliminary survey of the rice genome sequences**. *BMC genomics* 2004, **5**(1):18.
 56. Price AL, Jones NC, Pevzner PA: **De novo identification of repeat families in large genomes**. *Bioinformatics* 2005, **21 Suppl 1**:i351-358.
 57. Tarailo-Graovac M, Chen N: **Using RepeatMasker to identify repetitive elements in genomic sequences**. *Current protocols in bioinformatics* 2009, **Chapter 4**:Unit 4 10.
 58. Ouyang S, Buell CR: **The TIGR Plant Repeat Databases: a collective resource for the identification of repetitive sequences in plants**. *Nucleic acids research* 2004, **32**(Database issue):D360-363.
 59. Ou SJ, Jiang N: **LTR_retriever: A Highly Accurate and Sensitive Program for Identification of Long Terminal Repeat Retrotransposons**. *Plant Physiol* 2018, **176**(2):1410-1422.
 60. Han Y, Wessler SR: **MITE-Hunter: a program for discovering miniature inverted-repeat transposable elements from genomic sequences**. *Nucleic acids research* 2010, **38**(22):e199.
 61. Xu Z, Wang H: **LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons**. *Nucleic acids research* 2007, **35**(Web Server issue):W265-268.
 62. Storz G: **An expanding universe of noncoding RNAs**. *Science* 2002, **296**(5571):1260-1263.
 63. Nawrocki EP, Eddy SR: **Infernal 1.1: 100-fold faster RNA homology searches**. *Bioinformatics* 2013, **29**(22):2933-2935.
 64. Griffiths-Jones S, Bateman A, Marshall M, Khanna A, Eddy SR: **Rfam: an RNA family database**. *Nucleic acids research* 2003, **31**(1):439-441.

65. Korf I: **Gene finding in novel genomes**. *BMC bioinformatics* 2004, **5**:59.
66. Keilwagen J, Hartung F, Grau J: **GeMoMa: Homology-Based Gene Prediction Utilizing Intron Position Conservation and RNA-seq Data**. *Methods in molecular biology* 2019, **1962**:161-177.
67. Avram O, Kigel A, Vaisman-Mentesh A, Kligsberg S, Rosenstein S, Dror Y, Pupko T, Wine Y: **PASA: Proteomic analysis of serum antibodies web server**. *PLoS computational biology* 2021, **17**(1):e1008607.
68. Stanke M, Steinkamp R, Waack S, Morgenstern B: **AUGUSTUS: a web server for gene finding in eukaryotes**. *Nucleic acids research* 2004, **32**(Web Server issue):W309-312.
69. Holland MM, Parson W: **GeneMarker(R) HID: A reliable software tool for the analysis of forensic STR data**. *Journal of forensic sciences* 2011, **56**(1):29-35.
70. Majoros WH, Pertea M, Salzberg SL: **TigrScan and GlimmerHMM: two open source ab initio eukaryotic gene-finders**. *Bioinformatics* 2004, **20**(16):2878-2879.
71. Puma JE, Young M, Foerster S, Keller K, Bruno P, Franck K, Naja-Riese A: **The SNAP-Ed Evaluation Framework: Nationwide Uptake and Implications for Nutrition Education Practice, Policy, and Research**. *Journal of nutrition education and behavior* 2021, **53**(4):336-342.
72. Haas BJ, Salzberg SL, Wei Z, Pertea... M: **Automated eukaryotic gene structure annotation using EvidenceModeler and the Program to Assemble Spliced Alignments**. *Genome Biology* 2008, **9**(1):R7.
73. Minematsu K: **[Evidence-based medicine (EVM) for thrombolytic therapy during the ultra-acute stage of cerebrovascular diseases and its current status in Japan]**. *Nihon Naika Gakkai zasshi The Journal of the Japanese Society of Internal Medicine* 2004, **93**(9):1821-1826.
74. Xie C, Mao X, Huang J, Ding Y, Wu J, Dong S, Kong L, Gao G, Li CY, Wei L: **KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases**. *Nucleic acids research* 2011, **39**(Web Server issue):W316-322.
75. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R: **InterProScan: protein domains identifier**. *Nucleic acids research* 2005, **33**(Web Server issue):W116-120.
76. Syed A, Upton C: **Java GUI for InterProScan (JIPS): a tool to help process multiple InterProScans and perform ortholog analysis**. *BMC bioinformatics* 2006, **7**:462.
77. Tang H, Bowers JE, Wang X, Ming R, Alam M, Paterson AH: **Synteny and collinearity in plant genomes**. *Science* 2008, **320**(5875):486-488.
78. Li L, Stoeckert CJ, Jr., Roos DS: **OrthoMCL: identification of ortholog groups for eukaryotic genomes**. *Genome research* 2003, **13**(9):2178-2189.
79. Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA: **Circos: an information aesthetic for comparative genomics**. *Genome research* 2009, **19**(9):1639-1645.
80. Jiao Y, Wickett NJ, Ayyampalayam S, Chanderbali AS, Landherr L, Ralph PE, Tomsho LP, Hu Y, Liang H, Soltis PS *et al*: **Ancestral polyploidy in seed plants and angiosperms**. *Nature* 2011, **473**(7345):97-100.
81. Chen B, Silvestri GA, Dahne J, Lee K, Carpenter MJ: **The Cost-Effectiveness of Nicotine Replacement Therapy Sampling in Primary Care: a Markov Cohort Simulation Model**. *Journal of general internal medicine* 2022.
82. Vanneste K, Van de Peer Y, Maere S: **Inference of genome duplications from age**

- distributions revisited. *Molecular biology and evolution* 2013, **30**(1):177-190.
83. Nakamura T, Yamada KD, Tomii K, Katoh K: **Parallelization of MAFFT for large-scale multiple sequence alignments.** *Bioinformatics* 2018, **34**(14):2490-2492.
 84. Katoh K, Misawa K, Kuma K, Miyata T: **MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform.** *Nucleic acids research* 2002, **30**(14):3059-3066.
 85. Hohler D, Pfeiffer W, Ioannidis V, Stockinger H, Stamatakis A: **RAxML Grove: an empirical phylogenetic tree database.** *Bioinformatics* 2022, **38**(6):1741-1742.
 86. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A: **RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference.** *Bioinformatics* 2019, **35**(21):4453-4455.
 87. Stamatakis A: **RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.** *Bioinformatics* 2014, **30**(9):1312-1313.
 88. Yang C, Su XP, Liu DP, Guo ZW, Wang F, Lu YL: **A New Method of Aquatic Animal Personality Analysis Based on Machine Learning (PAML): Taking Swimming Crab Portunus trituberculatus as an Example.** *Front Mar Sci* 2020, **7**.
 89. Xu B, Yang ZH: **pamlX: A Graphical User Interface for PAML.** *Molecular biology and evolution* 2013, **30**(12):2723-2724.
 90. Kumar S, Stecher G, Suleski M, Hedges SB: **TimeTree: A Resource for Timelines, Timetrees, and Divergence Times.** *Molecular biology and evolution* 2017, **34**(7):1812-1819.
 91. Panda S: **The CaFe project: Optical <monospace>Fe II</monospace> and near-infrared <monospace>Ca II</monospace> triplet emission in active galaxies: simulated EWs and the co-dependence of cloud size and metal content.** *Astron Astrophys* 2021, **650**.
 92. Van der Straeten J, De Brouwer W, Kabongo E, Dresse MF, Fostier K, Schots R, Van Riet I, Bakkus M: **Validation of a PCR-Based Next-Generation Sequencing Approach for the Detection and Quantification of Minimal Residual Disease in Acute Lymphoblastic Leukemia and Multiple Myeloma Using gBlocks as Calibrators.** *Journal Of Molecular Diagnostics* 2021, **23**(5):599-611.
 93. Mahram A, Herbordt MC: **NCBI BLASTP on High-Performance Reconfigurable Computing Systems.** *Acm T Reconfig Techn* 2015, **7**(4).
 94. Wang YP, Li JP, Paterson AH: **MCScanX-transposed: detecting transposed gene duplications based on multiple colinearity scans.** *Bioinformatics* 2013, **29**(11):1458-1460.
 95. Nemati M, Tabari MMR, Hosseini SA, Javadi S: **A Novel Approach Using Hybrid Fuzzy Vertex Method-MATLAB Framework Based on GMS Model for Quantifying Predictive Uncertainty Associated with Groundwater Flow and Transport Models.** *Water Resour Manag* 2021, **35**(12):4189-4215.

Fig. 1 Plant morphology and Hi-C-assisted genome assembly of red clover. **(a)** Phenotype of the sequenced red clover plant. **(b)** Hi-C interaction heatmap showing 100-kb resolution super scaffolds.

Fig. 2 Venn analysis of five major databases(NR, Swiss-Prot, eggNOG, GO, KEGG) containing gene function annotation information.

Fig. 3 Gene family and phylogenetic tree analyses of red clover and other representative plant genomes. **(a)** Venn

diagram of the number of shared gene families. **(b)** A phylogenetic tree based on shared single-copy gene families (left), gene family expansions and contractions among red clover and seven other species (middle), and Gene family clustering in red clover and seven other plant genomes (right). **(c)** Genome-wide replication Ks distribution map of red clover and its related species. **(d)** Genome-wide replication Ks analysis of red clover .

Fig. 4 Features of the red clover and *M.truncatula* genome. **(a)** Length of each pseudochromosome (Mb). **(b)** Distribution of repetitive sequence. **(c)** Distribution of gene density. **(d)** Distribution of the GC content **(e)** Red clover and *M.truncatula* synteny analysis; the beginning of NC represents the chromosome of *M.truncatula*, while the beginning of CHR represents the chromosome of red clover.

Table 1 Summary statistic for the *Trifolium pratense* genome

Assembly		
Genome assembly	Estimated genome size	456Mbp
	Total length of assembly	423Mbp
	Number of contigs	194
	Contig N50	13Mbp
	Largest contig	34Mbp
	Number of scaffolds	96
	Scaffold N50	55Mbp
	Chromosome coverage (%)	97.32%
	GC content of genome	33.6%
Annotation		
Transposable elements	Total length	
	Total	224Mbp (52.81%)
	Retrotransposon	140Mbp (33.04%)
	DNA Transposon	37Mbp (8.69%)
Noncoding RNAs	Copies	
	rRNAs	13,053
	tRNAs	1,281
	miRNAs	477
	snRNAs	990
Gene models	Number of genes	44,588
	Mean gene length	3,620bp
	Mean coding sequence length	1,585bp

Table 2 The information of annotated gene models per species for all the species

Organism	Number of genes	Mean CDS length(bp)	Exons per transcript	Mean exon length(bp)	Mean intro length(bp)
<i>Vigna radiata</i>	29,006	1,430	7.6	293	449
<i>Glycine max</i>	54,881	1,391	8	295	413
<i>Trifolium medium</i>	119,102	306	1.4	219	172
<i>Cicer arietinum</i>	28,772	1,393	7.7	291	418
<i>Medicago truncatula</i>	36,079	1,428	6.9	324	393
<i>Trifolium pratense</i>	44,588	1,585	5	348	440

Figures

Figure 1

Plant morphology and Hi-C-assisted genome assembly of red clover. (a) Phenotype of the sequenced red clover plant. (b) Hi-C interaction heatmap showing 100-kb resolution super scaffolds.




Figure 2

Venn analysis of five major databases(NR, Swiss-Prot, eggNOG, GO, KEGG) containing gene function annotation information.

Figure 3

Gene family and phylogenetic tree analyses of red clover and other representative plant genomes. (a) Venn

diagram of the number of shared gene families. (b) A phylogenetic tree based on shared single-copy gene families

(left), gene family expansions and contractions among red clover and seven other species (middle), and Gene

family clustering in red clover and seven other plant genomes (right). (c) Genome-wide replication Ks distribution

map of red clover and its related species. (d) Genome-wide replication Ks analysis of red clover .

Figure 4

Features of the red clover and *M.truncatula* genome. (a) Length of each pseudochromosome (Mb). (b)

Distribution of repetitive sequence. (c) Distribution of gene density. (d) Distribution of the GC content (e) Red

clover and *M.truncatula* synteny analysis; the beginning of NC represents the chromosome of *M.truncatula*, while

the beginning of CHR represents the chromosome of red clover.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1TableS1.zip](#)
- [Additionalfile2TableS2.zip](#)
- [Additionalfile3TableS3.zip](#)
- [Additionalfile4TableS4.zip](#)

- [Additionalfile5FigureS1.zip](#)