



Unveiling the Diverse Medicinal Properties of *Murraya koenigii*

Shouvik Kumar Nandy  , Sattwik Das

[The author informations are in the declarations section. This article is published by ETFLIN in Sciences of Phytochemistry, Volume 2, Issue 2, 2023, Page 211-225. <https://doi.org/10.58920/sciphy02020107>]

Received: 17 August 2023

Revised: 17 September 2023

Accepted: 03 November 2023

Published: 07 November 2023

Editor: Samir Chtita

 This article is licensed under a Creative Commons Attribution 4.0 International License. © The author(s) (2023).

Keywords: *Murraya koenigii*, Phytochemistry, Pharmacological activity: Ethanobotany, Traditional medicine.

Abstract: Since the beginning of time, nature has provided medicines for treating fatal illnesses. Herbalists believe that certain plants have medicinal qualities. In tribal and rural communities, medicinal plants provide as an easily accessible source of treatment. The identification of various rejuvenating molecules that can halt or lessen the pathology of a variety of diseases will be regarded as a significant development of the present. There has been a scientific advancement in this area, and current studies on herbal medicines and traditional cures have attracted significant interest from all over the world. This plant, popularly known as the curry tree, is frequently used as herb and is also used to treat a variety of problems in traditional Indian medicine. About 80–85% of people worldwide rely on herbal products because they are thought to be efficient, secure, and cost-effective. Reviewing the plant taxonomy, ethnobotanical characteristics, folkloric or traditional uses, phytochemical, and pharmacological qualities of the *Murraya koenigii* plant was the goal of the current study. The leaves are used internally for dysentery as a carminative, tonic, stomachic, and inducer of vomiting. The usage of other parts includes treating piles, preventing helminthiasis, and reducing body heat, itchiness, and inflammation. Following several reports that this plant may treat a wide range of illnesses, scientists have worked to confirm the effectiveness of this plant by biological screening. A review of the literature suggests that *Murraya koenigii* has various medical benefits, including activity of antimicrobial, cardiac, antioxidative, antidiabetic and cholesterol reduction, cytotoxic action, antiulcer, and antidiarrhea.

Introduction

India is the habitat to more than 50,000 plant species, the majority of which are employed in folk and traditional herbalism. Many medicinal plants are used directly to treat illnesses or heal wounds, while some natural or pure compounds are consumed every day as a source of vital nutrients (1). For his basic requirements, such as food, clothing, and shelter, man uses plants in a variety of ways. For both urban and rural cultures, wild plants are the primary source of medicines, crafts, and cosmetics (2). Additionally, in rural regions, plants are the primary employers and sources of income (3). Plants have been utilised as remedies for thousands of years all across the world. According to WHO, around 80% of the population still relies on plant-based medications for primary care, especially in underdeveloped Nations. Siddha, Unani,

Ayurveda, and indigenous health civilizations are just a few of the medicinal systems used in India; they all make extensive use of herbs to cure human and animal illnesses (4). These therapeutic plants are crucial to us in another manner as well. India has a great variety of natural resources and a successful track record of traditional or old-style medicines; a sizable portion of the population still relies on plant-based medications for primary healthcare (5). The uses of medicinal plants are not only economical, but they also come with few or no adverse effects.

Murraya koenigii, commonly known as Curry, Kadi, Kari Patta, or Mitha Neem, is one such plant that is well-known in many nations, including India (6). *Murraya koenigii* is a native of Sri Lanka and India and is a member of the Rutaceae family. Curry leaves are a common ingredient in south Indian cuisine and have

been used to flavor food for a very long time. Utilizing medicinal plants has little to no negative effects and is also cost-effective. *Murraya koenigii* is also known by the common names Curry or Mitha Neem. Curry leaves are a common ingredient in south Indian cuisine and have been used to flavour food for a very long time (7). Vitamins A, B, C and E are abundant in this plant. Curry leaves are a good source of folic acid and iron, which fight anaemia (8). Curry leaves include a variety of phytochemicals that prevent cancer, treat liver damage, have neuro-protective capabilities, and fight against issues with the stomach, mouth, heart, and other organs (4, 8).

Table 1. Vernacular names of *Murraya koenigii* (8, 9, 10, 11).

Language	Commonly Known as
English	Curry leaves
Hindi	Karipatta, Mithanim
Bengali	Kartaphulli
Kannada	Karibevu
Gujarathi	Mitholimado
Tamil	Kariveppilai
Malayalam	Kariveppu
Marathi	Kadhilimb
Sanskrit	Girinimba
Telugu	Karepeku
Tulu	Bevusoppu
Portuguese	Folhas de caril
Russian	Listya karri
Spanish	Hojas de curry
Italian	Fogli di Cari
French	Feuilles de Cari
German	Curryblatter

Various names have been given to *Murraya koenigii*, including Karepaku in Andhra Pradesh, Kartaphulli in Bengal, Curry/Kari Patta in Hindi, and Curry Leaf in English. Karivempu in Tamilnadu, Kathnim and Karibevu in Karnataka, Narasingha in Assam, Gani, Gandhela and Gandla in Uttarakhand, Mitha Neemin Himachal Pradesh, Kariveppilei in Kerala, and Bhursangain from Orissa; Pindosine from Burmese; Gorenimbin, from Gujarat; Kerriebladeren from Dutch; Karry bald, from Danish; Curryblatter, from German; Daunkari, from Indonesia; Feuilles de curry from French; Hoja, from Spanish; and Fogli de Car, from Italian (Table 1). *Murraya koenigii* is one of the therapeutically significant herbs, and its taxonomy, ethnobotany, traditional use, and scientific significance are being evaluated (8). Therefore, the present review will describe and cover prior and current key works on *Murraya koenigii* linked to the themes chosen rather than covering a small number of carefully chosen

studies over a short period of time. The information will be methodically sorted, contrasted, and summarised, including phytochemical screening, identification, and pharmacological activity. We predicted that all of these efforts would result in a useful synthesis of pharmacological activity that would start off future views in the clearest possible way.

Taxonomy of Plant Kingdom

Murraya koenigii, a member of the Rutaceae family and commonly referred to as a "curry-leaf" tree, is a native of Sri Lanka, India, and other south Asian nations (10). It is native to practically all of India and has a distinctive perfume. It is a deciduous tree or shrub that may grow up to 6 meter tall and have trunk diameters of 15 to 40 cm with thin, smooth, brown or grey bark. (12). This plant's majority of parts emit a potent, off-putting odour. A species of tree known as *Murraya koenigii* is indigenous to the Asian tropical region, which extends from the Indian Himalayan foothills to Sri Lanka, via Indonesia, Myanmar, Southern China, and Hainan. The curry tree has bipinnately complex leaves that are 15–30 cm long, each carrying 11–25 leaflets alternate on rachis, and 2.5–3.5 cm long ovate lanceolate leaves.

The curry tree has grey colour bark with longitudinal striations, and beneath it, white colour bark is visible (13). Bisexual, funnel-shaped, white, sweetly scented, complete, stalked, irregular flowers with 2–3 mm long petioles and irregular edges have an average diameter when fully opened. 12 cm long terminal cymes with 60–90 blooms each. The ovoid, rough, or wrinkled fruits have glands, (Table 2) (10, 13).

Table 2. *Murraya koenigii*'s plant taxonomy (8, 10, 11).

Kingdom	Plantae
Subkingdom	Tracheobionta
Superdivision	Spermatophyta
Division	Magnoliophyta
Class	Magnoliopsida
Subclass	Rosidae
Order	Sapindales
Family	Rutaceae
Genus	<i>Murraya koenigii</i> ex L.
Species	<i>Murraya koenigii</i> L.

Ethnobotanical Use of *Murraya koenigii*

In Eastern Asia *Murraya koenigii* has countless important uses in the traditional system of medicine. Based on ethno-medicine, it is used as an anti-dysenteric, anti-diabetic and stimulant (14). In Indian

this plant is extremely valued for its leaves as an important ingredient to promote digestion and appetite. The leaves, roots and barks are tonic, carminative and stomachic (15). Leaves are used in dysentery also check vomiting. Steam distillate of leaves can be used as anti-anaemic, febrifuge stomachic and purgative.

Externally, leaves are applied to eruption and bruises. The leaves, roots are bitter in test, acid, cooling, analgesic, anti-helminthic, allays heat of the body, thirst, itching and inflammation, it also cures piles (16, 17). Leuco-derma and blood diseases can both benefit from it. To stop vomiting, use the toasted leaves. Root juice is beneficial for treating kidney-related discomfort. Fruits are regarded as astringent in Indo-China (18). Crushed leaves are applied topically to treat skin eruptions and soothe burns. To cure snake bites from deadly animals, leaf pastes that have been crushed are administered. For flavouring curries, egg, meat, and fish meals, traditional curry powder, etc., people mostly utilise fresh leaves, dried leaf powder, and essential oil (19). To create cosmetics and soaps, the aromatherapy sector uses essential oils (20). The nutritional value of the fruits is very high. The branches of this plant are frequently used as datun to clean teeth. In **Table 3**, the ethno-botanical profile is presented.

Phytochemical Study

Alkaloids, flavonoids, polyphenols, and terpenoids have been isolated from curry leaves, stems, bark, roots, and various plant extracts (31). The leaves contain 63.2% moisture, 8.8% protein, 39.4% carbohydrates,

1.15% total nitrogen, 6.15% fat, 18.92% total sugar, 14.6% starch, and 6.8% crude fiber (32, 33). Curry leaves are also rich in vitamins and minerals, including calcium, magnesium, sodium, and vitamin A (β -carotene) at 6.04 ± 0.02 mg/100 g, vitamin B3 (niacin) at 2.73 ± 0.02 mg/100 g, and vitamin B1 (thiamin) at 0.89 ± 0.01 mg/100 g (34).

The main chemical components of *Murraya koenigii*, along with their plant parts, are listed in Table 4 and Figure 1 (35, 36). Phytochemical screening of ethanol, petroleum ether, chloroform, aqueous, and ethyl acetate extracts confirmed the presence of alkaloids, carbohydrates, flavonoids, and sterols (37). Several experiments verified the phyto-constituents in curry tree extracts. When alkaloids were tested with chloroform, petroleum ether, alcohol, ethyl acetate, and water extracts, Mayer's reagent confirmed their presence by forming cream- or white-colored precipitates (38). Phenolic compounds were identified by adding lead acetate (5%) to an alcoholic root extract, which produced a white precipitate. Flavonoids were detected by dipping yellow filter paper into aqueous and alcoholic extracts with ammonia (39).

Saponins were indicated when the extract produced honeycomb-like foaming upon shaking with sodium bicarbonate. Free amino acids and proteins were confirmed using Biuret's, Millon's, and Ninhydrin's tests (36, 37). The hydro-alcoholic extract was mixed with CHCl_3 , acetic anhydride, and concentrated H_2SO_4 , producing a blue to brick-red color, confirming triterpenes and sterols (33). The plant exhibits a bitterness level of 2.5 units/g and possesses hemolytic properties. Carbohydrates and amino acids were detected in both aqueous and alcoholic extracts (39).

Table 3. Ethnobotanical use of *Murraya koenigii*.

No.	Used plant parts	Folk/ Ethnobotanical uses	References
1.	Leaves	Anti-anaemic, Anti-helminthic, Analgesic, Anti-ulcer, Anti-nociceptive Anti-amnesic, Anti-inflammation, cooling and itching, Stomachic, Purgative, Febrifuge, Hair tonic Stimulant of hair growth, Night blindness, Vomiting, Bruises and Eruption, Bites of poisonous animals, Hypercholesterolemia lightening, maintaining the natural skin, enhancing memory, lighting and rough skin improving, Pigmentation and showed skin, help to weight loss, to Enhance Appetite and digestion	(21, 22, 23)
2.	Whole plant	Stimulant, Blood-purifier, Hair tonic, Antidepressant, Anti-dysenteric, Antidiarrheal, Antifungal, Anti-inflammatory, Antiemetic, Febrifuge, Stomachic, Anti-periodic, Anti-diabetic, Prevent body aches, Kidney pain and Vomiting	(24, 25)
3.	Stem	Strengthen, Datum for cleaning, gums and teeth	(26, 27)
4.	Bark	Hair tonic, Carminative and Stomachic	(28)
5.	Fruits	Astringent	(11)
6.	Roots	Analgesic, Anti-helminthic, Cooling agent, Kidney pain, Blood disorders, Itching, Inflammation	(29, 30)




Figure 1. List of phytochemicals present in *Muraya koenigii* (16, 33, 40).

Folk/Traditional Use

Essential oils, fresh leaves, and dried leaf powder from *Muraya koenigii* are widely incorporated into various culinary preparations, including seafood, soups, meat dishes, eggs, and curries, enhancing both flavor and aroma. Beyond their role in cooking, these essential oils are highly valued in the soap and cosmetic aromatherapy industries for their therapeutic and fragrant properties (46). In hair care, curry leaves serve as a natural remedy for maintaining healthy hair and promoting hair growth, commonly used as a nourishing hair tonic (47). Traditionally, the leaves are boiled in coconut oil until a concentrated residue forms, which is then applied to the scalp to reduce hair fall, strengthen roots, and prevent premature graying.

Beyond their culinary and cosmetic benefits, *Muraya koenigii* holds a significant place in traditional medicine. Various parts of the plant have been used for centuries to treat a range of ailments, including nausea, vomiting, blood purification, depressive disorders, fungal infections, body aches, and diarrhea, as illustrated in Figure 2 (48). Indigenous communities have long relied on the bark and roots for treating

venomous bites, while the green leaves are consumed raw to alleviate morning sickness and gastrointestinal discomfort. Root juice is often applied to boils to promote healing, whereas leaf paste is used to relieve kidney pain (49). Additionally, the combination of raw green curry leaves and lime juice is a popular home remedy for diarrhea and nausea, believed to aid digestion and soothe the stomach (50).

In Ayurvedic medicine, dried curry leaf powder is mixed with honey and betel nut juice and consumed as an anti-periodic remedy to prevent recurring illnesses (51, 52). The plant is also a rich source of essential nutrients, including iron, vitamins A, B, B2, and C, making it particularly beneficial for individuals suffering from anemia or calcium deficiencies. Due to its diverse pharmacological properties, curry leaves are widely recognized for their antifungal, antidepressant, anti-diarrheal, anti-inflammatory, and blood-purifying effects. These healing properties make *Muraya koenigii* an essential component of various traditional and holistic health practices, where it is either used alone or combined with other medicinal plants to enhance its efficacy (53, 54).

Table 4. The major chemical constituents of *Murraya koenigii*.

Classification	Name of Components	Plant Part	References		
Alkaloids	Mahanine	Stem, bark, leaves and seeds	(16, 33, 40)		
	Mahanimbine	Stem, root, leaves and seeds			
	Murrayanol Koenimbine	Leaves, fruits and roots			
	O-Methylmurrayamine A Koenigicine	Leaves			
	Koenigine	Stem, bark and leaves			
	Murrayone (Coumarine) Mahanimbicine Bicyclomahanimbicine Phebalosin	Leaves			
	Isomahanimbine Koenimbidine	Leaves and roots		(11, 33, 40)	
	Euchrestine B Bismurrayafoline E	Leaves			
	Isomahanine	Leaves, fruits and seeds			
	Mahanimbinine	Leaves and seeds			
	Girinimbilol	Leaves			
	Pyrayafoline-D	Leaves, stem bark			
	Glycozoline Cyclomahanimbine Isomurrayazoline Mahanimboline Mukonicine Isolongifolene	Leaves			
	Mukonal Mukeic acid	Stems			
	9-Carbethoxy-3-methyl carbazole 9-Formyl-3-methyl carbazole	Roots and stems			
	Murrayazolinol Mahanimbinol Mukoeic acid Osthol Umbelliferone Murrayanine	Stems bark	(36, 37, 41)		
	Mukoenine-A Mukoenine-B	Roots, stem bark			
	Mukoline	Roots			
	Mukolidine (M)-murrastifoline-F	Stem bark and roots			(40, 41)
	3-Methyl-9H-carbazole-9-carbaldehyde	Roots			
	Bismahanine Bikoeniquinone A Bismurrayaquinone	Stem bark and roots	(36)		
	3-Methylcarbazole Murrayafoline A	Roots			
	Murrayakonine A Murrayakonine B Murrayakonine C Murrayakonine D	Stem and leaves	(40-41)		
	Girinimbine	Stem bark, roots and seeds			
	Murrayacine Murrayazoline	Stem			

Flavonoids	Quercetin Apigenin Kaempferol Rutin Catechin Myricetin 4-O-β-D-Rutinosyl-3-methoxyphenyl-1-propanone 1-O-β-D-Rutinosyl-2(R)-ethyl-1-pentanol 8-Phenylethyl-O-β-D-rutinoside		(42, 43)
Terpenoids	Blumenol A Icariside B1 Loliolide Blumenol A Icariside B1 (-)-Epiloliolide (-)-α-pinene (-)-β-pinene (+)-β-pinene (+)-sabinene	Leaves	(43, 44, 45)
	Squalene		(43)
	β-sitosterol	Bark and leaves	
Polyphenols	Selin-11-en-4α-ol		(44-45)
	2-hydroxy-4-methoxy-3,6-dimethylbenzoic acid	Bark	




Figure 2. Pharmacological activity and traditional use of *Murraya koenigii* (59, 62).

Pharmacological Activity

Researchers have discovered a number of pharmacological activities present in *Murraya koenigii*, as this plant contains several phytochemicals with varied compositions. Curry plant also shows pharmacological activities such as Neuro-protective activity through decreasing Glycemic levels, Anti-Nephrotoxicity activity against unilateral renal ischemia (55), it also exhibits decreased GI motility (Anti-Diarrheal Activity), anti-ulcer activity by working against lesion index, *Murraya koenigii* prevents tooth decay, stimulates digestive enzymes, lowers total cholesterol, acts as an anti-pyretic by reducing fever, acts as a memory enhancer, and promotes wound healing (56, 57). Curry tree exhibits insecticidal, phagocytic, anti-helminthic, hypercholesterolemic, antinociceptive, analgesic, and vasodilatory activities (see **Table 5** and **Figure 2**) (58).

In Vitro Studies

Antioxidative Activity

According to the literature, various solvents were used to study the antioxidative capabilities of the *Murraya koenigii* leaf extract. Natural anti-oxidants derived from plants have been touted as a viable therapy for the treatment and prevention of many illnesses, particularly cancer, cardiovascular disease, and other illnesses as well as neurological disorders (59). *Murraya koenigii* contains a number of naturally occurring bioactive substances that have outstanding antioxidant activities, including mahanine, mahanimbine, isolongifolene, koenimbine, girinimbine, isomahanine, koenoline, and O-methylmurrayamine (60). They were assessed based on their radical scavenging capacity against 1,1-diphenyl-2-picrylhydrazyl (DPPH) and the oil stability index (OSI) (61). In comparison to those of α -tocopherol, the OSI values of the methylene chloride (CH_2Cl_2) extract and the ethyl acetate (EtOAc) soluble portion of the 70% acetone extract were significantly prolonged. Three categories were used to group the 12 carbazoles (62). Based on the oil stability index (OSI) and their ability to scavenge radicals against 1, 1-diphenyl-2-picrylhydrazyl, the antioxidative capabilities of *Murraya koenigii* leaf extracts employing various solvents were assessed. Two carbazole alkaloids, mahanimbine and koenigine, were isolated from the leaves of *Murraya koenigii* and demonstrated antioxidant activity. Additionally, Koenigine demonstrated strong anti-radical effects (63).

Antimicrobial Activity

Several investigations have shown that the plant itself has a number of active compounds with antimicrobial properties, including Mahanimbine, Murrayanol, Mahanine, and Gurjunene (64). The stem bark of *Murraya koenigii* was used to isolate benzoisofuranone derivatives, three known steroids, and six known

carbazole alkaloids. These substances are proven to be efficacious at concentrations between 3.13 and 100 g/ml. *In vitro* antibacterial activity against Gram positive and Gram negative strains of bacteria was tested using methanolic extracts of 21 different plant species, according to a literature review. A study revealed that *Murraya koenigii* had the strongest antibacterial effect. Curry leaf plant greatly reduced *Staphylococcus epidermidis* (29, 65). Three carbazole alkaloids-mahanine, murrayanol, and mahanimbine were discovered in the acetone extract of Curry fresh leaves. Mahanimbine demonstrated antioxidant activity at 33.1 g/ml, whereas murrayanol had an IC₅₀ of 109 g/mL against hPGHS-1 and an IC₅₀ of 218 g/mL against hPGHS-2 in anti-inflammatory tests. These three carbazole alkaloids all had antibacterial and mosquito-killing properties as well as topoisomerase I and II inhibitory activities (66).

Anticancer Activity

Extracts from various plant parts have been shown through methodical study and pharmacological examination to have antiviral, anti-inflammatory, antioxidant, antidiabetic, anti-diarrheal, anti-leishmanial, and anticancer activity (67). Among a number of different chemical components found in plant extracts, the main bioactive component of the carbazole alkaloid mahanine has been identified (16). The chemopreventive as well as the therapeutic potential of this plant and its active ingredients against a variety of malignancies are demonstrated through the presentation of research articles in this review (68). By doing an *in vitro* cell line study, several studies reveal MK's anticancer properties. The *in vitro* anti-tumor promoting action of a pure chemical, girinimbine, which was isolated from the stem bark of *Murraya koenigii*, was demonstrated by measuring the percentage suppression of the induced early antigen EA of Epstein Barr virus EBV on the surface of Raji cells (69, 70). According to a study, a polyphenol-rich hydro-methanolic extract of MK leaves (CLE) reduced cell viability, growth kinetics, and arrested the S phase of the cell cycle in MCF-7 and MDA-MB-231 breast cancer cells, therefore inhibiting 26S proteasome proteolytic activity and inducing apoptosis. According to this study, MK leaves are a powerful source of proteasome inhibitors and may be helpful in the treatment of many malignancies (71). Despite the beneficial outcomes of the *in vitro* and *in vivo* research, very few pre-clinical trials have been carried out to investigate the anticancer effect of phytochemicals. Because cancer is a persistent disease, it is necessary to investigate how it affects people through clinical trials in order to introduce this product as an anticancer drug (72).

Antifungal Activity

Several investigations have reported *Murraya koenigii*'s antifungal activity by resisting the growth of gram-

positive and gram-negative bacteria. For instance, it has been claimed that the leaves' essential oil has antifungal properties (73). The presence of phytochemical components with complex molecular structures and a variety of action mechanisms, such as alkaloids, terpenoids, flavonoids, phenolics, tannins, and saponins, which are known for their antimicrobial properties, is what gives *Murraya koenigii* leaves their antifungal properties (74). Various studies back up the plant's historic use as an antifungal agent. Curry leaves' use in traditional medicine for the treatment of diarrhoea, dysentery, and skin eruptions may be explained by their *in vitro* antifungal action (75).

The ethanolic extract of *Murraya koenigii* had noticeable impacts on the morphology of the hyphae, including an increase in branching potential that led to the growth of brief, slender hyphal branches with inflated terminals (76). The current study comes to the conclusion that these plants contain a variety of chemical components that could be useful in pharmacology. It can be utilised to raise society's health level because it contains a variety of substances that are necessary for excellent health. Both plants' methanolic extracts exhibited a modest amount of antibacterial activity against the examined microorganisms (77).

In Vivo Studies

Antidiabetic Activity

Currently, the major goal of treating diabetes is to reduce hyperglycemia over time by using a combination of insulin, conventional therapy, and α -glucosidase inhibitors (78). However, the effectiveness of these chemicals is debatable due to unintended side effects, which increases the possibility for alternative treatments in the management of diabetes (79, 80). For this reason, it has been argued that plants represent a vast, yet largely untapped source of potentially effective antidiabetic medication. Statistically considerable hypoglycemia potential is present in *Murraya koenigii* in STZ-induced rats with diabetes (81). The *Murraya koenigii* extract appeared to be more efficient than the well-known drug antidiabetic drug glibenclamide. The petroleum ether extract of the dried plant was used to isolate mahanimbine, a chemical component of *Murraya koenigii*.

On streptozotocin-induced wistar rats, the anti-diabetic action was tested using a pure substance at doses of 50 mg/kg and 100 mg/kg. Mahanimbine may lower blood sugar levels by potentiating the effects of insulin, either by boosting pancreatic insulin secretion from beta cells of the islets of Langerhans or by enhancing peripheral glucose uptake (47). When compared to acarbose, mahanimbine had a noticeable α amylase inhibitory effect (82). However, given

that plant substances act more slowly than synthetic drugs and that higher dosages may have a plateau effect, which would be harmful to the treatment of diabetes, long-term research is necessary (3, 81). It will need more research to extract and pinpoint the precise active ingredients that give the examined plant materials their antidiabetic properties (11, 82).

Antiulcer Activity

One of the most prevalent gastrointestinal diseases is the peptic ulcer. *Murraya koenigii*'s aqueous extract has anti-ulcer properties (83). The antiulcer properties of *Murraya koenigii*'s aqueous and ether extracts were investigated in albino rat models of stomach ulceration brought on by reserpine (4). Using hot aqueous leaf extract at dosages of 250 and 400 mg/kg, the anti-ulcer efficacy was seen. The extract prevented stomach lesions brought on by pylorus ligation and anti-inflammatory, non-steroidal medicines. In a pylorus ligation model, the extract decreased gastric volume, ulcerative lesions, free and total acidity but increased the pH of gastric juice (79). The findings revealed that the extract has strong anti-ulcer properties (47). Extracts appeared to be as protective as ranitidine in treating stomach ulcers. The anti-ulcer activity of a crude aqueous extract of leaves was assessed using rat models of acute gastric lesions brought on by ethanol, aspirin, cold restriction stress, and pylorus ligation. These findings support previous research suggesting that an aqueous extract of *Murraya koenigii* leaves can be an effective antiulcer medication (84).

Anti-Inflammatory Activity

The leaf extracts of *Murraya koenigii* have anti-diabetic properties in addition to having some effects that control immunology in relation to oxidative stress metabolism. Expression of the cytokines interleukin (IL)-2, 4, 10, and tumour necrosis factor alpha (TNF-alpha) demonstrated this immune-modulatory and anti-inflammatory action (85). Various studies reveal that the anti-inflammatory and analgesic activity of methanol extract of dried *Murraya koenigii* Linn leaves given orally to healthy animals at doses of 100, 200, and 400 mg/kg body weight. Carrageenan-induced hind paw edoema in albino rats was used to test the extract's anti-inflammatory action. At various time points following injection of carrageenan (1% w/v), the mean increase in paw volume and inhibition in paw volume were determined plethysmometrically. The effectiveness of the extract as an analgesic was also examined in albino rats using the Eddy's hot plate method and the formalin-induced paw licking method. The carrageenan-induced paw edoema was significantly reduced (P 0.001) by the methanol extract, and analgesic effectiveness was demonstrated by an increase in reaction time using the eddy's hot plate method and a percentage increase in pain during

the formalin test. Comparing the methanol extract to the standard and reference medicine, diclofenac sodium (10mg/kg), the anti-inflammatory and analgesic effects were dose-dependent. Statistics showed that these inhibitions were significant ($P < 0.05$). Thus, our research implies that curry plant may be useful in the treatment of diseases that are related to inflammatory pain (2).

Antihelmintic Activity

When compared to the common medication Piperazine, the *Murraya koenigii* leaves' ethanolic and aqueous extract demonstrated anti-helmintic action, various research data reveals that it is thought that the polyphenolic component tannins, which are contained in the leaves, are what give them their antihelmintic properties. The methanolic extract has antihelmintic activity against Indian earthworm in a dose-dependent manner, such that it can paralyse the worm in 18 minutes and have a deadly effect in 45 minutes (86).

Wound Healing Activity

Mahanine, Mahanimbine, and Mahanimbicine are only a few of the phytochemicals found in *Murraya koenigii*'s leaves, stem, and bark's ethanol extract that

have wound-healing properties (45).

Anti-Amnestic Activity

After being treated with *Murraya koenigii* leaves, rats using behavioural models of scopolamine, aging-induced forgetfulness, and diazepam were assessed for memory improvement (87). Several groups were given plants-leaf powder mixed with wheat flour by mouth, and their behaviour was assessed using elevated plus-maze and Hebb-Williams maze trials. The memory scores of both young and old rats showed a significant dose-dependent improvement, and the induction of scopolamine (0.4 mg/kg, i.p.) and diazepam (1 mg/kg, i.p.) in the treatment groups may have contributed to the significant reduction in forgetfulness (88).

Antidiarrhoeal Activity

Three pure compounds of bioactive carbazole alkaloids kurryam, koenimbine, and koenine were isolated from the n-hexane extract of *Murraya koenigii* seeds under the guidance of a bioassay. Kurryam and koenimbine, two of the three substances, significantly inhibited rats' castor oil-induced diarrhoea and PGE2-induced enter pooling (89). Also, the substances significantly decreased Wister rats' gastro-intestinal motility in the test using charcoal meals (90).

Table 5. Pharmacological use of *Murraya koenigii*.

No.	Uses	Phytochemicals	Pharmacological action	Reference
1.	Anti-Diabetic	Koenimbidine, Murrayazoline, Murrayacine	Decreases Oxidative Stress	(91)
2.	Anti-Trichomona	Girinimbine, Mahanimbilo	Doing Against Trichomonas Gallinae	(69, 92)
3.	For Oral Health	Essential Oil	Stimulating Salivation Process	
4.	Vasodilation	Mahanimbilol, Murrayazoline	Work as negative Chronotropic Effect	
5.	Anti-Oxidant	Mahanimbine, Koenigine	Reduction in Hepatic Malondialdehyde In the Kidney	
6.	Anti-Cancer Activity	Girinimbine, Mahanine, Mahanimbine, Murrayafoline	Proliferation Death of Cancer Cell Proteasome Inhibitor	(90)
7.	Bronchial Disorders	Girinimbine, Mahanine	Block 5-Lipoxygenase Activity	
8.	Effect in Dental Caries	Isomahanine, Mahanine and Murrayanol	Inhibit of Cavity Formation	(92)
9.	Anthelmintic Activity	Mahanine, Koenimbidine	Cause Paralysis	(69, 92)
10.	Effect on Wound Healing	Mahanine, Mahanimbine, Essential Oil, Mahanimbicine	Work Against Inflammatory Cells And reduce Collagen Deposition	(93)
11.	Anti-Amnesic	Koenimbidine, Mahanimbicine	Protect Against the Neurodegenerative Diseases	(11)
12.	Improves Eyesight	Essential Oil and Vitamin A	Eye Sight Improvement	(92)
13.	Radiation Protection Activity	Mahanine, Murrayafoline	Increases Glutathione, Including Enzymes Levels and Decrease Chromosomal Damage	(83)
14.	Anti-Ulcer Activity	Mahanimbine And Essential Oil	Effect against Lesion Index, Area and Percentage of Lesion and On Ulcer	(6)
15.	Anti-Microbial Activity	Mahanimbine, Mahanine, Murrayanol	Inhibition of Topoisomerase I And II	(92)

16.	Anti-Diarrhoeal Activity	Kurryam, Koenine, Koenimbine	Prostaglandin E2-Induced enter pooling and reduction in gastrointestinal motility	(2)
17.	Chemo-protective Activity	Koenimbin	Reduce Chromosomal Damage	(94)
18.	Immunomodulatory Activity	Mahanimbine, Mahanine	Remove Carbon Partical From Blood	(95)
19.	Haema-tological Activity	Koenimbidine, Mahanimbicine	Not any Adverse Effect Against Food Efficiency Ratio	(92)
20.	Antipyretic Activity	Murrayacine, Murrayazoline	Murrayacine, Murrayazoline	(96)
21.	Nephro-protective Activity	Koenimbidine	Work against unilateral renal Ischemia	(1, 96)
22.	Cardio-Protective Activity	Girinimbine, Girinimbiol	Cadmium-Induced Oxidation Is Reduces	
23.	Anti-Cytotoxicity Activity	Girinimbine, Koenoline, Mahanine and Pyrafoline-D	Exhibiting Cell Death Resulted As The Mortality Of The Cell	
24.	Inotropic Activity	Girinimbiol	Positive Inotropic Effect	(97)
25.	Hepatoprotective Activity	Mahanimbine, Isomahanimbine, Girinimbine, Mahanine, Murrayazolidine, Murrayazoline	Oxidative Stress Inducer	(98)
26.	Anti-Lipase Activity	Mahanimbin, Koenimbin and Koenigicine	Reduced Total Cholesterol (Tc) And Triglyceride (Tg) Levels	(92)
27.	Anti-Alzheimer's Activity	Isomahanimbine, Murrayazolidine	Improves the Values of Protective Antioxidants	
28.	Anti-Analgesic Activity	Mahanine, Mahanimbine, Girinimbine, Isomahanimbine	Anti-Nociceptive Effects	
29.	Effect Digestive System	Mahanine, Murrayafoline	Stimulates Digestive Enzymes	(93)
30.	Neuro-Protective Activity	Koenimbin, Clausazoline-K, Koenigicine	Decreasing Glycemic Levels	(92)
31.	Anti-Inflammatory Activity	Mahanine, Mahanimbine, Girinimbine, Isomahanimbine	Cox-Inhibitory Activity	

Conclusion

All around the world, medicinal herbs are less expensive and freely accessible. As a result, we must increase the commercial usage of medicinal plants as a reliable supply of medication. *Murraya koenigii* is a readily available, multipurpose plant that has the ability to treat a wide range of common ailments as well as challenges we face every day. The therapeutic applications, phytochemistry, and pharmacological characteristics of *Murraya koenigii* are outlined in the current review. Several bioactive substances, such as alkaloids, polyphenols, terpenoids, and flavonoids, are found in *Murraya koenigii*. The pharmacological effects of *Murraya koenigii* and its derivatives, such as anticarcinogenic, proapoptotic, antiangiogenic, antimetastatic, immunomodulatory, and antioxidant capabilities, appear to be quite significant. The diverse roles that *Murraya koenigii* and its derivatives play in a variety of cell signalling pathways at various levels in different illnesses are the basis for the molecular mechanisms behind these activities. Oxidative stress, neurotoxicity, neuroinflammation, neuronal loss, and cognitive dysfunctions are all reduced by *Murraya koenigii* and its derivatives. However, like other polyphenols, *Murraya koenigii*'s actions are somewhat constrained by its bioavailability, and in such

circumstances, efficiency improvement should be carried out. Therefore, further experimental research on improving bioavailability and efficiency in clinical studies has to be included in future studies.

Declarations

Author Informations

Shouvik Kumar Nandy ✉

Affiliation: Department of Pharmacology, School of Pharmacy, Techno India University, Kolkata – 700091, India.

Contribution: Conceptualization, Data Curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing - Original Draft, Writing - Review & Editing.

Sattwik Das

Affiliation: Department of Pharmacology, School of Pharmacy, Techno India University, Kolkata – 700091, India.

Contribution: Conceptualization, Formal analysis, Funding acquisition, Methodology, Project administration, Resources, Supervision, Validation, Writing - Review & Editing.

Acknowledgment

The authors are thankful to Techno India University, Salt lake, Sector- V, Kolkata, West Bengal, for providing supplementary data and Chem draw software for structure making.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability

All data related to this review are available within the paper and its Supplementary Information.

Ethics Statement

Not applicable.

Funding Information

Not applicable.

Supplemental Material

Supplemental data contains two figures showing the compounds' structures identified in *Murraya koenigii*.

Supplemental Figure 1

(<https://etflin.com/file/document/202311061431321875603767.png>); Supplemental Figure 2

(<https://etflin.com/file/document/20231106143132321503236.png>).

References

1. Subramaniam G, XinYap K, Ong GH. The synergistic antimicrobial effects of crude plant extracts from *Plectranthus amboinicus*, *Azadirachta indica* and *Murraya koenigii* on *Staphylococcus* sp. 2015;58- 60.
2. Nirjara G, Sujatha P, Prabhudas P. Efficacy of pulverized *Punica granatum* (Lythraceae) and *Murraya koenigii* (Rutaceae) leaves against stored grain pest *Tribolium castaneum* (Coleoptera: Tenebrionidae). *International Journal of Agriculture and Biology*. 2010;12(4):616-20.
3. Kumar SR, Loveleena D, Godwin S. Medicinal property of *Murraya koenigii*-a review. *Int. res. j. biol* 2013; 2(9):80-3.
4. Handral HK, Pandith A, Shruthi SD. A review on *Murraya koenigii*: multipotential medicinal plant. *Asian J. Pharm. Clin. Res* 2012; 5(4):5-14.
5. Malwal M, Sarin R. Antimicrobial efficacy of *Murraya koenigii* (Linn.) Spreng. root extracts. 2011. 48-51.
6. TOMA M, LUCHIAN V, HOZA D. Avantguard of Romanian research: *Murraya koenigii* L.-an amazing flower and medicinal plant. *Scientific Papers-Series B, Horticulture*. 2020;64(1):717-28.
7. Dhongade H, Sawarkar H, Muley B, Deshmukh V, Pande A. Therapeutic potentials of *Murraya koenigii* Spreng (Rutaceae). *Am J Pharm Educ* 2013; 3(9).
8. Gahtori K, Aggarwal C, Tyagi M, Bajpai AB, Srivastava N, Singh S, Singh S, Gaurav N. Review on *Murraya koenigii*: Dietary Supplements and Highly Prosperous Plants of Pharmacological Value. *J. sci. temper* 2022;108.
9. Tiwari S, Talreja S. A Pharmaceutical Importance of *Murraya koenigii*-A Complete Study. *Indian J. Public Health* 2020;11(11):277.
10. Verma S. Overview study on *Murraya koenigii* (mithaneem): Rutaceae. *J. drug deliv. Ther.* 2018;8(4):90-2.
11. Gahlawat DK, Jakhar S, Dahiya P. *Murraya koenigii* (L.) Spreng: an ethnobotanical, phytochemical and pharmacological review. *J. pharmacogn. phytochem* 2014; 3(3):109-19.
12. Saraswat N, Wal P, Pal RS, Wal A, Pal Y, Pandey A. Pharmacognostic Evaluation and Standardization of the leaves of *Mimosa pudica*, leaves of *Murraya koenigii* and root of *Asparagus racemosus*. *Res J Pharm Technol* 2020; 13(12):5743-8.
13. Abeyasinghe DT, Alwis DD, Kumara KA, Chandrika UG. Nutritive importance and therapeutics uses of three different varieties (*Murraya koenigii*, *Micromelum minutum*, and *Clausena indica*) of curry leaves: An updated review. *Evidence-Based Complementary and Alternative Medicine*. 2021. 2021. <https://doi.org/10.1155/2021/5523252>.
14. Goyal P, Chhabra R, Vij L. Ethnobotany, Phytochemical, Pharmacological Potentials of *Murraya koenigii*, and Its Health Benefits-A Review. *Current Journal of Applied Science and Technology*. 2020 Sep 1;39(26):29-38.
15. Akula P, Sree AN, Santosh B, Sandeep B, Raviteja KB, Keerthi T. Evaluation of anti-microbial activity of leaf and bark extracts of *Murraya koenigii* (curry leaves). *Journal of Pharmacognosy and Phytochemistry*. 2016;5(3):101-5.
16. Samanta SK, Kandimalla R, Gogoi B, Dutta KN, Choudhury P, Deb PK, Devi R, Pal BC, Talukdar NC. Phytochemical portfolio and anticancer activity of *Murraya koenigii* and its primary active component, mahanine. *Pharmacological research*. 2018 Mar 1;129:227-36.
17. Vijayanand S. Evaluation of Antidiabetic activity of *Murraya koenigii* on Alloxan Induced Diabetic rats. *Int. J. Pharm. Sci. Res* 2015; 6:12.
18. Ghosh S, Mandi SS. Altitudinal effect in active principle content in *Murraya koenigii* (L) correlated with DNA fingerprinting study. *Journal of Medicinal Plants Studies*. 2018;6:20-6.
19. Singh AG. *Murraya koenigii* (L.) Spreng-Curry Leaves/Mitho Nim-A Miracle Plant. Butwal Campus

Journal. 2020, 3(1):125-30.

20. Batool S, Khera RA, Hanif MA, Ayub MA, Memon S. Curry leaf. In Medicinal Plants of South Asia. Elsevier sci 2020; (179-190).

21. Gupta S, George M, Singhal M, Sharma GN, Garg V. Leaves extract of *Murraya koenigii* linn for anti-inflammatory and analgesic activity in animal models. Journal of advanced pharmaceutical technology & research. 2010, 1(1):68.

22. Sharma P, Vidyasagar G, Bhandari A, Singh S, Ghule S, Agrawal A, Goyal S, Panwar MS. antiulcer activity of leaves extract of *Murraya koenigii* in experimentally induced ulcer in rats. Pharmacologyonline 2011; 2: 818-24.

23. Fiebig M, Pezzuto JM, Soejarto DD, Kinghorn AD. Koenoline, a further cytotoxic carbazole alkaloid from *Murraya koenigii*. Phytochemistry 1985; 24(12):3041-3.

24. Xie JT, Chang WT, Wang CZ, Mehendale SR, Li J, Ambihaipahar R, Ambihaipahar U, Fong HH, Yuan CS. Curry leaf (*Murraya koenigii* Spreng.) reduces blood cholesterol and glucose levels in ob/ob mice. Am. J. Chin. Med 2006; 34(02):279-84.

25. Rao BR, Rajput DK, Mallavarapu GR. Chemical diversity in curry leaf (*Murraya koenigii*) essential oils. Food Chem 2011; 126(3):989-94.

26. Bhandari PR. Curry leaf (*Murraya koenigii*) or cure leaf: review of its curative properties. Journal of medical nutrition and nutraceuticals 2012; 1(2):92.

27. Sarla GS. Complementary and alternative medicine. Res Rev J Pharm 2019; 6(3):1-5p.

28. Karthik S, Sahana KG, Babu A, HS HG, Meera V, Kumar R, Bharathi DR. *Murraya koenigii*: phytopharmacological, traditional and medicinal considerations. International Journal of Health Care and Biological Sciences. 2022 Dec 26:86-93.

29. Ilangovan SS, Krishna P, Koushika Das SS. A review on anti-microbial properties of *Murraya koenigii*. American Journal of Pharmaceutical Research. 2016;6(12).

30. Saini SC, Reddy GB. A review on curry leaves (*Murraya koenigii*): Versatile multi-potential medicinal plant. Am J Phytomed Clin Ther 2015; 3(4):363-8.

31. Waghmare AN, Tembhurne SV, Sakarar DM. Phytochemical analysis and in vitro antioxidant properties of *Murraya koenigii* (L.) fruits. Am. J. Phytomedicine Clin. Ther. 2015;3:403-16.

32. Shankar PR, Bhuminathan S, Rekha UV. Anti-Diabetic Activity of *Murraya koenigii*-A Comprehensive Review. Journal of Pharmaceutical Research International. 2021 Dec 16:462-70.

33. Rekha UV, Bhuminathan S, Shankar PR. Anti-Diabetic Activity of *Murraya koenigii*-A Comprehensive Review. J. Pharm. Res. Int 2021; 33(58B): 462-470.

34. Igara CE, Omoboyowa DA, Ahuchaogu AA, Orji NU, Ndukwe MK. Phytochemical and nutritional profile of *Murraya koenigii* (Linn) Spreng leaf. J. pharmacogn. phytochem 2016; 5(5):7.

35. Tripathi YC, Anjum N, Rana A. Chemical composition and in vitro antifungal and antioxidant activities of essential oil from *Murraya koenigii* (L.) Spreng. Leaves. Asian Journal of Biomedical and Pharmaceutical Sciences. 2018;8(65):6-13.

36. Balakrishnan R, Vijayraja D, Jo SH, Ganesan P, Su-Kim I, Choi DK. Medicinal profile, phytochemistry, and pharmacological activities of *Murraya koenigii* and its primary bioactive compounds. Antioxidants 2020; 9(2):101.

37. Vats M, Singh H, Sardana S. Phytochemical screening and antimicrobial activity of roots of *Murraya koenigii* (Linn.) Spreng. (Rutaceae). Brazilian Journal of Microbiology. 2011;42:1569-73.

38. Deepika T, Noorjahan CM. Phytochemical Screening and Thin Layer Chromatographic analysis for Antioxidant activity of *Murraya koenigii* (Curry leaf). Int. j. pharm. life sci 2016; 7(12).

39. Ghosh D, Firdaus SB, Mitra EL, Dey M, Chattopadhyay AI, Pattari SK, Dutta SA, Jana K, Bandyopadhyay DE. Hepatoprotective activity of aqueous leaf extract of *Murraya koenigii* against lead-induced hepatotoxicity in male wistar rat. Int J Pharm Pharm Sci. 2013;5(1):285-95.

40. Sindhu RK, Arora S. Phytochemical and Pharmacognostical Studies on *Murraya koenigii* (L.) spreng. Roots. Drug Discov. Today 2015; 7(1).

41. Ramsewak RS, Nair MG, Strasburg GM, DeWitt DL, Nitiss JL. Biologically active carbazole alkaloids from *Murraya koenigii*. J. Agric. Food Chem 1999; 47(2):444-7.

42. Srivastava SK, Srivastava SD. New Constituents and biological-activity of the roots of *Murraya koenigii*. ISSN International Centre 1993; 70(7):655-9.

43. Noolu B, Ajumeera R, Chauhan A, Nagalla B, Manchala R, Ismail A. *Murraya koenigii* leaf extract inhibits proteasome activity and induces cell death in breast cancer cells. BMC complementary and alternative medicine. 2013, 13(1):1-7.

44. Ma Q, Tian J, Yang J, Wang A, Ji T, Wang Y, Su Y. Bioactive carbazole alkaloids from *Murraya koenigii* (L.) Spreng. Fitoterapia 2013; 87:1-6.

45. Tan SP, Ali AM, Nafiah MA, Amna U, Ramli SA,

- Ahmad K. Terpenes and phenolic compounds of *Murraya koenigii*. Chem. Nat. Compd 2017. 53(5):980-1.
46. Devi RS, Rath SK, Kumar S. Medico-Biowealth of India. Vol. II. Odisha: APRF publisher. 2021.
47. Goel A, Sharma A, Kulshrestha S. A phyto-pharmacological review on *Murraya koenigii*: an important medicinal plant. Int J Pharm Sci Rev Res. 2020;62(2):113-9.
48. Paul SK, Goyal S, Yadav P, Yadav S, Parashari D. Phytochemistry and pharmacological aspects of *Murraya koenigii* Linn. Res J Pharm Technol 2013; 28;6:695-7.
49. Kumari B. Taxonomy and ethnobotany of *Murraya koenigii* (L.) Spreng: An exotic shrub in Rohilkhand region of Uttar Pradesh. Journal of Medicinal Plants. 2018;6(4):123-5.
50. Lakshmikandhan T. Green synthesis of zinc oxide nanoparticles using *Murraya koenigii* (curry leaf) leaf extract. Malaya J Matematik. 2020;2:4309-17.
51. Jayasinha P, Warnasuriya D, Dissanayake H. Karapincha: *Murraya koenigii*-a literature survey. Medicinal and aromatic plant series: No. 2. 2007.
52. Tan MA, Sharma N, An SS. Multi-Target Approach of *Murraya koenigii* Leaves in Treating Neurodegenerative Diseases. Pharmaceuticals. 2022;15(2):188.
53. Aswin Sakthivel M, Velmurugan S, Selvi BS, Senthil A. Studies on vegetative propagation in curry leaf (*Murraya koenigii*Spreng.). Pharma Innovation 2021.
54. Malode GP, Parbat AY, Shaikh AR, Panchale WA, Manwar JV, Bakal RL. Phytochemistry, pharmacology and botanical aspects of *Murraya koenigii* in the search for molecules with bioactive potential-A review. GSC Adv. Res. Rev 2021; 6(3):143-55.
55. ChV S, Meera I. Antioxidant and biological activities of three morphotypes of *Murraya koenigii* L. from Uttarakhand. J. Food Process. Technol 2013 ;4:1-7.
56. Mandlik DS, Namdeo AG. Immunomodulators and Phytodrugs. In Evidence Based Validation of Traditional Medicines. Singapore.Springer Sci. Rev 2021 (pp. 901-920).
57. Sharma R, Kumar U. Exploration and phytochemical estimation of *Murraya koenigii* leaves for pharmaceutical applications. Asian Journal of Pharmaceutical Research. 2019, 9(3):159-68.
58. Ansari MH, Mahapatra DK. A Short Overview on Anti-Diabetic Natural Products: Reviewing the Herbotherapeutic Potentials. Natural Products Pharmacology and Phytochemicals for Health Care 2021; 1-22.
59. Mitra PK, Maitra T, Mitra P, Paul B, Ghosh D, Guria M, Chowdhury D, Das AP. Antiulcer activity of an isolated compound (MK-1) from *Murraya koenigii* (Linnaeus) Sprengel leaf in rats. Pleione 2011; 5(1):49-55.
60. Rehana D, Mahendiran D, Kumar RS, Rahiman AK. In vitro antioxidant and antidiabetic activities of zinc oxide nanoparticles synthesized using different plant extracts. Bioprocess and biosystems engineering. 2017 Jun;40(6):943-57.
61. Gill NS, Sharma B. Study on antioxidant potential of *Murraya koenigii* leaves in Wistar rats. Pakistan Journal of Biological Sciences: PJSBS. 2014 Jan 1;17(1):126-9.
62. Ajay S, Rahul S, Sumit G, Paras M, Mishra A, Gaurav A. Comprehensive review: *Murraya koenigii* Linn. Asian J Pharm Life Sci. 2011, 2231:4423.
63. Yukari T, Hiroe K, Nordin H L and Nobuji N. Antioxidative Activity of Carbazoles from *Murraya koenigii* leaves. J Agri Food Chem 2001; 49:5589-5594.
64. Rao LJM, Ramalakshmi K, Borse BB, Raghavan B. Chemical composition of volatiles from coconut sap neera and effect of processing. Food Chemistry 2006; 100: 742-747.
65. Rahman MM, Gray AI. A benzoisofuranone derivative and carbazole alkaloids from *Murraya koenigii* and their antimicrobial activity. Phytochem 2005; 66 13 :1601-1606.
66. Manju Panghal, Vivek Kaushal, Yadav Jaya P. In vitro antimicrobial activity of ten medicinal plants against clinical isolates of oral cancer cases. Annals of Clinical Microbiology and Antimicrobials 2011; 10:21.
67. Kureel SP, Kapil RS, Popli SP. Terpenoid alkaloids from *Murraya koenigii* Spreng.-II. The constitution of cyclomahanimbine, bicyclomahanimbine & mahanimbidine. Tetrahedron Letters 1969; 44:3857-3862.
68. Anika A, Kaur S, Sadwal S. A Review of the Anti-Cancer Potential of *Murraya koenigii* (Curry Tree) and Its Active Constituents. Nutrition and Cancer. 2022, 74(1):12-26.
69. Nagappan T, Ramasamy P, Wahid ME, Segaran TC, Vairappan CS. Biological activity of carbazole alkaloids and essential oil of *Murraya koenigii* against antibiotic resistant microbes and cancer cell lines. Molecules. 2011, 16(11):9651-64.
70. Bonde, S.D.; Nemade, L.S.; Patel, M.R.; Patel, A.A. *Murraya koenigii* (Curry leaf): Ethnobotany, Phytochemistry and Pharmacology—A Review. Int. J. Pharm. Phytopharm. Res. 2011, 1, 23.
71. Noolu B, Gogulothu R, Bhat M, SYH Qadri S, Sudhakar Reddy V, Bhanuprakash Reddy G, Ismail A. In

vivo inhibition of proteasome activity and tumour growth by *Murraya koenigii* leaf extract in breast cancer xenografts and by its active flavonoids in breast cancer cells. *Anti-Cancer Agents Med. Chem* 2016; 16(12):1605-14.

72. Nadaf S, Desai R, More T, Shinde P, Dakare S, Killedar S. Antiproliferative and caspase-mediated apoptosis inducing effects of *Murraya koenigii* seeds against cancer cells. *South African Journal of Botany*. 2020, 132:328-37.

73. Lubis MF, Kaban VE, Aritonang JO, Satria D, Mulina AA, Febriani H. Acute toxicity and antifungal activity of the ointment *Murraya koenigii* ethanol extract. 2022, 256-261.

74. Yee KT, bin Ibrahim MI, Nwe TM, Thwin MM, Lwin MM, Aung KC, Oo MS, binti Raduan SZ, San Yi M. Bioactive compounds screening, antimicrobial activities of leave extract from two palatable plants: *Piper betle* and *Murraya koenigii* (Curry leaves). *Research Journal of Pharmacy and Technology*. 2023, 16(3):1452-8.

75. Kumar, N.S.; Mukherjee, P.K.; Bhadra, S.; Saha, B.P.; Pal, B.C. Acetylcholinesterase inhibitory potential of a carbazole alkaloid, mahanimbine, from *Murraya koenigii*. *Phyther. Res.* 2010, 24, 629-631.

76. James SA, Omwirhiren RE, Joshua IA, Dutse I. Anti-diabetic properties and phytochemical studies of ethanolic leaf extracts of *Murraya koenigii* and *Telfairia occidentalis* on alloxan-induced diabetic albino rats. *ornament*. 2016, 49.

77. Dhamane SP, Patil SA, Kulkarni AS, Potnis VV. Evaluation of antimicrobial activity of ethanolic extract of *Murraya koenigii* against *S. mutans*. *Journal of Pharmacognosy and Phytochemistry*. 2019, 8(4):1223-8.

78. Selvan DS, Kumar RS, Murugesan S, Shobana S, Rahiman AK. Antidiabetic activity of phytosynthesized Ag/CuO nanocomposites using *Murraya koenigii* and *Zingiber officinale* extracts. *Journal of Drug Delivery Science and Technology*. 2022, 67:102838.

79. Sharma A, Nagraik R, Venkidasamy B, Khan A, Dulta K, Kumar Chauhan P, Kumar D, Shin DS. In vitro antidiabetic, antioxidant, antimicrobial, and cytotoxic activity of *Murraya koenigii* leaf extract interceded ZnO nanoparticles. *Luminescence*. 2023, 38(7):1139-48.

80. Singh S, More PK, Mohan SM. Curry leaves (*Murraya koenigii* Linn. Sprengal)-a mircale plant. *Indian Journal of Scientific Research*. 2014;4(1):46-52.

81. Dineshkumar B, Mitra A, Mahadevappa M. Antidiabetic and hypolipidemic effects of mahanimbine carbazole alkaloid from *Murraya koenigii* Rutaceae leaves. *International Journal of Phytomedicine* 2010; 2:22-30.

82. Widayanti A, Srifiana Y, Efendi K. Antidiabetics activity of salam koja (*Murraya koenigii*) leaves tea bag. *Pharmaceutical Sciences and Research*. 2019;6(2):5.

83. Patidar DK. Anti-ulcer activity of aqueous extract of *Murraya koenigii* in albino rats. *Int J Pharma Bio Sci*. 2011, 2(1):524-9.

84. Nalla S, Dodoala S, Golla U. A review on pharmacognosy, phytochemistry and therapeutic importance of *Murraya koenigii*, *azadirachta indica* and *Piper nigrum*. *Journal of Integral Sciences*. 2020, 1-8.

85. Kaloni D, Chakraborty D, Tiwari A, Biswas S. In silico studies on the phytochemical components of *Murraya koenigii* targeting TNF- α in rheumatoid arthritis. *Journal of Herbal Medicine*. 2020, 24:100396.

86. Nishan M, Subramanian P. *Murraya koenigii* (curry leave)-A review on its potential. *Int. J. PharmTech Res*. 2015;7(4):566-72.

87. Mani V, Mohd Azahan NS, Ramasamy K, Lim SM, Abdul Majeed AB. Mahanimbine Improved Aging-Related Memory Deficits in Mice through Enhanced Cholinergic Transmission and Suppressed Oxidative Stress, Amyloid Levels, and Neuroinflammation. *Brain Sciences*. 2022;12(1):12.

88. Mandal S, Nayak A, Kar M, Banerjee SK, Das A, Upadhyay SN, Singh RK, Banerji A, Banerji J. Antidiarrhoeal activity of carbazole alkaloids from *Murraya koenigii* Spreng (Rutaceae) seeds. *Fitoterapia*. 2010;81(1):72-4.

89. Prabhu KA, Tamilanban T. Investigation of antidiabetic activity of stem of *Murraya koenigii*. *Int. j. res. pharmacol. pharmacother* 2012; 1(2):165-8.

90. Pagariya A, Maithili V. Anti-diarrhoeal activity of *Murraya koenigii* Linn root extracts. *Journal of Natural Remedies*. 2009;9(1):8-11.

91. Jain M, Gilhotra R, Singh RP, Mittal J. Curry leaf (*Murraya koenigii*): A spice with medicinal property. *MOJ Biol Med* 2017; 2(3):00050.

92. Darvekar VM, Patil VR, Choudhari AB. Anti-inflammatory activity of *Murraya koenigii*Spreng on experimental animals. *J Nat Prod Plant Resour* 2011; 1(1):65-9.

93. Paul S, Bandyopadhyay TK, Bhattacharyya A. Immunomodulatory effect of leaf extract of *Murraya koenigii* in diabetic mice. *Immunopharmacol. Immunotoxicol* 2011;33(4):691-9.

94. Rageeb MD, Usman MD, Barhate SD. Phytochemical evaluation and effect of antipyretic activity on *Murraya koenigii*Spreng. Leaves extract. *Int. j. pharm. chem. sci* 2012;1(1):231-6.

95. Kadam SH, Dombé S, Naikwadi P, Patil M. Cardiovascular effects of aqueous extract of *Murraya koenigii* on isolated perfused frog heart preparation. *J. Pharm. Res* 2011; 4(2):462-3.

96. Reddy BM, Dhanpal CK, Lakshmi BV. A review on curry leaves (*Murraya koenigii*): Versatile multi-potential medicinal plant. *Int. J. Adv. Pharm. Med. BioalliedSci* 2018; 6:31-41.

97. Sathaye SA, Amin PD, Mehta VB, Zala VB, Kulkarni RD, Kaur H, Redkar R. Hepatoprotective activity of *Murraya koenigii* against ethanol induced liver toxicity model in experimental animals. *Int. J. Pharma Bio Sci* 2012; 3(1):430-8.

98. Rana VS, Juyal JP, Blazquez MA. Chemical constituents of the volatile oil of *Murraya koenigii* leaves. *Int. J. Aromather* 2004; 14(1):23-5.

Publish with us

In ETFLIN, we adopt the best and latest technology in publishing to ensure the widespread and accessibility of our content. Our manuscript management system is fully online and easy to use.

Click this to submit your article:

<https://etflin.com/#loginmodal>



This open access article is distributed according to the rules and regulations of the Creative Commons Attribution (CC BY) which is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

How to cite: Nandy, S.K., Das, S.. Unveiling the Diverse Medicinal Properties of *Murraya koenigii*. *Sciences of Phytochemistry*. 2023; 2(2):211-225