



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Review Article

**JASMINUM GRANDIFLORUM L.: A COMPREHENSIVE
REVIEW OF ITS PHYTOCHEMISTRY AND THERAPEUTIC
APPLICATIONS**Ms. Akshaya. S.S ¹, Mr. Alan Antony¹, Mrs. Anu Pallavi. L¹,Mrs. Savitha Mol G M², Mrs. Sanitha M³, Dr. Kiran K J⁴, Dr. Prasobh G R ⁵¹ B. Pharm Students, Sree Krishna College of Pharmacy and Research Centre, Parassala,
Thiruvananthapuram, Kerala, India² Associate Professor, Department of Pharmacology, Sree Krishna College of Pharmacy and
Research Centre, Parassala, Thiruvananthapuram, Kerala, India³ Assistant Professor, Department of Pharmacology, Sree Krishna College of Pharmacy and
Research Centre, Parassala, Thiruvananthapuram, Kerala, India⁴ Vice Principal, Sree Krishna College of Pharmacy and Research Centre, Parassala,
Thiruvananthapuram, Kerala, India⁵ Principal, Sree Krishna College of Pharmacy and Research Centre, Parassala,
Thiruvananthapuram, Kerala, India**Abstract:**

Jasminum grandiflorum L., belonging to the family *Oleaceae*, is a perennial aromatic shrub or climber widely cultivated for its fragrant white flowers. The plant bears opposite pinnate leaves and star-shaped flowers, making it an important ornamental and commercial crop, especially in perfumery. Taxonomically, it is classified under Kingdom *Plantae*, Order *Lamiales*, Family *Oleaceae*, Genus *Jasminum*, and Species *grandiflorum*. It grows well in warm climates with full sunlight and well-drained loamy soil. Propagation is mainly through cuttings or layering, and regular pruning enhances flowering. Medicinally, the plant is valued in traditional systems like *Ayurveda* and *Unani*. Its flowers, leaves, and roots possess anti-inflammatory, antimicrobial, antioxidant, and mild sedative properties, and are used in the treatment of skin diseases, wounds, fever, and stress-related disorders.

Keywords: *Jasminum grandiflorum L.*, medicinal uses, Phytochemistry**Corresponding author:**

Ms. Akshaya .S.S , Mrs. Alan Antony , Mrs. Anupallavi .L

B.Pharm students,

Sree Krishna College of Pharmacy and Research Centre,

Parassala, Thiruvananthapuram, Kerala, India

E-Mail: akshayasanandakumar237@gmail.com ,alanantony0803@gmail.com, anupallavi628292@gmail.com

QR CODE



Please cite this article in press Akshaya .S.S et al., *Jasminum grandiflorum L.: a comprehensive review of its phytochemistry and therapeutic applications*, Indo Am. J. P. Sci, 2025; 12(12).

INTRODUCTION:**PLANT DESCRIPTION**

Jasminum grandiflorum is a large, evergreen, scrambling or twining shrub that grows up to 10–15 m in height. It is native to Asia, including Kashmir, Afghanistan, and Persia, and occurs at elevations of 700–2,700 m. The leaves are opposite and imparipinnate, 5.0–12.5 cm long, with 7–11 leaflets. The terminal leaflet is slightly larger, rhomboid-ovate to lanceolate, while the lateral leaflets are ovate, usually obtuse, with mucronate tips; the distal pair may be connate, and the proximal leaflets are shortly petiolated or sessile.

The flowers are highly fragrant, 3.0–3.8 cm across, white with a pink tinge externally, and borne in lax terminal or axillary cymes longer than the leaves. The calyx has five subulate lobes, 2–3 times longer than the short tube. The corolla tube measures 1.8–2.5 cm with five elliptic to obovate lobes. The ovary is bicarpellary. Fruits are black, elliptic berries when ripe. The species is cultivated in India and found wild in the sub-tropical north-west Himalayas, Western Ghats, Nilgiris, and other regions, as well as in France, Italy, China, Japan, Morocco, and Egypt.^[1,2]



Fig no:1 *Jasminum grandiflorum* L. plant

CULTIVATION AND COLLECTION

Jasminum grandiflorum is preferably cultivated in well-drained loamy soil with a pH range of 6.5–7.5, although it can also grow successfully in other soil types such as black soil, lateritic soil, and clay loam, provided an efficient drainage system is maintained. The plant is highly sensitive to waterlogging; therefore, proper soil aeration and drainage are essential for healthy growth. Propagation can be effectively carried out using the shoot tip culture method, which ensures uniform and disease-free plants.

Flowering generally begins within the first year of planting, indicating its early reproductive potential. The flowering season extends from May to December, during which flowers are harvested regularly. The leaves of *Jasminum grandiflorum* are of medicinal importance and are carefully collected by hand to avoid damage to the plant. After collection, the plant material is thoroughly cleaned and shade-dried to preserve its quality, making it suitable for further pharmacological and scientific studies.^[3]

CHEMICAL CONSTITUENTS

Jasminum grandiflorum L. contains a diverse range of bioactive phytochemicals, including alkaloids such as jasminine and nuciferine. It is also rich in flavonoids, notably quercetin, kaempferol, and rutin, which contribute to its antioxidant potential. In addition, the plant possesses important terpenoid compounds, including linalool and methyl jasmonate, which are largely responsible for its characteristic fragrance and therapeutic properties.^[4]

MEDICINAL USES

Jasminum grandiflorum exhibits a wide range of pharmacological activities, including significant anti-inflammatory and antimicrobial properties. It is also known for its wound-healing potential, promoting tissue repair and recovery. Additionally, the plant has demonstrated anti-diabetic activity, supporting its traditional use in the management of metabolic disorders.^[5]

SUMMARY

Jasminum grandiflorum L., commonly known as Spanish jasmine, is a medicinally important plant widely used in traditional systems of medicine. The plant is rich in bioactive constituents such as flavonoids, phenolic compounds, alkaloids, terpenoids, and glycosides, which contribute to its diverse pharmacological activities. Various studies have reported its antioxidant, anti-inflammatory, antimicrobial, antidiabetic, hepatoprotective, wound-healing. Different parts of the plant, especially the leaves and flowers, have been explored using aqueous, ethanolic, and methanolic extracts, showing significant therapeutic relevance. This review compiles and critically summarizes the botanical description, phytochemical profile, traditional uses, and reported pharmacological activities of *Jasminum grandiflorum* L.

CONCLUSION:

Based on the reviewed literature, *Jasminum grandiflorum* L. emerges as a valuable medicinal plant with broad pharmacological potential supported by its rich phytochemical composition. Scientific evidence validates many of its traditional uses, particularly its antioxidant and protective roles

against oxidative stress-related disorders. Despite promising experimental findings, further systematic studies including standardization of extracts, isolation of active compounds, toxicity evaluation, and clinical investigations are necessary. Future research may lead to the development of novel, safe, and effective plant-based therapeutic agents derived from *Jasminum grandiflorum* L.

REFERENCES:

- Ornek K, Karel F, Büyükbingöl Z. May nitric oxide molecule have a role in the pathogenesis of human cataract? *Exp Eye Res* 2003; 76: 23-27
- Olofsson EM, Marklund SL, Behndig A. Enhanced diabetes-induced cataract in copper-zinc superoxide dismutase-null mice. *Invest Ophthalmol Vis Sci* 2009; 50: 2913-2918
- Papadimitriou DT, Bothou C, Skarmoutsos F, Alexandrides TK, Papaevangelou V, Papadimitriou A. The autoimmune hypothesis for acute bilateral cataract in type 1 diabetes. *Diabetes Metab* 2016; 42: 386-387
- Klein BE, Klein R, Moss SE. Prevalence of cataracts in a population-based study of persons with diabetes mellitus. *Ophthalmology* 1985; 92: 1191-1196
- Nielsen NV, Vinding T. The prevalence of cataract in insulin-dependent and non-insulin-dependent diabetes mellitus. *Acta Ophthalmol (Copenh)* 1984; 62: 595-602
- Kothadia AD, Shenoy AM, Shabaraya AR, Rajan MS, Viradia UM, Patel NH. Evaluation of cataract preventive action of phycocyanin. *Int J Pharm Sci Drug Res* 2011; 3: 42-44.
- Kato A, Yasuko H, Goto H, Hollinshead J, Nash RJ, Adachi I. Inhibitory effect of rhetsinine isolated from *Evodia rutaecarpa* on aldose reductase activity. *Phytomedicine* 2009; 16: 258-261.
- Gupta SK, Kalaiselvan V, Sharma A, Srivastava S, Agrawal SS. Anti cataract potential of *Phyllanthus niruri* in galactose induced cataractogenesis of rat. *Int J Ophthalmol* 2009; 2: 286-290.
- Awasthi S, Srivastava SK, Piper JT, Singhal SS, Chaubey M, Awasthi YC. Curcumin protects against 4-hydroxy-2-transnonenal induced cataract formation in rat lenses. *Am J Clin Nutr* 1996; 64: 761-766.
- Truscott, R. J. W. (2005). Age-related nuclear cataract—oxidation is the key. *Experimental Eye Research*, 80(5), 709–725.
- Spector, A. (1995). Oxidative stress-induced cataract: Mechanism of action. *FASEB Journal*, 9(12), 1173–1182.
- Rao, P. V., & Rao, C. M. (2015). Oxidative stress and cataract formation: Role of antioxidants in cataract prevention. *Indian Journal of Ophthalmology*, 63(6), 448–454.
- Gupta, S. K., et al. (2020). Role of phytochemicals in the prevention and treatment of cataract: An overview. *Journal of Ophthalmic & Vision Research*, 15(4), 462–473.
- Michael, R., & Bron, A. J. (2011). *Philosophical Transactions of the Royal Society B*, 366, 1278–1292.
- Shiels, A., & Hejtmancik, J. F. (2017). Mutations and mechanisms in congenital and age-related cataracts. *Cold Spring Harbor Perspectives in Medicine*, 7(6), a025387.