


RESEARCH ARTICLE

UHPLC-QTOF-MS/MS-based metabolomic discovery of anticancer compounds in ethanolic extracts of *Ficus hispida* L. f.

Saengrawee Thammawithan^{1,2}, Jirattiporn Thanuma^{1,2}, Sirinya Sitthirak³, Chadapohn Panplu^{1,2}, Thapanee Pruksatrakul⁴, Piya Prajumwongs^{1,2}, Arporn Wangwiwatsin^{1,2}, Poramate Klanrit^{1,2}, Watcharin Loilome^{1,2}, Nisana Namwat^{1,2*}

1 Department of Systems Biosciences and Computational Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand, **2** Cholangiocarcinoma Research Institute, Khon Kaen University, Khon Kaen, Thailand, **3** School of Allied Health Sciences, Walailak University, Nakhon Si Thammarat, Thailand, **4** National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), Thailand

* nisana@kku.ac.th



OPEN ACCESS

Citation: Thammawithan S, Thanuma J, Sitthirak S, Panplu C, Pruksatrakul T, Prajumwongs P, et al. (2026) UHPLC-QTOF-MS/MS-based metabolomic discovery of anticancer compounds in ethanolic extracts of *Ficus hispida* L. f. PLoS One 21(3): e0343411. <https://doi.org/10.1371/journal.pone.0343411>

Editor: Ghulam Yaseen, University of Education, PAKISTAN

Received: October 21, 2025

Accepted: February 2, 2026

Published: March 3, 2026

Copyright: © 2026 Thammawithan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data availability statement: All relevant data are within the manuscript and its [Supporting Information](#) files.

Abstract

Ficus hispida L. f. (*F. hispida*) is commonly used in traditional medicine for various health problems. No comprehensive analysis of all its components has yet been undertaken, despite researchers having explored its chemical constituents and biological activities. Using untargeted metabolomics, we aimed to assess the chemical compositions and metabolic differences among the five parts of *F. hispida*: bark, fruits, leaves, twigs, and stalks. To discover the compounds that might be accountable for the noted efficacy, our study assessed the correlation between the identified metabolites and anticancer activities. We applied untargeted metabolomics using UHPLC-QTOF-MS/MS to confirm a total of 82 metabolites. These compounds were classified into six phytochemical groups with predominant accumulation in each part: fatty acids and their conjugates (twigs), terpenoids (stalks), phenylpropanoids (bark and twigs), alkaloids (bark and leaves), saccharides and their conjugates (bark), and amino acids and peptides (twigs and stalks). Multivariate analysis showed markedly distinct metabolic patterns across the five tested parts, especially leaf and bark extracts. The sulforhodamine assay (SRB) for the cytotoxicity test revealed that the bark and leaf extracts had the highest potential to inhibit the viability of cholangiocarcinoma (CCA) cells, with IC₅₀ values of 0.71 ± 0.17 µg/mL and 0.82 ± 0.22 µg/mL for KKU-213A, and 0.78 ± 0.13 µg/mL and 1.03 ± 0.21 µg/mL for KKU-055. Univariate analysis revealed that four metabolites, including catharanthine, cianidanol, procyanidin B2, and quinic acid, had significant correlation with these anticancer abilities. Subsequent cytotoxicity studies on these candidate metabolites revealed that catharanthine suppressed CCA cell viability at the IC₅₀ of 41.0 ± 0.99 µM (KKU-213A) and 47.4 ± 4.34 µM (KKU-055) over other candidates. Our research suggests that

Funding: 1. The coordinating center for the plant genetic conservation project under the royal initiation of her royal highness Princess Maha Chakri Sirindhorn (RSPG), Thailand 2. A grant from Faculty of Medicine (grant. no. IN68060), Khon Kaen University, Thailand The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

catharanthine, which is a terpene indole alkaloid found in the bark and leaves of *F. hispida*, is responsible for the anticancer efficacy against CCA cells.

1. Introduction

Ficus hispida L. f. (*F. hispida*) is a medicinal plant of the Moraceae family that is commonly used in traditional medicine in China, India, Sri Lanka, Myanmar, Thailand, and Australia [1]. The root, leaf, fruit, and bark of this plant have been applied to treat ulcers, jaundice, diabetes, bleeding, and dysentery, among other medical conditions [2]. Comprehensive research has revealed an extensive variety of phytochemicals in different parts of *F. hispida*, comprising phenols, flavonoids, glycosides, alkaloids, steroids, sterols, saponins, and terpenes [3,4]. Terpenoids, flavonoids, and alkaloids are recognized as the most prominent and prevalent biologically active substances, potentially having a role in antioxidant and anticancer effects, among other biological activities [4].

A metabolomic approach is a potent tool for a comprehensive analysis of plant metabolites with high-throughput analytical methods such as mass spectrometry [5]. Metabolomics offer key insights into the biochemical constitution of plant material by methodically profiling metabolites in plant tissues, extracts, and derivatives [6]. We have applied UHPLC-QTOF-MS/MS, a high-throughput technique, in metabolomics to identify metabolites and classify them into each phytochemical class. Ethanol was used as an extraction solvent to obtain moderately polar metabolites with antioxidant and anticancer effects, including phenolic acids, flavonoids, alkaloids, terpenoids, and small peptides. Statistical analyses, including principal component analysis (PCA), partial least squares discriminant analysis (PLS-DA), and hierarchical cluster analysis, were performed to generate metabolomics profiling that allowed for the comparison of metabolite changes in various parts of plants [7]. Moreover, specific significant metabolites can be evaluated for their correlation with biological activities such as antioxidant, antibacterial, and anticancer activities to identify specific bioactive molecules responsible for biological effects observed in an extract.

Globally, cancer remains a leading cause of death. Cholangiocarcinoma (CCA), or bile duct cancer, is an aggressive and highly lethal malignancy originating from bile duct epithelial cells that is generally associated with a poor prognosis [8]. Current treatments, such as surgery, radiation, and chemotherapy, frequently exhibited limited effectiveness against CCA due to the aggressive characteristics of this cancer and prevalent drug resistance [9,10]. Consequently, the study and development of effective natural therapeutic agents derived from medicinal plants may prove beneficial for CCA treatment.

F. hispida has been identified as a potential source of novel anticancer agents and has previously demonstrated anticancer effects on various cell lines [11]. Previous research has shown that stem extract has strong antineoplastic activity against breast cancer [12], that crude extracts of leaves and the bark have antiproliferative potency against glioblastoma (U87MG), and that methanolic fruit extract has cytotoxic activity against HL60, KB, HeLa, HT29, and HepG2 cells [13]. Additionally, a previous study

identified several compounds from fruits that may serve as promising lead candidates for androgen-dependent prostate cancer [14,15]. These results indicate that *F. hispida* is a significant source of therapeutically relevant phytochemicals that can be developed into natural chemopreventive anticancer agents.

Therefore, our study aimed to utilize an untargeted metabolomic approach to evaluate the chemical compositions and metabolic variations among five specific parts of *F. hispida*: the bark, fruits, leaves, twigs, and stalks. Furthermore, we aimed to assess the *in vitro* anticancer activity, specifically the cytotoxicity against CCA cells, and investigate the relationship between the identified metabolites and this anticancer activity to discover the compounds that might be responsible for the noted efficacy.

2. Materials and methods

2.1 Chemicals and reagents

Commercial-grade 99.7% ethanol was obtained from S.C. Science Co., Ltd., Thailand. Dimethyl sulfoxide (DMSO), sulforhodamine B (SRB), Dulbecco's Modified Eagle Medium (DMEM), (-)-quinic acid, (+)-Catechin (cyanidanol), and procyanidin B2 were purchased from Sigma-Aldrich (St. Louis, MO, USA). Catharanthine was purchased from MedChemExpress (Monmouth Junction, NJ, USA). Gemcitabine was purchased from Fresenius Kabi (Maharashtra, India). Trichloroacetic acid (TCA), penicillin-streptomycin, and trypsin-EDTA were obtained from Life Technologies (Grand Island, NY, USA).

2.2 Plant materials and sample preparation

The fresh *F. hispida* samples were collected from the Romklao Kallapruek Park, Khon Kaen University, Khon Kaen, Thailand (16°28'25.3"N 102°49'03.0"E), in January 2024. The plant material was identified by Mr. Piya Sukkharom, and a voucher specimen (S. Thammawithan 1) was deposited at the KKU Herbarium (KKU No. 27631), Khon Kaen, Thailand (check the details in [S1 File](#)). The collected samples were subsequently chopped and washed with water to eliminate the surface dirt and adhering particles. Subsequently, the samples were then dried in a hot air oven at 60°C. The dried samples were ground into fine powder and stored at 4°C until the subsequent step.

2.3 Sample extraction

The extraction process was performed according to a previous study with some modifications [16]. The powder of each sample was extracted with 99.7% ethanol. Briefly, 20 g of powder was added to 80 mL of ethanol. The mixture was subjected to sonication using an ultrasonic bath cleaner (Jeken, China) at 40 Hz for 30 min. Various previous studies [17,18] have used this sonication treatment, which effectively enhances the antioxidant compounds. The extraction was repeated three times for the high yield recovery and then combined and filtered with Whatman filter paper no. 1. A rotatory evaporator (Buchi/R-100, Buchi (Thailand) Ltd.) and SpeedVac (Labconco, Wertheim, Germany) were used for concentrating and evaporating the filtered extracts at 40°C under reduced pressure for 48 h and then kept at 4°C.

2.4 LC-MS/MS method for metabolomics analysis of *F. hispida* extracts

The metabolomics analysis was carried out by using ultra-high-pressure liquid chromatography (UHPLC) coupled to a QTOF impact II mass spectrometer (Bruker Daltonics, Bremen, Germany). Briefly, ethanolic extracts of the plant were diluted to a concentration of 5 mg/mL and filtered using a 0.2 µm syringe filter. Thereafter, the extracts were separated by an Intensity Solo 2 C18 chromatographic column (100 x 2.1 mm, 2.1 mm). The flow rate was set at 0.4 mL/min, with a 5 µL injection volume. The temperature of the autosampler was set at 10°C and a column oven temperature of 40°C. Mobile phases A and B were used: A was a mix of ultrapure water and methanol (99:1) with 5 mM ammonium formate and 0.01% formic acid, while B was methanol with 5 mM ammonium formate and 0.01% formic acid. The LC gradient program started with 0.1% B from 0 to 1 min, followed by a linear increase to 99.9% B at 10 min for the separation of plant metabolites.

The mobile phase composition was maintained for 12 min for the column washing step, then returned to the initial conditions at 12.1 min and held until 15 min for column re-equilibration prior to the next injection. The mass spectra were acquired through data-dependent acquisition (DDA) in positive and negative polarity modes. The scan range was from 50 to 1300 m/z with a high-resolution accurate mass at 120,000. The precursor ions were fragmented at collision energies of 20, 25, and 30 eV, with a resolution of 30,000 for fragmented ions.

2.5 Anticancer activities of *F. hispida* extracts

Cell culture. The CCA cell lines KKKU-213A and KKKU-055 were obtained from the Japanese Collection of Research Bioresources (JCRB) Cell Bank, Osaka, Japan (check the details in S2 in [S1 File](#)), and cultured in Dulbecco's modified Eagle medium (DMEM) mixture supplemented with 10% fetal bovine serum and 100 IU/mL of penicillin-streptomycin. The cultured cell lines were incubated at 37°C in a humidified incubator with a 5% CO₂ atmosphere.

Cytotoxicity test by sulforhodamine B (SRB) method. The cell lines were plated into a 96-well plate at a density of 2.5×10^3 cells per well. After culturing for 12–13 h, the various concentrations of crude extract were added, and the cells were cultured for a further 48 h. Gemcitabine served as a positive control. A plate containing only cell suspension for a no-growth control (day 0) was set, and cell viability was measured by the Sulforhodamine B (SRB) method [19]. Briefly, the cells were fixed in 10% trichloroacetic acid for 1 h and washed three times with DI water. Then, fifty microliters of 0.4% SRB were exposed and incubated for 45 min. After unbound dye was washed with 1% acetic acid, SRB was solubilized with 200 μ l of 10 mM Tris-base pH 10.5, and the optical density was read at 540 nm with a microplate reader (Tecan, Switzerland).

2.6 Statistical analysis and data analysis

Statistical analysis of the cytotoxic test was conducted by using GraphPad Prism software. All assays were performed in three independent experiments, and data were reported as mean values with standard deviations (mean $\pm$ SD). The data were analyzed using the one-way analysis of variance (ANOVA) to evaluate the differences between the samples. A p -value of less than 0.05 was considered statistically significant. The analysis of the complex data, which included methods such as principal component analysis (PCA), partial least squares discriminant analysis (PLS-DA) score plots, variable importance in projection (VIP) score plot analysis, a heatmap plot, and univariate analysis, was done using MetaboAnalyst 6.0.

3. Results

3.1 Metabolite identification and classification among five parts of *F. hispida* extract by UHPLC-QTOF-MS/MS analysis

The chemical compositions of the bark, fruits, leaves, twigs, and stalks of *F. hispida* extract were investigated using UHPLC-QTOF-MS/MS in the positive and negative ionization modes to identify various components. A comparison of chromatograms among the five parts of *F. hispida* extract is shown in S1 in [S1 File](#). The chromatogram in positive mode revealed that the categories of metabolites among five parts were similar, while the content of many metabolites varied (S1A Fig in [S1 File](#)). It was noted that the chromatogram showed peak intensity at 325 m/z in all parts. This characteristic was annotated to be denatonium, a synthetic compound, which is used for denaturing ethanol. Therefore, denatonium was excluded from our analysis. In contrast, the types and the content of metabolites were different between the bark and other parts in negative mode (S1B Fig in [S1 File](#)). The ethanol extract of *F. hispida* yielded metabolites in the positive rather than the negative mode. Tentative metabolite identifications were evaluated by comparing MS/MS fragment ions, retention time, and comparability with available literature data and open-source software, including MZmine, GNPS, and SIRIUS. We confirmed a total of 82 metabolites under both ionization modes, belonging to various classes, including fatty

acids and conjugates, terpenoids, phenylpropanoids, alkaloids, saccharides and conjugates, amino acids, peptides, and analogues (S1 Table). Of all the metabolites, we found 68 and 16 in the positive and negative modes, respectively. Two metabolites, procyanidin B2 and chlorogenic acid, were detected in both modes. The total number of compounds varied among all groups. There were 26, 14, 37, 47, and 45 metabolites identified in the bark, fruits, leaves, twigs, and stalks, respectively.

Fig 1 illustrates the phytochemical classes present in five parts of *F. hispida*. The identified compounds encompassed various fatty acids and conjugates, terpenoids, phenylpropanoids, alkaloids, saccharides and conjugates, amino acids, peptides, and analogues, which accounted for 28.4%, 28.4%, 23.7%, 10.1%, 6.51%, and 2.96% of the total (Fig 1A),




Fig 1. Classification of phytochemical classes in different parts of *F. hispida*. (A) The number of differential metabolites that accumulated predominantly in each part of *F. hispida*. (B) % Composition of metabolite in each phytochemical class in five parts of *F. hispida*.

<https://doi.org/10.1371/journal.pone.0343411.g001>

respectively. When comparing the five parts of the extract, the maximum numbers of metabolites in fatty acids and conjugates and terpenoids were accumulated in the twigs and stalks. The phenylpropanoid content was found in the bark and twig extracts, respectively. In contrast, alkaloids, saccharides and conjugates, and amino acids, peptides, and analogues were predominantly accumulated in leaves, bark, and twigs together with stalks, respectively. Comparing the phytochemical classes in each part of the plant, the bark extract contained the greatest phenylpropanoid content, with 38.5% (Fig 1B). The fruit and twig extracts had higher amounts of fatty acids and conjugates than other classes, with 42.9% and 38.3%, respectively. The leaf and stalk extracts comprised the highest amount of terpenoids, with 29.7% and 33.3%, respectively. The results indicated that the metabolite content varied among the five parts of the *F. hispida* extract for each chemical class.

3.2 The metabolic variations among the five parts of *F. hispida*

To differentiate the various metabolite profiles across the five parts of the ethanolic *F. hispida* extract, multivariate analyses were conducted, including unsupervised principal component analysis (PCA) score plots and supervised partial least squares discrimination analysis (PLS-DA) score plots, utilizing UHPLC-QTOF-MS/MS data in both positive and negative modes. Results are shown in Fig 2. The PCA, unsupervised multivariate analysis, was employed to evaluate natural sample variance, clustering, and outlier identification. In the positive mode, PC1 and PC2 explained 44.3% and 39.1% of the variance, respectively (Fig 2A). The initial two components accounted for 83.4% of the overall data variance. As a result, metabolite profiles among five parts were grouped into 3 distinct clusters, namely bark, leaves, and overlapping of fruits, twigs, and stalks. Bark, fruits, twigs, and stalks exhibited a close correlation, whereas these four parts were separated from leaf extract. On the other hand, PCA score plots for the negative data analysis demonstrated a distinct division between the bark extract and other groups (Fig 2B). PC1 and PC2 accounted for 67.1% and 14.2% of the variance, respectively. These first two factors explained 81.3% of the variation in the data. In addition, PCA score plots for both modes showed that all samples in the score plots were within the 95% Hotelling's T-squared ellipse. These results indicated no outlier among the analyzed samples. To further investigate differences in metabolic variance among the five parts, a supervised analysis of PLS-DA was employed. In positive mode, the PLS-DA revealed that the samples could be divided into five distinct clusters (Fig 2C). As a result, there were differences in the metabolic variations of the five parts in positive mode. In particular, leaf extract was discriminated from other groups. The PLS1 explained 35.8% of the variation, while PLS2 explained 14.0%, according to the PLS-DA score plots. The quality metrics for PLS-DA were indicated by R^2 and Q^2 to evaluate modeling and predictive performance. The values of R^2 and Q^2 were 0.734 and 0.660, respectively. This model was considered reliable and showed no indications of overfitting. The PLS-DA score plot in negative mode showed clear differences in the metabolite patterns between the bark extract and the others (Fig 2D). PLS1 and PLS2 explained 66.8% and 12.9%, respectively. The model showed R^2 and Q^2 values were 0.758 and 0.701, respectively. Accordingly, R^2 and Q^2 confirmed that the model was effective with satisfactory predictable accuracy.

3.3 The distribution of metabolites in the five parts of *F. hispida*

The hierarchical cluster analysis and heatmap plot were performed to cluster and visualize the content profile of the metabolites across five parts of *F. hispida* extract. Fig 3 displayed the distribution of different metabolites among the five parts. In positive mode, the hierarchical cluster showed that bark, fruits, leaves, twigs, and stalks were separated into five clusters (Fig 3A). The leaf extract was sorted out from the other groups. Following the deeper colors on the heatmap plot (Fig 3B), each part of the extract exhibited different metabolite expression levels. The twig extract had the greatest metabolite expression, followed by the leaf and stalk extracts, whereas bark and fruit extracts exhibited lower concentration levels. Conversely, differential metabolites in negative mode exhibited a clear clustering pattern into two groups, comprising bark and other components (Fig 3C). The heatmap plot distinctly illustrated two variations in metabolite distribution characterized by high and low expression levels (Fig 3D). The main distribution of metabolites was found in the bark extract, while extracts from fruits, leaves, twigs, and stalks exhibited minimal distribution and contents of metabolites.




Fig 2. Multivariate data analysis of the metabolites. Principal component analysis (PCA) scores plots and partial least square-discriminant analysis (PLS-DA) scores plots of bark, leaf, fruit, twig, and stalk extracts. PCA score plots in positive mode (A) and negative mode (B). PLS-DA scores plot for positive mode (C) and negative mode (D). Shaded areas are the 95% confidence regions of each group.

<https://doi.org/10.1371/journal.pone.0343411.g002>

3.4 The significantly different metabolites among the five parts

The metabolites that significantly differed among the five parts were determined by an ANOVA statistical assay with p -values less than 0.05 and PLS-DA based on the values of variable importance in the projection (VIP) more




Fig 3. Hierarchical cluster and heat map illustrating the distribution of metabolites across the five parts of *F. hispida*. (A) Hierarchical cluster analysis between five parts and (B) heat map of metabolite distribution in positive mode. (C) Hierarchical cluster analysis between five parts and (D) heat map of metabolite distribution in negative mode. Each sample is represented by a column, and each metabolite is represented by a row. The blue and red colors refer to low and high abundance, respectively.

<https://doi.org/10.1371/journal.pone.0343411.g003>

than 1. Thirteen metabolites significantly differed among the five parts of *F. hispida* extract, including phytosphingosine, 9-Oxo-10E,12Z- octadecadienoic acid, delta-9-tetrahydrocannabinol, catharanthine, 2-[[[(2R,3S)-2-(3,4-dihydroxyphenyl)-3,5-dihydroxy-3,4-dihydro-2H-chromen-7-yl]oxy]oxane-3,4,5-triol, 13S-hydroxy-9Z,11E, 15Z-octadecatrienoic acid, pheophorbide A, 1-hexadecanoyl-sn-glycerol, harmol, quinic acid, embelin, procyanidin B2, and cianidanol (Fig 4A and 4B). Among these, phytosphingosine, 9-Oxo-10E,12Z-octadecadienoic acid, 2-[[[(2R,3S)-2-(3,4-dihydroxyphenyl)-3,5-dihydroxy-3,4-dihydro-2H-chromen-7-yl]oxy]oxane-3,4,5-triol, 13S-hydroxy-9Z,11E,15Z-octadecatrienoic acid, pheophorbide A, and 1-hexadecanoyl-sn-glycerol had the highest expression in twigs. Delta-9-tetrahydrocannabinol and catharanthine showed maximum expression in stalks and leaves, respectively, while harmol, quinic acid, embelin, procyanidin B2, and cianidanol exhibited maximum accumulation in bark.

3.5 Cytotoxicity of crude extract against CCA cell lines

The viability of KKU-213A and KKU-055 cells was expressed in Fig 5. The results showed that all crude extracts reduced a percentage of cell viability on the two CCA cell lines in a dose-dependent manner. Bark and leaf extracts significantly inhibited CCA cell viability more than fruit, twig, and stalk. The IC_{50} values (S2 in S1 Table) of bark, leaves, fruits, twigs, stalks, and gemcitabine on KKU-213A were 0.71 ± 0.17 , 0.82 ± 0.22 , 7.96 ± 3.29 , 7.53 ± 4.24 , 8.51 ± 1.59 , and 0.02 ± 0.00 $\mu\text{g/mL}$, respectively. The IC_{50} values of bark, leaves, fruits, twigs, stalks, and gemcitabine on KKU-055 were 0.78 ± 0.13 , 1.03 ± 0.21 , 11.5 ± 3.18 , 9.98 ± 7.02 , 11.9 ± 1.72 , and 0.05 ± 0.00 $\mu\text{g/mL}$, respectively. These revealed that all crude extracts showed potential inhibition of CCA cell lines. In particular, the bark and leaf extracts had the lowest IC_{50} values on the two CCA cell lines. Compared to gemcitabine, the reference drug, the cytotoxicity of those extracts was lower.




Fig 4. VIP score plot and significantly differential metabolites among the five parts of *F. hispida*. VIP scores are derived from PLS-DA analysis at different parts (bark; B, fruits; F, leaves; L, stalks; ST, twigs; TW) of *F. hispida* in (A) positive mode and (B) negative mode. The column to the right of each figure indicates variations in metabolite peak intensities. * is added to the significant metabolites with VIP scores >1 and $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0343411.g004>




Fig 5. Cytotoxic effect of various parts of *F. hispida* on CCA cell viability. (A and B) KKU-213A and (C and D) KKU-055 after treatment with each extract for 48 h (* and ** indicate statistical significance; $p < 0.05$ and $p < 0.001$, respectively).

<https://doi.org/10.1371/journal.pone.0343411.g005>

3.6 Identification of key metabolites correlated to the anticancer activities of *F. hispida* extract

The univariate analysis with Pearson's correlation coefficient was determined between thirteen of the significant metabolites and the IC_{50} values. This correlation can evaluate the extent to which one variable may impact another, potentially indicating a positive or negative correlation. A heatmap of three colors was plotted to observe the correlation data: the shades of red represent positive correlations ($0 < r < 1$), whereas shades of blue represent negative correlations ($-1 < r < 0$). A significant correlation was considered at a p -value less than 0.05. Among high-level metabolites, catharanthine showed a strong negative correlation as well as p -values less than 0.01 with the IC_{50} values ($r = -0.615$ (KKU-213A), -0.634 (KKU-055)) (Fig 6 and S3 in S1 Table). Moreover, there were 4 metabolites that showed a moderate negative correlation ($r = (-0.40) - (-0.59)$) as well as p -values less than 0.05 with the cytotoxicity test (IC_{50} value) in the following order: cianidanol > procyanidin B2 > quinic acid > embelin (Fig 6 and S3 in S1 Table).

3.7 Cytotoxic effect of key metabolites on CCA cell lines

The cytotoxic effect of key metabolites on CCA cells was performed using the SRB assay. The IC_{50} values were assessed after KKU-213A and KKU-055 cells were treated with catharanthine, cianidanol, procyanidin B2, and quinic acid for 72 h. The percentage of cell viability is shown in Fig 7. Catharanthine, cianidanol, and procyanidin B2 decreased the cell viability of KKU-213A and KKU-055 in dose-dependent manners. The IC_{50} values of catharanthine, cianidanol, and procyanidin B2 on KKU-213A were 41.0 ± 0.99 , 812 ± 166 , and $>500 \mu M$ (S4 in S1 Table), respectively. The IC_{50} values of




Fig 6. Correlation patterns of metabolites and anticancer activities. Correlation map of the metabolites analyzed by UHPLC-QTOF-MS/MS with cytotoxicity (IC₅₀). The shades of red represent positive correlations ($0 < r < 1$), whereas shades of blue represent negative correlations ($-1 < r < 0$). Each square indicates Pearson's correlation coefficient of a pair of metabolites and IC₅₀. **, * indicates p -values less than 0.05 with a strong correlation ($r = (-0.6) - (-0.79)$) and a moderate correlation ($r = (-0.40) - (-0.59)$), respectively.

<https://doi.org/10.1371/journal.pone.0343411.g006>

catharanthine, cianidanol, and procyanidin B2 on KKU-055 were 47.4 ± 4.34 , 474 ± 155 , and >500 μ M. Nevertheless, quinic acid at the concentration of 0–1000 μ M did not inhibit the viability of KKU-213A and KKU-055. The result indicated that catharanthine and cianidanol exhibited a cytotoxicity effect against KKU-213A and KKU-055 cells.

4. Discussion

Our study revealed a metabolite profile of five parts of *F. hispida* (bark, fruits, leaves, twigs, and stalks) using UHPLC-QTOF-MS/MS in both positive and negative ionization modes, demonstrating 82 metabolites across various phytochemical classes, including fatty acids and conjugates, terpenoids, phenylpropanoids, alkaloids, saccharides and conjugates, amino




Fig 7. Cytotoxic effect of key metabolites on CCA cell viability. (A) KKU-213A and (B) KKU-055 after treatment with 0, 1, 10, 100, 500, and 1,000 μM of catharanthine, cianidanol, procyanidine B2, and quinic acid for 72 h.

<https://doi.org/10.1371/journal.pone.0343411.g007>

acids, peptides, and their analogues. The majority of metabolites in positive ionization mode corresponded to previous findings indicating that ethanol extracts of *Ficus* species prefer the ionization of non-polar or moderately polar compounds, such as terpenoids and fatty acids [4,20]. Consistent with earlier reports, terpenoids, phenylpropanoids, and flavonoids were identified as the main chemical components contributing to the antioxidant and anti-inflammatory properties of *F. hispida* [2]. Bark extract contained a high proportion of phenylpropanoids (38.5%). It is rich in phenolic compounds that may contribute to its traditional use in skin and liver diseases. Terpenoids, which are known as photoprotective and cytoprotective agents, were abundant in leaves and stalks [21]. The abundance of fatty acids and conjugates in twigs and fruits, such as linoleic, linolenic, and 13-hydroxy-octadecatrienoic acids, indicates their functions in structural and energy storage metabolism, supporting the tissue-specific specialization observed in other *Ficus* taxa [22,23]. In addition, the discovery of alkaloids, such as catharanthine and harmol, in various regions enriches the chemical profile of *F. hispida* and aligns with evidence suggesting that phenanthroindolizidine alkaloids exhibit antiproliferative effects [24,25]. Our results demonstrate more chemical diversity and clearer patterns of accumulation for each part compared to previous studies that primarily examined single parts or isolated compound groups. This comprehensive profiling highlights *F. hispida* as a chemically versatile medicinal species, revealing previously unrecognized insights into its tissue-specific metabolite biosynthesis and affirming its ethnomedicinal significance for antioxidant, anti-inflammatory, and anticancer applications.

Thereafter, we used PCA, PLS-DA, the hierarchical cluster, and heatmaps to evaluate the similarities and differences in metabolite profiles between five parts of *F. hispida* extract. In positive mode, the results indicated distinct clustering patterns among the five groups, with the leaf extract separated from other groups. For the negative mode, there was some overlapping of metabolite profiles, but bark extract separated from the other four parts. The hierarchical cluster and heatmaps of metabolites confirmed separation of each group. The results showed that the types and the content of metabolites were different in each group. The total number and the contents of compounds were found to be the highest in the ethanolic twig extract for positive mode, while negative mode indicated the highest metabolite distribution only in bark extract. Furthermore, the fruit, leaf, twig, and stalk extracts exhibited a very low number of metabolites (1–3) in the negative mode. This phenomenon explains why the metabolite profiles in the four different parts overlapped in negative mode.

Subsequently, the VIP score highlighted significant metabolites based on relative variables, facilitating the identification of potential metabolite markers. This study identified thirteen metabolites that vary significantly in content across the five parts of the *F. hispida* extract. Among thirteen metabolites, 9-oxo-10E,12Z-octadecadienoic acid, delta-9-tetrahydrocannabinol, catharanthine, 2-[[[(2R,3S)-2-(3,4-dihydroxyphenyl)-3,5-dihydroxy-3,4-dihydro-2H-chromen-7-yl] oxy]oxane-3,4,5-triol, 13S-hydroxy-9Z,11E,15Z-octadecatrienoic acid, 1-hexadecanoyl-sn-glycerol, harmol, and embelin are plant metabolites that have not been reported in any *Ficus* species. These metabolites were initially discovered in *F. hispida*, which may contribute to the medicinal properties of this plant. In contrast, phytosphingosine, pheophorbide A, and procyanidin B2 have not been reported in *F. hispida*; however, they are found in other *Ficus* species [26–29]. Of these, quinic acid and cianidanol were reported in *F. hispida* extract in a previous study [3].

The results showed that five parts of *F. hispida* have potent anticancer activity to CCA cell lines of KKU-213A and KKU-055 with IC_{50} values of 0.71 ± 0.17 – 11.9 ± 1.72 $\mu\text{g/mL}$. This value conformed to the anticancer potency set by the National Cancer Institute standard for the plant extracts with an $IC_{50} < 30$ $\mu\text{g/mL}$ [30]. Previous studies on the cytotoxicity test of bark and leaf extracts against human cancer cell lines (U87MG, A549, HT-29) showed that the IC_{50} values were higher than those in this study [31]. The cytotoxicity of fruit extract was in agreement with a previous study by Jie Zhang et al., 2018. They reported the IC_{50} values of fruit extract on human cancer cell lines (HL60, A549, SKBR3, KB, Hela, HT29, HepG2, and LO2) were in the range of 0.1–100 $\mu\text{g/mL}$ [13]. Furthermore, our results revealed that bark and leaf extracts had the greatest potential cytotoxicity against the tested CCA cells. The results related to the variation of metabolites is that the main metabolite in bark extract consists of phenolic compounds. Many studies define phenolic compounds as potential anticancer agents against several types of cancer [32]. The leaf extract showed the maximum number of metabolites in the alkaloids class compared to other extracts. Numerous research studies have confirmed the antiproliferative and anticancer effects of the alkaloids [33]. Therefore, phenolic compounds and alkaloid metabolites may play an important role in anticancer activities of bark and leaf extracts.

After evaluation of the cytotoxicity assay, univariate analysis was performed to identify key metabolites that are responsible for anticancer capability. As shown in previous analyses, thirteen metabolites were identified that significantly differed among five parts. The correlation analysis between significant metabolites with cytotoxicity (IC_{50} values) revealed a strong negative correlation to catharanthine that exhibited high accumulation in bark and leaf extracts. Catharanthine is a pharmaceutically important antitumor alkaloid that is found in *Catharanthus roseus* [34]. It was reported to have an anticancer effect on liver carcinoma cells (HepG2) [35], leukemia cells (HL60, K562) [36], and colorectal carcinoma cells (HCT 116) [37]. Moreover, a moderate negative correlation appeared in four metabolites, including cianidanol, procyanidin B2, quinic acid, and embelin, that exhibited the highest accumulation in bark extract. Cianidanol, or (+)-catechin, a polyphenolic antioxidant plant metabolite, occurs in green tea, red wine, chocolate, and cocoa [38]. The metabolite shows various medicinal effects, such as antioxidant, antimicrobial, anticancer, and anti-inflammatory properties [39]. Procyanidin B2, a flavonoid compound, is a potent natural antioxidant that is

present in grape seeds, apples, and cacao beans [40]. It has properties that fight cancer, viruses, inflammation, and free radicals [41]. Additionally, prior studies indicated that procyanidin B2 suppressed the proliferation of liver, breast, and gastric cancer cells [42–44]. Quinic acid is a naturally occurring phenolic compound found in many plants, such as coffee, bilberries, prunes, cranberries, and kiwifruit [45–47]. These compounds exhibited various biological activities, such as antioxidant, antidiabetic, anticancer, antimicrobial, antiviral, aging-protective, antinociceptive, and analgesic effects [39]. Embelin is a plant-based benzoquinone alkaloid that displays various biological activities, including antioxidant, antidiabetic, analgesic, antibacterial, antitumor, and anti-inflammatory activities [48,49]. These compounds exhibited various biological activities, such as antioxidant, antidiabetic, antimicrobial, antiviral, aging-protective, antinociceptive, and anticancer effects [50].

Subsequently, we confirmed the anticancer effect of the metabolites on CCA cells. Among four potential metabolites, including catharanthine, cianidanol, procyanidin B2, and quinic acid, our result showed that catharanthine, cianidanol, and procyanidin B2 had cytotoxicity on CCA cells. Nonetheless, catharanthine exhibited the most significant inhibitory effect on CCA cell viability. Catharanthine is a precursor to the potent chemotherapeutic vinca alkaloids, such as vinblastine and vincristine [51]. This finding aligns with previous research, indicating that alkaloids play a role in the anticancer activity of *F. hispida* (bark and leaves) [18]. A previous study has shown that hispiloscine, a phenanthroindolizidine alkaloid isolated from the stems, bark, and leaves of the Malaysian *F. hispida*, showed appreciable antiproliferative activities against MDA-MB-231, MCF-7, A549, HCT-116, and MRC-5 cell lines [25]. O-methyltylophorinidine was isolated from the leaves and stalks of *F. hispida* and showed a potent cytotoxic effect on Lu1 (lung) and Col2 (colon) cancer cells [52]. A comparison of IC₅₀ values to the crude extracts, catharanthine (41–47 μ M or 13–16 μ g/mL) was found to be higher than the bark and leaf extracts (0.7–1 μ g/mL). Due to plants containing plenty of phytochemical substances and targeting several biological pathways, they can occasionally work synergistically to increase overall potency and solubility. Thus, these substances could sometimes be more potent than the pure substance [53,54]. In addition, extracts work in concert to enhance activity, potentially to lessen side effects and drug resistance, while pure compounds offer precision in targeting cancer cells.

We have shown, for the first time, that the anticancer effect attributed to catharanthine on CCA cells is present in the bark and leaves of *F. hispida*. Because there are no previous reports on the use of catharanthine against CCA, we can only compare the IC₅₀ value to reported results for other cancers. We found that the IC₅₀ value of catharanthine for CCA cells in this study was lower than that observed for HepG2 cells (IC₅₀ at 130–135 μ M) in the past. They demonstrated that catharanthine-triggered apoptosis was linked to increased autophagy signaling pathways by interacting with caspase-8, inhibiting the PI3K/Akt/mTOR pathway, upregulating Beclin-1, LC3, and ULK1, and enhancing SIRT1 expression [35]. In CCA cells, they may be cytotoxically affected by catharanthine via a synergistic anti-cancer effect through a multimodal mechanism of action typical of vinca alkaloids. Hence, it is necessary to further clarify the biological mechanisms of catharanthine that underlie its anti-CCA activity. Furthermore, in comparison to chemotherapy drugs, such as gemcitabine, the cytotoxic potency of catharanthine is typically lower *in vitro* and requires higher dosages to impede growth as much. Despite this, its different mechanisms of action, such as activating autophagy signaling pathways, highlight the potential of catharanthine for development as an adjuvant or combination therapy. This could potentially enhance the efficacy of treatment or help overcome drug resistance in cancer cells. Nonetheless, our study has limitations, as all findings were derived from *in vitro* experiments, which fail to adequately represent the complexity of tumor-stromal interactions or pharmacokinetic constraints *in vivo*. In addition to this, data on the stability, bioavailability, and metabolic pathway of catharanthine are lacking, limiting clinical interpretation. Concurrently with clarifying the mechanisms of action in CCA cells, further studies should focus on validating results in animal models, analyzing pharmacokinetics, and evaluating synergistic effects with standard chemotherapeutic agents to more accurately determine the therapeutic potential of catharanthine in cholangiocarcinoma.

5. Conclusion

Our findings show that *F. hispida* contains a wide range of metabolites and has a pharmacological potential that provides a valuable source for natural product development. The identification of catharanthine as a major cytotoxic metabolite provides new information about the bioactive compounds in this species, and establishes a promising basis for future mechanistic and therapeutic advancements against CCA. Further research should focus on elucidating the exact anti-cancer mechanisms by which catharanthine exerts anti-CCA effects, paving the way for the development of alternative chemotherapeutic agents.

Supporting information

S1 File. Plant material. Detailed information on plant identification.
(PDF)

S1 Table. Metabolites identified and phytochemical classes in various parts of *F. hispida* extract by UHPLC-QTOF-MS/MS analysis.
(ZIP)

Acknowledgments

We would like to acknowledge Professor Ross H Andrews for editing this manuscript.

Author contributions

Conceptualization: Saengrawee Thammawithan, Nisana Namwat.

Data curation: Saengrawee Thammawithan, Thapanee Pruksatrakul.

Formal analysis: Saengrawee Thammawithan, Sirinya Sitthirak, Thapanee Pruksatrakul.

Funding acquisition: Watcharin Loilome, Nisana Namwat.

Investigation: Saengrawee Thammawithan, Jirattiporn Thanuma, Piya Prajumwongs.

Methodology: Saengrawee Thammawithan, Jirattiporn Thanuma, Chadapohn Panplu.

Resources: Arporn Wangwiwatsin, Poramate Klanrit, Watcharin Loilome, Nisana Namwat.

Supervision: Nisana Namwat.

Validation: Saengrawee Thammawithan.

Visualization: Saengrawee Thammawithan, Sirinya Sitthirak.

Writing – original draft: Saengrawee Thammawithan.

Writing – review & editing: Saengrawee Thammawithan, Arporn Wangwiwatsin, Watcharin Loilome, Nisana Namwat.

References

1. Ripu MK, Bussmann R. *Ficus* (Fig) species in Nepal: A review of diversity and indigenous uses. *J Ecol Appl*. 2006;11:85–97.
2. Ali M, Chaudhary N. *Ficus hispida* Linn.: A review of its pharmacognostic and ethnomedicinal properties. *Pharmacogn Rev*. 2011;5(9):96–102. <https://doi.org/10.4103/0973-7847.79104> PMID: 22096323
3. Kumar S, Singh A, Kushwaha AK, Tiwari R, Chaudhary LB, Srivastava M, et al. The UPLC–ESI–QqQ LIT–MS/MS method for quantitative determination of phytochemicals in ethanolic extracts of different parts of eight *Ficus* species: Development and validation. *International Journal of Food Properties*. 2018;21(1):328–44. <https://doi.org/10.1080/10942912.2017.1414838>
4. Cheng J-X, Zhang B-D, Zhu W-F, Zhang C-F, Qin Y-M, Abe M, et al. Traditional uses, phytochemistry, and pharmacology of *Ficus hispida* L.f.: A review. *J Ethnopharmacol*. 2020;248:112204. <https://doi.org/10.1016/j.jep.2019.112204> PMID: 31669442

5. Shen S, Zhan C, Yang C, Fernie AR, Luo J. Metabolomics-centered mining of plant metabolic diversity and function: Past decade and future perspectives. *Mol Plant*. 2023;16(1):43–63. <https://doi.org/10.1016/j.molp.2022.09.007> PMID: 36114669
6. Manickam S, Rajagopalan VR, Kambale R, Rajasekaran R, Kanagarajan S, Muthurajan R. Plant Metabolomics: Current Initiatives and Future Prospects. *Curr Issues Mol Biol*. 2023;45(11):8894–906. <https://doi.org/10.3390/cimb45110558> PMID: 37998735
7. Plumb RS, Gethings LA, Rainville PD, Isaac G, Trengove R, King AM, et al. Advances in high throughput LC/MS based metabolomics: A review. *TrAC Trends in Analytical Chemistry*. 2023;160:116954. <https://doi.org/10.1016/j.trac.2023.116954>
8. Brindley PJ, Bachini M, Ilyas SI, Khan SA, Loukas A, Sirica AE, et al. Cholangiocarcinoma. *Nat Rev Dis Primers*. 2021;7(1):65. <https://doi.org/10.1038/s41572-021-00300-2> PMID: 34504109
9. Tepsiri N, Chaturat L, Sripa B, Namwat W, Wongkham S, Bhudhisawasdi V, et al. Drug sensitivity and drug resistance profiles of human intrahepatic cholangiocarcinoma cell lines. *World J Gastroenterol*. 2023;11(18):2748–53. <https://doi.org/10.3748/wjg.v11.i18.2748> PMID: 15884115
10. Marin JJG, Lozano E, Herraiz E, Asensio M, Di Giacomo S, Romero MR, et al. Chemoresistance and chemosensitization in cholangiocarcinoma. *Biochim Biophys Acta Mol Basis Dis*. 2018;1864(4 Pt B):1444–53. <https://doi.org/10.1016/j.bbadis.2017.06.005> PMID: 28600147
11. Cabanlit KL, Torres MAJ, Demayo CG. *Ficus* spp. (Figs) and their anticancer potential: A systematic review of laboratory studies and traditional uses in the Philippines. *J Herbmec Pharmacol*. 2025;14(2):133–52. <https://doi.org/10.34172/jhp.2025.52854>
12. Pratumnit B, Srisapoomi T, Worawattananon P, Oparatiattikul N, Jiratchariyakul W, Kummalue T. In vitro antineoplastic effect of *Ficus hispida* L. plant against breast cancer cell lines. *J Med Plant Res*. 2009;3(4):255–61. <https://doi.org/10.5897/JMPR.9000746>
13. Zhang J, Zhu W-F, Xu J, Kitdamrongtham W, Manosroi A, Manosroi J, et al. Potential cancer chemopreventive and anticancer constituents from the fruits of *Ficus hispida* L.f. (Moraceae). *J Ethnopharmacol*. 2018;214:37–46. <https://doi.org/10.1016/j.jep.2017.11.016> PMID: 29197545
14. Rahman H, Rahman A, Zahan S, Biswas P, Sohag AAM, Islam S, et al. A Pharmacoinformatics and molecular dynamics approach to the phytochemical screening of *Ficus hispida* fruits for prostate cancer therapy. *Curr Comput Aided Drug Des*. 2025;:10.2174/0115734099409352251103111907. <https://doi.org/10.2174/0115734099409352251103111907> PMID: 41276981
15. Tabrez S, Hoque M, Suhail M, Khan MI, Zughaibi TA, Khan AU. Identification of anticancer bioactive compounds derived from *Ficus* sp. by targeting Poly[ADP-ribose]polymerase 1 (PARP-1). *Journal of King Saud University - Science*. 2022;34(5):102079. <https://doi.org/10.1016/j.jksus.2022.102079>
16. Mutungi MM, Muema FW, Kimutai F, Xu Y-B, Zhang H, Chen G-L, et al. Antioxidant and Antiproliferative Potentials of *Ficus glumosa* and its bioactive polyphenol metabolites. *Pharmaceuticals (Basel)*. 2021;14(3):266. <https://doi.org/10.3390/ph14030266> PMID: 33804242
17. Annegowda HV, Bhat R, Min-Tze L, Karim AA, Mansor SM. Influence of sonication treatments and extraction solvents on the phenolics and antioxidants in star fruits. *J Food Sci Technol*. 2012;49(4):510–4. <https://doi.org/10.1007/s13197-011-0435-8> PMID: 23904662
18. Sérgio PL, Tiago BA, Thiago RB, Michel R dos RS, Claudia ALC, Laiza CK, et al. Ultrasonic assisted extraction of bioactive compounds from different parts of *Hancornia Speciosa* Gomes. *J Med Plants Res*. 2020;14(7):300–8. <https://doi.org/10.5897/jmpr2020.6944>
19. Vichai V, Kirtikara K. Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nat Protoc*. 2006;1(3):1112–6. <https://doi.org/10.1038/nprot.2006.179> PMID: 17406391
20. Cruz JMDA, Corrêa RF, Lamarão CV, Kinupp VF, Sanches EA, Campelo PH, et al. *Ficus* spp. fruits: Bioactive compounds and chemical, biological and pharmacological properties. *Food Res Int*. 2022;152:110928. <https://doi.org/10.1016/j.foodres.2021.110928> PMID: 35181099
21. Karinchai J, Buacheen P, Rodthayoy D, Yodweeraopong T, Imsumran A, Wongnoppavich A, et al. Anti-inflammatory and anti-adipocyte dysfunction effects of *Ficus lindsayana* latex and root extracts. *Pharmaceuticals (Basel)*. 2024;17(3):287. <https://doi.org/10.3390/ph17030287> PMID: 38543073
22. Knothe G, Razon LF, de Castro MEG. Fatty acids, triterpenes and cycloalkanes in ficus seed oils. *Plant Physiol Biochem*. 2019;135:127–31. <https://doi.org/10.1016/j.plaphy.2018.11.030> PMID: 30529978
23. Hasnat H, Alam S, Akter Shompa S, Saha T, Richi FT, Hossain MH, et al. Phyto-pharmacological wonders of genus *Ficus*: Ethnopharmacological insights and phytochemical treasures from natural products. *Saudi Pharm J*. 2024;32(12):102211. <https://doi.org/10.1016/j.jsps.2024.102211> PMID: 39697478
24. Jia X-Y, Wu Y-M, Li J-Y, Lei C, Hou A-J. Alkaloid constituents of *Ficus hispida* and their antiinflammatory activity. *Nat Prod Bioprospect*. 2020;10(1):45–9. <https://doi.org/10.1007/s13659-020-00233-5> PMID: 32072482
25. Yap VA, Loong B-J, Ting K-N, Loh SH-S, Yong K-T, Low Y-Y, et al. Hispidacine, an unusual 8,4'-oxyneolignan-alkaloid with vasorelaxant activity, and hispidosine, an antiproliferative phenanthroindolizidine alkaloid, from *Ficus hispida* Linn. *Phytochemistry*. 2015;109:96–102. <https://doi.org/10.1016/j.phytochem.2014.10.032> PMID: 25468714
26. Dongfack MDJ, Lallemand M-C, Kuete V, Mbazoa CD, Wansi J-D, Trinh-van-Dufat H, et al. A new sphingolipid and furanocoumarins with antimicrobial activity from *Ficus exasperata*. *Chem Pharm Bull (Tokyo)*. 2012;60(8):1072–5. <https://doi.org/10.1248/cpb.c12-00279> PMID: 22863713
27. Murni A, Hanif N, Kita M, Darusman LK. Methyl 10-epi-pheophorbide a from mcf-7 Cells Active Layer Of The Indonesian *Ficus Deltoidea* jack leaves. *Int J Pharm Pharm Sci*. 2017;9(8):183. <https://doi.org/10.22159/ijpps.2017v9i8.18750>
28. Lata S, Koli P, Singh S, Bhadoria BK, Chand U, Yadav DK, et al. Proanthocyanidins isolated from the leaves of *Ficus glomerata* evaluated on the activities of rumen enzymes: in vitro and in silico studies. *Front Chem*. 2024;12:1359049. <https://doi.org/10.3389/fchem.2024.1359049> PMID: 38380397
29. Sonibare O, Sojinu S. Chemical composition of leaf lipids of angiosperms: origin of land plant-derived hydrocarbons in sediments and fossil fuel. *Eur J Sci Res*. 2009;25(2):192–9.

30. Suffness M, Pezzuto JM, Hostettmann K. Methods in plant biochemistry: assays for bioactivity. In: Hostettmann K, editor. *Methods in Plant Biochemistry*. 1990. p. 33–71.
31. Abubakar IB, Lim K-H, Loh H-S. Alkaloid extracts of *Ficus* species and palm oil-derived tocotrienols synergistically inhibit proliferation of human cancer cells. *Nat Prod Res*. 2015;29(22):2137–40. <https://doi.org/10.1080/14786419.2014.991927> PMID: 25515603
32. Bakrim S, El Omari N, El Hachlafi N, Bakri Y, Lee L-H, Bouyahya A. Dietary phenolic compounds as anticancer natural drugs: recent update on molecular mechanisms and clinical trials. *Foods*. 2022;11(21):3323. <https://doi.org/10.3390/foods11213323> PMID: 36359936
33. Mondal A, Gandhi A, Fimognari C, Atanasov AG, Bishayee A. Alkaloids for cancer prevention and therapy: Current progress and future perspectives. *Eur J Pharmacol*. 2019;858:172472. <https://doi.org/10.1016/j.ejphar.2019.172472> PMID: 31228447
34. Ferreres F, Pereira DM, Valentão P, Oliveira JMA, Faria J, Gaspar L, et al. Simple and reproducible HPLC-DAD-ESI-MS/MS analysis of alkaloids in *Catharanthus roseus* roots. *J Pharm Biomed Anal*. 2010;51(1):65–9. <https://doi.org/10.1016/j.jpba.2009.08.005> PMID: 19720492
35. Gholami F, Seyedalipour B, Heidari-Kalvani N, Nabi-Afjadi M, Yaghoubzad-Maleki M, Fathi Z, et al. Catharanthine, an anticancer vinca alkaloid: an *in silico* and *in vitro* analysis of the autophagic system as the major mechanism of cell death in liver HepG2 cells. *Naunyn Schmiedeberg Arch Pharmacol*. 2024;397(11):8879–92. <https://doi.org/10.1007/s00210-024-03191-8> PMID: 38856913
36. Wang X-D, Li C-Y, Jiang M-M, Li D, Wen P, Song X, et al. Induction of apoptosis in human leukemia cells through an intrinsic pathway by catha-chunine, a unique alkaloid isolated from *Catharanthus roseus*. *Phytomedicine*. 2016;23(6):641–53. <https://doi.org/10.1016/j.phymed.2016.03.003> PMID: 27161405
37. Siddiqui MJ, Ismail Z, Aisha AFA, Abdul Maji AMS. Cytotoxic Activity of *Catharanthus roseus* (Apocynaceae) crude extracts and pure compounds against human colorectal carcinoma cell line. *International J of Pharmacology*. 2009;6(1):43–7. <https://doi.org/10.3923/ijp.2010.43.47>
38. Bernatoniene J, Kopustinskiene DM. The role of catechins in cellular responses to oxidative stress. *Molecules*. 2018;23(4):965. <https://doi.org/10.3390/molecules23040965> PMID: 29677167
39. Bae J, Kim N, Shin Y, Kim S-Y, Kim Y-J. Activity of catechins and their applications. *biomed dermatol*. 2020;4(1). <https://doi.org/10.1186/s41702-020-0057-8>
40. Chen J, Zhong K, Jing Y, Liu S, Qin S, Peng F, et al. Procyanidin B2: A promising multi-functional food-derived pigment for human diseases. *Food Chem*. 2023;420:136101. <https://doi.org/10.1016/j.foodchem.2023.136101> PMID: 37059021
41. Lee Y. Cancer chemopreventive potential of procyanidin. *Toxicol Res*. 2017;33(4):273–82. <https://doi.org/10.5487/TR.2017.33.4.273> PMID: 29071011
42. Li Y, Lu X, Tian P, Wang K, Shi J. Procyanidin B2 induces apoptosis and autophagy in gastric cancer cells by inhibiting Akt/mTOR signaling pathway. *BMC Complement Med Ther*. 2021;21(1):76. <https://doi.org/10.1186/s12906-021-03225-1> PMID: 33627124
43. Avelar MM, Gouvêa CMCP. Procyanidin b2 cytotoxicity to mcf-7 human breast adenocarcinoma cells. *Indian J Pharm Sci*. 2012;74(4):351–5. <https://doi.org/10.4103/0250-474X.107070> PMID: 23626391
44. Liu G, Shi A, Wang N, Li M, He X, Yin C, et al. Polyphenolic Proanthocyanidin-B2 suppresses proliferation of liver cancer cells and hepatocellular carcinogenesis through directly binding and inhibiting AKT activity. *Redox Biol*. 2020;37:101701. <https://doi.org/10.1016/j.redox.2020.101701> PMID: 32863234
45. Ryan JJ, Dupont JA. Identification and analysis of the major acids from fruit juices and wines. *J Agric Food Chem*. 1973;21(1):45–9. <https://doi.org/10.1021/jf60185a018>
46. Coppola ED, Conrad EC, Cotter R. High pressure liquid chromatographic determination of major organic acids in cranberry juice. *J Assoc Off Anal Chem*. 1978;61(6):1490–2. <https://doi.org/10.1093/jaoac/61.6.1490> PMID: 730655
47. Uleberg E, Rohloff J, Jaakola L, Tröst K, Junttila O, Häggman H, et al. Effects of temperature and photoperiod on yield and chemical composition of northern and southern clones of bilberry (*Vaccinium myrtillus* L.). *J Agric Food Chem*. 2012;60(42):10406–14. <https://doi.org/10.1021/jf302924m> PMID: 23033879
48. Sheng Z, Ge S, Gao M, Jian R, Chen X, Xu X, et al. Synthesis and biological activity of embelin and its derivatives: an overview. *Mini Rev Med Chem*. 2020;20(5):396–407. <https://doi.org/10.2174/1389557519666191015202723> PMID: 31644404
49. Danquah M. Embelin and its derivatives: design, synthesis, and potential delivery systems for cancer therapy. *Pharmaceuticals (Basel)*. 2022;15(9):1131. <https://doi.org/10.3390/ph15091131> PMID: 36145352
50. Nguyen V, Taine EG, Meng D, Cui T, Tan W. Chlorogenic acid: a systematic review on the biological functions, mechanistic actions, and therapeutic potentials. *Nutrients*. 2024;16(7):924. <https://doi.org/10.3390/nu16070924> PMID: 38612964
51. Mendonça KC, Palani N, Rajadesingu S, Radhakrishnan K, Ayyar M, Priya LS. Pharmacological potential of bioactive compounds in *Catharanthus roseus* extract: A comprehensive review. *Toxicol Rep*. 2025;14:101998. <https://doi.org/10.1016/j.toxrep.2025.101998> PMID: 40213418
52. Peraza-Sánchez SR, Chai H-B, Shin YG, Santisuk T, Reutrakul V, Farnsworth NR, et al. Constituents of the leaves and twigs of *Ficus hispida*. *Planta Med*. 2002;68(2):186–8. <https://doi.org/10.1055/s-2002-20257> PMID: 11859479
53. Rasoanaivo P, Wright CW, Willcox ML, Gilbert B. Whole plant extracts versus single compounds for the treatment of malaria: synergy and positive interactions. *Malar J*. 2011;10 Suppl 1(Suppl 1):S4. <https://doi.org/10.1186/1475-2875-10-S1-S4> PMID: 21411015
54. Aiyelaagbe OO, Hamid AA, Fattorusso E, Taglialatela-Scafati O, Schröder HC, Müller WEG. Cytotoxic activity of crude extracts as well as of pure components from *jatropha* species, plants used extensively in african traditional medicine. *Evid Based Complement Alternat Med*. 2011;2011:134954. <https://doi.org/10.1155/2011/134954> PMID: 21754941