

Ethnomedical knowledge of plants used in alternative medicine for the treatment of hemorrhoids in Lubumbashi, Haut-Katanga Province, Southern Democratic Republic of Congo

Bashige Chiribagula Valentin

`bashige.chiribagula@unilu.ac.cd`

University of Lubumbashi

Okusa Ndjolo Philippe

University of Lubumbashi

Muhona Melman

University of Lubumbashi

Manya Mboni Henry

University of Lubumbashi

Bakari Amuri Salvius

University of Lubumbashi

Lumbu Simbi Jean Baptiste

University of Lubumbashi

Research Article

Keywords: Ficus stuhlmannii, Khaya nyasica, Phyllanthus amarus, Lubumbashi, Traditional healers, hemorrhoids.

Posted Date: July 22nd, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-4608274/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

Ethnomedical knowledge of plants used in alternative medicine for the treatment of hemorrhoids in Lubumbashi, Haut-Katanga Province, Southern Democratic Republic of Congo

Bashige Chiribagula Valentin^{*1}, Okusa Ndjolo Philippe¹, Muhona Melman ¹, Many Mboni Henry¹, Bakari Amuri Salvius², Lumbu Simbi Jean Baptiste³

¹ Department of Pharmacology - Laboratory of Therapeutic Chemistry and Analysis of Natural Substances - Faculty of Pharmaceutical Sciences - University of Lubumbashi (UNILU) - 27, av. Kato, Commune Kampemba, Lubumbashi - DR Congo.

² Department of Pharmacology - Laboratory of Pharmacognosy – Faculty of Pharmaceutical Sciences – University of Lubumbashi (UNILU)- 27, av Kato, Commune Kampemba, Lubumbashi - DR Congo.

³ Department of Chemistry– Faculty of Sciences–University of Lubumbashi (UNILU) N°1 Maternity Avenue, Commune of Lubumbashi, Lubumbashi - DR Congo.

Corresponding author: Bashige Chiribagula Valentin: bashige.chiribagula@unilu.ac.cd; bachival@gmail.com

Author's ORCID

Bashige Chiribagula Valentin (0000-0002-1371-0517), **Okusa Ndjolo Philippe** (0000-0002-5082-4097), **Muhona Melman** (0009-0006-0938-8373), **Many Mboni Henry** (0000-0001-9236-7288), **Bakari Amuri Salvius** (0000-0001-6360-8660), **Lumbu Simbi Jean Baptiste** (0000-0002-8889-3397).

Abstract

Background: A variety of medicinal plants are used in traditional medicine in Lubumbashi for the management of hemorrhoidal diseases. However, no investigation has been conducted to gather the knowledge required for this type of management in the region. The present study was conducted to inventory the plants used in Lubumbashi to treat hemorrhoidal diseases and to relate their ethnomedical characteristics.

Methods: This study was conducted between March 2022 and February 2023 using semi-structured interviews with households (n=1520), herbalists (n=25), and traditional healers : THs (n=59).


Results: The 1,604 respondents (sex ratio M/F = 0.9; mean age: 56 ± 3 years; experience: 12 ± 3 years) provided information on 100 taxa, 84 of which are used against internal hemorrhoids, *Phyllanthus amarus* being the most cited (Citation Index, CI: 0.76). Most of them are trees (38%) or shrubs (32%), belonging to 90 genera and 45 families dominated by the Fabaceae (10%) and Asteraceae (9%). They are indicated in 76 other pathologies, dominated by gastrointestinal disorders (GID), wounds and sexually transmitted infections (CI > 0.57). From these 100 taxa, 117 anti-hemorrhoidal formulations were derived, 11 of which combined more than one plant. In all these recipes, the leaf is the most commonly used part (>60%) and the liniment (>45%) is the most popular form of application. For the first time, this study reports 14 taxa as plants used in the treatment of hemorrhoids for which some evidence of hemorrhoidal activity has been reported. Among these taxa, *Ficus stuhlmannii*, *Ficus laurifolia*, and *Ocimum centraliafricanum* are listed as medicinal plants for the first time. *Khaya nyasica*, and *Syzygium cordatum*, each with 11 uses, have the highest traditional medicinal value.

Conclusion: The findings of this study indicate that a significant number of medicinal plants are used in traditional medicine in Lubumbashi for the treatment of hemorrhoidal diseases. Some of these plants are endemic to the biodiversity area, while

others are shared with other cultures and regions. A series of pharmacological studies is currently underway with the objective of validating the anti-hemorrhoidal properties of these plants and in order to identify phytochemical compounds responsible of this activity.

Keywords: *Ficus stuhlmannii*, *Khaya nyasica*, *Phyllanthus amarus*, Lubumbashi, Traditional healers, hemorrhoids.

Graphical Abstract



Background

Hemorrhoids are a network of arterial and venous vessels that are present in the normal state at the level of the anal canal. They contribute to the maintenance of continence [1]. Hemorrhoids may be located either under the skin of the anal orifice (external hemorrhoids) or higher up under the mucosa of the anal canal (internal hemorrhoids) [2]. Normally, hemorrhoids may exhibit a slight degree of swelling during the act of defecation. However, when they remain permanently dilated, they constitute pathological hemorrhoids [3,4].

Hemorrhoidal pathologies are defined by the presence of symptoms such as irritation, extreme pain, and itching around the anus, a painful or itchy lump or swelling near the anus, fecal leakage, painful stools, and the presence of blood on the tissue after defecation [5–7]. A number of studies have identified a number of risk factors for the development of hemorrhoids, including chronic constipation [8,9], diarrhea, insufficient dietary fiber, exertion and prolonged sitting, particularly during defecation [10], abdominal obesity, depression, pregnancy [11], hypertension, smoking [12], as well as advanced age and a sedentary lifestyle [13]. The severity of hemorrhoidal disease is typically evaluated according to the Goligher classification system, which is based on the degree of prolapse through the anus. The Goligher classification system defines four grades of hemorrhoidal disease. Grade 1 is defined as a hemorrhoid that bleeds but does not prolapse. Grade 2 is defined by the capacity of the hemorrhoid to prolapse upon exertion, which may or may not be accompanied by spontaneous reduction or return to the anal canal. Grade 3 is defined by the ability of the hemorrhoid to prolapse on exertion, which can be manually pushed back into the anal canal. Grade 4 is defined by the permanent prolapse of the hemorrhoid, which cannot be manually reduced [13–15].

Hemorrhoidal disease is one of the most frequently encountered anorectal conditions in general practice [16]. However, its true prevalence in the general population is not well known, in part because many patients do not seek treatment in hospital facilities [17]. A multicenter study conducted in Europe and Brazil reported a prevalence of 11% [16]. Partial prevalences have been reported in various countries around the world, but these studies are not truly representative of their countries. In the literature, we find that 16.6% of cases occur in Korea [12]; 48.1% in Eastern India [18]; 16% in Saudi Arabia [19]; and 38.9% in Australia [20]. In Africa, the reported incidence ranges from 7% to 13.1% in Ethiopia [9,21], 56.2% in Bamako [22], 2.3% in Brazzaville [23], 14.6% in Ouagadougou [24], and 54% in Nigeria [25]. To date, no studies have been conducted on the prevalence of hemorrhoids in the Democratic Republic of the Congo, particularly in Lubumbashi.

The treatment of hemorrhoids is aimed at correcting the underlying cause, which may be a variety of factors. In cases where the condition is relatively straightforward, it is advisable to adopt a diet high in fiber, drink plenty of water, and apply local anesthetic creams to relieve itching and pain. In symptomatic cases, treatments containing analgesics, non-steroidal anti-inflammatories, or anesthetics are employed to address hemorrhoidal attacks. Another potential treatment option is the application of a ligature to the hemorrhoidal artery. In the event of disease progression, surgical intervention is the definitive treatment for severe and thrombosed external hemorrhoids [26–28].

In Lubumbashi, as in most developing countries, the use of traditional medicine is a primary approach to addressing health concerns such as hemorrhoidal pathologies [29,30]. A recent study indicated that 79.4% of the Lubumbashi population sought traditional medicine for various pathologies and social issues. This form of medicine is largely based on the use of medicinal plants. The study additionally revealed that hemorrhoidal pathologies represent the primary reason for consulting traditional medicine in Lubumbashi [31]. However, neither national nor local studies have been conducted on the plants used to treat hemorrhoidal pathologies.

Several ethnomedicinal studies have been conducted in Africa on the use of medicinal plants to treat hemorrhoidal pathologies. For illustrative purposes, ethnobotanical studies have been conducted in Nigeria [32,33], Cameroon [34], Tanzania [35], and Kenya [36].

In Lubumbashi, the majority of ethnobotanical studies have focused on the targeted pathologies. Some of these studies, although oriented towards other pathologies, have yielded insights into the treatment of hemorrhoidal pathologies by medicinal plants that have been inventoried on this occasion. This is exemplified by *Vernonia shirensis* [synonym of *Baccharoides adoensis* var. *mossambiquensis* (Steetz) Isawumi, El-Ghazaly & B Nord, Asteraceae], whose macerated leaves are used locally against hemorrhoids [37]. Another example is the macerated leaves or stem bark of *Aframomum*

angustifolium (Sonn.) K.Schum., a member of the Zingiberaceae family [38]. Furthermore, the decoction of the roots of *Bridelia atroviridis* Müll Arg, Phyllanthaceae [39]; decoction from the leaves of *Asparagus africanus* Lam, Asparagaceae [40]; The decoction derived from the leaves of *Pericopsis angolensis* (Baker) Meeuwen, Fabaceae [41], and the decoction derived from the roots of *Piliostigma thonningii* Schumacher Milne-Redh, Fabaceae [42] are employed as an anti-hemorrhoidal agent in local non-conventional medicine. Nevertheless, no study has yet focused on the plants utilized in traditional medicine in Lubumbashi to treat hemorrhoidal pathologies. In light of the above, the objective of this study was to conduct an ethnobotanical survey with the aim of compiling a list of plants used in traditional medicine in Lubumbashi to treat hemorrhoidal pathologies. The list was to be compiled from both the general population and among traditional medicine professionals in the region. The objective was to identify the most commonly utilized plants in the region for the management of hemorrhoidal pathologies and to highlight the region's characteristic taxa.

1. Material and methods.

1.1. Experimental Framework

The study was conducted in the seven communes of the city of Lubumbashi: Annexe, Kampemba, Katuba, Kamalondo, Kenya, Lubumbashi, and Rwashi, in the province of Haut-Katanga in the Democratic Republic of Congo (Fig 1).




Fig.1. Lubumbashi's city in Haut-Katanga province, Democratic Republic of Congo

Lubumbashi is situated between 11°26' and 11°55' north latitude and 27°15' and 27°40' east longitude at an altitude of 1,230 meters. The climate is tropical, with an average annual temperature of 22.4 °C and an average annual rainfall of 512.7 mm³. The study area experiences two distinct seasons, with a relatively shorter rainy season spanning from November to April. The Miombo clear forest is the most prevalent vegetation type [42].

1.2. Ethnomedical data collection

Ethnomedical data were collected from a variety of sources, including households, herbalists, and traditional healers (THs), in Lubumbashi between March 2021 and August 2022. The sample size of household respondents was determined using two-stage sampling: a stratified sample and a cluster sample. For each commune considered a stratum, the corresponding commune office provided the population size (Table 1). The size of the representative sample at this level was determined by the following formula:

$$n = \frac{p \times (1-p) \times N \times t_p^2}{t_p^2 \times p \times (1-p) + (N-1) \times y^2} \quad (1);$$

Where, n= sample size, N= actual population size, p = expected proportion of respondents (estimated p at 0.5), tp = sampling confidence interval (tp = 1.96 given that we set a 95% confidence interval) and y = margin of sampling error: we set this at 0.05. This yields a household population of 2,690. (Table 1).

Table 1 Household sample size

Commune	N	n_r	n_e	D (%)
Annexe	749142	384.0	390	0.2
Kamalondo	27911	379.0	379	0.0
Kampemba	526 926	383.9	385	0.1
Katuba	469228	383.8	384	0.0
Kenya	480974	383.9	384	0.0
Lubumbashi	481563	383.9	384	0.0
Ruashi	410 034	383.8	384	0.0
Total	3 145 778	2 683	2 690	0.3

Legend – N (Target population), n_r(Calculated representative sample), n_e(Population surveyed in the field), D: Difference.

All individuals were interviewed in their own homes. Of the 2,690 people interviewed, 264 did not give consent to participate in the study, 407 said they had never suffered from hemorrhoids, and 499 said they had never used traditional medicine to treat hemorrhoids. However, the results of this study, based on data collected from households, concern only 1,520 households.

The sample was divided into clusters, with each cluster representing a specific avenue. This was achieved by counting every three households. In each household, we included any adult (≥ 18 years old) who had ever used plants to treat hemorrhoids, provided they were sufficiently unfamiliar with the interviewers.

Due to the lack of reliable data on the size of the herbalist population in Lubumbashi, the objective was to interview as many herbalists as possible, accessible through markets and other public spaces. This stage enabled us to meet 25 herbalists, all of whom consented to participate in the study.

The respondents in the Traditional Healer (TH) category were identified from a list of 50 Traditional Practitioners compiled during a recent survey conducted in the study area [42]. This stage yielded 35 traditional healers (THs), all of whom consented to participate. Subsequently, snowball sampling was employed to expand the sample size, resulting in an additional 30 Traditional Healers (THs). Of the 50 THs identified in the initial survey, 24 agreed to participate in the study, while four were already included on the initial list. The survey of the THs was thus conducted on 59 respondents.

2.3. Data processing and analysis

The names of the plant species inventoried were collected in the local languages. Subsequently, the plant species were converted into a scientific name at the Kipopo herbarium, where they were deposited, in order to assign scientific names to the herbaria collected with the help of resource persons. The names were updated by comparison with the following databases: Plants of the World Online (<https://powo.science.kew.org/>), The World Flora Online (<http://www.worldfloraonline>) or the African Plant Database (<https://africanplantdatabase.ch/>).

In light of the current controversy surrounding ethnobotanical indexes [43], and in order to provide quantitative support for this study, we have employed only headcounts and three ratios.: citation index (CI), medicinal use index (MUI) and medicinal capability index (MCI). The plant citation index (CI) was calculated using the formula:

$$CI = \frac{\text{Number of informants citing the plant}}{\text{Total number of informants who participated in the study}} \quad (2)$$

The medicinal use index (MUI) was calculated by the formula:

$$\text{MUI} = \frac{\text{Number of uses reported for a taxon}}{\text{Total number of uses reported for the entire study}} \quad (3)$$

The medicinal capability index (MCI) was calculated by the formula:

$$\text{MCI} = \frac{\text{NNumber of taxa cited to manage a given pathology}}{\text{Total number of taxa inventoried during the survey}} \quad (4)$$

In this study, the CI allows us to determine the number of times a plant has been cited in the management of hemorrhoids in the region, and consequently to identify the level of consensus within the population on the curative use of a given species in the management of hemorrhoids. The MUI highlights the level of medicinal use of a plant among the population studied. Finally, the MCI enables us to measure the potential of a given community to manage a well-defined pathology.

2.4. Ethics approval and consent to participate

The project proposal and Ethical were reviewed and provided by the Department of Pharmacology, Faculty of Pharmaceutical Sciences, University of Lubumbashi, Democratic Republic of the Congo (FSPUNILU-DP-BD-022022).

Prior to the commencement of data collection, In accordance with the ethical standards of the University of Lubumbashi, all 59 THs, 25 herbalists, and 1,520 households were asked to provide voluntary verbal consent. All respondents were informed that the data collected would be used exclusively for academic purposes. Informants were made aware that our goals were scientific rather than commercial.

3 Results

3.1 General knowledge of hemorrhoidal pathologies among the interviewees.

The informants interviewed for this study (herbalists: n=25, traditional practitioners: n=59, and households: n=2690) demonstrated a clear distinction between internal and external hemorrhoidal pathologies. However, only 1,520 individuals, representing 56.5% of the initial population, were responsible for the ethnobotanical knowledge of this category of the survey population.

In fact, of the 2690 individuals interviewed, 264 had not provided informed consent, 407 had never suffered from hemorrhoidal pathologies, and 499 had never utilized plants during a hemorrhoidal crisis. Consequently, the proportion of the population who had ever suffered from hemorrhoids was 2019 out of 2426, or 83.2%, and the proportion who had ever used plants during a hemorrhoidal crisis was 1520, or 62.6%.

The survey population diagnosed hemorrhoidal pathologies, locally known as "*Kilonda ntumbu*," based on nine symptoms: itching, pain in the anal region, fever, irritation, bleeding during or after bowel movements, internal burning sensations, and protuberances protruding from the anus. These symptoms were reported by over 50% of the subjects consulted, regardless of their category (Table 2).

Table 2 Healers' knowledge of hemorrhoidal disease

Sub-parameter	Herbalists (n=25)		Traditional healers (n=59)		Households (n=1520)	
	Nc	CI	Nc	CI	Nc	CI
Consequences of hemorrhoidal pathologies						
Anemia	16	0.64	39	0.66	1102	0.73
Pain and discomfort	25	1.00	59	1.00	1520	1.00
Sexual dysfunction	15	0.60	38	0.64	1101	0.72
Frequent unsuccessful bowel movements	20	0.80	52	0.88	1508	0.99
Anal fissures	19	0.76	51	0.86	1506	0.99
Discomfort	17	0.68	49	0.83	1501	0.99
Type of hemorrhoids found						
Internal	20	0.80	59	1.00	1401	0.92
External	21	0.84	54	0.92	1447	0.95
Number of people treated for hemorrhoids one month prior to the study (year)						
20-30	5	0.2	60	1.0	NA	NA
30-40	25	1.0	205	3.5	NA	NA
40-50	2	0.1	105	1.8	NA	NA
50-60	7	0.3	10	0.2	NA	NA
≥60	0	0.0	61	1.0	NA	NA
Symptoms						
Itching	20	0.8	52	0.9	1508	0.99
Pain in the anal region	12	0.5	31	0.5	997	0.66
Discharge of mucus from the anus	9	0.4	20	0.3	340	0.22

Sub-parameter	Herbalists (n=25)		Traditional healers (n=59)		Households (n=1520)	
Fever	16	0.6	39	0.7	1102	0.73
Irritation	19	0.8	51	0.9	1506	0.99
Bleeding during or after bowel movements	17	0.7	49	0.8	1501	0.99
Sensation of internal burning	15	0.6	38	0.6	1101	0.72
Sensation of inflammation in the rectum	10	0.4	21	0.4	965	0.63
Sensitive protuberances emerge from the anus	25	1.0	59	1.0	1520	1.00
Causes of hemorrhoidal disease						
Natural childbirth	5	0.2	17	0.3	164	0.11
Foods low in fresh vegetables and fruit	1	0.0	10	0.2	51	0.03
Frequent constipation	21	0.8	50	0.8	1410	0.93
Pregnancy	18	0.7	34	0.6	1189	0.78
Habitual spicy dishes	22	0.9	53	0.9	1418	0.93
Carrying heavy loads on a daily basis	23	0.9	57	1.0	1508	0.99
Frequent sitting for long periods	24	1.0	58	1.0	1510	0.99
Frequent sodomy	4	0.2	15	0.3	10	0.01
Frequent cycling	7	0.3	7	0.1	30	0.02
Genetic predisposition	0	0.0	3	0.1	10	0.01
Sedentary lifestyle	7	0.3	8	0.1	35	0.02
Overweight	5	0.2	4	0.1	37	0.02

Legend NA: Not applicable; N: Number citations (of people surveyed); CI: citation index.

The 12 factors identified as responsible for hemorrhoidal pathologies include frequent constipation, pregnancy, habitual spicy food, carrying heavy loads on a daily basis, and long and frequent sitting positions. These factors were reported by over 60% of interviewees in all categories (Table 2).

These informants are aware that hemorrhoidal pathologies can have a number of consequences, including pain and discomfort, as well as the frequent desire to have a bowel movement with no result. These consequences were reported by over 80% of them. The experience of the traditional medical practitioners consulted (herbalists and traditional practitioners) indicates that the majority of individuals who consult them for hemorrhoids are between 30 and 50 years of age (Table 2).

3.2. General Characteristics of Inventoried Plants

Ethnobotanical surveys have revealed that 100 plant species belonging to 90 genera and 45 botanical families are used to treat hemorrhoidal pathologies in Lubumbashi (Table 3). The majority of these medicinal taxa are trees (38%), followed by shrubs (32%) (Fig. 2). These taxa are native to seven geographical regions, with the TA (Tropical Africa) and TA-SA (South Africa) regions accounting for over 60% of the taxa inventoried (Fig. 3). The taxa are named in 16 local ethnic languages, with Swahili, Luba-Kat, and Bemba accounting for over 50% of the names of the taxa collected (Fig. 4).

Table 3 General characteristics on plants inventoried during the survey

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T_{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Acmella caulirhiza</i> Delile ^c (Asteraceae)	KIP000 111011	Mchapia tumbili ^v (Swahili)	0.027	WP ^{y3}	Pilage ^l	Hemorrhoid	0.013	NR	NR
<i>Adansonia digitata</i> L. ^c (Malvaceae)	KIP000 111012	Mutshi munene ^o (Luba-kas)	0.003	Sb ^w	Decoction ^l	Hemorrhoid, Clogged trumpet	0.026	Niger [44]	<i>In vivo</i> -Rat: ED ₅₀ = 145.3 ± 7.6 mg/Kg [45]
<i>Aframomum daniellii</i> (Hook.f.) K. Schum. ^c (Zingiberaceae)	KIP000 111013	Matundu ^u (Luba-kas)	0.004	Sd ^{y3}	Decoction ^l	Hemorrhoid	0.013	Cameroon [46]	NR
<i>Aframomum melegueta</i> K Schum (Zingiberaceae)	KIP000 001341	Masusu ^u (Bemba)	0.007	R ^{y3}	Decoction ^E	Hemorrhoid, wound, stomachache, diarrhea	0.052	Cameroon [34]	<i>In vivo</i> -Rat: ED ₅₀ =1000 mg/Kg [47]
<i>Ageratum conyzoides</i> L. ^c (Asteraceae)	KIP000 111014	Riyani ^m (Swahili)	0.011	WP ^{y1}	Pilage ^l	Hemorrhoid, Diarrhea, Hepatitis, Epilepsy and gonorrhea	0.079	Ivory Coast [48]	<i>In vivo</i> -Rat: ED ₅₀ =200 mg/Kg- Lv [49]
<i>Aidia micrantha</i> (K.Schum.) Bullock ex F.White ^c (Rubiaceae)	KIP000 111015	Mutshi wa buanga ^u (Luba-kas)	0.013	Sb ^w	Decoction ^l	Hemorrhoid, Sexual weaknesses	0.039	NR	NR
<i>Albizia adianthifolia</i> (Schum) W Wight ^a (Fabaceae)	KIP000 001031	Kapetanzovu ^u (Luba-kat)	0.064	Rb ^w	Decoction ^E	Asthenias, Tooth decay, Sexual dysfunction, Hemorrhoids, High blood pressure, Stomachache, Sinusitis, Coughs	0.105	Nigeria and South Africa [50]	<i>In vitro</i> -Cox1: IC ₆₀ - 80 : 100 µg/mL – Rb [50]

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Albizia lebbbeck</i> (L.) Benth. ^a (Fabaceae)	KIP000 111016	Kitutumie °(Luba-Kat)	0.095	R ^w	Maceration ^E	Hemorrhoid	0.011	DRC: [51]	<i>In vivo</i> -Rat: ED ₅₀ =400 mg/Kg -Sb [52]
<i>Alchornea cordifolia</i> (Schumach. & Thonn.) Müll. Arg. ^c (Euphorbiaceae)	KIP000 111017	Tshibeji ^u (Luba-kas)	0.032	Lv ^x	Pilage ^l	Hemorrhoid, Diarrhea, Wounds, Diabetes, Anemia, Dysentery	0.078	DRC [53]	<i>In vivo</i> -Rat: DE ₅₀ =1000 mg/Kg-Sb [54]
<i>Aloe vera</i> (L.) Burn.f. ^a (Asphodelaceae)	KIP452 120002	Aloé ^q (Swahili)	0.634	Lv ^x	Maceration ^l	Hemorrhoid, Acne, tooth decay, constipation, burns, gastric ulcer	0.078	Cameroon [55]	<i>In vivo</i> -Rat: DE ₅₀ =100 mg/Kg-Lv [56]
<i>Alstonia boonei</i> De Wild. ^a (Apocynaceae)	KIP000 111018	Mutshi wa bukula ^u (Luba-kas)	0.009	Sb ^w	Maceration ^l	Hemorrhoid, diarrhea and typhoid	0.039	Cameroon [46]	<i>In vivo</i> -Rat: ED ₇₀₋₉₀ =50-200 mg/Kg-Sb [57]
<i>Amaranthus hypochondriacus</i> L. ^a (Amaranthaceae)	KIP000 111019	Tshitekuteku ^r (Luba-kas)	0.006	Ry ¹	Pilage ^l	Hemorrhoid	0.013	NR	NR
<i>Amaranthus spinosus</i> L. ^a (Amaranthaceae)	KIP000 111020	Tshiteku ^v (Luba-kas)	0.686	Lv ^{y1}	Exudation ^l	Hemorrhoid	0.013	DRC [53]	<i>In vivo</i> -Rat: ED ₇₀₋₉₀ =50-250 mg/Kg-Lv [58]
<i>Anacardium occidentale</i> L. ^c (Anacardiaceae)	KIP000 111021	Kabanga banga ^v (Luba-kas)	0.004	Rb ^x	Decoction ^l	Hemorrhoid, leprosy, diarrhea	0.039	Burkina Faso: [59]	<i>In vivo</i> -Rat: ED ₅₀ =100 mg/Kg-Sb [60]
<i>Anchomanes difformis</i> (Blume) Engl. ^c (Araceae)	KIP000 111022	Sikili ^u (Swahili)	0.003	Rz ^w	Pilage ^l	Hemorrhoid	0.013	Cameroon [46]	<i>In vivo</i> -Rat: ED ₅₀ =800 mg/Kg: Lv [61]

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti- hemorrhoidal uses	Previous anti- inflammatory activity
<i>Annona senegalensis</i> Pers. ^c (Annonaceae)	KIP452 120003	Elo ^o (Luba-kat)	0.003	Rb ^x	Infusion ^l	Hemorrhoid, abdominal pain, STI, Gastritis	0.052	DRC [62] Niger [44]	<i>In vivo</i> -Rat: ED ₅₀ =100 mg/Kg: Lv [63]
<i>Anonidium mannii</i> (Oliv). Engl. & Diels ^c (Annonaceae)	KIP000 111023	Nkundo ^u (Lingala)	0.004	Rb ^w	Decoction ^l	Hemorrhoid	0.013	NR	NR
<i>Antidesma venosum</i> auct. ^b (Phyllanthaceae)	KIP293 612638	Kifubia ^v (Luba-kat)	0.011	Sb ^x	Decoction ^l	Hemorrhoid, STI, bilharzia	0.039	NR	<i>In vitro</i> -Cox: ED ₅₀ - 100=250 µg/mL - Lv [64]
<i>Antirrhinum majus</i> L. ^a (Plantaginaceae)	KIP000 111024	Lueni ^p (Swahili)	0.011	R ^x	Maceration ^l	Hemorrhoid	0.013	NR	<i>In vitro</i> -Cox2: ED ₁₀₀ =100 µg/mL: Lv [65]
<i>Antrocaryon klaineianum</i> Pierre ^b (Anacardiaceae)	KIP000 111025	Mbota ya nyama ^u (Luba-kas)	0.071	Rb ^w	Decoction ^l	Hemorrhoid, STI, stomach pain, painful menstruation	0.052	Cameroon [46]	<i>In vivo</i> -Rat: ED ₉₀ =600 mg/Kg - Sb [66]
<i>Asparagus africanus</i> Lam. ^b (Asparagaceae)	KIP000 111026	Mukoma ^v (Luba-kat)	0.014	Lv ^x	Decoction ^l	hemorrhoid, STI, cough, diarrhea, rheumatism, sore throat	0.079	DRC [67]	<i>In vivo</i> -Rat: ED ₂₅ =250 mg/Kg - Rb [64]
<i>Aspilia kotschyi</i> (Sch.Bip. ex Hochst.) Oliv. ^c (Asteraceae)	KIP000 111027	Tshipukapuka ^u (Luba-kas)	0.012	WP ^{y1}	Decoction ^l	Hemorrhoid, Haemorrhagia, wound	0.039	DRC [68]	NR
<i>Baccharoides adoensis</i> (Sch.Bip. ex Walp.) H.Rob ^c (Asteraceae)	KIP000 111028	Alete ^v (Hunde)	0.007	Rb ^w	Decoction ^l	Hemorrhoid, fever, diabetes, flu.	0.052	NR	NR

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Baillonella toxisperma</i> Pierre ^c (Sapotaceae)	KIP000 111029	Muabi ^u (Luba-kas)	0.009	Rb ^w	Decoction ^l	Hemorrhoid, backache, kidney failure, inflammation	0.052	Cameroon [46]	NR
<i>Balanites aegyptiaca</i> (L) Delile (Zygophyllaceae)	KIP000 111030	Mubambangoma ^r (Bemba)	0.002	Rb ^w	Decoction ^E	Hemorrhoid, fever, diabetes, jaundice	0.052	India [69]	<i>In vivo</i> -Rat: ED ₈₀ =600 mg/Kg-Rb [70]
<i>Bridelia ferruginea</i> Benth (Phyllanthaceae)	KIP000 001040	Inkuka-i-nsii ^u (Yanzi)	0.009	Rb ^w	Decoction ^{&} & Maceration ^E	Hemorrhoid, wound, TBC, Gonorrhea, Sickle cell disease, hypertension, dental caries	0.078	Nigeria [71]	<i>In vitro</i> -Cox2: ED ₈₀ =200 µg/mL-Rb [72]
<i>Bridelia micrantha</i> (Hochst.) Baill. ^c (Euphorbiaceae)	KIP000 001041	Muteja ^v (Swahili)	0.011	Rb ^w	Decoction ^l	Hemorrhoid	0.013	DRC [39]	<i>In vivo</i> -Rat: ED ₅₀ =200 mg/Kg-Lv [73]
<i>Canarium schweinfurthii</i> Engl. ^d (Burseraceae)	KIP000 111031	Elemi ^u (Luba-kat)	0.039	Rb ^w	Decoction ^l	Hemorrhoid, renal failure	0.026	DRC [40]	<i>In vitro</i> -Lox: IC ₅₀ =62,6 µg/L - Resin [74]
<i>Capsicum frutescens</i> L. ^c (Solanaceae)	KIP000 001044	Ushinda ^r (Luba-kat)	0.024	Lv ^x	Expression ^l	Hemorrhoid, rheumatism, muscle and joint pain	0.052	DRC [75]	NR
<i>Carica papaya</i> L. ^a (Caricaceae)	KIP041 463406	Kipapayi ^u (Luba-kat)	0.015	Sd ^x	Maceration ^l	Hemorrhoid	0.013	DRC [68]	<i>In vitro</i> -Lox: IC ₅₀ =48.0 µg/mL-Seed [76]
<i>Chromolaena odorata</i> (L.) R.M. King & H. Rob. ^{c,a} (Asteraceae)	KIP000 001049	Mokili-mobimba ^v (Lingala)	0.013	Lv ^x	Decoction ^l	Hemorrhoid	0.013	NR	<i>In vivo</i> -Rat: ED ₅₀ =400 mg/Kg- Lv [77]

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti- hemorrhoidal uses	Previous anti- inflammatory activity
<i>Citrus limon</i> (L.) Burm.f. ^f (Rutaceae)	KIP000 111032	Limau ^o (Swahili)	0.627	Fr ^w	Fermentation ^l	Hemorrhoid, Tired, stomachache	0.039	DRC [78,79]	<i>In vivo</i> -Rat: ED ₄₀ =400 mg/Kg - Fr [80]
<i>Clerodendrum bungei</i> Steud. ^c (Lamiaceae)	KIP000 111033	Sinawolo ^u (Kimbala)	0.017	Lv ^w	Decoction ^l	Hemorrhoid	0.013	DRC [79]	<i>In vitro</i> - lipopolysaccharide (LPS): Clerodenone B IC ₅₀ =6.3. µM - Lv [81]
<i>Cola chlamydantha</i> K. Schum. ^c (Malvaceae)	KIP000 111034	Karoj ^u (Rund)	0.007	Rb ^w	Decoction ^l	Hemorrhoid, anemia	0.026	Cameroon [46]	NR
<i>Coleus kilimandschari</i> Gürke ^b (Lamiaceae)	KIP000 111035	Mulavumba ^y (Swahili)	0.009	Lv ^x	Infusion ^l	Hemorrhoid, asthma, thrombosis, weight loss	0.052	DRC [67]	NR
<i>Costus afer</i> Ker Gawl. ^c (Costaceae)	KIP000 111036	Tangawisi sapiral ^u (Swahili)	0.029	Sy ³	Decoction ^l	Hemorrhoid, constipation, diarrhea, rheumatism, malaria	0.056	Cameroon [46]	<i>In vivo</i> -Rat: ED ₄₀ =400 mg/Kg - Lv [82]
<i>Crossopteryx febrifuga</i> (Afzel. ex G.Don) Benth ^c (Rubiaceae)	KIP000 001056	Okane ^u (Kimbala)	0.007	Rb ^w	Maceration ^l	Hemorrhoid, diabetes, chickenpox, measles, GID, malaria, constipation, rheumatism	0.089	[62]	NR
<i>Croton oligandrus</i> Pierre ex Hutch. ^c (Euphorbiaceae)	KIP000 111037	Tuki ^u (Swahili)	0.011	Rb ^w	Decoction ^l	Hemorrhoid, Cystitis	0.026	Cameroon [46]	NR
<i>Cyathula prostrata</i> (L.) Blume ^c (Amaranthaceae)	KIP000 111038	Tshilawulawu ⁿ (Luba-kas)	0.016	Lv ^{y1}	Decoction ^l	Hemorrhoid, asthma, diarrhea, colic, abdominal pain	0.056	Cameroon [46]	<i>In vivo</i> -Rat: DE ₆₀ =50 mg/Kg: Lv [83]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Cylicodiscus gabunensis</i> Harms ^c (Mimosaceae)	KIP000 111039	Okane ^u (Kimbala)	0.030	Rb ^w	Decoction ^l	Hemorrhoid, itching, swollen foot	0,039	NR	<i>In vivo</i> -Rat: DE ₅₀ =200 mg/Kg - Lv [84]
<i>Cymbopogon citratus</i> ^c (DC.) Stapf (Poaceae)	KIP000 111040	Tshipuka ^l (Luba-kas)	0.029	Lv ^{y3}	Decoction ^l	hemorrhoid, swollen foot	0.026	Cameroon [46]	<i>In vivo</i> -Rat: DE ₈₀ =68.24 mg/Kg: Lv [85]
<i>Cyperus articulatus</i> L. ^a (Cyperaceae)	KIP000 111041	Sokasoka ^m (Luba-Kat)	0.014	R ^x	Maceration ^l	hemorrhoid, tired, dizziness	0.039	DRC [79]	<i>In vitro</i> -Cox2: DE ₁₅₋₃₅ : 10 µM - Rb [86]
<i>Cyphostemma adenocaula</i> (Steud. Ex A. Rich.) Desc. ex Wild & R.B. Drumm. ^c (Vitaceae)	KIP000 111042	Muteja ya pori ^u (Swahili)	0.011	R ^{y3}	Infusion ^l	hemorrhoid	0.013	DRC [79]	NR
<i>Dalbergia lactea</i> Vatke ^c (Fabaceae)	KIP000 111043	Obwat ^u (Kimbala)	0.012	Lv ^x	Decoction ^l	hemorrhoid, pneumonia, diarrhea, dental caries, diabetes	0.056	DRC [79]	NR
<i>Diplorhynchus condylocarpon</i> (Mull. Arg.) ^c (Apocynaceae)	KIP361 753009	Mwembe ^u (Bemba)	0.013	Lv ^x	Decoction ^E	hemorrhoid, wound, GID, fever, snakebite	0.056	NR	NR
<i>Droogmansia munamensis</i> De Wild ^c (Fabaceae)	KIP000 001350	Mununganunga ^h (Bemba)	0.009	Lv ^x	Decoction ^E	hemorrhoid, diabetes, dysentery, wounds	0.056	NR	NR
<i>Entada abyssinica</i> Steud. Ex A. Rich. ⁹ (Fabaceae)	KIP000 001353	Tshitefu ⁿ (Luba-Kas)	0.618	R ^x	Decoction ^l	hemorrhoid, malaria, diabetes, low back pain, headaches	0.056	DRC: [38]	<i>In vivo</i> -Rat: DE ₅₀ =200 mg/Kg - Lv [87]

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Erigeron bonariensis</i> L. ^a (Asteraceae)	KIP000 111044	Riyani moya ^f (Swahili)	0.623	Lv ^{y1}	Exudation ^l	hemorrhoids, rheumatism, stretch marks	0.039	NR	NR
<i>Erigeron sumatrensis</i> Retz. ^f (Asteraceae)	KIP000 111045	Kitukituki ^m (Luba- kat)	0.644	Ry ¹	Maceration ^l	hemorrhoids, rheumatism, stretch marks	0.039	DRC [79]	NR
<i>Euphorbia hirta</i> L. ^e (Euphorbiaceae)	KIP000 001354	Kisanga ^m (Kikongo)	0.642	Sb ^{y1}	Decoction ^l	hemorrhoid, diarrhea, gonorrhea, dysentery	0.056	Cameroon [46]	<i>In vivo</i> -Rat: DE ₇₅ =100 mg/Kg - Lv [88]
<i>Ficus laurifolia</i> Lam (Moraceae) ^f (Moraceae)	KIP501 400542	Kikuyu ^v (Luba-kat)	0.635	Rb ^w	Pilage ^{l,E}	hemorrhoid, wound, peptic ulcer, diabetes, STI, tuberculosis	0.078	NR	NR
<i>Ficus stuhlmannii</i> Warb ^f (Moraceae)	KIP000 111046	Chilemba ^s (Bemba)	0.635	Rb ^w	Pilage ^{l,E}	hemorrhoids, Anemia, Abdominal pain, Gonorrhea (STI), Diarrhea, Syphilis, Coughs	0.078	NR	NR
<i>Ficus sur</i> Forssk ^f (Moraceae)	KIP539 378804	Mupulamako ^u (Bemba)	0.013	Rb ^w	Pilage ^l	hemorrhoids, anemia, female sterility, cough, syphilis, jaundice, amoebiasis, dysentery and helminthiasis	0.100	NR	<i>In vitro</i> -Cox2: DE ₅₀ : 80 µg/mL - Lv [89]
<i>Garcinia punctata</i> Oliv. ^c (Clusiaceae)	KIP000 111047	Nkodo ^u (Lingala)	0.006	Lv ^w	Decoction ^l	hemorrhoid, sexual weakness, infections	0.039	NR	NR
<i>Gardenia ternifolia</i> subsp. jovis-tonantis	KIP000 001366	Mankeba ^u (Bemba)	0.007	Rb ^x	Pilage ^E	hemorrhoid, malaria, diarrhea,	0.067	DRC [90]	<i>In vitro</i> -Cox1: IC ₉₀ : 100 µg/mL - Lv [91]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
(Welw) VeDRC ^c (Rubiaceae)						hypertension, dental caries, schistosomiasis			
<i>Gymnanthemum amygdalinum</i> (Delile) Sch.Bip ^c (Asteraceae)	KIP312 286606	Mubilishi ^v (Kirega)	0.006	R ^x	Decoction ^l	hemorrhoid, gonorrhea, malaria, diabetes	0.056	DRC [92]	<i>In vivo</i> -Rat: ED ₉₅ =200 mg/Kg - Lv [93]
<i>Hibiscus cannabinus</i> L. ^c (Malvaceae)	KIP000 111048	Ngayi ^v (Kirega)	0.011	Lv ^{y1}	Maceration ^l	hemorrhoid, obesity, indigestion	0.039	DRC [79]	<i>In vivo</i> -Rat: DE ₆₅ =400 mg/Kg - [94]
<i>Hymenocardia acida</i> Tul. ^b (Phyllanthaceae)	KIP464 414491	Ambalanga ^u (Hemba)	0.012	R ^x	Decoction ^l	hemorrhoid	0.011	DRC [40]	<i>In vivo</i> -Rat: DE ₇₀ =200 mg/Kg - Lv [95]
<i>Indigofera capitata</i> Kotschy ^b (Fabaceae)	KIP000 111049	Onkai ^u (Teke)	0.009	Fy ¹	Maceration ^l	hemorrhoid, sexual dysfunction, bronchitis, cough, dyspnea	0.066	NR	NR
<i>Khaya nyasica</i> Stapf ex Baker f. ^b (Meliaceae)	KIP000 001372	Bamba ^j (Luba-kat)	0.012	F ^w	Pilage ^{l,E}	hemorrhoid, anemia, diarrhea and dysentery, amebiasis, ascariasis, cholera, ovarian cyst, malaria, wound, dermatosis	0.145	Nigeria [71]	NR
<i>Landolphia owariensis</i> P.Beauv ^b (Apocynaceae)	KIP000 111050	Mazangu ^u (Kikongo)	0.066	Lv ^z	Pilage ^E	hemorrhoid, spasms	0.026	DRC [79]	<i>In vivo</i> -Rat: DE ₅₅ =400 mg/Kg - Lv [96]

Taxon^{source} (Family)	NH	Local name (ethnicity)^{GT}	CI (n=1604)	UP^{MT}	Preparation^T HD	Medicinal uses	MUI (n=76)	Previous anti- hemorrhoidal uses	Previous anti- inflammatory activity
<i>Luffa cylindrica</i> (L.) M.Roem. ^c (Cucurbitaceae)	KIP000 111051	Tseke ⁿ (Kiyombe)	0.011	Sb ^z	Maceration ^l	hemorrhoid, bronchitis, sinusitis, rhinitis	0.056	Nigeria [97]	<i>In vivo</i> -Rat: DE ₃₀ =50 mg/Kg - Lv [98]
<i>Mangifera indica</i> L. ^e (Anacardiaceae)	KIP000 111052	Mwembe ^y (Shi)	0.009	Sb ^w	Decoction ^l	hemorrhoid, diarrhea, diabetes, colic, dysentery	0.066	DRC [62] Cameroon [34]	<i>In vitro</i> -Human Red Blood Cell (HRBC) Membrane Stabilization Method: IC ₅₀ : 100 µg/mL - Lv [99]
<i>Markhamia tomentosa</i> (Benth.) K.Schum. ex Engl. ^c (Bignoniaceae)	KIP000 111053	Esate ou onz ^u (Kimbala)	0.007	Lv ^x	Decoction ^l	hemorrhoid, skin infection, wound, rheumatism	0.066	DRC [79]	<i>In vivo</i> -Rat: DE ₆₆ =200 mg/Kg -Lv [100]
<i>Memecylon flavovirens</i> Baker ^c (Melastomaceae)	KIP000 001376	Musunda ^u (Bemba)	0.006	Lv ^{y1}	Pilage ^E	hemorrhoid, cough, helminthiasis, nasal congestion	0.056	NR	NR
<i>Mesosphaerum suaveolens</i> (L.) Kuntze ^f (Lamiaceae)	KIP000 111054	Mifisha ^u (Bemba)	0,704	Lv ^{y1}	Pilage ^E	hemorrhoid, malaria, fever, hernias, wounds	0.066	NR	<i>In vitro</i> -TNF-α: IC ₉₀ : 100 µg/mL - Lv [101]
<i>Monodora myristica</i> (Gaertn.) Dunal ^c (Annonaceae)	KIP000 111055	Mpeya ^u (Kikongo)	0.080	Lv ^w	Decoction ^l	hemorrhoid, stomachache, headache, diarrhea, hypertension	0.066	DRC [67]	<i>In vivo</i> -Rat: DE ₆₀ =200 mg/Kg: Seed [102]
<i>Moringa oleifera</i> Lam. ^c (Moringaceae)	KIP000 111056	Konse-konse ^l (Bemba)	0.064	Sb ^w	Maceration ^l	hemorrhoid, diarrhea, abdominal pain, sexual impotence	0.066	Niger [44]	<i>In vivo</i> -Rat: DE ₆₀ =250 mg/Kg-Lv [103]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Musa x paradisiaca</i> L. ^f (Musaceae)	KIP000 111057	Mugomba ^u (Swahili)	0.013	Lv ^w	Maceration ^E	hemorrhoids, wounds, dysentery, bronchitis, diarrhea	0.066	Nigeria [71]	<i>In vitro</i> -Human Red Blood Cell Membrane Stabilization Method: IC ₄₃ = 1000 µg/mL – Sb [104]
<i>Myrianthus arboreus</i> P.Beauv. ^a (Urticaceae)	KIP000 111058	Mbubu ^u (Luba-Kat)	0.032	R ^w	Decoction ^l	hemorrhoid, diarrhea, constipation, abdominal pain	0.056	Cameroon [46]	<i>In vivo</i> -Rat: DE ₅₉ =250 mg/Kg - Lv [105]
<i>Newbouldia laevis</i> (P.Beauv.) Seem. ex Bureau ^c (Bignoniaceae)	KIP000 111059	Mpese ^u (Kikongo)	0.026	Lv ^w	Pilage ^l	hemorrhoid, epilepsy, convulsion	0.039	Ghana Cameroon Nigeria [106]	<i>In vivo</i> -Rat: DE ₉₀ =400 mg/Kg: Rb [107]
<i>Ocimum centraliafricanum</i> R.E.Fr. ^a (Lamiaceae)	KIP000 111060	Gish ^u (Rund)	0.006	Lv ^{y3}	Decoction ^l	hemorrhoid	00.13	NR	NR
<i>Ocimum gratissimum</i> L. ^a (Lamiaceae)	KIP000 001383	Ditsususu ^o (Kiyombe)	0.004	Lv ^{y3}	Maceration ^l	hemorrhoid, cough, cold, dysentery	0.056	DRC [38]	<i>In vitro</i> -Cox-Lox: IC ₅₀ : 125-144 µg/mL Lv [108]
<i>Parinari curatellifolia</i> subsp ^c (Chrysobalanaceae)	KIP000 001385	Mpundu ⁿ (Bemba)	0.009	Rb ^w	Pilage ^E	Hemorrhoid, malaria, diarrhea, hypertension	0.056	DRC [41]	<i>In vivo</i> -Rat: DE ₂₅ =400 mg/Kg - Lv [109]
<i>Passiflora foetida</i> L. ^c (Passifloraceae)	KIP000 111061	Maua ^v (Swahili)	0.012	Sb ^{y2}	Decoction ^l	hemorrhoid, asthma, nervousness, digestive disorder, muscle spasm	0.066	NR	<i>In vitro</i> -NO: IC ₅₀ : 30-50 µg/mL - Lv, Fr [110]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Pentaclethra macrophylla</i> Benth. ^c (Fabaceae)	KIP000 111062	Ngunga ^u (Swahili)	0.017	R ^w	Infusion ^l	hemorrhoid, inflammation, parasites	0,039	DRC [67]	<i>In vivo</i> -Rat: DE ₂₀ =150 mg/Kg: Sb [111]
<i>Pentadiplandra brazzeana</i> Baill. ^c (Pentadiplandraceae)	KIP000 111063	Mutitatita ^u (Kirega)	0.031	Sb ^x	Maceration ^l	hemorrhoid, pneumonia, edema	0,039	[112]	<i>In vitro</i> -Albumin denaturation method: IC ₅₂ : 50 µg/mL-Rb [113]
<i>Pericopsis angolensis</i> (Baker) ^c (Fabaceae)	KIP000 111064	Mubanga ^h (Bemba)	0.032	Rb ^w	Decoction ^l	hemorrhoids, fever, diarrhea	0.039	NR	<i>In vitro</i> -NO assay: No activity -Sb [114]
<i>Phyllanthus amarus</i> Schumach. & Thonn. ^f (Phyllanthaceae)	KIP000 111065	Kivumbwa ⁿ (Hemba)	0.756	R ^{y1}	Decoction ^l	hemorrhoid, diabetes, flu, inflammation, liver cirrhosis	0.066	DRC [115]	<i>In vitro</i> -pro-inflammatory cytokines assay: IC ₅₀ = 60 µg/mL -WP [116]
<i>Phyllanthus muellerianus</i> (Kuntze) Exell ^c (Phyllanthaceae)	KIP277 202454	Mupetwa lupe ^u (Bemba)	0.002	Lv ^x	Pilage ^l	hemorrhoid, constipation, wound, hepatitis	0.056	DRC [67]	<i>In vivo</i> -Rat: DE ₄₇ =300 mg/Kg -Lv [117]
<i>Phytolacca dodecandra</i> L'Hér. ^c (Phytolaccaceae)	KIP000 111066	Loko ^o (Kirega)	0.002	R ^x	Pilage ^l	hemorrhoids	0.013	DRC [51]	<i>In vivo</i> -Rat: DE ₇₄ =600 mg/Kg -Lv [118]
<i>Plumbago zeylanica</i> L. ^c (Plumbaginaceae)	KIP000 111067	Beji ya risasi (Swahili)	0.003	Sb ^x	Decoction ^l	hemorrhoid, vomiting, rheumatism	0.039	DRC [79]	<i>In vivo</i> -Rat: DE ₄₅ =100 mg/Kg: Rb [119]
<i>Psidium guajava</i> L. ^a (Myrtaceae)	KIP000 301861	Mapele ^r (Swahili)	0.019	Lv ^x	Maceration ^l	hemorrhoids, hemorrhage, diarrhea	0.039	DRC [62]	<i>In vivo</i> -Rat: DE ₅₅ =250 mg/Kg -Lv [120]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Psorospermum febrifugum</i> Spach ^a (Hypericaceae)	KIP510 193316	Kafifi ^u (Bemba)	0.021	Rb ^x	Decoction ^E	hemorrhoid, diarrhea, amoebae, malaria	0.056	DRC [37]	<i>In vivo</i> -Rat- xylene-induced ear oedema assay: DE ₅₅ =400 mg/Kg: Sb [120]
<i>Pterocarpus angolensis</i> DC. ^e (Fabaceae)	KIP000 001393	Mulambwa ^v (Luba-kat)	0.646	Sb ^w	Pilage ^l	hemorrhoid, epilepsy, abdominal pain, ascariasis	0.056	DRC [40]	<i>In vitro</i> -Cox assay: IC ₉₀ = 250 µg/mL -Sb [121]
<i>Radlkofera calodendron</i> Gilg ^c (Sapindaceae)	KIP000 111068	Luet ^u (Kikongo)	0.007	Lv ^w	Decoction ^l	hemorrhoid	0.013	DRC [79]	NR
<i>Senna alata</i> (L.) Roxb. ^c (Fabaceae)	KIP000 111069	Tunda ⁿ (Swahili)	0.009	Lv ^x	Decoction ^l	hemorrhoids, fever, diarrhea, dysentery, sexual weakness	0.056	Thailand [122]	<i>In vitro</i> -HRBC membrane stabilizing method: IC ₅₀ = 9.93 µg/mL -Lv [123]
<i>Sida rhombifolia</i> L. ^c (Malvaceae)	KIP000 111070	Lumvumvu ^v (Kikongo)	0.013	Lv ^x	Decoction ^l	hemorrhoid, depression, bronchitis, inflammation, fever	0.066	DRC [79]	<i>In vitro</i> -NO assay: IC ₅₀ = 9.93 µg/mL -Lv [124]
<i>Solanum lycopersicum</i> L. ^a (Solanaceae)	KIP000 111071	Tomate ^m (Luba-kat)	0.016	Lv ^{y3}	Exudation ^l	Hemorrhoids	0.011	DRC [51]	<i>In vivo</i> -Rat: DE ₅₀ =50 mg/Kg: Fr [125]
<i>Strychnos spinosa</i> Lam (Malvaceae)	KIP000 111072	Lumvumvu ^v (Kikongo)	0.127	Lv ^x	Pilage ^l	hemorrhoids, depression, bronchitis, inflammation, fever	0.066	DRC [79]	<i>In vitro</i> -NO assay: IC ₅₀ = 88.43 µg/mL -Lv [126]
<i>Symphytum officinale</i> L. ^c (Boraginaceae)	KIP000 111073	Duyemba ^t (Tshisala)	0.009	Sb ^{y3}	Decoction ^l	hemorrhoid, inflammation, dry cough	0.039	NR	<i>In vivo</i> -Rat: DE ₅₈ =500 mg/Kg - R [127]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Syzygium cordatum</i> Subsp. ^a (Myrtaceae)	KIP000 111074	Ciyambaganyi ⁱ (Shi)	0.006	Lv ^x	Exudation ^l	hemorrhoid, malaria, diabetes, convulsion, pains, Abdominal, epilepsy, meningitis, Typhoid fever, Gonorrhea, tooth decay	0.145	DRC [51]	<i>In vitro</i> -Cox assay: IC ₇₅ = 250 µg/mL -Lv [128]
<i>Terminalia mollis</i> M.A.Lawson ^f (Combretaceae)	KIP452 120033	Kibobo ^u (Luba-kat)	0.728	Lv ^w	Pilage ^l	Hemorrhoid, constipation, diarrhea, abdominal pain	0.056	DRC [31]	<i>In vitro</i> -NO assay: IC ₅₀ = 40 µg/mL -Lv [129]
<i>Tetradenia riparia</i> (Hochst.) Codd ^c (Lamiaceae)	KIP370 001297	Mutuzo ^v (Shi)	0.011	WP ^w	Pilage ^l	Hemorrhoid, cough, diarrhea, constipation	0.056	DRC [78]	<i>In vitro</i> -NO assay: IC ₅₀ = 2.05 ± 1.18 µg/mL -Lv [130]
<i>Thonningia sanguinea</i> Vahl ^c (Balanophoraceae)	KIP000 111075	Bisusu ^u (Luba-kat)	0.015	Sb ^{y3}	Decoction ^l	Hemorrhoid, Osteomyelitis	0.026	DRC [79]	<i>In vivo</i> -Rat-Albumin induced paw oedema: Non active - Lv [131]
<i>Uapaca guineensis</i> Müll.Arg. ^c (Phyllanthaceae)	KIP000 111076	Ngwaka ^u (Kirega)	0.032	Lv ^w	Infusion ^l	hemorrhoids, muscular weakness, sexual weakness	0.039	NR	<i>In vivo</i> -Rat: DE ₉₀ =124 mg/Kg: R [132]
<i>Vernonia shirensis</i> Oliv. & Hiern ^b (Asteraceae)	KIP000 111076	Kilulunkunja ^v (Swahili)	0.009	Lv ^{y3}	Decoction ^l	hemorrhoid, headache, diarrhea, diabetes, fever	0.066	DRC [40]	NR
<i>Zanha africana</i> (Radlk.) Exell ^b (Sapindaceae)	KIP000 001397	Kalayi ^h (Bemba)	0.012	Rb ^w	Pilage ^l	hemorrhoid, malaria, diabetes, convulsion	0.056	NR	<i>In vivo</i> -Rat: IC ₅₀ =0.47 mM-Lv: Zanhasaponin A [133]

Taxon ^{source} (Family)	NH	Local name (ethnicity) ^{GT}	CI (n=1604)	UP ^{MT}	Preparation ^T _{HD}	Medicinal uses	MUI (n=76)	Previous anti-hemorrhoidal uses	Previous anti-inflammatory activity
<i>Zanthoxylum chalybeum</i> Engl. ^c (Rutaceae)	KIP000 001016	Pupwe kyulu ^u (Luba-kat)	0.019	R ^x	Decoction ^l	hemorrhoid	0.013	NR	<i>In vitro</i> -Cox assay: IC ₉₅ = 500 µg/mL -Rb [64]
<i>Zingiber officinale</i> Roscoe ^a (Zingiberaceae)	KIP000 111077	Tangawisi ^k (Swahili)	0.548	Rz ^{y3}	Decoction ^l	hemorrhoid, constipation, diarrhea, hepatitis	0,056	DRC [62]	In vivo -Rat: DE ₈₀ = 100 mg/Kg - Rz [134]

Legend: Abbreviations: NH: herbarium number; CI: citation index; GT: geographical type; UP: used part; MT: morphological type; FU: form of use; MUI: medicinal use index; THD: type of hemorrhoid diseases. **Data sources:** Traditional practitioners & herbalists (d), Traditional practitioners (c), Households (a), Herbalists (b), Traditional practitioners & herbalists & Households (e), Traditional practitioners & Households (f). **Morphological types:** Tree (w), Shrub (x), Annual herb (y1), Biennial herb (y2), Perennial herb (y3) and Liana (z). **Hemorrhoidal types:** I (internal hemorrhoid), E (external hemorrhoid). **Geographical types:** CA : (h) CA-SA (i) EA (j) MA (k); MA-NA-TA (l); MA-NA-TA-SA (m); MA-TA (n) ; MA-TA-SA (o); NA (p); NA-TA (q); NA-TA-SA (r) W (s); NN (t); TA (u); TA-SA (v) ; CA (Central Africa); SA (Southern Africa); MA(Madagascar); EA (East Africa); NA (North Africa); WE (West Africa). **Used part** : Whole plant, WP; Root bark, Rb; Stem Bark, Sb; Leaves, Lv; Seed, Sd; Fr, Fruit; Rz: Rhizome.




Fig.2. Morphological types (n=100).




Fig. 3 Geographical type.

Legend: TA(Tropical Africa); NEA(North-East Africa); CA (Central Africa); SA(Southern Africa); MA(Madagascar); EA(East Africa); NA(North Africa); WE(West Africa).




Fig. 4 Plant naming languages.

Seventeen families have more than two taxa each. The Fabaceae (10 species), Asteraceae (9 plant species), Lamiaceae (7 species), Phyllanthaceae (6 species), Anacardiaceae (5 species), Euphorbiaceae (4 species), Malvaceae (4 species) and Sapindaceae (4 species) families are the eight most important, collectively accounting for 49% of all plants cited (Fig.5).




Fig. 5 Family with taxa $n \geq 2$

The most frequently cited taxa are *Phyllanthus amarus* Schumach. & Thonn. (Phyllanthaceae), *Terminalia mollis* MA Lawson (Combretaceae), and *Mesosphaerum suaveolens* (L.) Kuntze (Lamiaceae). These taxa were claimed by over 1123 resource persons ($CI \geq 0.728$) in total, with each taxon being cited individually (Table 2).

3.3. Medicinal plants used to treat hemorrhoidal diseases

A total of 100 taxa are utilized in 117 anti-hemorrhoidal recipes within the study area. Of these, 101 employ a single plant, while 17 utilize mixtures of two or three plants. Three of the single-plant recipes are employed for mixed hemorrhoidal pathologies, 14 for external hemorrhoids, and the remaining 84 for internal hemorrhoids. In contrast, all the mixed-plant recipes are utilized for mixed hemorrhoids. The leaves of the plant are the most commonly used part in the preparation of herbal remedies, representing 69% of the cases, followed by root barks, which represent 36% of the cases (Fig 6).




Fig. 6 Parts of plants used to treat hemorrhoids.

The most prevalent method of preparing herbal remedies is decoction (51%) in mono-herbal recipes and pounding (44%) in polyherbal recipes (Fig. 7). Topical administration of herbal remedies is employed in five forms, with liniment (CI: 0.49-0.38) and sitz bath (CI: 0.30-0.25) representing over 50% of anti-hemorrhoidal recipes (Fig. 8).




Fig. 7 Mode of anti-hemorrhoidal recipe's preparation.




Fig. 8 Form of uses of recipe.

The respondents indicated that they utilize a variety of utensils, including teaspoons, tablespoons, cups, and bottles, to adjust the doses according to the patient's age (child or adult) and/or physiological state (e.g., pregnancy). In order to contextualize the results of this ethnobotanical survey, we sought to estimate the quantities of solutions (volumes) and solids (mass or powdered plant parts) as reported by respondents. This was done in reference to a

previous study [135], which had attempted to do the same. The informants indicated that the treatment of hemorrhoidal pathologies typically lasts between two and five days.

The information provided by the resource persons interviewed indicates that liniments are prepared by mixing three parts of palm oil with one part of solution (exudates, expression, decoction, macerated, or infused) or vegetable matter reduced to paste or powder, as the case may be. Ointments are prepared by mixing two parts of beeswax that have been melted in a water bath with one part of hot palm oil. Subsequently, plant extracts (30-40%) or dry matter are added to the aforementioned mixture.

With regard to the 16 recipes resulting from mixtures of two (12 recipes) or three taxa (3 recipes), these involve 27 taxa. It is notable that *Euphorbia hirta* (4 recipes) plays a significant role in several recipes, accounting for 25% of the total. The next most prevalent taxa are *Ageratum conyzoides*, *Albizia adianthifolia*, *Bridelia micrantha*, *Khaya nyasica*, *Monodora myristica*, and *Parinari curatellifolia*, which are each involved in two recipes. The remaining nine taxa are reported in only one polyherbal recipe. The most frequently cited polyherbal recipes are those based on the mixture of *P. angolensis* (leaves) and *P. thonningii* (root bark) (121 citations) and the mixture of *A. cordifolia* (root bark) and *A. hybridus* (roots) (107 citations) (Table 4).

Table 4 Characteristics of herbal healing recipes used in blends

N°	Taxon1	Taxon 2	Taxon 3	UP (Ratio)	Form	CI (n=2774)	Preparation
1	<i>A melegueta</i>	<i>E hirta</i>	NA	Lv-Lv (1÷2)	Ointment	0.009	Expression
2	<i>A conyzoides</i>	<i>A adianthifolia</i>	NA	R-R (1÷1)	Sitz bath	0.013	Decoction
3	<i>A cordifolia</i>	<i>A hybridus</i>	NA	R-Rb (1÷2)	Drink	0.039	Maceration
4	<i>A senegalensis</i>	<i>B aegyptiaca</i>	NA	Lv-R (2÷1)	Liniment	0.027	Spraying
5	<i>E hirta</i>	<i>A conyzoides</i>	NA	Lv-Lv (2÷1)	Ointment	0.028	Expression
6	<i>B micrantha</i>	<i>C kilimandschari</i>	NA	Rb-Rb (2÷1)	Sitz bath	0.032	Decoction
7	<i>C febrifuga</i>	<i>D condylocarpon</i>	NA	Lv-R (1÷1)	Drink	0.032	Decoction
8	<i>F sur</i>	<i>K nyasica</i>	NA	R-Sb (1÷1)	Liniment	0.011	Pulverization
9	<i>L owariensis</i>	<i>M indica</i>	NA	Lv-Lv (2÷1)	Ointment	0.014	Pulverization
10	<i>M tomentosa</i>	<i>M myristica</i>	NA	Lv-Rb (1÷1)	Liniment	0.016	Pulverization
11	<i>O fimbriatum</i>	<i>P curatellifoli</i>	NA	Rb-R (2÷1)	Liniment	0.026	Spray
12	<i>P angolensis</i>	<i>P thonningii</i>	NA	Lv-Rb (2÷1)	Liniment	0.044	Spray

N°	Taxon1	Taxon 2	Taxon 3	UP (Ratio)	Form	CI (n=2774)	Preparation
13	<i>P angolensis</i>	<i>S cordatum</i>	<i>M myristica</i>	Lv-Rb-Rb (1÷2÷1)	Liniment	0.013	Pulverization
14	<i>T riparia</i>	<i>G amygdalinum</i>	<i>E hirta</i>	Lv-Lv-Lv (1÷2÷1)	Ointment	0.032	Expression
15	<i>P curatellifolia</i>	<i>A adianthifolia</i>	<i>P guajava</i>	Lv-Lv-Lv (1÷2÷2)	Sitz bath	0.033	Maceration
16	<i>B micrantha</i>	<i>K nyasica</i>	<i>E hirta</i>	Lv-Lv-Lv (1÷2÷2)	Sitz bath	0.035	Decoction

Legend L: Leave, R: Root, Rb: Root bark, Sb: Stem bark. The unit of raw material (plant organ) intake is the handshake. NA : Not applicable, UP: used part.

3.4. Medicinal uses of plants surveyed

The taxa reported in this study are known and utilized by householders, herbalists, and traditional healers for the treatment of hemorrhoids (Fig 9). The taxa are employed to treat internal hemorrhoids (83 taxa), external hemorrhoids (14 taxa), or both types concomitantly (3 taxa). Nevertheless, these taxa are also involved in the management of 75 other pathologies, signs, or symptoms, where gastrointestinal disorders are particularly prevalent. The most frequently cited indications were gastrointestinal disorders (CI: 0.622) and wounds (CI: 0.68) (Table 4).




Fig. 9 Distribution of plants inventoried across the 3 categories of people surveyed

The five pathologies for which the resource persons in this study have the most taxa, in addition to hemorrhoids, are GID (31 taxa), diabetes (15 taxa), sexually transmitted infections (STI; 14 taxa), malaria (13 taxa), and wounds (11 taxa) (Table 5).

Table 5 Other pathologies treated by plants inventoried during surveys

Indications	Nt (n=76)	MCI (n=100)	Nc	CI (n=1604)	Taxon
Acne	1	0.01	48	0.030	<i>Aloe vera</i>
Amoebiasis	4	0.04	12	0.007	<i>Ficus sur</i> , <i>Khaya nyasica</i> , <i>Pentaclethra macrophylla</i> , <i>Psorospermum febrifugum</i> .
Anemia	4	0.04	1	0.001	<i>Alchornea cordifolia</i> , <i>Ficus stuhlmannii</i> , <i>Ficus sur</i> , <i>Khaya nyasica</i> .
Asthma	3	0.03	6	0.004	<i>Coleus kilimandschari</i> , <i>Cyathula prostrata</i> , <i>Passiflora foetida</i>
Bleeding	2	0.02	18	0.011	<i>Aspilia kotschy</i> , <i>Psidium guajava</i> .
Bronchitis	5	0.05	2	0.001	<i>Indigofera capitata</i> , <i>Luffa cylindrica</i> , <i>Musa paradisiaca</i> , <i>Sida rhombifolia</i> , <i>Strychnos spinosa</i> .
Burns	1	0.01	48	0.030	<i>Aloe vera</i>
Chicken pox	1	0.01		0.000	<i>Crossopteryx febrifuga</i>
Cholera	1	0.01	1	0.001	<i>Khaya nyasica</i>
Cirrhosis of the liver	1	0.01	1	0.001	<i>Phyllanthus amarus</i>
Clogged tubes	1	0.01	1	0.001	<i>Adansonia digitata</i>
Colic	9	0.09	439	0.274	<i>Annona senegalensis</i> , <i>Cyathula prostrata</i> , <i>Ficus stuhlmannii</i> , <i>Mangifera indica</i> , <i>Moringa oleifera</i> , <i>Myrianthus arboreus</i> , <i>Pterocarpus angolensis</i> , <i>Syzygium cordatum</i> , <i>Terminalia mollis</i> .
Constipation	9	0.09	247	0.154	<i>Aloe vera</i> , <i>Costus afer</i> , <i>Myrianthus arboreus</i> , <i>Phyllanthus muellerianus</i> , <i>Crossopteryx febrifuga</i> , <i>Psidium guajava</i> , <i>Terminalia mollis</i> , <i>Tetradenia riparia</i> , <i>Zingiber officinale</i> .
Convulsion	3	0.03	1	0.001	<i>Newbouldia laevis</i> , <i>Syzygium cordatum</i> , <i>Zanha africana</i> .
Cough	9	0.09	126	0.079	<i>Albizia adianthifolia</i> , <i>Asparagus africanus</i> , <i>Ficus stuhlmannii</i> , <i>Ficus sur</i> , <i>Indigofera capitata</i> , <i>Memecylon flavovirens</i> , <i>Ocimum gratissimum</i> , <i>Symphytum officinale</i> , <i>Tetradenia riparia</i> .
Cyst	2	0.02	7	0.004	<i>Croton oligandrus</i> , <i>Khaya nyasica</i> .
Depression	2	0.02	1	0.001	<i>Sida rhombifolia</i> , <i>Strychnos spinosa</i> .
Diabetes	15	0.15	721	0.450	<i>Alchornea cordifolia</i> , <i>Baccharoides adoensis</i> , <i>Balanites aegyptiaca</i> , <i>Crossopteryx febrifuga</i> , <i>Dalbergia lactea</i> , <i>Droogmansia</i>

Indications	Nt (n=76)	MCI (n=100)	Nc	CI (n=1604)	Taxon
					<i>munamensis</i> , <i>Entada abyssinica</i> , <i>Ficus brachypoda</i> , <i>Gymnanthemum amygdalinum</i> , <i>Mangifera indica</i> , <i>Phyllanthus amarus</i> , <i>Syzygium cordatum</i> , <i>Vernonia amygdalina</i> , <i>Vernonia shirensis</i> , <i>Zanha africana</i> .
Dysentery	9	0.09	134	0.084	<i>Alchornea cordifolia</i> , <i>Euphorbia hirta</i> , <i>Droogmansia munamensis</i> , <i>Ficus sur Forssk</i> , <i>Khaya nyasica</i> , <i>Mangifera indica</i> , <i>Musa paradisiaca</i> , <i>Ocimum gratissimum</i> , <i>Senna alata</i> .
Dyspnea	1	0.01	4	0.002	<i>Indigofera capitata</i>
Edema	1	0.01	1	0.001	<i>Pentadiplandra brazzeana</i>
Epilepsy	4	0.04	12	0.007	<i>Ageratum conyzoides</i> , <i>Newbouldia laevis</i> , <i>Syzygium cordatum</i> , <i>Pterocarpus angolensis</i> .
Tiredness	3	0.03	124	0.077	<i>Citrus limon</i> , <i>Albizia adianthifolia</i> , <i>Cyperus articulatus</i> .
Fever	9	0.09	494	0.308	<i>Baccharoides adoensis</i> , <i>Balanites aegyptiaca</i> , <i>Diplorhynchus condylocarpon</i> , <i>Mesosphaerum suaveolens</i> , <i>Pericopsis angolensis</i> , <i>Senna alata</i> , <i>Sida rhombifolia</i> , <i>Strychnos spinosa</i> , <i>Vernonia shirensis</i> .
Flu	2	0.02	13	0.008	<i>Baccharoides adoensis</i> , <i>Phyllanthus amarus</i> .
Gastric ulcer	2	0.02	48	0.030	<i>Aloe vera</i> , <i>Ficus brachypoda</i>
Gastritis	2	0.02	13	0.008	<i>Aframomum melegueta</i> , <i>Albizia adianthifolia</i> , <i>Annona senegalensis</i> .
Gastrointestinal disorders (GID)	31	0.31	997	0.622	<i>Aframomum melegueta</i> , <i>Ageratum conyzoides</i> , <i>Alchornea cordifolia</i> , <i>Alstonia boonei</i> , <i>Anarcadium occidentale</i> , <i>Asparagus africanus</i> , <i>Costus afer</i> , <i>Crossopteryx febrifuga</i> , <i>Cyathula prostrata</i> , <i>Dalbergia lactea</i> , <i>Diplorhynchus condylocarpon</i> , <i>Euphorbia hirta</i> , <i>Ficus brachypoda</i> , <i>Ficus stuhlmannii</i> , <i>Gardenia ternifolia</i> , <i>Khaya nyasica</i> , <i>Mangifera indica</i> , <i>Monodora myristica</i> , <i>Moringa oleifera</i> , <i>Musa paradisiaca</i> , <i>Myrianthus arboreus</i> , <i>Parinari curatellifolia</i> , <i>Pericopsis angolensis</i> , <i>Plumbago zeylanica</i> , <i>Psidium guajava</i> , <i>Psorospermum febrifugum</i> , <i>Senna alata</i> , <i>Terminalia mollis</i> , <i>Tetradenia riparia</i> , <i>Vernonia shirensis</i> , <i>Zingiber officinale</i>
Headach	3	0.03	25	0.016	<i>Entada abyssinica</i> , <i>Monodora myristica</i> , <i>Vernonia shirensis</i> .

Indications	Nt (n=76)	MCI (n=100)	Nc	CI (n=1604)	Taxon
Hepatitis	3	0.03	108	0.067	<i>Ageratum conyzoides</i> , <i>Phyllanthus muellerianus</i> , <i>Zingiber officinale</i> .
Hernias	1	0.01	10	0.006	<i>Mesosphaerum suaveolens</i>
Hypertension	5	0.05	159	0.099	<i>Albizia adianthifolia</i> , <i>Bridelia ferruginea</i> , <i>Gardenia ternifolia</i> , <i>Monodora myristica</i> , <i>Parinari curatellifolia</i>
Indigestion	1	0.01	5	0.003	<i>Hibiscus cannabinus</i>
Infections	1	0.01	10	0.006	<i>Garcinia punctata</i>
Inflammations	6	0.06	217	0.135	<i>Baillonella toxisperma</i> , <i>Pentaclethra macrophylla</i> , <i>Phyllanthus amarus</i> , <i>Sida rhombifolia</i> , <i>Strychnos spinosa</i> , <i>Symphytum officinale</i> .
Intestinal worms	5	0,05	134	0,084	<i>Ficus sur</i> , <i>Pterocarpus angolensis</i> , <i>Khaya nyasica</i> , <i>Memecylon flavovirens</i> , <i>Pentaclethra macrophylla</i> .
Itching	1	0.01	1	0.001	<i>Cylicodiscus gabunensis</i>
Jaundice	2	0.02	10	0.006	<i>Balanites aegyptiaca</i> , <i>Ficus sur</i> .
Joint pain	1	0.01	1	0.001	<i>Capsicum frutescens</i>
Kidney failure	2	0.02	2	0.001	<i>Baillonella toxisperma</i> , <i>Canarium schweinfurthii</i> .
Leprosy	1	0.01	3	0.002	<i>Anacardium occidentale</i>
Lumbago	2	0.02	1	0.001	<i>Baillonella toxisperma</i> , <i>Entada abyssinica</i>
Malaria	13	0.13	258	0.161	<i>Costus afer</i> , <i>Crossopteryx febrifuga</i> , <i>Entada abyssinica</i> , <i>Gardenia ternifolia</i> , <i>Gymnanthemum amygdalinum</i> , <i>Khaya nyasica</i> , <i>Mesosphaerum suaveolens</i> , <i>Parinari curatellifolia</i> , <i>Pentaclethra macrophylla</i> , <i>Psorospermum febrifugum</i> , <i>Syzygium cordatum</i> , <i>Vernonia amygdalina</i> , <i>Zanha africana</i> .
Measles	1	0.01	4	0.002	<i>Crossopteryx febrifuga</i>
Meningitis	3	0.03	3	0.002	<i>Antrocaryon klaineianum</i> , <i>Citrus lemon</i> , <i>Monodora myristica</i> .
Meningitis	1	0.01	5	0.003	<i>Syzygium cordatum</i>
Muscle pain	1	0.01	1	0.001	<i>Capsicum frutescens</i>
Muscle spasm	2	0.02	1	0.001	<i>Landolphia owariensis</i> , <i>Passiflora foetida</i>
Muscular weakness	1	0.01	3	0.002	<i>Uapaca guineensis</i>
Nasal congestion	1	0.01	5	0.003	<i>Memecylon flavovirens</i>
Nervousness	1	0.01	1	0.001	<i>Passiflora foetida</i>
Obesity	1	0.01	2	0.001	<i>Hibiscus cannabinus</i>
Osteomyelitis	1	0.01	10	0.006	<i>Thonningia sanguinea</i>
Painful periods	1	0.01	1	0.001	<i>Antrocaryon klaineianum</i>
Parasites	1	0.01	1	0.001	<i>Pentaclethra macrophylla</i>
Pneumonia	2	0.02	2	0.001	<i>Dalbergia lactea</i> , <i>Pentadiplandra brazzeana</i>

Indications		Nt (n=76)	MCI (n=100)	Nc	CI (n=1604)	Taxon
Rheumatism		8	0.08	38	0.024	<i>Asparagus africanus</i> , <i>Capsicum frutescens</i> , <i>Crossopteryx febrifuga</i> , <i>Costus afer</i> , <i>Erigeron bonariensis</i> , <i>Erigeron sumatrensis</i> , <i>Markhamia tomentosa</i> , <i>Plumbago zeylanica</i>
Rhinitis		2	0.02	1	0.001	<i>Luffa cylindrica</i> , <i>Ocimum gratissimum</i>
Schistosomiasis		2	0.02	1	0.001	<i>Antidesma venosum</i> , <i>Gardenia ternifolia</i>
Sexual dysfunction		9	0.09	478	0.298	<i>Aidia micrantha</i> , <i>Albizia adianthifolia</i> , <i>Ficus sur</i> , <i>Garcinia punctata</i> , <i>Indigofera capitata</i> , <i>Indigofera capitata</i> , <i>Moringa oleifera</i> , <i>Senna alata</i> , <i>Uapaca guineensis</i> .
Sexual	Transmitted	14	0.14	917	0.572	<i>Ageratum conyzoides</i> , <i>Annona senegalensis</i> , <i>Antidesma venosum</i> , <i>Antrocaryon klaineianum</i> , <i>Asparagus africanus</i> , <i>Bridelia ferruginea</i> , <i>Euphorbia hirta</i> , <i>Ficus brachypoda</i> , <i>Ficus stuhlmannii</i> , <i>Ficus sur</i> , <i>Garcinia punctata</i> , <i>Gymnanthemum amygdalinum</i> , <i>Syzygium cordatum</i> , <i>Vernonia amygdalina</i> .
Infections						
Sickle cell disease		1	0.01	1	0.001	<i>Bridelia ferruginea</i>
Sinusitis		2	0.02	1	0.001	<i>Albizia adianthifolia</i> , <i>Luffa cylindrica</i>
Skin infections		1	0.01	6	0.004	<i>Markhamia tomentosa</i>
Snake bite		1	0.01		0.001	<i>Diplorhynchus condylocarpon</i>
Sore throat		1	0.01	3	0.002	<i>Asparagus africanus</i>
stretch mark		2	0.02	31	0.019	<i>Erigeron bonariensis</i> , <i>Erigeron sumatrensis</i>
Swollen feet Edema		2	0.02	2	0.001	<i>Cylicodiscus gabunensis</i> , <i>Cymbopogon citratus</i>
Thrombosis		1	0.01	4	0.002	<i>Coleus kilimandschari</i>
Tooth decay		6	0.06	49	0.031	<i>Albizia adianthifolia</i> , <i>Aloe vera</i> , <i>Bridelia ferruginea</i> , <i>Dalbergia lactea</i> , <i>Gardenia ternifolia</i> , <i>Syzygium cordatum</i> .
Tuberculosis		2	0.02	1	0.001	<i>Bridelia ferruginea</i> , <i>Ficus brachypoda</i>
Typhoid		2	0.02	1	0.001	<i>Alstonia boonei</i> , <i>Syzygium cordatum</i>
Vertigo		1	0.01	21	0.013	<i>Cyperus articulatus</i>
Weight loss		1	0.01	4	0.002	<i>Coleus kilimandschari</i>
Wounds		11	0.11	963	0.600	<i>Aframomum melegueta</i> , <i>Alchornea cordifolia</i> , <i>Aspilia kotschyi</i> , <i>Bridelia ferruginea</i> , <i>Diplorhynchus condylocarpon</i> , <i>Droogmansia munamensis</i> , <i>Ficus brachypoda</i> , <i>Markhamia tomentosa</i> , <i>Mesosphaerum suaveolens</i> , <i>Musa paradisiaca</i> , <i>Phyllanthus muellerianus</i> .

Legend: Nt: Number of taxa; Nc : Number of citations ; MCI : Medical Capability Index. The Medical Capability Index (MCI) refers to a community's potential to manage a disease; AHT: Arterial hypertension; HIV: human immunodeficiency virus; STI : Sexually transmitted infections.

According to the informants consulted, the taxa with the greatest number of medicinal uses in the study area are *Khaya nyasica* Stapf ex Baker f.b (Meliaceae), with 11 uses; *Syzygium cordatum* Subsp.a (Myrtaceae), with 11 uses; and *Ficus sur* Forssk f (Moraceae), with 9 uses. The next most frequently cited taxa are *Albizia adianthifolia* (Schum.) W. Wight (Fabaceae) (8 uses) and *Crossopteryx febrifuga* (Afzel. ex G. Don) Benth. (Rubiaceae) (8 uses). Collectively, they are implicated in over 60% of the pathologies mentioned in this study (Table 2).

3.5. Socio-demographic profile of persons consulted

The individuals consulted in this study were either household contacts (94.8%), herbalists (1.6%), or practitioners of traditional medicine (3.8%), with the majority being women (57.9%). The age range of the individuals consulted was 50 to 60 years old, with the youngest being 21 and the oldest 73 years old. The most respondents had received no formal education (37.6%) or had completed only primary school (36.7%). The informants were distributed across the 44 districts that comprise the seven communes of the city of Lubumbashi in similar proportions, ranging from 12.9% to 22.4%. However, in the Kamalondo and Kampemba communes, the proportion was slightly lower, at 7.9% and 9.9%, respectively. The respondents were engaged in six occupational categories, with housework (40.8%) and liberal (21.2%) representing the most prevalent types. In the majority of cases, the informants have accumulated over 11 years of experience in the use of medicinal plants (Table 6).

Table 6 Socio-demographic characteristics of people consulted

Category	Household (n=1520)		Herbalists (n=25)		THs (n=59)		Total (n=1604)
	Nc	%	Nc	%	Nc	%	%
Age							
20-30	280	18.40	7	28	11	18.64	18.58
30-40	672	44.20	11	44	21	35.59	43.89
40-50	410	26.97	5	20	14	23.72	26.75
50-60	153	10.06	2	8	8	13.55	10.16
>60	5	0.33	0	0	5	8.47	0.62
Experience (Year range)							
0-5	149	9.8	6	24	10	16.9	10.3
6-10	105	6.9	7	28	9	15.3	7.5
11-15	1006	66.2	3	12	20	33.9	64.2
16-20	17	1,1	8	32	11	18.6	2.2

Category	Household (n=1520)		Herbalists (n=25)		THs (n=59)		Total (n=1604)
	Nc	%	Nc	%	Nc	%	%
21-25	143	9.4	1	4	9	15.3	9.5
Profession							
Retailer	304	20	25	100	5	8.47	20.82
Civil servant	191	12.57	0	0	0	0	11.91
Liberal	320	21.05	0	0	40	67.79	21.2
Housekeeper	655	43.09	0	0	0	0	40.84
THs	50	3.29	0	0	14	23.72	3.99
Herbalist	907	59.67	17	68	4	6.78	57.86
Gender							
Female	907	59.7	17	68	4	6.8	57.9
Male	613	40.3	8	32	55	93.2	42.1
Residence							
Annex	210	13.8	5	20	15	25.4	14.3
Kamalondo	157	10.3	0	0	2	3.4	9.9
Kampemba	112	7.4	3	12	11	18.6	7.9
Katuba	243	16.0	3	12	7	11.9	15.8
Kenya	198	13.0	4	16	5	8.5	12.9
Lubumbashi	347	22.8	4	16	9	15.3	22.4
Ruashi	253	16.6	6	24	10	16.9	16.8
Level of education							
None	587	38.62	8	32	8	13.6	37.59
Primary	542	35.66	12	48	35	59.32	36.72
Vocational	138	9.08	2	8	5	8.93	9.04
Secondary	239	15.72	3	12	11	18.64	15.77
University	18	1.18	0	0	0	0	1.12

Legend: THs: traditional healers; NC: Number of citation ; %: Percentage

4. Discussion

4.1. Plants used to treat hemorrhoidal diseases.

Until now, traditional medicine has been the primary means of treating hemorrhoidal pathologies in Katanga, and various plants are used for this purpose. This study reports for the first time 24 taxa of the 100 plants reported in our surveys as plants used in the management of hemorrhoidal pathologies, in view of the available literature. These are *Acmella caulirhiza* Delile (Asteraceae), *Aidia micrantha* (K.Schum.) Bullock ex F.White (Rubiaceae), *Amaranthus hybridus* L. (Amaranthaceae), *Anonidium mannii* (Oliv). Engl. & Diels (Annonaceae), *Antirrhinum*

majus L. (Plantaginaceae), *Baccharoides adoensis* (Sch.Bip. ex Walp.) H.Robc (Asteraceae), *Chromolaena odorata* (L.) R.M. King & H. Rob. (Asteraceae), *Crossopteryx febrifuga* (Afzel. ex G.Don) Benthc (Rubiaceae), *Diplorhynchus condylocarpon* (Mull. Arg.) (Apocynaceae), *Droogmansia munamensis* De Wild (Fabaceae), *Erigeron bonariensis* L. (Asteraceae), *Ficus laurifolia* Lam.f (Moraceae), *Ficus stuhlmannii* Warb (Moraceae), *Ficus sur* Forssk (Moraceae), *Garcinia punctata* Oliv. (Clusiaceae), *Indigofera capitata* Kotschyb (Fabaceae), *Mesosphaerum suaveolens* (L.) Kuntze (Lamiaceae), *Ocimum centraliafricanum* R.E.Fr. (Lamiaceae), *Passiflora foetida* L. (Passifloraceae), *Pericopsis angolensis* (Baker) (Fabaceae), *Symphytum officinale* L. (Boraginaceae), *Uapaca guineensis* Müll.Arg.c (Phyllanthaceae), *Zanha africana* (Radlk.) Exellb (Sapindaceae) and *Zanthoxylum chalybeum* Engl. (Rutaceae) (Table 2). The most cited by more than 60% of interviewees are: *Mesosphaerum suaveolens* (CI: 0.704), *Ficus laurifolia* (CI: 0.635), *Ficus stuhlmannii* (CI: 0.635) and *Erigeron bonariensis* (CI: 0.623).

Of the 24 taxa, 14 have not previously been studied for their anti-hemorrhoidal activity. In order of importance of citation, the taxa are as follows: *Crossopteryx febrifuga*, *Ficus stuhlmannii*, *Ficus laurifolia*, *Indigofera capitata*, *Diplorhynchus condylocarpon*, *Droogmansia munamensis*, *Baccharoides adoensis*, *Aidia micrantha*, *Erigeron bonariensis*, *Garcinia punctata*, *Acmella caulirhiza*, *Amaranthus hypochondriacus*, *Anonidium mannii*, and *Ocimum centralafricanum*. According to Plants of the World Online (<https://powo.science.kew.org>), all of these taxa are native to the DRC, with the exception of *Erigeron bonariensis* and *Amaranthus hypochondriacus*, which are introduced taxa.

Of the 14 taxa initially reported by this study as plants with anti-hemorrhoidal uses in Katanga, a literature search revealed that only eight taxa, namely *Acmella caulirhiza* [136], *Aidia micrantha* [79], *Amaranthus hypochondriacus* [137], *Anonidium mannii* [138], *Crossopteryx febrifuga* [40,41], *Diplorhynchus condylocarpon* [37,42,139], *Droogmansia munamensis* [40], *Garcinia punctata* [140,141] have previously been documented as medicinal plants in Congolese traditions.

Although not reported as medicinal plants in the Democratic Republic of the Congo (DRC), three of the fourteen taxa, *Indigofera capitata*, *Baccharoides adoensis*, and *Erigeron bonariensis*, are reported as medicinal plants in other African regions. In Nigeria [142] and, in Uganda [143], powder from *Indigofera capitata* is employed as a

remedy for snakebites, while in the Republic of Congo, the dried infrutescences of this plant are utilized for their hallucinogenic properties [144]. In Zimbabwe, *Baccharoides adoensis* is traditionally employed for the treatment of fever, upper respiratory tract infections, and currently for tuberculosis [145]. In Kenya, the root decoction is utilized for the treatment of sexually transmitted diseases, heart and kidney problems, while the leaves are employed in the treatment of malaria and tuberculosis [146]. In South Africa, the decoction of *Erigeron bonariensis* [synonym: *Conyza bonariensis* (L.)] is a folk remedy that is widely used for the treatment of tuberculosis. The leaves of this plant are employed as folk medicine for the treatment of a number of ailments, including rheumatism, gout, cystitis, nephritis, dysmenorrhea, toothache, and headaches [147].

To the best of our knowledge, although *Ficus stuhlmannii*, *Ficus laurifolia*, and *Ocimum centraliafricanum* are considered native plants to the country, no medicinal use has yet been reported for any of them. This study, therefore, presents the first account of the related ethnomedicinal knowledge. The fact that *Ocimum centraliafricanum* was reported only by the household population indicates that ethnomedicinal knowledge in our study area is not the exclusive domain of traditional healers. This is further corroborated by the fact that ethnomedicinal knowledge in our area is a family legacy, as evidenced by the findings of previous studies [37,40,42,148]. Although these plants are not reported as having anti-hemorrhoidal uses, some taxa of the same genus are used in traditional medicine to treat hemorrhoidal ailments. This is exemplified by *Ocimum gratissimum*, whose leaves are utilized in Benin as a decoction against typhoid fever [149]. The same is true of the taxon *Ocimum basilicum* L described in the Persian pharmacopoeia, which has been demonstrated to possess anti-inflammatory and myorelaxant properties in vivo, thus confirming its indication in the management of hemorrhoids [150]. In the *Ficus* genus, several taxa have been reported to be used in traditional medicine to treat hemorrhoids, including *Ficus vallischooudae* delile [151], *Ficus exasperata* Vahl [152], *Ficus carica* L [150], *Ficus umbellata* (Vahl) [153], *Ficus benghalensis* [154], *Ficus sycomorus* L, *Ficus palmata* Forssk, and *Ficus vasta* Forssk [30]. The two taxa of the *Ficus* genus reported in our study contribute to the growing list of

numerous species of this genus used in various cultures for the management of hemorrhoidal pathologies.

The majority of the medicinal plants mentioned by the respondents belong to the Fabaceae family, with 10 taxa identified (Fig. 3a). The prevalence of Fabaceae in this study can be attributed to the fact that it is the most dominant family of medicinal plants in the region. Ethnobotanical studies conducted in the Miombo region (Democratic Republic of the Congo, Tanzania, Angola, Zambia, Malawi, Zimbabwe, and Mozambique) have demonstrated that Fabaceae is the most prevalent family of medicinal plants. A case in point is the study conducted in the Sikonge district of Tanzania, which found that Fabaceae accounted for seven out of 28 taxa [35]. Another example is the study conducted in the province of Bié in Angola, which found that Fabaceae accounted for 12 out of 87 taxa [155]. A similar pattern was observed in the provinces of West, Copperbelt, Central, and Northern Zambia, where Fabaceae accounted for seven out of 35 taxa [156]. A similar pattern emerges from studies conducted in Zomba, Malawi, where Fabaceae accounted for 10 out of 54 taxa [157], and in Gokwe, Zimbabwe, where Fabaceae represented 13 out of 127 taxa [158]. A bibliographical review of medicinal plants in Mozambique also highlighted the preponderance of Fabaceae, with 95 out of 731 taxa [159]. In the Democratic Republic of the Congo (DRC), the Miombo forest covers the provinces of Lualaba, Tanganyika, and Haut-Katanga. In Haut-Katanga province, where the city of Lubumbashi is located, no study has inventoried and characterized all the medicinal plants in the region. However, several ethnomedicinal studies targeting a specific pathology have been carried out, and in most cases, Fabaceae have taken the lead. These include the study of plants used to treat gastrointestinal disorders in Kamina and Kanyama (n=10 taxa, Fabaceae =2 taxa [160], the study of anti-cariogenic plants in Lubumbashi (n=14 taxa, Fabaceae = 3 taxa [161], and the study of anti-malarial plants in the Lubumbashi area and its surroundings (n= 19, Fabaceae: 11 taxa [162], n= 96, Fabaceae = 22 taxa [41]). The study on anti-diabetic plants from Lubumbashi and its surroundings yielded 45 taxa, with Fabaceae accounting for 11 taxa [40]. The study on plants used against urogenital

schistosomiasis yielded 61 taxa, with Fabaceae accounting for 17 taxa [37]. The study on plants used in the management of sexual dysfunctions in the Kampemba commune (Lubumbashi) yielded 21 taxa, with Fabaceae accounting for 7 taxa [163] is followed by the study on plants used in Lubumbashi and surrounding areas to treat gastritis (n=14, Fabaceae = 3 taxa [164]. Finally, the study on plants used in Lubumbashi to treat typhoid fever (n=54, Fabaceae = 20 taxa) [42] is presented.

Each of the plants reported in this study is named in one of the DRC's local languages, with Swahili, Luba, and Bemba being the most prevalent (Fig. 3b). This result is consistent with previous ethnobotanical studies conducted in the region [37,40,163,165], which have reported the predominance of Bemba and Luba in the practice of traditional medicine in Katanga. It is noteworthy that the names of ethnic groups not endemic to Katanga, such as Lingala and Kikongo, which are languages spoken in the west of the country, also appear in the list. This may be indicative of a certain degree of cultural cross-fertilization that traditional Katangese medicine has undergone [42].

A citation index: CI of 0.7 or greater was observed in three taxa, which were the most frequently cited in the overall study. The most frequently cited taxa were *Phyllanthus amarus* Schumacher & Thonn f. (Phyllanthaceae), *Terminalia mollis* MA Lawson f. (Combretaceae) and *Mesosphaerum suaveolens* (L) Kuntze (Lamiaceae). These taxa were reported by both traditional healers and the general population. According to Plants of the World Online (<https://powo.science.kew.org/>), only *Terminalia mollis* is native to the DRC, while the two other taxa are introduced plants. To the best of our knowledge, only the first two taxa are reported as medicinal plants in the DRC.

In southern Katanga, the infusion of *Phyllanthus amarus* leaves is employed to treat asthma, diabetes, typhoid fever, hepatitis, bronchial infections, and peptic ulcer [42], while decoction of the whole plant is utilized to treat dysentery [39]. In Central-Kongo, infusion of the plant's leaves is employed to promote diuresis, while maceration of the roots is utilized to treat amoebiasis and snail fever [138]. Its potential use in the

treatment of haemorrhoidal pathologies may be based not only on its anti-inflammatory properties, as demonstrated *in vitro* [116], but also on the numerous flavonoids reported in its leaves [166], which have been associated with strengthening blood vessel walls, increasing venous tone and lymphatic drainage, and normalizing capillary permeability at the anal level [167,168].

The root bark of *Terminalia mollis* is employed in Katanga as a decoction for a variety of ailments, including dysentery, intestinal worms, cancer, lice, diarrhea, abdominal pain [139,169], typhoid fever [42], sexual dysfunction [163], diabetes, and syphilis [40]. In powder form, they are employed as a treatment for jaundice. The anti-inflammatory activity observed *in vitro* and the presence of tannins in its root barks may be the basis for its use against hemorrhoids.

To the best of our knowledge, *Mesosphaerum suaveolens* has never been reported as a medicinal plant in the Democratic Republic of the Congo (DRC). However, the plant is a species that has been traditionally used for the treatment of a wide range of ailments, including stomach pain, cough, verminosis, ulcer, liver disease, fever, influenza, nasal congestion, and inflammation in various countries, such as Brazil, Benin, India, Nigeria, and Togo [170]. The use of the plant against hemorrhoids is linked to the presence of numerous phenolic compounds isolated from its leaves and to its anti-inflammatory activity *in vitro* [101].

The majority of the taxa identified in this study are either trees or shrubs (Fig. 5), which is consistent with previous ethnobotanical studies conducted in the region [39,42,162,171]. Previous ethnomedicinal studies targeting anti-hemorrhoidal plants have yielded varying results regarding the predominant morphology of the taxa used, depending on the study. In a study conducted in southwestern Nigeria, the majority of taxa inventoried were trees [71]. This is also the case for the study carried out in southeast Cameroon [34] or in Tabora, Tanzania [[35]. The results were similar to those reported in our study. In contrast, in studies conducted in Madurai district in India [172] or in north-central Nigeria [32], grass was in the majority. The morphological aspect of

taxa is therefore not a specific characteristic of plants with a reputation for anti-hemorrhoidal action. However, the preferential use of tree or shrub morphological types offers the advantage of year-round availability of plant material, providing the possibility of uninterrupted pathology management throughout the year. However, this practice increases the risk of extinction of a taxon in great demand, especially in the current context of global warming [173–175]. It would be beneficial for traditional healers to receive training in plant taxon conservation techniques, particularly *in situ* and *ex vivo* methods and biotechnological approaches. Additionally, it is crucial for the country to regulate harvesting practices and the use of plant resources for the intensive management and conservation of medicinal plants, with the objective of perpetuating the practice of traditional medicine while eradicating the risk of extinction of interesting taxa.

4.2. Anti-hemorrhoidal Recipes

Of the 117 anti-hemorrhoidal recipes, over 60% are based on leaves, which have been prepared in 7 different ways, with decoction being the most common method (Figure 8). The preponderance of leaves and decoction as the part used and mode of extraction of the active ingredient are reported in various ethnobotanical studies carried out in Katanga [37,41,161,162]. However, it should be noted that these modes of extraction are not universally consistent. In other ethnobotanical studies conducted in the same region, the root was found to be the main organ used [40,176]. This diversity in the organ preferentially used is also observed in ethnobotanical studies focusing on plants with anti-hemorrhoidal uses. In some cases, the leaf is the most commonly used organ [30,32,177], while in others, the root [35,178–180], stem bark [34], or aerial parts [181] are the preferred choice. The lack of consensus on the most important organ for the use of plants in the management of hemorrhoidal pathologies can be justified by the diversity of compounds likely to contribute to the management of hemorrhoidal disorders. The utilization of leaves as a source of raw material offers a number of advantages. (i) Leaves are generally more accessible and abundant than other plant

parts such as roots or bark, which may require damaging the plant to harvest [182]; (ii) they can contain a high concentration of bioactive compounds because they are the site of photosynthesis and other metabolic processes, which can lead to the accumulation of beneficial phytochemicals [183]. (iii) They are more readily dried and ground than other plant parts, which renders them more convenient for preparing extracts and medicines [183]. (iv) They are more sustainable as they can be harvested without killing the plant, allowing for continuous use and conservation of plant species [182].

In traditional medicine, the preparation of a medicinal recipe through the use of a decoction offers several advantages. Firstly, it is an optimal method for extracting active compounds that are stable at high temperatures. Secondly, the process does not necessitate the use of complex or expensive equipment, thereby facilitating its widespread applicability. Thirdly, the process is relatively straightforward and does not necessitate the involvement of a trained operator, which is advantageous in traditional settings. Fourthly, it is a cost-effective method that can be readily implemented in a domestic setting. Nevertheless, decoctions are unsuitable for the extraction of heat-sensitive constituents, as the boiling process may result in their degradation [184,185]. Nevertheless, it is necessary to await the results of biological experiments before a proper assessment of the choice of organ and method of preparation used for each recipe can be made.

In contrast to previous ethnobotanical studies conducted in the region, which predominantly reported oral administration [40–42], our study primarily focused on topical applications. This is consistent with previous studies on plants purported to have anti-hemorrhoidal properties [30,32,34]. This is particularly evident when considering the localized nature of hemorrhoidal pathologies.

The majority (n=101) of the recipes documented in this study utilize a single plant, a finding consistent with several ethnobotanical studies conducted in Katanga [37,38,40]. However, there are also recipes resulting from mixtures of two or three taxa (n=16), a

pattern observed in other ethnobotanical studies conducted in the region [42,163]. While the combination of plants may be based on the search for a certain synergy, there is a risk of increasing the risk of drug interactions and collateral effects. Nevertheless, it is of the utmost importance to remain vigilant during biological experiments in order to ensure that each recipe is duly evaluated.

4.3. Knowledge of Hemorrhoidal Diseases

In this survey, respondents referred to hemorrhoidal pathologies in Swahili as "*Kilonda ntumbu*," which translates to "the wound of the belly." This term is linked to the perception among the local population that hemorrhoidal pathologies originate from a specific type of diet, which is reflected in the allusion to the stomach.

The fact that more than 65% of subjects mentioned symptoms typically associated with hemorrhoidal pathologies (Table 6) suggests that the people interviewed were familiar with the pathology for which they mentioned healing plants. This finding is consistent with previous studies that have demonstrated the prevalence of hemorrhoids in Lubumbashi [31].

4.4. Other pathologies treated by inventoried plants

In the Democratic Republic of the Congo (DRC), the 10 most deadly pathologies are in order of importance, as follows: malaria, tuberculosis, lower respiratory infections, neonatal disorders, diarrheal diseases, stroke, ischemic heart disease, road injuries, hypertensive heart disease, and cirrhosis and other chronic liver diseases [174]. Among the most deadly pathologies, the plants inventoried in this study were cited in the treatment of diarrhea (31 taxa), malaria (10 taxa), lower respiratory infections (7 taxa), and tuberculosis (2 taxa) (Table 4). It can be reasonably concluded that the resource persons consulted during this study have plant resources capable of coping with at least four of the ten most deadly pathologies in the Democratic Republic of the Congo. This provides indirect evidence of the capacity of traditional medicine in Lubumbashi to address the major health challenges facing the DRC.

In addition to hemorrhoidal pathologies, the resource persons have identified at least ten taxa for each of the following five pathologies: In descending order of importance, these are gastrointestinal disorders, diabetes, sexually transmitted infections, malaria, and wounds (Table 4). The availability of multiple taxa to treat a given pathology indicates the community's capacity to address the condition in question. Given that traditional medicine is culturally specific, this can provide insight into the particular management approaches of a given community in the context of specific pathologies.

4.5 The socio-demographic characteristics of the interviewees.

In contrast to previous ethnobotanical studies conducted in Lubumbashi, this study interviewed a greater proportion of women than men [40–42,163]. This discrepancy can be attributed to the fact that, unlike the present study, the aforementioned studies were conducted among practitioners of traditional medicine, predominantly men. In contrast, the present study's account was conducted with the general population, which, like the national population in the 40-60 age range, is predominantly female according to Index Mundi in 2023 (<https://www.indexmundi.com>). Furthermore, the fact that the majority of informants were met in households may also explain the preponderance of women. Indeed, in households, the probability of encountering women during surveys is higher than that of encountering men, particularly given that in our society, household tasks are typically performed by women. Almost two-thirds of those surveyed had experience with the use of medicinal plants to treat their ailments, which suggests that traditional medicine plays a role in the study area. These data are consistent with a study conducted in the region on the practice of traditional medicine, which found that 79.4% of respondents had utilized traditional medicine to address their health or social concerns [31]. Moreover, several of the plants referenced in this study have been identified as anti-hemorrhoidal agents in other regions, and some have been demonstrated to possess indirect anti-hemorrhoidal properties (Table 2). These findings should enhance the credibility of the information presented in the present study.

5. Conclusion

The findings of this study indicate that the population of Lubumbashi (DRC) employs a diverse range of taxa to treat hemorrhoidal diseases, particularly internal hemorrhoids. However, these plants offer considerable potential for managing a wide range of other pathologies, including gastrointestinal disorders, wounds, sexually transmitted infections, and diabetes. Among the taxa employed, some are endemic to the study area and specific to the region. Currently, studies are being conducted to assess the efficacy of these plants in the management of hemorrhoidal pathologies. Nevertheless, further ethnopharmacological studies are required to validate the other properties of these plants. Such studies could result in the development of improved traditional medicines or the identification of new drug candidate molecules. These findings provide compelling evidence to motivate self-sufficient communities to explore sustainable methods of conserving these plants for the collective benefit of biodiversity conservation in the Miombo.

6. Abbreviations

TA(Tropical Africa); **NEA**(North-East Africa); **CA** (Central Africa); **SA**(Southern Africa); **MA**(Madagascar); **EA**(East Africa); **NA**(North Africa); **WE**(West Africa); **MCI** : Medical Capability Index ; **CI**: citation index; **GT**: geographical type; **UP**: used part; **MT**: morphological type; **FU**: form of use; **MUI**: medicinal use index; **TH**: type of hemorrhoid; **WP**: Whole plant ; **Rb**: Root bark; **Sb**: Stem Bark; **Lv** : Leaves; **Sd** : Seed; **Fr**: Fruit; **Rz**: Rhizome; **NH**: herbarium number; **FU**: form of use; **MUI**: medicinal use index; **THD**: type of hemorrhoid diseases; **Nc**: Number of citations; **Nt**: Number of taxa.

7. Acknowledgements

The authors would like to thank the Kipopo Herbarium for the identification of plant species. They also thank the traditional healers, herbalists, and households of Lubumbashi for the acquisition of ethnobotanical data.

8. Authors' contributions

BCV designed, coordinated this research and drafted the manuscript. MM, MBH and BCV carried out experiments and data analysis. BCV, BAS, ONP and LSJB conceived of the study, and participated in research coordination. The all authors read and approved the final manuscript.

9.Funding Statement

No funding was received for this research.

10.Data Availability

The data used in this study are included within the article.

11. Declarations

11.1. Ethics approval and consent to participate

Ethical review was provided by the Department of Pharmacology – Faculty of Pharmaceutical Sciences – University of Lubumbashi (Ethics approval number: FSPUNILU-DP-BD-022022).

11.2. Consent for publication

There is no third party data. It is our original research data; With regard to the people interviewed, Oral informed consent was obtained before the initiation of data collection in all cases.

All concerned individual authors have agreed to the publication of the outcome of this study as stated above under ethical consideration and authors 'contribution.

11.3. Conflicts of Interest

The authors declare that they have no conflicts of interest.

11.4. Author details

¹ Department of Pharmacology - Laboratory of Therapeutic Chemistry and Analysis of Natural Substances - Faculty of Pharmaceutical Sciences - University of Lubumbashi (UNILU) - 27, av. Kato, Commune Kampemba, Lubumbashi - DR Congo.

² Department of Pharmacology - Laboratory of Pharmacognosy – Faculty of Pharmaceutical Sciences – University of Lubumbashi (UNILU)- 27, av Kato, Commune Kampemba, Lubumbashi - DR Congo.

³ Department of Chemistry– Faculty of Sciences–University of Lubumbashi (UNILU) N°1 Maternity Avenue, Commune of Lubumbashi, Lubumbashi - DR Congo.

12. References

- [1] Fiðere I, Groma V, Goldiðð NR, Gardovskis A, Gardovskis J. Worldwide disease — Haemorrhoids . How Much Do We Know? *Proc Latv Acad Sci Sect B* 2021;**75**(1):1–10.
- [2] Mahmoud A, Abd Elmoniem S, Ghonaem S, Mohammed W. Effect of Nursing Guidelines on Hemorrhoids Symptoms among Pregnant Women. *Egypt J Nurs Heal Sci* 2023;**4**(2):4–28.
- [3] Gupta S, Singh TG, Baishnab S, Garg N, Kaur K, Satija S. Review article recent management of Hemorrhoids: A pharmacological and surgical perspective. *Plant Arch* 2020;**20**:3828–3837.
- [4] Lalisang TJ. Hemorrhoid: Pathophysiology and Surgical Management Literature review. *New Ropanasuri J Surg* 2016;**1**(1):31–36.
- [5] Sinha P, Singh MP, Rajpoot BK. Review article on guda arsha (haemorrhoids): causes, sign and symptoms, prevention & treatment. *world J Pharm Res* 2021;**9**(8):336–344.
- [6] Soeseno SW, Wahyudi PAE, Febyan F. Diagnosis and Management of Internal Hemorrhoids: A Brief Review. *Eur J Med Heal Sci* 2021;**3**(5):1–5.
- [7] Haider R. Hemorrhoids the Clinical Practice. *J Surg Case Reports Images* 2023;**6**(5):e166.
- [8] Sandler RS, Peery AF. Rethinking What We Know About Hemorrhoids. *Clin Gastroenterol Hepatol* 2019;**17**(1):8–15.
- [9] Kibret AA, Oumer M, Moges AM. Prevalence and associated factors of hemorrhoids among adult patients visiting the surgical outpatient department in the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia. *PLoS One* 2021;**16**(4):1–11.
- [10] Ravindranath GG, Rahul BG. Prevalence and risk factors of hemorrhoids: a study in a semi-urban centre. *Int Surg J* 2018;**5**(2):496.
- [11] Lee JH, Kim HE, Kang JH, Shin JY, Song YM. Factors associated with hemorrhoids in Korean adults: Korean national health and nutrition examination survey. *Korean J Fam Med* 2014;**35**(5):227–236.
- [12] Hong YS, Jung KU, Rampal S, Zhao D, Guallar E, Ryu S, et al. Risk factors for hemorrhoidal disease among healthy young and middle-aged Korean adults. *Sci Rep* 2022;**12**(1):1–9.
- [13] Khan MA, Chowdri NA, Parray FQ, Wani RA, Mehraj A, Baba A, et al. "PNR-Bleed" classification and Hemorrhoid Severity Score—a novel attempt at classifying the

hemorrhoids. *J Coloproctology* 2020;**40**(4):398–403.

- [14] Picciariello A, Tsarkov P, Papagni V, Efetov S, Markaryan D, Tulina I, et al. Classifications and clinical assessment of haemorrhoids: the proctologist's corner. *Rev Recent Clin Trials* 2021;**16**(1):10–16.
- [15] Wang L, Ni J, Hou C, Wu D, Sun L. Time to change? Present and prospects of hemorrhoidal classification. *Front Med* 2023;**10**(4):1252468.
- [16] Sheikh P, Goron F, Salmat G, Republic C. The prevalence , characteristics and treatment of hemorrhoidal disease: results of an international web-based survey. *J Comp Eff Res* 2020;**9**(17):1219–1232.
- [17] Rørvik HD, Davidsen M, Gierløff MC, Brandstrup B, Olaison G. Quality of life in patients with hemorrhoidal disease. *Surg Open Sci* 2023;**12**:22–28.
- [18] Komal A, Majumadar D. Hemorrhoids: Incidence & Risk Factors in a Corporate Hospital of Eastern India. *IAR J Med Surg Res* 2021;**2**(1):9–14.
- [19] Al-masoudi RO, Shosho R, Alquhra D, Alzahrani M, Hemdi M. Prevalence of Hemorrhoids and the Associated Risk Factors Among the General Adult Population in Makkah , Saudi Arabia. *Cureus* 2024;**16**(1):e51612.
- [20] Jain A, Lew C, Aksakal G, Hiscock R, Mirbagheri N. Laser hemorrhoidoplasty in the treatment of symptomatic hemorrhoids: a pilot Australian study. *Ann Coloproctol* 2024;**40**(1):52–61.
- [21] Tilahun A, Teklemariam AB, Tilahun Z, Baye ND, Amsalu A, Mengist B, et al. Prevalence and Associated Factors of Hemorrhoids and Other Perianal Complications During the Puerperium Among Mothers Who Gave Birth at Debre Tabor Referral Hospital, Debre Tabor, Ethiopia, 2022. *Qeios* 2024;**1**:1–10.
- [22] Cheickna T, Issaka D, Modibo S, Modibo T, Soma CL. Study of Hemorrhoid Disease in the Department of General Surgery of the Cs Ref of the Commune I Bamako. *Surg Sci* 2024;**15**:36–47.
- [23] Etiennette M, Note J, Ossibi PE, Ngami RS, Miabaou DM, Philestine C, et al. Surgical Treatment of Hemorrhoidal Disease at Brazzaville University Hospital Center. *Surg Sci* 2024;**15**:159–168.
- [24] Zamane H, Coulibaly A, Kain DP, Kiemtore S, Mireille S, Sanon K, et al. External Hemorrhoidal Thrombosis in Recent Births at the Maternity Ward of Saint Camille Hospital in Ouagadougou. *Open J Obstet Gynecol* 2023;**13**:1487–1497.
- [25] Okafor HC, Okafor JN, Okafor RI. Hemorrhoids Among People of Sifawa Community in Sokoto State Hemorrhoids Among People of Sifawa Community in Sokoto State. *J Fam Med Heal Care* 2023;**9**(2):28–33.
- [26] Ma W, Guo J, Yang F, Dietrich CF, Sun S. Progress in endoscopic treatment of hemorrhoids. *J Transl Intern Med* 2020;**8**(4):237–244.
- [27] Stratta E, Trompetto M. Conservative treatment of hemorrhoidal disease. *Rev Recent Clin Trials* 2021;**16**(1):87–90.

- [28] Devi V, Deswal G, Dass R, Chopra B, Kriplani P, Grewal A, et al. Hemorrhoid Disease: A Review on Treatment, Clinical Research and Patent Data. *Infect Disord Targets* 2023;**23**(6):15–35.
- [29] Boakye Y, Mensah D, Agyei E, Agyen R, Agyare C. Role of Medicinal Plants in the Treatment of Hemorrhoids. In: *Herbs*. Apple Academic Press: Florida, 2022: 45–59.
- [30] Getachew M, Belayneh A, Kebede B, Alimaw Y. Medicinal plants used for management of hemorrhoids in Ethiopia: A systematic review. *Heliyon* 2022;**8**:e10211.
- [31] Mutombo CS, Bakari SA, Ntabaza VN, Nachtergaeel A, Lumbu JS, Duez P, et al. Perceptions and use of traditional African medicine in Lubumbashi , Haut-Katanga province (DR Congo): A cross-sectional study. *PLoS One* 2022;**17**(10):1–25.
- [32] Chukwu MN, Ayodele SM, Edoke BP. Ethnobotanical Survey of Plants used in Treatment of Haemorrhoids in North-Central Nigeria. *Univ Thi-Qar J Sci* 2023;**10**(2):131–136.
- [33] Ugwu CO, Asogwa FK, Ali JI, Ozoani HA, Asadu C, Nwokedi KC, et al. Anti- hemorrhoids potential of selected medicinal plants used in treatment of Hemorrhoids: A review. *GSC Adv Res Rev* 2024;**18**(3):36–42.
- [34] Mvogo Ottou PB, Ngotta Biyon JB, Mokake SE, Bissemb PO, Owono Fouda LR, Ngouondjou Foze T, et al. Knowledge of Tradi-Practitioners on Hemorrhoidal Disease and Anti-Hemmoroidal Plants in the Southeast Region of Cameroon: Pharmacology and Preliminary Phytochemistry. *Saudi J Med Pharm Sci* 2020;**06**(04):321–333.
- [35] Kacholi DS, Mvungi Amir H. Herbal remedies used by traditional healers to treat haemorrhoids in Tabora region, Tanzania. *Pharm Biol* 2022;**60**(1):2182–2188.
- [36] Charles NN, Bonareri NL. A survey of medicinal plants used by abagusii traditional healers of South Western Kenya in the treatment of haemorrhoids. *Int J Adv Acad Stud* 2020;**2**(3):244–252.
- [37] Muya K, Tshoto K, Cioci CC, Aseho MM, Kalonji M, Byanga K, et al. Survol ethnobotanique de quelques plantes utilisées contre la schistosomiase urogénitale à Lubumbashi et environs. *Phytothérapie* 2014. doi:10.1007/s10298-014-0877-z.
- [38] Bashige-Chiribagula V, Many-Mboni H, Ntabaza-Ndage V, Numbi Ilunga E, Bakari-Amuri S, Kalonda Mutombo E, et al. Étude ethnobotanique, biologique et chimique de plantes réputées anticariogènes à Lubumbashi – RD Congo. *Phytotherapie* 2017;**15**(1):2–9.
- [39] Mbayo KM, Kalonda ME, Tshisand TP, Kisimba KE, Mulamba M, Richard MK, et al. Contribution to ethnobotanical knowledge of some Euphorbiaceae used in traditional medicine in Lubumbashi and its surroundings (DRC). *J Adv Bot Zool* 2016;**4**(2):1–16.
- [40] Amuri B, Maseho M, Lumbu S, Pierre D, Byanga K. Ethnobotanical survey of herbs used in the management of diabetes mellitus in Southern Katanga Area/DR Congo. *Pan African Med Journal* 2018;**30**(2018). doi:10.11604/pamj.2018.30.218.11718.
- [41] Kalonji M, Muya RK, Mutombo EK, Cimanga BCC, Numbi EWI, Kahumba JB. Aperçu ethnobotanique de plantes réputées antipaludéennes utilisées dans la ville de Lubumbashi et ses environs , dans le Haut-Katanga en RD Congo. *Ethnopharmacologia*

2019;**61**(6):75–84.

- [42] Bashige C V, Pierre KI, Many MH, Kasali mushagalusa F, Okusa NP, Lumbu SJ. Ethnobotanical study of plants used by traditional healers in Lubumbashi (Democratic Republic of Congo) in the management of typhoid fever. *GSC Biol Pharm Sci* 2022;**21**(01):265–286.
- [43] Leonti M. The relevance of quantitative ethnobotanical indices for ethnopharmacology and ethnobotany. *J Ethnopharmacol* 2022;**288**:1–19.
- [44] Jazy MA, Karim S, Morou B, Sanogo R, Mahamane S. Enquête Ethnobotanique Auprès Des Tradipraticiens De Santé Des Régions De Niamey Et Tillabéri Au Niger: Données 2012-2017. *Eur Sci J* 2017;**13**(33):276–304.
- [45] Anna KQ, Isaac A, Emmanuel O, Nana AM-G, Yakubu J, Phoebe EG. In-vivo anti-inflammatory activity of different parts of *Adansonia Digitata*. *J Med Plants Res* 2021;**15**(6):252–259.
- [46] Dibong SD, Brice P, Ottou M, Vandí D, Christelle R, Tchamaha FM, et al. Ethnobotanique des plantes médicinales anti hémorroïdaires des marchés et villages du Centre et du Littoral Cameroun. *J Appl Biosci* 2015;**96**:9072–9093.
- [47] Ilic NM, Dey M, Poulev AA, Logendra S, Kuhn PE, Raskin I. Anti-inflammatory activity of grains of paradise (*Aframomum Melegueta Schum*) extract. *J Agric Food Chem* 2014;**62**(43):10452–10457.
- [48] Sidio S, N'guessan K, Abrou J, Kouadio V. Plants used in traditional medicine against haemorrhoidal pathology by Beté of the Ouragahio sub-prefecture, department of Gagnoa (Côte d'Ivoire). *J Appl Biosci* 2020;**150**:5403–15418.
- [49] Andissa NO, Moussoungou A, Koloungous B, Abena A. Topical Anti-inflammatory effect of aqueous extract ointment of *Ageratum conyzoides* L. in wistar rat. *Int J Phytopharm* 2015;**5**(3):37–41.
- [50] Maroyi A. *Albizia Adianthifolia*: Botany , Medicinal Uses , Phytochemistry , and Pharmacological Properties. *Sci World J* 2018;**18**:7463584.
- [51] Ngunde S, Liyongo CI, Kowozogono R, Gozo TZ, Kwezi BM, Iteku J. Etudes ethnobotanique des plantes utilisées en médecine traditionnelle à Gini (Yakoma , Nord-Ubangi , République Démocratique du Congo). *Int J Appl Res* 2021;**7**(1):36–43.
- [52] Avoseh ON, Mtunzi FM, Ogunwande IA, Ascrizzi R, Guido F. *Albizia lebbeck* and *Albizia zygia* volatile oils exhibit anti-nociceptive and anti-inflammatory properties in pain models. *J Ethnopharmacol* 2021;**268**:113676.
- [53] Kwezi BM, Munkolo PM, Ayingweu CL, Luyeye FL. Utilisation des plantes par la population du groupement Nguma au plateau des Bateke dans la Commune de Maluku à Kinshasa (RD Congo). *J Appl Biosci* 2023;**186**:19556–19570.
- [54] Oruka O, Achuba FI. In vitro Antioxidant and Anti-Inflammatory Activities of Aqueous Leaf Extract of *Alchornea cordifolia*. *J Appl Sci Environ Manag* 2023;**27**(2):299–304.
- [55] Tiomela Japhette D, Yimta F, Jonas K, Tientcheu Jean Philippe D, Nguele Marc Olivier O, Joseph Renaud M, et al. Ethnobotanical survey of anti-hemorrhoidal plants in the

- Bamboutos division, West region of Cameroon. *J Pharmacogn Phytochem* 2019;**8**(5):2287–2292.
- [56] Devaraj A, Karpagam T. Evaluation of anti-inflammatory activity and analgesic effect of aloe vera leaf extract in rats. *Int Res J Pharm* 2011;**2**(3):103–110.
- [57] Olanlokun JO, Olowofolahan AO, Bodede O, Adegbuyi AT, Prinsloo G, Steenkamp P, et al. Anti-inflammatory potentials of the n-hexane fraction of alstonia boonei stem bark in lipopolysaccharide-induced inflammation in wistar rats. *J Inflamm Res* 2021;**14**:3906–3920.
- [58] Paswan S, Srivastava S, Rao C. Incision Wound healing, Anti-inflammatory and Analgesic activity of Amaranthus spinosus in Wistar rats. *Res J Pharm Technol* 2020;**13**(5):2439–2444.
- [59] Yoni M, Senou I, Yonli EY. Caractérisation des Parcs Agroforestiers à Anarcadium occidentale L. et Services Ecosystémiques dans la Forêt Classée de Dindéresso à l'Ouest du Burkina Faso. *Eur Sci Journal, ESJ* 2023;**19**(27):203.
- [60] De Araújo Vilar MS, De Souza GL, De Araújo Vilar D, Leite JA, Raffin FN, Barbosa-Filho JM, et al. Assessment of phenolic compounds and anti-inflammatory activity of ethyl acetate phase of Anacardium occidentale L. bark. *Molecules* 2016;**21**(8):1087.
- [61] Adebayo AH, John-Africa LB, Agbafor AG, Omotosho OE, Mosaku TO. Anti-nociceptive and anti-inflammatory activities of extract of Anchomanes difformis in rats. *Pak J Pharm Sci* 2014;**27**(2):265–270.
- [62] Ngbolua K, Inkoto C, Mongo N, Ashande MC, Masens Y, Mpiana PT. Étude ethnobotanique et floristique de quelques plantes médicinales commercialisées à Kinshasa, République Démocratique du Congo. *Rev Marocaine des Sci Agron Vétérinaires* 2019;**7**(1):118–128.
- [63] Megwas AU, Akuodor GC, Chukwu LC, Aja DO, Okorie EM, Ogbuagu EC, et al. Analgesic, anti-inflammatory and antipyretic activities of ethanol extract of Annona senegalensis leaves in experimental animal models. *Int J Basic Clin Pharmacol* 2020;**9**(10):1477.
- [64] Elgorashi EE, McGaw LJ. African plants with in vitro anti-inflammatory activities: A review. *South African J Bot* 2019;**126**:142–169.
- [65] Jang M, Hwang I, Hwang B, Kim G. Anti-inflammatory effect of Antirrhinum majus extract in lipopolysaccharide-stimulated RAW 264.7 macrophages. *Food Sci Nutr* 2020;**8**(9):5063–5070.
- [66] Fongang ALM, Laure Nguemfo E, Djouatsa Nangue Y, Bogning Zangueu C, Fouokeng Y, Azebaze AGB, et al. Antinociceptive and anti-inflammatory effects of the methanolic stem bark extract of Antrocaryon klaineianum Pierre (Anacardiaceae) in mice and rat. *J Ethnopharmacol* 2017;**203**:11–19.
- [67] Kasali FM, Kadima JN, Peter EL, Mtewa AG, Ajayi CO, Tusiimire J, et al. Antidiabetic Medicinal Plants Used in Democratic Republic of Congo: A Critical Review of Ethnopharmacology and Bioactivity Data. *Front Pharmacol* 2021;**12**(October):1–40.
- [68] Fundiko MC, Mandango MA, Mutambala BM. Etude Ethnobotanique des plantes

- utilisées dans le traitement des maladies de l'appareil digestif à Kinshasa et ses environs, République Démocratique du Congo [Ethnobotany of plants used in the treatment of diseases of the digestive system in Kinshasa. 2017;**31**(1):194–203.
- [69] Bhoyar PS, Deshpande SA, Amle PN, Gomkale KM, Agrawal RN, Sunchuwar UP, et al. Hemorrhoids: A Review on Herbal Treatments and Models for Pharmacological Evaluation. *Int J Pharm Sci Rev* 2020;**63**(27):160–167.
- [70] Kadiatou Tata Traoré, Noufou Ouédraogo, Lazare Belemnaba, Gilchrist Abdoul Laurent Boly, Leïla Marie Esther W. Kabré, Constantin B. Atchadé, et al. Anti-inflammatory and analgesic activities of extracts from *Balanites aegyptiaca* L. Delile (Balanitaceae) root bark: Plant used against liver diseases in Bukina Faso. *African J Pharm Pharmacol* 2019;**13**(18):322–329.
- [71] Soladoye MO, Adetayo MO, Chukwuma EC, Adetunji AN. Ethnobotanical Survey of Plants Used in the Treatment of Haemorrhoids in South-Western Nigeria. *Ann Biol Res* 2010;**1**(4):1–15.
- [72] Olajide OA, Aderogba MA, Okorji UP, Fiebich BL. *Bridelia ferruginea* produces antineuroinflammatory activity through inhibition of nuclear factor-kappa B and p38 MAPK signalling. *Evidence-based Complement Altern Med* 2012;**2012**:ID 546873.
- [73] Nwaehujor Chinaka O, Igile Godwin O, Ode Julius O, Udegbumam Rita I. Anti-inflammatory activities of methanol leaf extract of *Bridelia micrantha* (Hochst) Baill. (Euphorbiaceae) in Wistar rats. *J Appl Pharm Sci* 2014;**4**(6):068–073.
- [74] Jazet Dongmo PM, Tchoumboungang F, Ndongson B, Agwanande W, Sandjon B, Amvam Zollo PH, et al. Chemical characterization, antiradical, antioxidant and anti-inflammatory potential of the essential oils of *Canarium schweinfurthii* and *Aucoumea klaineana* (Burseraceae) growing in Cameroon. *AgricBioUNorth Am* 2010;**1**:606–610.
- [75] Pathy KK, Flavien NB, Honoré BK, Vanhove W, Patrick VD. Ethnobotanical characterization of medicinal plants used in Kisantu and Mbanza-Ngungu territories, Kongo-Central Province in DR Congo. *J Ethnobiol Ethnomed* 2021;**17**(1):1–15.
- [76] Singh PG, S. B M, Shailasreesekhar S, TS G, Basalingappa KM, BV DS. In Vitro Antioxidant, Anti-Inflammatory And Anti-Microbial Activity of *Carica Papaya* Seeds. *Glob J Med Res* 2020;**20**(February):19–38.
- [77] Elion, Itou R, Etou, Ossibi A, Epa C, Nsondé, Ntandou G, Bokia C, Ouamba J, et al. Anti-inflammatory and analgesic effects of leaves of *Chromolaena odorata* L. (King and Robinson). *African J Pharm Pharmacol* 2017;**11**(17):217–223.
- [78] Iragi GK, Imani B, Nfizi IB, Masumbuko CN, Gendusa PA, Matendo A, et al. Ethnomedicinal study of plants used in the Uvira Territory (Democratic Republic of Congo) Ethnomedicinal study of plants used in the Uvira Territory (Democratic Republic of Congo). *Forest Sci Technol* 2021;**17**(3):144–154.
- [79] Konda K, Kabakura M, Mbembe B, Mahuku K, Mafuta M, Mpoyi K, et al. *Plantes Médicinales de Traditions - Province de l'Equateur – R.D. Congo*. Institut d. Ministère de la Recherche Scientifique et Technologie: Kinshasa, RDC, 2015.
- [80] Bekkouch O, Zengin G, Harnafi M, Touiss I, Khoulati A, Saalaoui E, et al. Anti-

- Inflammatory Study and Phytochemical Characterization of Zingiber officinale Roscoe and Citrus limon L. Juices and Their Formulation. *ACS Omega* 2023;**8**(30):26715–26724.
- [81] Wu Y, Wu D, Li H, Huang H, Hu Y, Zhang Q, et al. Cyclohexylethanoid derivative and rearranged abietane diterpenoids with anti-inflammatory activities from *Clerodendrum bungei* and *C. inerme*. *Arab J Chem* 2024;**17**(1):105338.
- [82] Omoyiola OL, Ayomikun LT, Rafiu SD, Olufunso AB, Samuel OO, Abraham OK, et al. Anti-Inflammatory Property of *Costus afer* Ker Gawl Ethanol Leaf Extract in STZ-Induced Diabetic Rats. *Sch Int J Biochem* 2023;**6**(10):129–137.
- [83] Ibrahim B, Sowemimo A, Van Rooyen A, Van De Venter M. Antiinflammatory, analgesic and antioxidant activities of *Cyathula prostrata* (Linn.) Blume (Amaranthaceae). *J Ethnopharmacol* 2012;**141**(1):282–289.
- [84] Fotso AF, Longo F, Djomeni PDD, Kouam SF, Spiteller M, Dongmo AB, et al. Analgesic and antiinflammatory activities of the ethyl acetate fraction of *Bidens pilosa* (Asteraceae). *Inflammopharmacology* 2014;**22**(2):105–114.
- [85] Garcia R, Ferreira JP, Costa G, Santos T, Branco F, Caramona M, et al. Evaluation of anti-inflammatory and analgesic activities of *Cymbopogon citratus* in vivo-polyphenols contribution. *Res J Med Plant* 2015;**9**(1):1–13.
- [86] Mittas D, Mawunu M, Magliocca G, Lautenschläger T, Schwaiger S, Stuppner H, et al. Bioassay-Guided Isolation of Anti-Inflammatory Constituents of the Subaerial Parts of *Cyperus articulatus* (Cyperaceae). *Molecules* 2022;**27**(18):1–14.
- [87] Olajide OA, Alada ARA. Studies on the anti-inflammatory properties of *Entada abyssinica*. *Fitoterapia* 2001;**72**(5):492–496.
- [88] Uwaya D, Okakwu R, Omozuwa O. In-Vivo and In-Vitro Anti-Inflammatory Activities of the Aqueous Extract of Di-Herbal Formulation (*Euphorbia hirta* and *lactuca virosa*). *J Appl Sci Environ Manag* 2021;**24**(11):1979–1985.
- [89] Mikailu S, Obomate NL, Ugochukwu OP, Ekenna IC. Anti-Inflammatory, Fibrinolytic and Anti-Oxidant Activities of the N-Hexane Extract of *Ficus sur* Forssk (Moraceae) Leaves. *Haya Saudi J Life Sci* 2022;**7**(2):44–55.
- [90] Agbodjento E, Klotóé JR, Dramane G, Doughton TV, Ategbo JM. *Gardenia ternifolia* Schumach. & Thonn.: revue sur les aspects ethnobotanique, ethnopharmacologique, phytochimique et toxicologique. *Int J Biol Chem Sci* 2019;**12**(6):2922.
- [91] Larsen BHV, Soelberg J, Jäger AK. COX-1 inhibitory effect of medicinal plants of Ghana. *South African J Bot* 2015;**99**:129–131.
- [92] Mandjo BL, Ifulu JB, Dande PA. Etude ethnobotanique des plantes médicinales vendues aux marchés de Kinshasa (RD Congo). *Int J Innov Appl Stud* 2021;**34**(4):858–869.
- [93] Setiani LA, Rusli Z. Anti-inflammatory potential of african leaf water extract. *J Agric Appl Biol* 2020;**1**(2):46–53.
- [94] Shaikh S, Joshi YM, Kadam V. Comparative study of anti-inflammatory activity of aqueous and methanolic extracts of *Hibiscus cannabinus* leaf (Malvaceae). *Int J Pharm Pharm Sci* 2016;**8**(4):64–68.

- [95] Sofidiya MO, Odukoya OA, Adedapo AA, Mbagwu HOC, Afolayan AJ, Familoni OB. Investigation of the anti-inflammatory and antinociceptive activities of *Hymenocardia acida* Tul. (Hymenocardiaceae). *African J Biotechnol* 2010;**9**(49):8454–8459.
- [96] Ibekwe NN, Ibekwe NN, John-Africa LB. Anti-inflammatory Effect of the Alkaloid-rich Fraction of *Landolphia owarensis*. *Niger J Pharm Res* 2021;**17**(1):39–44.
- [97] Akinwumi K, Eleyowo O, Oladipo O. A Review on the Ethnobotanical Uses, Phytochemistry and Pharmacological Effect of *Luffa cylindrica*. In: Hany A El-Shemy, editor. *Natural drugs from plants*. London,UK, 2021: 83–93.
- [98] Onyegbule FA, Okoye CI, Chukwunwejim CR, Umeokoli BO, Eze PM. Evaluation of antioxidant, anti-inflammatory, and antimicrobial activities of the leaf extracts of *Luffa cylindrica*. *J Heal Sci* 2018;**8**(2):101–109.
- [99] Sravani D, Aarathi K, Sampath Kumar NS, Krupanidhi S, Vijaya Ramu D, Venkateswarlu TC. In vitro anti-inflammatory activity of *Mangifera indica* and *Manilkara zapota* leaf extract. *Res J Pharm Technol* 2015;**8**(11):1477–1480.
- [100] Sowemimo A, Samuel F, Fageyinbo MS. Anti-inflammatory activity of *Markhamia tomentosa* (Benth.) K. Schum. Ex Engl. ethanolic leaf extract. *J Ethnopharmacol* 2013;**149**(1):191–194.
- [101] Mohanta O, Ray A, Jena S, Sahoo A, Panda SS, Das PK, et al. Mesosphaerum suaveolens Essential Oil Attenuates Inflammatory Response and Oxidative Stress in LPS-Stimulated RAW 264.7 Macrophages by Regulating NF- κ B Signaling Pathway. *Molecules* 2023;**28**:5817.
- [102] Ishola IO, Ikumawoyi VO, Afolayan GO, Olorife OJ. Antinociceptive and anti-inflammatory properties of hydroethanolic seed extract of *Monodora myristica* (Annonaceae) in rodents. *West African J Pharm* 2016;**27**(1):22–32.
- [103] Mahdi HJ, Abdul N, Khan K, Mahmud R, Zaini M, Asmawi B. LC / MS , GC / MS screening and in vivo anti-inflammatory activity of Malaysian *Moringa oleifera* Lam leaf extracts and fractions against carrageenan -induced paw oedema in rats. *J Innov Pharm Biol Sci* 2017;**4**(3):48–54.
- [104] Khairinisa G, Mahargyani W, Agnia G. Anti-Inflammatory Activity of the Peel Extract of Ambon Bananas (*Musa Paradisiaca* L .) Examined With the Human Red Blood Cell Membrane Stabilization Method. *4th Int Semin Glob Heal* 2022;**22**:216–223.
- [105] Oluwole OG, Ologe O, Alabi A, Yusuf GT, Umukoro S. Anti-inflammatory effects and anti-oxidant capacity of *Myrathius arboreus* (Cecropiaceae) in experimental models. *J Basic Clin Physiol Pharmacol* 2017;**28**(6):521–529.
- [106] Okagu IU, Ndefo JC, Agbo MO. Trado-Medical Uses , Chemical Constituents and Biological Activities of *Newbouldia laevis* (Bignoniaceae): A Review. *Tabriz Univ Med Sci* 2022;**28**(1):51–75.
- [107] Olayemi O, John-Africa L, Abdullahi R, Isimi C. Effect of emulgel containing *Newbouldia laevis* stem bark extract on croton oil-induced hemorrhoids in mice. *J Res Pharm* 2023;**27**(4):1488.

- [108] Sahouo GB, Tonzibo Z, Boti B, Chopard C, Mahy J, N'guessan YT. Anti-Inflammatory and analgesic activities: chemical constituents of essential oils of *Ocimum gratissimum*, *Eucalyptus citriodora* and *Cymbopogon giganteus* inhibited lipoyxygenase 1-1 and cyclooxygenase of pghs. *Bull Chem Soc Ethiop* 2003;**17**(2):191–197.
- [109] Fakai IM, Abdulhamid A, Dada AY. Anti-Inflammatory and Analgesic Evaluation of *Parinari curatellifolia* Methanol Leaf Extract on Albino Rats. *Asian J Biochem Genet Mol Biol* 2021;**10**(1):9–18.
- [110] Dharmasiri PGNH, Ranasinghe P, Jayasooriya PT, Samarakoon KW. Antioxidant , Anti-Diabetic and Anti-Inflammatory Activities of *Passiflora foetida* grown in Sri Lanka. *Trop Agric Res* 2024;**35**(1):24–33.
- [111] Nnamani PO, Diovu EO, Peter IE, Onwuka AM, Ogbuanwu CV, Abonyi OE, et al. Anti-Inflammatory and Antinociceptive Activity of Herbal Lipospheres of *Pentaclethra macrophylla* (Fabaceae) Stem Bark Extract. *Processes* 2023;**11**:2557.
- [112] Chung J, Kong J, Choi H, Kong K. Antioxidant, anti-inflammatory, and anti-allergic activities of the sweet-tasting protein brazzein. *Food Chem* 2017;**267**:163–169.
- [113] Foe FMN, Flore T, Tchingang K, Nyegue AM, Abdou J, Joel A, et al. Chemical composition , in vitro antioxidant and anti-inflammatory properties of essential oils of four dietary and medicinal plants from Cameroon. *BMC Complement Altern Med* 2016;**16**:117.
- [114] Sigidi MT, Anokwuru CP, Zininga T, Tshisikhawe MP, Shonhai A, Ramaite IDI, et al. Comparative in vitro cytotoxic , anti- inflammatory and anti-microbiological activities of two indigenous Venda medicinal plants. *Transl Med Commun* 2016;**1**:1–7.
- [115] Mbayo MK, Kalonda EM, Muya RK, Tshisand PT, Kanangila AB, Maseho FM, et al. Test d'activité antimittotique et étude chimique préliminaire de quelques Euphorbiaceae du Katanga méridional (RDC). *Phytotherapie* 2018;**16**(2):74–86.
- [116] Harikrishnan H, Jantan I, Haque A, Kumolosasi E. Anti-inflammatory effects of *Phyllanthus amarus* Schum . & Thonn . through inhibition of NF- κ B , MAPK , and PI3K-Akt signaling pathways in LPS-induced human macrophages. *BMC Complement Altern Med* 2018;**18**:224.
- [117] Boakye YD, Agyare C, Ko W, George P, Poku P, Ossei S. Anti-in fl ammatory activity of aqueous leaf extract of *Phyllanthus muellerianus* (Kuntze) Exell . and its major constituent , geraniin. *J ofEthnopharmacology* 2016;**187**:17–27.
- [118] Nakalembe L, Kasolo JN, Nyatia E, Lubega A, Bbosa GS. Analgesic and Anti-Inflammatory Activity of Total Crude Leaf Extract of *Phytolacca dodecandra* in Wistar Albino Rats. *Neurosci Med* 2019;**10**:259–271.
- [119] Kumar D, Ganguly K, Hegde H V, Patil PA, Roy S, Kholkute SD. Activity of *Plumbago zeylanica* Linn . root and *Holoptelea integrifolia* Roxb . bark pastes in acute and chronic paw infl ammation in Wistar rats. *J Ayurveda Integr Med* 2014;**5**(1):33–37.
- [120] Souri MS, Oktavia S, Ifora I. Potential anti-inflammatory effects of *Psidium guajava* L.: A review. *Asian J Pharm Res Dev* 2021;**9**(2):47–52.
- [121] Mulaudzi RB, Ndhlala AR, Kulkarni MG, Finnie JF, Staden J Van. Anti-inflammatory and

- mutagenic evaluation of medicinal plants used by Venda people against venereal and related diseases. *J Ethnopharmacol* 2013;**146**(1):173–179.
- [122] Maneenoon K, Khuniad C, Teanuan Y, Saedan N, Prom-in S, Rukleng N, et al. Ethnomedicinal plants used by traditional healers in Phatthalung Province , Peninsular Thailand. *J Ethnobiol Ethnomed* 2015;**11**(43):1–20.
- [123] Oladeji OS, Adelowo FE, Oluyori AP, Bankole DT. Ethnobotanical Description and Biological Activities of *Senna alata*. *Evidence-based Complement Altern Med* 2020;**2020**:ID 2580259.
- [124] Mah SH, Teh SS, Ee GCL. Anti-inflammatory, anti-cholinergic and cytotoxic effects of *Sida rhombifolia*. *Pharm Biol* 2017;**55**(1):920–928.
- [125] Li H, Deng Z, Loewen S, Tsao R. Bioaccessibility , in vitro antioxidant activities and in vivo anti-inflammatory activities of a purple tomato (*Solanum lycopersicum* L .). *Food Chem* 2014;**159**:353–360.
- [126] Isa A, Dzoyem J, Adebayo S, Suleiman M, Eloff J. Nitric Oxide Inhibitory Activity Of *Strychnos Spinosa* (Loganiaceae) Leaf Extracts And Fractions. *African J Tradit Complement Altern Med* 2016;**13**(6):22–26.
- [127] Vostinaru O, Conea S, Mogosan C, Toma CC, Borza CC, Vlase L. Anti-inflammatory and antinociceptive effect of *Symphytum officinale* root. *Rom Biotechnol Lett* 2017;**2**(6):14160–7.
- [128] Maroyi A. *Syzygium cordatum* hochst. Ex krauss: An overview of its ethnobotany, phytochemistry and pharmacological properties. *Molecules* 2018;**23**(1084):1–18.
- [129] Muganga R, Bero J, Quetin-Leclercq J, Angenot L, Tits M, Mouithys-Mickalad A, et al. In vitro Antileishmanial, Antitrypanosomal, and Anti-inflammatory-like Activity of *terminalia mollis* root bark. *Planta Med* 2021;**87**(9):724–731.
- [130] Ghuman S, Ncube B, Finnie JF, McGaw LJ, Njoya EM, Cooposamy RM, et al. Antioxidant , anti-in fl ammatory and wound healing properties of medicinal plant extracts used to treat wounds and dermatological disorders. *South African J Bot* 2019;**126**:232–240.
- [131] Jigam AA, Abdulrazaq UT, Mahmud HA, Tijani FO, Jigam AA, Abdulrazaq UT, et al. Efficacy of *Thonningia Sanguinea* Vahl . (*Balanophoraceae*) Root Against *Plasmodium Berghei* , *Plasmodium Chabaudi* , Inflammation and Nociception in Mice. *J Appl Pharm Sci* 2012;**02**(01):47–51.
- [132] Nkeh-chungag BN, Temdie JR, Sewani-rusike C, Yolande M, Mbafor JT, Iputo JE. Analgesic , anti-inflammatory and antiulcer properties of the extract of *Uapaca guineensis* (*Euphorbiaceae*). *J Med Plants Res* 2009;**3**(9):635–640.
- [133] Maroyi A. *Zanha africana* (*Radlk .*) Exell: review of its botany , medicinal uses and biological activities. *J Pharm Sci Res* 2019;**11**(8):2980–2985.
- [134] Zammel N, Saeed M, Bouali N, Elkahoui S, Alam JM, Rebai T, et al. Antioxidant and Anti-Inflammatory Effects of *Zingiber officinale* roscoe and *Allium subhirsutum*: In Silico, Biochemical and Histological Study. *Foods* 2021;**10**:1383.
- [135] Bashige C V, Bakari AS, Ndjolo PO, Kahumba BJ, Duez P, Lumbu SJ. Ethnobotanical study

- of plants used as antimalarial in traditional medicine in Bagira in Eastern RD Congo. *J Pharmacogn Phytochem* 2020;**9**(4):1–14.
- [136] Pacifique MI, Jean de Dieu MM, Gentil IK, René M. Etude des Plantes Médicinales Utilisées par des Femmes Autochtones Pygmées (Batwa) Enceintes Dans l' hinterland du Parc National de Kahuzi-Biega (Rift albertin , RD . Congo). *Eur Sci J* 2020;**16**(27):107–132.
- [137] Ngbolua K-N, Innocent Z, Medard M, Colette MA, Ruphin DD, Honore Y, et al. Effect of Agricultural Residues Based-Compost on the Yield of *Amarantus hybridus* L. (Amaranthaceae) in Gbado-Lite City, Nord-Ubangi (Democratic Republic of the Congo). *Budapest Int Res Exact Sci J* 2019;**1**(4):53–61.
- [138] Latham P, Konda A. *Useful Plants of Kongo Central Province , Democratic Republic of Congo*. DFID: Blairgowrie, UK, 2017.
- [139] Vwakyankazi M, Petit P. Bunganga ya Mici: Guérisseurs et plantes médicinales à Lubumbashi. 2004;:125.
- [140] Ngbolua K, Mpiana PT, Mwanza FB, Tshibangu DST, Tshilanda DD, Ashande MC. Antisickling and antibacterial activities of *Garcinia punctata* Oliv . (Clusiaceae) and *Tetradenia riparia* (Hochst .) Codd (Lamiaceae) from Democratic Republic of the Congo. *J Adv Med Life Sci* 2016;**4**(1):1–8.
- [141] Kunyima PK, Muganza DM, Maloueki U, Mwabonkolo MM, Lami JN, Mbomba ANB, et al. Antimalarial efficacy and toxicity evaluation of 80 % ethanol extracts from the stem bark of *Enantia olivacea* , *Garcinia punctata* and *Massularia acuminata*. *Int J Biol Chem Sci* 2018;**12**(5):2093–2100.
- [142] Ibrahim B, Abdulkadir S. Effects of petroleum ether, ethanol and aqueous extracts of *Indigofera capitata* on oedema induced activity of *Naja nigricollis* venom in envenomed rats. *Sci World J* 2020;**15**(1):126–128.
- [143] Omara T, Nakiguli CK, Naiyl RA, Opondo FA, Otieno SB, Ndiege ML, et al. Medicinal plants used as snake venom antidotes in east african community: Review and assessment of scientific evidences. *J Med Chem Sci* 2021;**4**(2):107–144.
- [144] Kimpouni V, Koubouana F, Apani E, Motom M, Makita-Madzou JP. Contribution à l'inventaire et à l'utilisation des plantes à effets psychotropes et toxiques au Congo (Brazzaville). *Phytotherapie* 2012;**10**(1):19–24.
- [145] Mabhiza D, Chitemerere T, Mukanganyama S. Antibacterial Properties of Alkaloid Extracts from *Callistemon citrinus* and *Vernonia adoensis* against *Staphylococcus aureus* and *Pseudomonas aeruginosa* . *Int J Med Chem* 2016;**2016**:1–7.
- [146] Swamy TA, Obey J, Mutuku NC. Phytochemical analysis of *Vernonia adoensis* leaves and roots used as a traditional medicinal plant in Kenya. *Int J Pharm Biol Sci* 2013;**3**(3):46–52.
- [147] Thabit RAS, Cheng X-R, Al-Hajj N, Rahman RT, Le G-W, Le -Antioxidant G-W. Antioxidant and *Conyza bonariensis*: A Review. *Eur Acad Res* 2014;**2**(6):8454–8474.
- [148] Mbuyi KS, Kalunga MR, Kalonda M, Cimanga CC, Numbi WI, Kahumba BJ, et al. Aperçu ethnobotanique de plantes réputées antipaludéennes utilisées dans la ville de

Lubumbashi et ses environs , dans le Haut-Katanga en RD Congo. *Ethnopharmacologia* 2019;**61**:75–83.

- [149] Kpetehoto W, Johnson R, Houeto E, Hounkpatin A, Boni G, Mignanwande Z, et al. Ethnomedicinal study of *Ocimum gratissimum* Lin (Lamiaceae) in Cotonou. *J Complement Med Res* 2020;**11**(1):72.
- [150] Hashempur MH, Khademi F, Rahmanifard M, Zarshenas MM. An Evidence-Based Study on Medicinal Plants for Hemorrhoids in Medieval Persia. *J Evidence-Based Complement Altern Med* 2017;**22**(4):969–981.
- [151] Lawan A, Katsayal U, Yaro A. Anti-inflammatory and anti-nociceptive effects of the methanolic extract of the stem bark of *Ficus vallis-choudae* delile (Moraceae). *J Pharm Pharmacol* 2008;**2**(10):200–203.
- [152] Ahmed F, Mueen Ahmed K, Abedin M, Karim A. Traditional uses and pharmacological potential of *Ficus exasperata* vahl. *Syst Rev Pharm* 2012;**3**(1):15–23.
- [153] René YBGG, Bienvenu G, Fidèle AM, Huguette AB, Djidénu A, Fifamin AAJ, et al. Phytochemistry, metabolites quantification and antioxidant activity of *Calotropis procera* (Ait.) and *Ficus umbellata* (Vahl.), plants traditionally used against hemorrhoids in Benin. *Int J Curr Res Chem Pharm Sci* 2021;**8**(12):12–26.
- [154] Chatterjee R, Patil S, Sahu AN. Phytochemical Estimation And Anti- Hemorrhoidal Activity Of *Ficus Benghalensis* Linn . Prop Root Extract. *Res J Pharm Life Sci* 2020;**1**(1):10–21.
- [155] Novotna B, Polesny Z, Pinto-Basto MF, Van Damme P, Pudil P, Mazancova J, et al. Medicinal plants used by 'root doctors', local traditional healers in Bié province, Angola. *J Ethnopharmacol* 2020;**260**:112662.
- [156] Nyirenda J, Chipuwa M. An ethnobotanical study of herbs and medicinal plants used in Western, Copperbelt, Central and Northern provinces of Zambia. *Phytomedicine Plus* 2024;**4**(1):100514.
- [157] Chikowe I, Mnyenyembe M, Jere S, Mtewa A, Mponda J, Lampiao F.). An ethnomedicinal survey of indigenous knowledge on medicinal plants in the traditional authority Chikowi in Zomba, Malawi. *Curr Tradit Med* 2020;**6**(3):225–241.
- [158] Shopo B, Mapaya R, Maroyi A. Ethnobotanical study of medicinal plants traditionally used in Gokwe South District, Zimbabwe. *South African J Bot* 2022;**149**:29–48.
- [159] Siteo E, Wyk B Van. An inventory and analysis of the medicinal plants of Mozambique. *J Ethnopharmacol* 2024;**319**(P2):117137.
- [160] Okombe V, Simbi JL, Stévigny C, Vandenput S, Pongombo C, Duez P. Traditional plant-based remedies to control gastrointestinal disorders in livestock in the regions of Kamina and Kaniama (Katanga province , Democratic Republic of Congo). *J Ethnopharmacol* 2014;**153**(3):686–693.
- [161] Bashige C V, Many MH, Ntabaza N V, Numbi IE, Bakari AS, Kalonda ME, et al. Étude ethnobotanique, biologique et chimique de plantes réputées anticariogènes à Lubumbashi – RD Congo. *Phytothérapie* 2017;**15**(1):2–9.

- [162] Kalonda E, Mbayo M, Muhume S, Kasereka M, Mulamba J, Manya H, et al. Ethnopharmacological survey of plants used against malaria in Lubumbashi city (D.R. Congo). *J Adv Bot Zool* 2014;**1**(2):1–8.
- [163] Bashige C V, Alombong AG, Kamwimba MA, Bakari AS, Okusa NP. Ethnobotanical study of medicinal plants used in the treatment of sexual dysfunctions in traditional medicine in Kampemba-Lubumbashi, DR Congo. *World J Adv Res Rev* 2020;**7**(3):016–032.
- [164] Maloba MJ, Muyumba NW, Kasanya KJ, Mutombo SC, Mbayo KM, Ngoy KE, et al. Ethnobotanical survey and preliminary chemical screening of 14 plants reputed antigastitis in Lubumbashi and surroundings (DR Congo). *Int Eur Ext Enablement Sci Eng Manag* 2020;**8**(12):62–69.
- [165] Kalunga M, Mbuyi K, Ali K, Manya M, Numbi WI, Mbayo K, et al. Criblage chimique de quelques plantes utilisée contre la schistosomiase urogénitale dans le Haut-Katanga, RD Congo. *Phytochem BioSub J* 2019;**13**(2):86–101.
- [166] Mansi Gupta, J S Vaghela. Recent Advances in Pharmacological and Phytochemistry Studies on *Phyllanthus amarus*. *Pharm Biosci J* 2019;**7**(1):01–08.
- [167] Singh S, Strydom M, Balmith M, Megaw C, Nell M. The role of flavonoids in the pharmacological treatment of haemorrhoids. *South African Gen Pract* 2024;**5**(1):36–44.
- [168] Chiaretti M, Fegatelli DA, Pappalardo G, Venti MDS, Chiaretti AI. Comparison of Centella with Flavonoids for Treatment of Symptoms in Hemorrhoidal Disease and After Surgical Intervention: A Randomized Clinical Trial. *Sci Rep* 2020;**10**(1):1–14.
- [169] Shengo L, Mundongo T. A Survey of the Antibacterial Activity of Three Plants Used in the Congolese Herbal Medicine Practiced by the Healers in the City of Lubumbashi: Recent Advancement. In: Chaitanya M, editor. *Trends in Pharmaceutical Research and Development Vol. 1*. Book Publisher International, 2020: 81–91.
- [170] Abebe M, Asres K, Bekuretsion Y, Woldkidan S, Debebe E, Seyoum G. Teratogenic Effect of High Dose of *Syzygium guineense* (Myrtaceae) Leaves on Wistar Albino Rat Embryos and Fetuses. *Evidence-Based Complement Altern Med* 2021;**2021**:6677395.
- [171] Bashige C v, Kahumba BJ, Bakari AS, Mbuyi KS, Lumbu SJ. Étude ethnobotanique , phytochimique et évaluation de l ' activité antiplasmodiale de 13 plantes réputées antipaludéennes dans la commune du Kenya (Lubumbashi , RDC). *Phytothérapie* 2017;;1–10.
- [172] Sundaram SS, Suresh K, Sundaram SP. Indigenous knowledge on medicinal plants used to treat hemorrhoids in selected areas of Madurai district, Tamil Nadu, India. *J Med Plants Stud* 2019;**7**(4):24–27.
- [173] Gusain P, Uniyal D, Joga R. Conservation and sustainable use of medicinal plants. In: Elsevier, editor. *Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine*. Academic Press: Cambridge, Massachusetts, USA, 2020: 409–427.
- [174] Abbafati C, Abbas KM, Abbasi-Kangevari M, Abd-Allah F, Abdelalim A, Abdollahi M, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*

2020;**396**(10258):1204–1222.

- [175] Kaur N, Kaur N, Saggoo M. Conservation Strategies for Medicinal Plants in the Face of Environmental Challenges. In: Aftab T, editor. *Environmental Challenges and Medicinal Plants. Environmental Challenges and Solutions*. Springer, Cham: Gewerbestrasse, Switzerland, 2022: 461–485.
- [176] Bashige C V, Bakari AS, Manyah MH, Kahumba BJ, Okusa NP, Lumbu SJ. In vivo antiparasmodial and toxicological studies of *Dialium angolense* Welw. Ex Oliv. (Fabaceae) leaves extracts, a medicinal plant from Eastern Congo. *World J Biol Pharm Heal Sci* 2020;**04**(02):32–42.
- [177] Astana PRW, Nisa U, Triyono A, Ardiyanto D, Fitriani U, Zulkarnain Z, et al. Medicinal plants used by traditional healers for hemorrhoid treatment in Borneo island: Ethnopharmacological study RISTOJA. *IOP Conf Ser Earth Environ Sci* 2021;**913**(1):0–8.
- [178] Euloges KB, Clément MZ, Mâli K, Arsène H, Nogogna Z, Hervé H, et al. Connaissances , Attitudes et Pratiques des Tradipraticiens de Santé de Bobo Dioulasso à propos de la Maladie Hémorroïdaire Knowledge , attitude and practice of native doctors about hemorrhoidal disease at Bobo. *West African J Pharm* 2020;**21**.
- [179] Oyeyemi IT, Akinseye KM, Adebayo SS, Oyetunji MT, Oyeyemi OT. South African Journal of Botany Ethnobotanical survey of the plants used for the management of malaria in Ondo State , Nigeria. *South African J Bot* 2019;**124**:391–401.
- [180] Dhiman S, Nadda RK, Bhardwaj P. Medicinal herbs from Western Himalayas for hemorrhoids treatment: A review correlating traditional knowledge with modern therapeutics. *Pharmacol Res - Mod Chinese Med* 2023;**9**:100334.
- [181] Erbay MŞ, Sarı A. Plants used in traditional treatment against hemorrhoids in Turkey. *Marmara Pharm J* 2018;**22**(2):110–132.
- [182] Gagliano J, Anselmo-Moreira F, Sala-Carvalho W, Furlan C. What is known about the medicinal potential of bamboo? *Adv Tradit Med* 2022;**22**(3):467–495.
- [183] Shah F, Kim S. Therapeutic role of medicinal plant extracts and bioactive compounds in osteoarthritis. *Adv Tradit Med* 2022;**22**(4):837–844.
- [184] Mohammed Golam Rasul. Conventional Extraction Methods Use in Medicinal Plants, their Advantages and Disadvantages. *Int J Basic Sci Appl Comput* 2018;**2**(6):10–14.
- [185] El Maaiden E, Bouzroud S, Nasser B, Moustaid K, El Mouttaqi A, Ibourki M, et al. A Comparative Study between Conventional and Advanced Extraction Techniques: Pharmaceutical and Cosmetic Properties of Plant Extracts. *Molecules* 2022;**27**(7):2074.