

Applicability of entomopathogenic fungi and essential oils against the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

Mahmoud Fakeer

Gamal Hamam

Jin Joo

Khalid Hussein (✉ khussein@aun.edu.eg)

Research Article

Keywords: Fall armyworm, natural products, *Metarhizium anisopliae*, *Beauveria bassiana*, biocontrol

Posted Date: May 31st, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2959941/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) is a key polyphagous pest because of the damage it causes to maize crops. The entomopathogenic fungi *Metarhizium anisopliae* AUMC2605 and *Beauveria bassiana* AUMC3563 were evaluated to manage *S. frugiperda* in the lab and field. Additionally, the toxicity of essential oils from *Prunus amygdalus*, *Linum usitatissimum*, *Simmondsia chinensis*, and *Nigella sativa* were detected against larvae *S. frugiperda* in the field. Using the drenches technique, the tested *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 isolates were pathogenic to *S. frugiperda* and caused mortality ranging from 10.0 to 80.33%. The laboratory results indicated that *B. bassiana* AUMC3563 was more deadly than *M. anisopliae* AUMC2605. In the fifth instar larval stage *S. frugiperda* was not highly susceptible to the tested fungal isolates. However, the earlier instars larval of *S. frugiperda* were more susceptible to *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605. *B. bassiana* AUMC3563 caused the highest mortality of first, second, and third instar larvae at 5.6×10^7 conidia ml⁻¹ in the laboratory. The field trial results also indicated that the entomopathogenic fungus *B. bassiana* AUMC3563 was more effective than *M. anisopliae* AUMC2605. These results assert the potentiality of entomopathogenic fungi and natural products as effective tools in sustainable and integrated pest management.

Introduction

The fall armyworm, *Spodoptera frugiperda* is a serious insect pest that can infect more than one hundred-eighty six plant species belonging to forty-two families of different crops and cause much damages (Montezano et al., 2018). It is a native insect pest to tropical and subtropical crops such as maize, vegetable crops, rice, sorghum, and cotton in the American continent (Nagoshi et al. 2017; Otim et al. 2018; Padhee and Prasanna 2019). Recently, the fall armyworm was reported in West African countries such as Nigeria, Ghana, and Togo on the maize crop (Goergen et al., 2016; Abrahams et al., 2017). The larvae of *S. frugiperda* could cause damage ranged from 10 to 80% in the barley crops and 30 to 90% in the wheat crops (Resendiz et al., 2017; Yang, et al., 2020). The *S. frugiperda* larvae were also detected to feed on growing and young organs in the peanut crop causing 65 and 78% loss (He et al., 2020).

Pesticide application is not economical, leads to the evolution of insect resistance, dangerous to farmers, and harmful to natural enemies (Cai et al., 2017; Early et al., 2018). There is a pivotal need to develop alternative techniques to synthetic pesticides; these techniques should be environmentally friendly, safe, and costly-effective to reduce agricultural insect pests. Many pathogens, including viruses, bacteria, fungi, protozoa, and nematodes have been associated with the fall armyworm pest but only a few cause epizootics (Sharanabasappa et al., 2018; Ganiger et al., 2018). Despite *Entomophaga aulicae*, *Erynia radicans*, and *Nomuraea rileyi* fungi can cause high levels of mortality in the insect population, the disease appears too late to alleviate high levels of damage. The hyphomycetous entomopathogenic fungi e.g. *Beauveria bassiana*, are common soil dwellers (Ramos et al., 2020). Hence, they can be developed as biocontrol agents against the soil-inhabiting pests without risk to non-target insects (Dannon et al., 2020). *M. anisopliae* and *B. bassiana* are aerobic pathogenic fungi that attack insect

hosts. They have been extensively studied and widely applied as alternative eco-friendly pest management agents through the biological control of forestry and agricultural crop pests (Balla et al., 2021). These fungi not only infect insects at different life-cycle stages but also exert lasting sustained activity on the next generation (Kaczmarek and Boguś, 2021). Host invasion by entomopathogenic fungi occurs via germinating conidia that invade the insect's cuticle and reach the hemolymph (Mannino et al. 2019). Environmental factors, including relative humidity, temperature, nutrient composition, and light, can influence fungal pathogenicity (Mishra et al., 2015). Particularly, both humidity and temperature significantly affect the growth, germination, virulence, and survival of such mycopathogens (Sharma and Sharma, 2021). On the other hand, several lines of evidence suggested that natural products contribute to insect pests' management; however, the mechanisms of these insect-natural product interactions need more extensive investigations. There is also a strong tendency in the application of entomopathogenic fungi to gradually drive out chemical fertilizers (Gustianingtyas et al. 2020; Waqas et al. 2021).

Identifying potential natural products and deploying entomopathogenic fungi for insect pest management is one of the fundamental approaches for sustainable agriculture and adopting the next generation. While entomopathogenic fungi are widely suggested for pest biocontrol and sustainable agriculture, natural products e.g. essential oils also seem promising because of their rich content of bioactive compounds and low environmental impact to a greater extent (Fadda et al., 2022; Xu et al., 2019). Natural products of plant origin e.g. essential oils and entomopathogenic fungi such as *Metarhizium anisopliae* and *Beauveria bassiana* are likely candidates as biocontrol agents to manage multiple arthropod pests. The objective of this study was to detect their efficacy on the polyphagous pest *Spodoptera frugiperda* (Lepidoptera: Noctuidae) under laboratory and field conditions.

Materials and Methods

Insects and fungi

The eggs of fall armyworm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) were obtained from infested maize field in Assiut, Egypt (Fig. 1), and transferred to the laboratory in the Entomology Institute at Assiut, University. Eggs were placed in a Petri dish (15 cm × 0.9 cm) lined with filter paper and examined daily until hatching, the newly hatched larvae were provided with fresh maize leaves. Several pupae were kept in plastic cylindrical cages till adult emergence for a new colony. The newly appeared adults were transferred into adult cage. The ceiling of the cage was supplied with a sterile cotton ball saturated with 10% honey solution. The cotton ball was exchanged daily to guarantee identical egg ages. The fall armyworms colony was maintained in an acclimatized growth chamber at 25°C, 65% relative humidity, and photoperiod of 12 hours Day Light (Wanjiru and Sunday 2019). Two entomopathogenic isolates *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 were obtained from Assiut University Mycology Center (AUMC). The entomopathogenic fungi were refreshed on Difco™ Sabouraud dextrose agar (SDA) and incubated at 25°C for one week. The newly fungal hyphae were grown on malt extract agar (MEA) for two weeks at 25°C under static conditions. Conidia were scratched from the surface of the cultures and suspended in a sterile distilled H₂O containing 0.1% Tween 80 (Sigma Chemical Co. Ltd.,

UK). The spore suspension was mixed properly using a magnetic stirrer for 30 minutes. The conidial suspension was filtered through clinical gauze to detach the trapped hyphae. The numbers of spores in the stock conidial suspension were counted using the cell counting chamber (Marienfeld, Neubauer-improved, Germany). A serial dilution (5.0×10^7 to 1.0×10^7 conidia ml^{-1}) was prepared.

Insect fungi-exposure bioassay

The dipping technique was followed during this bioassay; ten larvae of the third instar per replicate were drenched in the conidial suspensions of *B. bassiana* AUMC3563 or *M. anisopliae* AUMC2605. After soaking in the conidial suspension, each treatment was placed in a Petri dish lined with wet sterile filter paper. A solution of 0.1% Tween 80 served as the un-treated (control). All Petri dishes were incubated at 25°C; fresh corn leaf pieces (15 cm) were served daily (Cordero et al., 2014). The larvae were examined daily for fifteen days, and the dead larvae were taken to another sterile moist chamber. The investigations were conducted in triplicate, average values were used, and the LC_{50} was calculated. Under field conditions, an experimental field of 10 x 10 m² cultivated with maize, in the Faculty of Agriculture, Asyut University, was divided and adopted for the field experiment. In this investigation, two concentrations of each entomopathogenic fungus were applied (5.6×10^7 and 9.25×10^7 conidia/ ml^{-1}) using spraying technique. The suspensions were sprayed onto corn seedling leaves with (1 ml per leaf) using a Precision Spray Tower (Potter Manufacturing, UK). One week later, the numbers of alive and dead were examined. The mortality value was detected according to the following formula:

$$\text{Mortality (\%)} = \frac{\text{Number of infected (dead)}}{\text{Total number of insects}} \times 100$$

Insect essential oil-exposure bioassay

The test was conducted under field conditions using four different essential oils (Flax oil Jojoba oil, Almond oil, and Nigella). These pure essential oils (100%) were products of The National Research Center in Dokki, Egypt. Under field conditions, a maize plant field (100 m²) was divided and adopted for this investigation in the experimental agriculture land of Assiut University, Egypt. The population of *S. frugiperda* was recorded from fifty randomly selected plants and sprayed with different concentrations (0%, 1%, 3%, and 5%) for each oil. The oils were sprayed using a 10-L precision tower sprayer (Potter Manufacturing, UK). To calculate the percentages of reduction in larvae of *S. frugiperda*, the numbers of larvae were recorded before and after the oil application according to Henderson and Tilton (1955) formula:

$$\text{Reduction \%} = 1 - \left(\frac{n \text{ in Co before treatment} \times n \text{ in T after treatment}}{n \text{ in Co after treatment} \times n \text{ in T before treatment}} \right) \times 100$$

Where; n = pest population, T = treated, Co = control

Statistical Analysis

The reduction percentages of natural oil and the entomopathogenic fungi in the field were calculated according to Henderson-Tilton's formula (Henderson and Tilton, 1955). Data were statistically analyzed using Statistical Package of the Social Sciences (SPSS) software windows (version 20.0). One-way analysis of variance (ANOVA) was done where appropriate. The results were considered statistically significant at $P \leq 0.05$.

Results

Mycosis of the fall armyworm larvae

In this study, *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 strains were evaluated, as a suspension drenches, against the *S. frugiperda* at different larvae stages (Table 1). Treatment efficacies under laboratory conditions were detected by assessing the mortality during development from the 1st instar through to pupae. Both of *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 were pathogenic to the larvae of *S. frugiperda* but overall mortality ranged between 10.0 and 80.0%, and 0.0% recorded for the treated and control experiments, respectively. In the treated experiment, larvae mortality after exposure to *B. bassiana* AUMC3563 increased from 10.66–80.33%. However, mortality by *M. anisopliae* AUMC2605 increased from 10–70.66% as the dose was increased from 1.05×10^7 conidia ml^{-1} to 5.6×10^7 conidia ml^{-1} . In the laboratory bioassay, the selected isolates evinced high virulence against the first and second instars of the *S. frugiperda* larvae. The isolate *B. bassiana* AUMC3563 realized 80% mortality at concentrations of 1.05×10^7 , 2.1×10^7 , 4.2×10^7 and 5.6×10^7 conidia ml^{-1} to the first and second instars of the insect (Fig. 2). Moreover, *B. bassiana* AUMC3563 at a concentration of 5.6×10^7 caused 80% mortality to the third instar. The entomopathogen strain *B. bassiana* AUMC3563 exhibited higher virulence against the fall armyworm than *M. anisopliae* AUMC2605. *B. bassiana* AUMC3563 isolate against *S. frugiperda* larvae caused mortality ranging from 10.0 to 80.33%. However, *M. anisopliae* AUMC2605 caused mortality ranging from 10.0 to 70.66% using the drenches technique. The first instar of *S. frugiperda* larvae showed the highest mortality for the two entomopathogenic fungi. However, the fifth instar of *S. frugiperda* larvae was markedly more resistant to the two entomopathogenic fungi.

Insecticidal indices of *B. bassiana* and *M. anisopliae* in the field

In this study, the virulence of the entomopathogenic fungi strains *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 was evaluated in the field. The investigation under field conditions was detected by assessing the mortality during insect development. At a concentration of 9.25×10^7 conidia/ml, *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 demonstrated mortality of 81.56% and 71.51%, respectively. The results seem to be less effective than those under laboratory conditions. However, at a concentration of 5.6×10^7 , the two applied entomopathogenic fungi achieved 66.23% and 53.57% mortality, respectively (Table 2). Either under the field or laboratory conditions *B. bassiana* AUMC3563 strain seems to be more virulent than the *M. anisopliae* AUMC2605 strain.

Insecticidal properties of essential oil

The used four essential oils exhibited significant insecticidal properties against the fall armyworm larvae. The essential oils at concentrations of 3% and 5% were very effective against *S. frugiperda* larvae (Table 3). The highest insecticidal activity was noticed at 3% v/v of the Jojoba oil (100% mortality), and the lowest insecticidal effect was detected at 1% v/v *Nigella sativa* oil (0% mortality). The insecticidal effect of the *Nigella sativa* oil achieved 100% mortality only after 48 h at 5% (v/v) of the oil. However, the same concentration showed 88.88% mortality after 24 h of application. Bitter almond oil at 5% (v/v) caused 87.5% mortality to the larvae of *S. frugiperda*, 80% mortality at 3% (v/v), and only 22.22% mortality at 1% (v/v). However, *Nigella* oil at 1% (v/v) didn't show any mortality. Jojoba oil at 3% (v/v) showed complete inhibition (100% mortality) to the *S. frugiperda* larvae in the field after 24 h, and only 63.63% mortality at 1% (v/v) on the second day. Jojoba oil at 3% (v/v) revealed 100% mortality against *S. frugiperda* larvae on the first day. Flax oil at very low concentration (1% (v/v)) could inhibit the *S. frugiperda* larvae in the corn field by mortality of 90%. Jojoba oil at 3% (v/v) revealed 100% mortality against *S. frugiperda* larvae on the first day and only 63.63% mortality at 1% (v/v) on the second day. The flax oil at 3% (v/v) revealed complete death (100% mortality) on the second day. The flax oil showed a relatively high insecticidal effect (88.88% mortality) at 1% (v/v) of the oil (Fig. 2). Both Jojoba and flax oils showed their highest insecticidal effect against the fall armyworms larvae (100% mortality) on the first day at 3% (v/v) and 5% (v/v) of the oil, respectively. However, 3% (v/v) of Flax oil showed 100% mortality in the next day. After one day application, the flax oil showed 88.88%, 90.9%, and 100% mortality at 1% (v/v), 3% (v/v), and 5% (v/v), respectively. However, the lowest insecticidal effect was exhibited by *Nigella* oil at 1% (v/v) (0% mortality) even after the second day treatment.

Discussion

The fall armyworm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae), is one of the most aggressive pests worldwide, and causes considerable losses in host crops. The fall armyworm comprises two strains, the rice strain, which feeds upon rice and other grass plants, and the maize strain, which refers to maize and sorghum (Abbas et al., 2022; Kumar et al., 2022). When fall armyworm attacks in huge numbers, it causes a long-term threat and damaging to many crops (Baudron et al., 2019; Makgoba et al., 2021; Mutyambai et al., 2022). The synthetic chemical insecticides are still the most effective and regularly utilized technique; however, these chemicals are risky to the ecosystem and have adverse

effects, e.g. the development of resistance in the insect pests, which increased the inevitability of search for safe and long-term, target-specific, and cost-effective alternatives. Several studies reported the resistance of the fall armyworm insect to most of the active synthetic insecticides; therefore biological control is considered an interesting option for insect pest management (Ahissou et al., 2021). Several species of the fall armyworm natural enemies were reported. Sisay et al. (2018) encountered 150 parasitoid species attack the fall armyworm. It is also attacked by diverse taxa of natural enemies (Harrison et al., 2019; Koffi et al., 2020) and entomopathogens such as viruses, bacteria, and fungi (Shylesha et al., 2018). The entomopathogenic fungi are thought to be an essential bio-control agent for this invasive pest. Recently, some entomopathogens were isolated from dead fall armyworm insect, in Africa and Asia, such as *Spodoptera frugiperda* Nuclear Polyhydrosis Virus, *Bacillus thuringiensis*, *B. bassiana*, and *M. anisopliae* (Chinwada, 2018; Shylesha et al., 2018; Sharanabasappa et al., 2019; Firake & Behere, 2020). Microbial insecticides neither induce resistance in their host insects nor deposit toxic residues in the environment (Qadri et al., 2020; Sato et al., 2021). The public awareness for healthful environment increased the attention on microbial bio-control agents (Paddock et al., 2021).

Mortality of the fall armyworm caused by mycopathogens

In the current investigation, the used entomopathogenic isolates showed high virulence against the early instars of the fall armyworm larvae. However, the fifth instar of the fall armyworm larvae showed the lowest sensitivity to the two entomopathogenic fungi. Idrees et al. (2021) evaluated the effect of five fungal isolates against immature instars of *S. frugiperda* larvae. The results showed significant effects on larvae mortality as *B. bassiana* ZK-5 reduced the feeding activity of first to third stage of *S. frugiperda* larvae. Sutanto et al. (2021) reported that the larvae hatched from treated eggs were decreased by the entomopathogenic fungi at 1×10^9 conidia ml^{-1} . The hatchability was significantly reduced by 26.84% in *B. bassiana* and 46.48% in *M. anisopliae*. Conidial concentrations of *M. anisopliae*, *B. bassiana*, and *Isaria fumosorosea* which ranged from 1×10^5 to 1×10^8 conidia ml^{-1} demonstrated virulence on the *S. litura* eggs with mortality of 48.2–71.6% (Afandhi et al., 2020). Among the five tested fungal strains, they found that *C. tenuissimum* SE-10, *B. bassiana* ZK5, and *P. citrinum* CTD24 showed significant effects on the insect mortality (Idrees et al., 2021). In this study, conidial concentrations of *B. bassiana* AUMC3563 and *M. anisopliae* AUMC2605 demonstrated virulence on the *S. frugiperda* larvae with mortality ranged from 10.0 to 80.0% under lab conditions.

B. bassiana AUMC3563 strain at conidial concentrations ranged from 1.05×10^7 to 5.6×10^7 conidia ml^{-1} achieved 80% mortality in the early instars of the fall armyworm larvae. Also, using dipping technique the fungal pathogen strain *M. anisopliae* AUMC2605 was effective on the fall armyworm with larvae mortality of 70.66%. Indeed, entomopathogenic fungi are common in maize fields and naturally involved in the suppression of several crop pests (Vega, 2018). After the outbreak of the fall armyworm, many larvae infected with mycopathogens were found in maize fields (Chinwada, 2018). *B. bassiana* is one of the most common used biological control agents globally (Dannon et al., 2020). Recently, *B. bassiana* demonstrated brilliant pathogenicity against eggs and early instar larvae of the fall armyworm (Gao et al., 2022). Some commercial products, based on *B. bassiana* or *M. anisopliae* are already available and in

use West Africa (Rwomushana et al., 2018). Ullah et al. (2022) demonstrated that *M. anisopliae* was more virulent to the fall armyworm larvae than *B. bassiana*. However, *B. bassiana* isolate was more lethal to the *Myzus persicae* nymphs than *M. anisopliae*. At concentration of 1.0×10^9 conidia ml⁻¹ *M. anisopliae* caused 88% mortality in *S. frugiperda* and 65% mortality in *M. persicae*. *B. bassiana* exhibited 76% mortality in *S. frugiperda* and 94% mortality in *M. persicae* at the same concentration. Rajula et al. (2021) isolated the entomopathogenic fungus *M. rileyi* during field survey of the fall armyworm *S. frugiperda*. This fungal species has demonstrated 95% mortality of the third larvae instar of the fall armyworm larvae. This finding could be evidence enough to start production of entomopathogenic fungus as a potential bio-pesticide and gain the farmers conviction for application to reduce the insect pest damage (Rajula et al., 2021). Recently, many insecticidal microbial strains have been isolated from soil and effectively used by limited number of farmers as bio-insecticides spray (Kumar et al., 2021). The use of the entomopathogenic fungi *B. bassiana* and *M. anisopliae* in the integrated pest management of the fall armyworm is promising because these fungi can establish endophytic habitat in maize plants (Ramos et al. 2020). Their colonization as endophytes in maize plant caused 100% mortality on the second instar larvae, while 75% and 87% mortality were detected on the forth instar larvae inoculated with *M. anisopliae* and *B. bassiana*, respectively (Ramos et al. 2020). These two entomopathogenic fungi used, caused lethal infection to the different life stages of *S. frugiperda*. In this present data, the fifth instar of *S. frugiperda* larvae showed the lowest susceptibility to the entomopathogenic fungi. The early larval instars particularly the first instar was the most susceptible to *B. bassiana* and *M. anisopliae* (Sutanto et al., 2021). The thick cuticle of pupae may serves as barrier to fungal invasion; therefore, this stage of insect life is seldom being infected by fungal pathogens (Elya et al., 2021). Thus, the virulence of a microbial pathogen not only depends on the insect species but also their developmental stage. Bio-control approaches based on *B. bassiana* and *Metarhizium anisopliae* don't just depend upon the interactions between pathogen and host but also on the ambient environmental conditions to which mycopathogens are exposed (Xu et al. 2016).

Botanical Insecticidal indices of in the field

Several plant species can produce a wide variety of secondary compounds that possess pesticidal properties and toxic to insect pests (Jaoko et al., 2020). Yarou et al. (2017) listed twenty plants that comprise insecticidal properties to control arthropod pests of vegetable crops. These bio-pesticides are expected to be more environmentally friendly with more diverse range of bioactive substances compared to synthetic chemical insecticides (Jaoko et al., 2020). Several studies reported that these botanicals were effective against fall armyworm (Siazemo et al., 2020; Kardinan et al., 2021; Rioba et al, 2020). Botanicals derived from plant extracts in Africa were effective in controlling fall armyworm with similar efficacy like chemical pesticides and also increased the production (Akeme et al., 2021). Sisay et al. 2019 reported seven potential plant extracts that were used in the control of the fall armyworm with mortality higher than 75%. Neem was probably the most effective among these pesticidal plants (Yarou et al., 2017). Moreover, the highest larval mortalities were detected with *Nicotiana tabacum* L. The aqueous extracts of *Cassia nigricans* reduced the fall armyworm infestation in maize by only 13% (Kambou &

Millogo, 2019). Botanical insecticides based on capsaicin, neem, orange oil, and other aromatic herbs are commercially available in several countries (Bateman et al., 2018). Some botanical pesticides caused high mortality under laboratory conditions. For example, neem extracts demonstrated 70% reduction in the fall armyworm (Matova et al., 2020). *Eucalyptus urograndis* was found to possess properties to protect maize from pests (Andrade et al., 2016). The *Carica papaya* seed powder was detected as an efficient bio-insecticide (Ogbonna et al., 2021). Neem oil at concentration of 0.17–0.33% reduced the fall armyworm damage in maize (Abbas et al., 2022). Botanical insecticides are non-hazardous, target-specific, and environmentally safe for natural enemies (Ahmed et al., 2021). In this investigation, Jojoba oil was probably the most effective among the pesticidal plant oils; the highest larval mortalities were detected at 3% v/v of the Jojoba oil (100% mortality). However, the lowest effective botanical insecticide was *Nigella sativa* oil. The botanical pesticide efficacy is most likely due to the secondary metabolites in the plant such as amides, isobutyl amides, natural lipophilic, and piperine which act as an anti-feedant, deterrent, and neurotoxin (Akeme et al., 2021). Presently, the pesticides market is dominated by the synthetic chemicals despite the development of various botanical pesticides and entomopathogens (Ngegba et al., 2022). Actually, alternative strategy to chemical insecticides still plays a little role in the sustainable agriculture and in integrated pest management. Extracts of neem plant biopesticides are not available on a large commercial scale due to the lack of financial support, materials, and appropriate equipment (Yarou et al., 2017). Our study provides important information of natural products for the field application of biological control of the fall armyworm. Nevertheless, more researches are needed to confirm the active property of these natural products as insecticides responsible for protection. Thus, the potential effective botanical insecticides should be communicated and made available for sustainable controlling of the invasive fall armyworm (Deguine et al. (2021). Furthermore, tests must be conducted in greenhouse and field experiments to assess the efficacy of these microbial and botanical pesticides, which can be detected by the reduction in damage and yield increase.

Conclusion

The fall armyworm *S. frugiperda* has become one of the most aggressive pests of maize and other economic crops in Africa and Asia. Among the alternatives to the chemical insecticides, bioinsecticides are an inevitable choice that should be explored. This investigation emphasizes the importance of the entomopathogenic fungi and natural products as biopesticides as biological control agents to reduce the damage of invasive insect pests and increase the productivity of plant crops. The first instar of *S. frugiperda* larvae was markedly more susceptible to the entomopathogenic fungi. Also, the results indicate that *B. bassiana* AUMC3563 strain was more virulent than *M. anisopliae* AUMC2605 strain. Jojoba oil extract achieved the highest reduction of the fall armyworm on maize plant. Thus, natural resources are important source for agro-ecological crop protection using plant biopesticides and biological control agents. This data could therefore be used to propose and produce new botanical insecticide that could also reduce the potential threat to the environment.

Declarations

Competing interests

The authors declare that they have no competing interests.

Author Contributions

Mahmoud Fakeer: Practetical work, Writing, Revising.

Gamal H. Hammam: Practetical work, Writing, Revising.

Jin Ho Joo: , Conceptualization, Visualization, Writing - Review & Editing.

Khalid Abdallah Hussein: Supervision, Conceptualization, Visualization, Writing - Review & Editing.

Funding

Funding is by the authors.

Availability of data and materials

All data and materials are available.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Abbas A, Ullah F, Hafeez M, Han X, Dara MZN, Gul H, Zhao CR (2022) Biological Control of Fall Armyworm, *Spodoptera frugiperda*. *Agronomy* 12:2704
2. Abrahams PT, Beale M, Cock N, Corniani R, Day J, Godwin S, Murphy G, Richards J (2017) Fall Armyworm: Status Impacts and control options in Africa. CABI Preliminary Evidence Note. (April 2017)
3. Afandhi A, Pratiwi VR, Hadi MS, Setiawan Y, Puspitarini RD (2020) Suitable Combination Between *Beauveria bassiana* (Balsamo) Vuillemin and Four Plant Leaf Extracts to Control *Spodoptera litura* (Fabricius). *AGRIVITA J Agric Sci* 42:341

4. Ahissou BR, Mathieu W, Bokonon-Ganta AH, Somda I, Verheggen F (2021) Integrated pest management options for the fall armyworm *Spodoptera frugiperda* in West Africa: Challenges and opportunities. A review. *BASE* 25:192–207
5. Ahmed N, Alam M, Saeed M, Ullah H, Iqbal T, Awadh Al-Mutairi K (2022) Botanical Insecticides Are a Non-Toxic Alternative to Conventional Pesticides in the Control of Insects and Pests [Internet]. *Global Decline of Insects*. IntechOpen; 2022. Available from: <http://dx.doi.org/10.5772/intechopen.100416>
6. Akeme C, Ngosong C, Sumbele S, Aslan A, Tening A, Krah C, Moses K, Justin O (2021) Different controlling methods of fall armyworm (*Spodoptera frugiperda*) in maize farms of small-scale producers in Cameroon. *IOP Conference Series: Earth and Environmental Science*. 911:012053
7. Andrade J, Sa V, Lodi S, Godoy B (2016) Age of *Eucalyptus urograndis* plantations and occurrence of pest insects. *Revista Árvore* 40:885–892
8. Balla A, Allaoua S, Hafsa C, Ali C, Warren K, Justyna AN, Tomasz O, Farida B, Lassaad B (2021) The Threat of Pests and Pathogens and the Potential for Biological Control in Forest Ecosystems. *Forests* 12:1579
9. Bateman ML, Luke B, Edgington S, Kuhlmann U, Cock JW (2018) Assessment of potential biopesticide options for managing fall armyworm (*Spodoptera frugiperda*) in Africa. *J Appl Entomol* 142:805–819
10. Baudron F, Abdou Zaman-Allah M, Chaipa I, Chari N, Chinwada P (2019) Understanding the factors influencing fall armyworm (*Spodoptera frugiperda* J.E. Smith) damage in African smallholder maize fields and quantifying its impact on yield. A case study in Eastern Zimbabwe. *Crop Prot* 120:141–150
11. Cai P, Gu X, Yao M, Zhang H, Huang J, Idress A, Ji Q, Chen J, Yang J (2017) The optimal age and radiation dose for *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae) eggs as hosts for mass-reared *Fopius arisanus* (Sonan) (Hymenoptera: Braconidae). *Biol Control* 108:89–97
12. Chinwada P (2018) Fall Armyworm Prevalence Assessment in Madagascar. In: *FAO Mission Report*. 32 pp
13. Cordero P, Príncipe A, Jofré E, Mori G, Fischer S (2014) Inhibition of the phytopathogenic fungus *Fusarium proliferatum* by volatile compounds produced by *Pseudomonas*. *Arch Microbiol* 196:803–809
14. Dannon HF, DANNON AE, Douro-Kpindou OK (2020) Toward the efficient use of *Beauveria bassiana* in integrated cotton insect pest management. *J Cotton Res* 3:24
15. Deguine JP, Aubertot JN, Flor RJ (2021) Integrated pest management: good intentions, hard realities. A review. *Agron Sustain Dev* 41:38
16. Early R, González-Moreno P, Murphy ST, Day R (2018) Forecasting the global extent of invasion of the cereal pest *Spodoptera frugiperda* the fall armyworm. *NeoBiota* 40:25–50
17. Elya C, De Fine HH (2021) The genus *Entomophthora*: bringing the insect destroyers into the twenty-first century. *IMA Fungus* 12:34

18. Fadda A, Daniele S, El Hassan S, Said G, Maurizio M, Silvia M, Nese SY, Asli CK, Celale KG, Massimo L, Ginevra LB, Zorita D, Alessandra D (2022) Innovative and Sustainable Technologies to Enhance the Oxidative Stability of Vegetable Oils. *Sustainability* 14:849
19. Firake DM, Behere GT (2020) Natural mortality of invasive fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) in maize agroecosystems of northeast India. *Biol Control* 148:1–11
20. Ganiger PC, Yeshwanath HM, Muralimohan K, Vinay N, Kumar ARV, Chandrashekara K (2018) Occurrence of the new invasive pest fall army worm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in the maize fields of Karnataka India. *Curr Sci* 115:621–623
21. Gao YP, Luo M, Wang XY, He XZ, Lu W, Zheng XL (2022) Pathogenicity of *Beauveria bassiana* PfBb and Immune Responses of a Non-Target Host, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Insects* 13:914
22. Goergen GPL, Kumar S, Sankung B, Abou T, Manuele T (2016) First Report of Outbreaks of the Fall Armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae) a New Alien Invasive Pest in West and Central Africa. *PLoS ONE* 11(10):e0165632
23. Gustianingtyas M, Herlinda S, Suwandi S, Hamidson H, Setiawan A, Verawaty M, Elfita A (2020) Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. *Biodiversitas* 21:1839–1849
24. Harrison RD, Thierfelder C, Baudron F, Midega C, Schaffner U, Van den Berg J (2019) Agro-ecological options for fall armyworm (*Spodoptera frugiperda* JE Smith) management: providing low-cost, smallholder friendly solutions to an invasive pest. *J Environ Manage* 243:318–330
25. He LM, Zhao SY, Wu KM (2020) Study on the damage of fall armyworm *Spodoptera frugiperda* to peanut. *Plant Prot* 46:28–33
26. Henderson CF, Tilton EW (1955) Tests with acaricides against brown wheat mite. *J Econ Entomol* 48:157–161
27. Idrees A, Qadir ZA, Akutse KS, Afzal A, Hussain M, Islam W, Waqas MS, Bamisile BS, Li J (2021) Effectiveness of Entomopathogenic Fungi on Immature Stages and Feeding Performance of Fall Armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Larvae. *Insects* 12:1044
28. Idrees A, Qadir ZA, Qasim M, Akutse KS, Afzal A, Hussain M, Islam W, Waqas MS, Bamisile BS, Li J (2021) Effectiveness of Entomopathogenic Fungi on Immature Stages and Feeding Performance of Fall Armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Larvae. *Insect* 12:1044
29. Jaoko V, Nji TT, Backx S, Mulatya J, Van den AJ, Magomere T, Olubayo F, Mangelinckx S, Werbrouck SPO, Smagghe G (2020) The Phytochemical Composition of *Melia volkensii* and Its Potential for Insect Pest Management. *Plants* 9:143
30. Kaczmarek A, Boguś MI (2021) Fungi of entomopathogenic potential in Chytridiomycota and Blastocladiomycota, and in fungal allies of the Oomycota and Microsporidia. *IMA Fungus* 12:29
31. Kambou G, Millogo AM (2019) Biological efficiency of natural substances aqueous extracts (*Cassia nigricans* Vahl., *Parkia biglobosa* (Jacq.) R. Br. Ex G. Donf., *Capsicum annum* L., *Cleome viscosa* L.)

- against *Spodoptera frugiperda* J.E. Smith on corn production and their effects on a fer. J Environ Sci Comput Sci Eng Technol 8:36–51
32. Kardinan A, Maris P (2021) Effect of botanical insecticides against Fall Armyworm *Spodoptera frugiperda* J. E. Smith (Lepidoptera: Noctuidae). IOP Conference Series: Earth and Environmental Science. 653:012060
 33. Koffi D, Kyerematen R, Eziah VY, Agboka K, Adom M, Goergen G, Meagher L (2020) Natural enemies of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in Ghana. Fla Entomol 103:85–90
 34. Kumar J, Ramlal A, Mallick D, Mishra V (2021) An Overview of Some Biopesticides and Their Importance in Plant Protection for Commercial Acceptance. Plants (Basel) 10:1185
 35. Kumar RM, Gadratagi BG, Paramesh V, Kumar P, Madivalar Y, Narayanappa N, Ullah F (2022) Sustainable Management of Invasive fall armyworm *Spodoptera frugiperda*, vol 12. Agronomy, p 2150
 36. Makgoba MC, Tshikhudo PP, Nnzeru LR, Makhado RA (2021) Impact of fall armyworm (*Spodoptera frugiperda*) (J.E. Smith) on small-scale maize farmers and its control strategies in the Limpopo province. South Afr Jamba 13:1016
 37. Mannino MC, Huarte-Bonnet C, Davyt-Colo B, Pedrini N (2019) Is the Insect Cuticle the only Entry Gate for Fungal Infection? Insights into Alternative Modes of Action of Entomopathogenic Fungi. J Fungi Basel 5:33
 38. Matova PM, Kamutando CN, Magorokosho C, Kutywayo D, Gutsa F, Labuschagne M (2020) Fall-armyworm invasion, control practices and resistance breeding in Sub-Saharan Africa. Crop Sci 60:2951–2970
 39. Mishra S, Kumar P, Malik A (2015) Effect of temperature and humidity on pathogenicity of native *Beauveria bassiana* isolate against *Musca domestica* L. J Parasit Dis 39:697–704
 40. Montezano DG, Specht A, Sosa-Gómez DR, Roque-Specht VF, Sous Silva JC, Paula-Moraes SV, Peterson JA, Hunt TE (2018) Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Americas Afr Entomol 26:286–301
 41. Mutyambai D, Munyao SN, Paul-André C, Sevgan S (2022) Agronomic Factors Influencing Fall Armyworm (*Spodoptera frugiperda*) Infestation and Damage and Its Co-Occurrence with Stemborers in Maize Cropping Systems in Kenya. Insects 13:266
 42. Nagoshi RN, Fleischer S, Meagher RL, Hay-roe M, Silvie P, Vergara C, Westbrook J, Khan A, Muru MG, Silvie P, Vergara C, Westbrook J (2017) Fall armyworm migration across the Lesser Antilles and the potential for genetic exchanges between North and South American populations. PLoS ONE 12:1–18
 43. Ngegba PM, Gaofeng C, Muhammad ZK, Guohua Z (2022) Use of Botanical Pesticides in Agriculture as an Alternative to Synthetic Pesticides. Agriculture 12:600
 44. Ogbonna C, Okonkwo N, Nwankwo E, Ezemuoka L, Anorue C, Irikannu K, Egbuche C (2021) Carica papaya seed oil extract in the management of insect pest of cabbage plant both in the laboratory and field. Int J Entomol Res 6:12–21

45. Otim MH, Tay WT, Walsh TK, Kanyesigye D, Adumo S, Abongosi J, Ochen S, Sserumaga J, Alibu S, Abalo G, Asea G, Agona A (2018) Detection of sister-species in invasive populations of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) from Uganda. PLoS ONE 13:1–18
46. Paddock KJ, Pereira AE, Finke DL, Ericsson AC, Hibbard BE, Shelby KS (2021) Host resistance to *Bacillus thuringiensis* is linked to altered bacterial community within a specialist insect herbivore. Mol Ecol 30:5438–5453
47. Padhee AK, Prasanna BM (2019) The emerging threat of fall armyworm in India. Indian Farm 69:51–54
48. Qadri M, Short S, Gast K, Hernandez J, Wong A (2020) Microbiome Innovation in Agriculture: Development of Microbial Based Tools for Insect Pest Management. Front Sustain Food Syst 4:547751
49. Rajula J, Pittarate S, Suwannarach N, Kumla J, Ptaszyńska A, Thungrabeab M, Mekchay S, Krutmuang P (2021) Evaluation of Native Entomopathogenic Fungi for the Control of Fall Armyworm (*Spodoptera frugiperda*) in Thailand: A Sustainable Way for Eco Friendly Agriculture. J Fungi 7:1073
50. Ramos Y, Taibo AD, Jiménez JA, Portal O (2020) Endophytic establishment of *Beauveria bassiana* and *Metarhizium anisopliae* in maize plants and its effect against *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) larvae. Egypt J Biol Pest Control 30:1–6
51. Resendiz RZ, Lopez JA, Estrada SB, Osorio E, Pecina JA, Mendoza C (2017) Efectos geneticos de la resistencia a *Spodoptera frugiperda* en lineas de maiz derivadas de germoplasma nativo de Tamaulipas. Revista Mexicana de Ciencias agrícolas 8:1329–1341
52. Rioba NB, Philip CS (2020) Opportunities and Scope for Botanical Extracts and Products for the Management of Fall Armyworm (*Spodoptera frugiperda*) for Smallholders in Africa. Plants 9:207
53. Sato Y, Jang S, Takeshita K (2021) Insecticide resistance by a host-symbiont reciprocal detoxification. Nat Commun 12:6432
54. Sharanabasappa et al (2019) Natural enemies of *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae), a recent invasive pest on maize in South India. Fla Entomol 102:619–623
55. Sharanabasappa CM, Asokan R, Mahadeva HM, Maruthi MS, Pavithra HB, Hegde K, Navi S, Prabhu ST, Goergen G (2018) First report of the fall armyworm *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) an alien invasive pest on maize in India. Pest Manage Hortic Ecsyst 24:23–29
56. Sharma R, Sharma P (2021) Fungal entomopathogens: a systematic review. Egypt J Biol Pest Control 31:57
57. Shylesha AN, Jalali SK, Gupta A, Varshney R, Enkatesant V, Shetty P, Ojha R (2018) Studies on new invasive pest *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) and its natural enemies. J Biol Control 32:1–7
58. Siazemo Martin, Simfukwe P (2020) An Evaluation of the Efficacy of Botanical Pesticides for Fall Armyworm Control in Maize Production. OALib 07:1–12

59. Sisay B, Simiyu J, Malusi P, Likhayo P, Mendesil E, Elibariki N, Wakgari M, Ayalew G, Tefera T (2018) First report of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) natural enemies from Africa. J Appl Entomol 142:800–804
60. Sutanto KD, Husain M, Rasool KG, Al-Qahtani WH, Aldawood AS (2021) Pathogenicity of local and exotic entomopathogenic fungi isolates against different life stages of red palm weevil (*Rhynchophorus ferrugineus*). PLoS ONE 16:e0255029
61. Ullah S, Raza ABM, Alkafafy M (2022) Isolation, identification and virulence of indigenous entomopathogenic fungal strains against the peach-potato aphid *Myzus persicae* Sulzer (Hemiptera: Aphididae) and the fall armyworm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). Egypt J Biol Pest Control 32:2
62. Vega F (2018) The use of fungal entomopathogens as endophytes in biological control: A review. Mycologia 110:4–30
63. Wanjiru J, Sunday K (2019) Ovicidal effects of entomopathogenic fungal isolates on the invasive fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae). J Appl Entomol 143:626–634
64. Waqas MS, Shi Z, Yi TC, Xiao R, Shoaib AAZ, Elabasy ASS, Jin DC (2021) Biology, ecology, and management of cotton mealybug *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae). Pest Manag Sci 77:5321–5333
65. Xu YJ, Luo F, Li B, Shang Y, Wang C (2016) Metabolic Conservation and Diversification of *Metarhizium* Species Correlate with Fungal Host-Specificity. Front Microbiol 7:2020
66. Xu P, Zhang D, Wang J, Wu K, Wang X, Wang X, Ren G (2019) The host preference of *Spodoptera frugiperda* on maize and tobacco. Plant Prot 45:61–64
67. Yang X, Sun X, Zhao S, Li J, Chi X, Jiang Y, Wu K (2020) Population occurrence spatial distribution and sampling technique of fall armyworm *Spodoptera frugiperda* in wheat fields. Plant Prot 46:10–16
68. Yarou B, komlan A, Tossou E, Mensah AC, Simon S, Verheggen FJ, Francis F (2017) Efficacy of basil-cabbage intercropping to control insect pests in Benin West Africa. Comm Appl Biol Sci 2:157–166

Tables

Table 1 to 3 are available in the Supplementary Files section.

Figures



Figure 1

Shows the damages of fall army-worm larvae to the leaves and ears of maize plant crop.




Figure 2

Shows the mortality (%) of the fall armyworm *S. frugiperda* caused by essential oils at different dilutions and the entomopathogenic fungi *B. bassiana* and *M. anisopliae* at conc. of 5.6×10^7 conidia/ml under the field conditions.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.docx](#)