



Evaluation of Different Dahlia (*Dahlia variabilis* L.) Cultivars for Tuber Yield

Khiromani Nag and S K Tamrakar

Department of Floriculture and Landscape Architecture,
College of Agriculture, IGKV, Raipur (Chhattisgarh) 492012

ABSTRACT

An experiment was conducted in the Department of Floriculture and Landscape Architecture, CoA, IGKV, Raipur from 2019-20 to 2020-21. The experiment was laid out in RBD with three replications and 20 cultivars with a view to find out the overall evaluation of different cultivars of dahlia. Result showed that the tuber yield parameters were observed *i.e.* number of tubers/plant (7.96), number of tuber/ plot (113.70), weight of single tuber (86.29 g), tuber yield/plant (86.95 g), tuber yield/plot (1.78 kg) and tuber yield (2.52 t/ ha) was observed in cv. T₄ (Blackout Black).

Key Words: Blackout Black, Dahlia, Kenya Yellow, Tuber, Yield.

INTRODUCTION

Dahlia (*Dahlia variabilis* L.) is one of the most popular flowers grown in many parts of the world for its beautiful ornamental blooms having varying shades of different colours, being useful as cut flowers besides for the beautification of gardens (Vikas *et al*, 2015). They are extensively used for garden display, home decoration, cut flowers of pompon and miniature types stay fresh in flower vases for many days and also make moderately good garlands (Gupta, 2015). Tubers of this plant contain significant amount of insulin and fructose and small quantities of medicinally active compound such as pythin and benzoic acid.

According to the National Horticulture Board, India's floriculture production area was 305 thousand ha in 2019-20, with a cut flower production of 762 thousand t. Floriculture is now commercially cultivated in several states like Andhra Pradesh (19.1%), Tamil Nadu (16.6%), Madhya Pradesh (11.99%), having gone ahead of other producing states like Karnataka, West Bengal, Mizoram, Gujrat, Orissa, Jharkhand, Haryana, Assam and Chhattisgarh. In India commercially grown major flowers such as Rose, Tuberose,

Gladiolus, Anthurium, Carnation and Marigold etc., but dahlia cultivation is limited only in hills and plains of Eastern India (Anonymous, 2019). In the Chhattisgarh state area under floriculture is 13,089 ha⁻¹ with the production of 2,29868 MT. approximately in the 2020-21. In Chhattisgarh, commercially grown major flowers like Marigold, Tuberose, Gladiolus, Roses, Gaillardia and Orchids but there is no exact scale production of dahlia is available (Anonymous, 2020).

As Dahlia is still called an uncut diamond for India, evaluation is the stepping stone in order to utilize any crop to its full potential. Since the performance of each genotype varies with region, season and growing environment, therefore testing the performance of the available genotypes for suitability and adaptability take prime importance, as the research works on this line is meagre in Chhattisgarh plains.

Dahlia is commercially propagated through rooted cutting which are prepared through tuber. Quality and disease-free tubers are required to fulfil the demand of rooted cuttings in Chhattisgarh state. The present investigation was carried out with the

Corresponding Author's Email: nagkhiromani@gmail.com

Evaluation of Different Dahlia (*Dahlia variabilis* L.) Cultivars for Tuber Yield

Table 1. Evaluation of Dahlia cultivars for tuber yield Characteristics (pooled data of three years). (pooled data of three years, 2019-20 to 2020-21)

Treatment	Treatment	Number of tubers/ plant	Number of tubers/ plot	Weight of tuber (g)	Tuber yield/ plant (g)	Tuber yield/plot (kg)	Tuber yield (t/ha ⁻¹)
T ₁	Gajanan Yellow	4.69	74.34	68.65	69.87	1.45	2.13
T ₂	Babananda Yellow	4.68	64.33	60.59	61.42	1.26	1.78
T ₃	Kenya Yellow	6.58	92.33	54.05	59.46	1.21	1.55
T ₄	Blackout Black	7.96	113.70	86.29	86.95	1.78	2.52
T ₅	Ankita Pink	4.08	99.47	57.53	57.85	1.18	1.53
T ₆	Pu Sinha Pink	4.22	89.00	63.13	63.38	1.29	1.90
T ₇	Raja New Blue	4.59	79.83	64.47	64.29	1.31	1.93
T ₈	Kenya Blue	7.45	71.03	66.47	67.36	1.38	2.03
T ₉	Kenya Original	6.92	75.43	74.67	75.29	1.54	2.25
T ₁₀	Anita Orange Red with Gold	3.45	58.33	51.57	51.72	1.07	1.36
T ₁₁	Jishu Red	6.15	93.64	79.13	79.46	1.63	2.46
T ₁₂	Sachin Red	6.85	82.66	78.71	78.61	1.60	2.43
T ₁₃	Red New	5.46	61.27	61.57	60.07	1.23	1.61
T ₁₄	Matangini White	4.38	95.44	62.44	62.70	1.28	1.86
T ₁₅	S. P. Romia Roy	4.09	78.61	56.92	66.83	1.37	2.00
T ₁₆	Suparna New Bicolor	4.26	71.53	74.89	71.82	1.48	2.18
T ₁₇	Prabhujee Bicolor	4.13	89.43	61.33	55.66	1.14	1.42
T ₁₈	Prime Minister Bicolor	6.51	69.34	66.45	69.30	1.42	2.10
T ₁₉	Raja Bicolor	6.14	98.53	66.66	61.45	1.26	1.71
T ₂₀	Glory of India Bicolor	5.98	98.02	57.69	65.24	1.33	1.95
SEm (±)		0.32	3.13	2.96	2.12	0.07	0.10
CD (P=0.05%)		0.92	8.97	8.50	6.09	0.21	0.29

objective to find out suitable genotypes/cultivars for quality planting materials with higher number of tuber yield under the evaluation of different Dahlia (*Dahlia variabilis* L.) cultivars for production of tuber yield under Chhattisgarh plains,

MATERIALS AND METHODS

The experiment was conducted at the Horticultural Research cum Instructional Farm, Department of Floriculture and Landscape

Architecture, CoA, IGKV, Raipur, during 2019-20 to 2020-21. The soil of the research field was clay-loam and was neutral in response with the pH 7.1. For studying the evaluation and association analysis of the experimental during the years all the twenty cultivars were grown in Randomized Block Design with three replications. Standard package of practices of dahlia cultivation were followed for growing a healthy crop. Observations were recorded on five randomly selected plants, from the border

Evaluation of Different Dahlia (*Dahlia variabilis* L.) Cultivars for Tuber Yield

plants were avoided. The experimental plots were irrigated at an interval of one week at initial stages and 10-12 days at later stages to maintain optimum moisture throughout the experimentation.

RESULTS AND DISCUSSION

Yield Characteristics

From the data (Table 1), cv. Blackout Black (T_4) resulted the significantly maximum number of tubers/plant *i.e.* 7.96, number of tubers/plot (113.70) which was noted *at par* with cv. Kenya Blue (T_8) in number of tuber/plant. Whereas minimum number of tubers/plant was observed in T_{10} (Anita Orange Red with Gold) *i.e.* 3.45, minimum number of tubers/plot (58.33). The tuber yield/plant may be directly related to production of more plant height, number of branches, there by synthesis of more photosynthesis and ultimately good tuber yield. The similar results observed in dahlia by Ahmed and Gul (2002) and Sree Devi (2020).

Among the different cultivars studied, the cultivar Blackout Black (T_4) recorded in maximum weight of tuber (86.29 g which was statistically *at par* with Jishu Red (T_{11}), Sachin Red (T_{12}). However, the weight of tuber (51.57 g) was minimum in cv. (T_{10}) Anita Orange Red with Gold. Variation in the bulb weight among the different dahlia cultivars under trial might due to the genetic different and local environmental condition soil structure and texture might have been responsible for the increase in weight of tubers. The variation in tuber weight may be due to the genotypic expression of the cultivars. This is in accordance with results of Gupta *et al* (2015), Manjula *et al* (2017), Bajaraya *et al* (2018), Singh and Singh (2018) in Dahlia.

The cultivars exhibited that the maximum tuber yield/plant (86.95 g) was registered in cultivar T_4 (Blackout Black). However, the minimum tuber yield/plant (51.72 g) was registered in cv. Anita Orange Red with Gold (T_{10}). Maximum tuber yield/plot was reported in cultivar T_4 (Blackout Black) 1.78 kg) which was statistically *at par* T_{11} (Jishu

Red) and T_{12} (Sachin Red) whereas, the minimum tuber yield/plot (1.07 kg) was observed in cultivar T_{10} (Anita Orange Red with Gold). The tuber yield (t/ha) of the experiment was noted that significantly highest tuber yield (2.52 t/ha) was observed in cultivar T_4 (Blackout Black) which was statically *at par* with cultivar T_9 (Kenya Original), T_{11} (Jishu Red) and T_{12} (Sachin Red). While the lowest tuber yield (t/ha) was found in cultivar T_{10} (Anita Orange Red with Gold) 1.36 t/ha.

Tuber yield/plant (g)/ plot(kg)/ (t/ha) was obtained in cultivars Blackout Black and least was obtained in Anita Orange Red with Gold. Tuber yield /plant (g)/ plot (kg)/ (t/ha) may be directly related to production of more number of branches, there by synthesis of more photosynthates and ultimately good tuber yield. The similar results observed in dahlia by Verma and Kulkarni (2017), Mahawer *et al* (2010) and Shukla *et al* (2018).

CONCLUSION

It may be concluded that cv. Red Blackout Black was most suitable for higher tuber yield with more number for production of higher tuber number as well as yield of tuber.

ACKNOWLEDGEMENT

Special thanks to the Dr. R. SadhuKhan, Professor in Genetics and Plant Breeding, BCKV, West Bengal for supporting with planting material.

REFERENCES

- Ahmed J and Gul S (2002). Evaluation of exotic cultivars of dahlia (*D. coccinea*) under Rawalakot conditions. *Asian J Plant Sci* 1(5): 565-566.
- Anonymous (2019). National Horticulture Board, Gurgaon. Database.
- Anonymous (2019). Directorate Horticulture and Farm Forestry, Chhattisgarh.
- Bajaraya B, Kanawjia A, Jaysawal N, Dubey A, Parveen S and Pawaiya (2018). Performance of different cultivars of Dahlia (*D. variabilis* L.) under agro-climatic conditions of Gwalior. *J Pharmacog Phytochem* 7(6): 98-102.

Evaluation of Different Dahlia (*Dahlia variabilis* L.) Cultivars for Tuber Yield

- Gupta A K, Jaiswal N K and Saravanan S. (2015). Varietal evaluation of different hybrids of dahlia (*D. variabilis* L.) under Allahabad agro-climatic conditions. *Int J Agric Sci Res* **5**(1) 55-58.
- Manjula B S, Nataraj S K, Hegde P P, Anitha G and Ayesha N (2017). Evaluation of Dahlia genotypes (*D. variabilis* L.) for growth, yield and quality traits under hill zone of Karnataka. *Environ & Ecol* **35** (4C): 3158—3161.
- Sree Devi M, Seetharamu G, Patil B C, Hanchinamani C N, Kukanoor L, Satish and Nishani S (2020). Assessment of genetic variability among dahlia (*D. variabilis* L.) genotypes for productivity and quality traits. *J Pharmacogn Phytochem* **9**(4): 3134-3137.
- Shukla P, Prasad V, Ainarkar A A (2018). Evaluation of dahlia hybrids (*D. variabilis* L.) under allahabad agro-climate conditions. *J Pharmacogn Phytochem* **7**(5): 1109-1113.
- Verma J and Kulkarni B S (2017). Evaluation of Dahlia genotypes for growth and yield characters under dry zone of Karnataka, India. *Int J Curr Microbiol App Sci* **6**(11): 402-409.
- Vikas D M, Patil V S and Dorajeerao A V D (2015). Evaluation of dahlia genotypes based on vegetative and quality characters. *Pl Archives* **15**(1): 283-286.

Received on 3/10/2022

Accepted on 31/3/2023