

Synergistic effect of foliar co-application of Bradyrhizobium japonicum and Nanoparticles on Growth, and Productivity of Soybean (Glycine max L.) in Sandy Soil Nano-Bradyrhizobium co-application Enhances Soybean in Sandy Soil

Abdou A. Soaud

Cairo University

Hend A. S. Ahmed

Cairo University

Heba B. A. Kandil

Abdel Wahab M. Mahmoud

Cairo University

Shimaa E Ibrahim Hassanien

shimaa@azhar.edu.eg

Al-Azhar University (Girls Branch), Yossuf Abbas st

Research Article

Keywords: Sandy soil, soybean, Nanoparticles, Rhizobium, Nutrient viability, Seed quality

Posted Date: June 2nd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-9581265/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Synergistic effect of foliar co-application of *Bradyrhizobium japonicum* and Nanoparticles on Growth, and Productivity of Soybean (*Glycine max* L.) in Sandy Soil

Nano-*Bradyrhizobium* co-application Enhances Soybean in Sandy Soil

Abdou A. Soaud^a, Hend A. S. Ahmed^a, Abdel Wahab M. Mahmoud^b, Heba B. A. Kandil^c, Shimaa El-Sayed Ibrahim Hassanien^{*}

^a Department of soil science, Faculty of Agriculture, Cairo University, 12613, Giza, Egypt. a2soaud@yahoo.com

^b Plant Physiology Division, Department of Agricultural Botany, Faculty of Agriculture, Cairo University, 12613, Giza, Egypt. Mohamed.mahmoud@agr.cu.edu.eg.

^c Agric. Microbiol. Res. Dep., Soils, Water and Environ. Res. Inst. (SWERI), Agric. Res. Center (ARC), Giza, Egypt. heba.belal@arc.sci.eg.

^{*} Botany and Microbiology Department, Faculty of Science, Al-Azhar University (Girls Branch), Yossuf Abbas st., P.O. 11754, Nasr City, Cairo, Egypt. shiamaa@azhar.edu.eg.

Abstract:

Many strategies, including the application of nanoparticles (NPs), that boost the growth and productivity of plant in sandy soil. The impacts of co-application of N, P, Zn, Fe nanoparticles (NPs) and *Bradyrhizobium japonicum* on soybean were investigated by pot experiment conducted in Faculty of Agriculture, Cairo University. The plants were exposed to 14 treatments (T1-T14) of 100ppm of N,P NPs and 40ppm of Fe, Zn NPs in exchange with *B. japonicum* inoculation and control (T0 non-treated plants). *B. japonicum* was grown at various concentration of NPs but the high growth rate was observed at 100ppm of N,P NPs and 40ppm of Fe, Zn NPs. And the optimum temperature and pH for its growth was (30-35) and (6.5-7), respectively. *B. japonicum* showed high nitrogenase activity, exopolysaccharides, IAA, (P, Fe, Zn mobilization) with 100ppm of N,P NPs and 40ppm of Fe, Zn NPs. The co-application of *B. japonicum* and nanomaterials were essential for boosting the viability of macro and micro elements in sandy soil, leaves and seed. The co-application of N, P, Zn, Fe NPs and *B. japonicum* (T7) showed an increase in Chlorophyll (3.82-9.54mg/g FW), total Carbohydrate (15.54- 20.67%) and Leaf area (115.41-122.74 cm²) in leaves. Furthermore, P, Zn, Fe NPs, and *B. japonicum* (T11) recorded the highest oil content(17.63%), where T7 increased protein content (32.43%) and the weight of 100 seed (8.07gm) compared to T0 t. The obtained results provide strong evidence that the foliar application of NPs improves the symbiotic performance between *B. japonicum* and the soybean plant.

Key words: Sandy soil, soybean, Nanoparticles, Rhizobium, Nutrient viability, Seed quality.

Acknowledgments: We would like to show our appreciation and express our deepest gratitude towards Cairo University, Faculty of Agriculture, Plant Physiology department, nanotechnology project for sample analyses and Department of Soil, as well Botany and Microbiology department, Faculty of science, Al-Azhar university for facilities supplies.

Consent for publication: No unlawful and defamatory materials.

Funding: No external fun

Conflicts of Interest: The authors declare no conflicts of interest.

Author Contributions: A. M. M., I. & Sh. E.I. H. – Conceptualization, data collection, methodology, data analysis ; Sh. E.I. H – manuscript writing and editing, resources; H. A. S. A.- methodology, resources, manuscript writing; H. B. A. K. – data collection; A. A. S.-

visualization, supervision, data analysis. All authors agreed for submission of manuscript to the journal.

1. Introduction

Soybeans (*Glycine max*) grow well in a variety of climates, temperate tropical, and subtropical regions. It is essential crop, has nutritional value and known as “poor man’s meat,” **Tafida et al.(2022)**. In Egypt, soybean farming has emerged as a practical agricultural solution; face the threats such as import reliance, food security and economic diversity. Soybean farming introduces to Egypt in the twentieth century and become a major crop, especially in the Valley and Nile Delta areas. It has several and important uses either as spanning animal feed, human food products or in industrial applications **FAO (2021)**. Egypt is regarded as the leading producer of soybeans in North Africa due to its climate, which includes ideal temperature ranges, exposure to sunlight, and irrigation techniques **Mahmoud et al. (2024)**. In the year 2022, Egypt's soybean yield was 2.92 t/ha. Globally, Egypt is arranged as the seventh-largest importer of soybeans coming after China, but in 2019 it became the highest importer reaching about 4257.30271 thousand tons for 1663.182 million USD **FAO (2021)**. Despite prosperity of soybean cultivation in Egypt, water scarcity, and limited availability of cultivated land and pest management are challenges facing soybean cultivation and productivity **Salama et al. (2023)**.

Sandy soil, is characterized by its large pores and low organic matter content, poses a particular challenge to crop productivity due to their quick water drainage and poor nutrient-holding capacities (**Ayman et al., 2020**) and (**Musei et al., 2024**). Sandy soils are widely distributed around the world, covering approximately 900 million hectares and in Egypt it represents more than 70% of total area **FAO (2021)**. In dry lands, where the sandy soils are widespread, the anticipated variability and change in climate are expected to aggravate food security problems for farmers who depend on these regions. Sandy lands reclamation through strategies that alleviate water and nutrient percolation has been developed for increasing crop productivity and assure food security (**Aoda et al., 2021**).

Nanotechnology is a recent technique for the enhancement of the agricultural sector because it introduces new ways to alleviate different soil stresses and boost crop yield (**Singh et al., 2023** and **Vardumyan et al., 2024**). Nano-fertilizer is an important part of modern agricultural techniques, enhancing productivity **Zahra et al. (2022)**, able to penetrate the

epidermis with great efficiency, regulate nutrient uptake, reduce wastes, and improvement of nutrients utilization **Ninama, et al.(2023)** and **Mahmoud et al. (2023)**. Nanotechnology is one of powerful strategy for improving the properties of sandy soil and protecting plants from its poverty of due to the very high surface area of nanomaterials, surface charges, and their morphological characteristics(**Ayman et al., 2020**) and (**Dawood, 2021**).(**Figure 1**)showed a graphical illustration the role of nanomaterials in improving plant growth in sandy soil.




Figure (1): showing a graphical illustration the role of nanomaterials in improving plant growth in sandy soil design by Biorender.com.

The over using and applying of chemical fertilizers in agriculture to enhance crop yields and soil productivity, it creates environmental problems, including reduced soil fertility and biodiversity **Khalid et al. (2023)** and (**Mahmoud and Swaefy 2020**). To overcome on the harmful impacts of synthetic fertilizers, different techniques are recently being used to boost plant growth and yield, such as fallowing and green manure cultivation (**Ma et al., 2021**), **Shang et al.(2021)** and promotion of biofertilizers (**Castro et al., 2020**) and (**Mahmoud et al. 2017**). Rhizobia, are Gram-negative bacteria have a positive impact on crop productivity and nutrient utilization such as boosting phosphorus and iron uptake, that stimulate crop growth, reduce and prevent the harm of pests and plant pathogens, and stimulate the growth of benefited microbes (**Ahemad and Kibret, 2014**). Enhancing nitrogen fixation by bacteria in soybean plant is essential to the sustainable boosting of agriculture by rhizobia biofertilization. The species of rhizobia isolated from soybean plants act as biofertilizers that enhancing N-fixation and decrease

emissions (Soumare *et al.*, 2020). Rhizobia produce plant growth regulating hormones like gibberellin that has essential role in the growth of nodules in legume plants (Nett *et al.*, 2022). Additionally, rhizobia are eco-friendly environment, bio-fertilization and has been reported to increase the productivity of most plants, like soybeans (Kolapo *et al.*, 2025). Investigating novel techniques for the cooperative interactions between microbes, nanomaterials, and plants may assist in reducing various stressors and promote plant development. This three-way relationship improves plant health and provides a viable method for guaranteeing agricultural approaches are sustainable (Campos *et al.*, 2023).

Upon previous introduction the current investigation was planned to determine the effects of different concentrations of N, P, Zn, Fe nanoparticles (NPs) on *Bradyrhizobium japonicum* growth and its plant growth regulating traits and studied the effect of foliar co-application of N, P, Zn, Fe nanoparticles (NPs) and *Bradyrhizobium japonicum* inoculum on soybean growth and the productivity of this important legume in sandy soil.

2. Material and Methods

2.1. Effect of the synthesized nanoparticles on rhizobium growth

To observe the effect of various nanomaterials on the growth of rhizobia, the strain of rhizobia (*Bradyrhizobium japonicum*) was obtained from biofertilizer production unit, Soils, Water and Environment Research Institute, Agriculture Research Center (ARC), Giza, Egypt. *Rhizobium* was cultivated in yeast mannitol broth medium (Vincent, 1970) only as control, and with N nanoparticles at concentrations 100, 150, 200 ppm, P NPs at concentrations 50, 100 and 150ppm and also with each Fe, Zn nanoparticles at concentrations 35, 40 and 50 ppm then incubated at 28°C for 72 hours under shaking conditions and the total viable count and the size of colony was determined by agar plate method.

2.2. Effect of different temperature on rhizobia growth in the presence of nanoparticles

Test tubes was contained with 5 ml yeast mannitol broth medium mixed with previous nanoparticles at its concentrations inoculated with 1 ml of rhizobia 4×10^8 cfu / mL and incubated at different degrees of temperature 15, 25, 30, 35, 40, 45 °C in triplicates for three days then the rhizobia growth was visualized.

2.3. Effect of different pH values on rhizobial growth in the presence of nanoparticles

Test tubes containing 5ml of yeast mannitol broth medium and the pH was adjusted as 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5 and 9.0 pH values and after their sterilization they were inoculated with *Bradyrhizobium japonicum* 4×10^8 cfu / mL culture with the previous nanoparticles with their concentrations in triplicates and incubated at 30°C for three days then the rhizobia growth was observed.

2.4. Detection of plant growth regulating (PGP) activities of rhizobium

The quantitative assay of IAA was carried out in broth YM medium as reported by **Brick et al. (1991)**. The YM broth medium was distributed in flasks 100ml capacity each contain 50ml supplemented with l-tryptophane (0.5%), then inoculated with 1 ml of rhizobia 4×10^8 cfu / mL and the flasks were treated with different concentration of four NPs and incubated for three days at 30°C for three days in shaking incubator at 120 r/min. After incubation, the culture medium undergo to centrifugation at 4000r/min for 20min and 1 ml of supernatant was mixed with 2ml of Salkowsky reagent (2% 0.5 M FeCl₃ in 35% perchloric acid) and incubated in dark conditions for 30 min. Pink or red color absorbance was measured at 530nm. The amount of IAA in supernatant was estimated using a standard curve of pure IAA. Rhizobium strain was screened to estimate EPS production according to **(Damery and Alexander, 1969)**. *Bradyrhizobium japonicum* allowed to grow on 50ml YM liquid medium with 1% glucose with various concentrations of N, P, Zn and Fe nanoparticles and inoculated with 1 ml of rhizobia 4×10^8 cfu / mL and incubated in rotary shaker at 200rpm for 3 days. The culture medium was centrifuged at 3500 rpm for 30 minutes after that the supernatant was combined with ow volume of acetone .Consequently, centrifugation was carried out at 3500rpm for 30, the crude of polysaccharides was washed with dis. Water and acetone alternately and filtered by filter paper and weighted after drying at 105° C overnight . A simple method for estimating nitrogenase activity by *Bradyrhizobium japonicum* through reduction of acetylene on agar plates with 2 to 2.5 mM glutamine as sole nitrogen is described by **Wilcockson and Werner (1978)**. The ammonium phosphate molybdate blue color method was used to quantitatively estimation of phosphate solubilization. The blue color absorbance was read at 660nm using KH₂PO₄ as standard and the amount of soluble phosphate were expressed as mg/l over control (**Jackson, 1973**).Quantitative estimation of zinc solubilizing potential of rhizobium on YM broth medium with 0.1% ZnO based on method described by **Francis et al. (1988)**. *Bradyrhizobium japonicum*

was grown on 5ml of YM broth medium supplemented with various concentrations of N, P, Zn and Fe nanoparticles and incubated for 72 hours at 30°C. After three days the culture medium was centrifuged 3000 rpm for 20 min then 0.5 mL of supernatant mixed with 0.5 mL of Chromo Azurol Sulfonate (CAS) solution and 10 μ L of sulfosalicylic acid solution to make final volume and kept at room temperature for 2hr, the absorbance was calculated at 630nm using UV-spectrophotometer **Bandemer and Shamble (1944)**. All experiments were carried out in triplicates with the same treatments. Tukey test ($p \leq 0.05$) was used to compare the treatment means using the high range statistical domain (HSD).

2.5. Synthesis of Nitrogen, Phosphorus, Iron and Zinc nanomaterials

The top to bottom molecular chemical technique was used to synthesis nanomaterials. From its precursor, Nitrogen, phosphorus, Iron and Zinc nanomaterials were prepared and synthesized (ammonium nitrate salt NH_4NO_3 , urea ($\text{CO}(\text{NH}_2)_2$), Calcium phosphate ($\text{Ca}(\text{H}_2\text{PO}_4)_2 \cdot \text{H}_2\text{O}$), magnetite (Fe_3O_4) and zinc chloride (ZnCl_2)) the used precursor and other reagents were bought from Sigma Chemical Company (Saint-Louis, Missouri, US).

2.5.1. Synthesis of Nitrogen, Phosphorus nanoparticles

Nitrogen and phosphorus nanomaterials (Figures 2 a and b) were prepared according to (**Mahmoud and Swaefy, 2020**) with modified method by using chitosan solution as a carrier to polymerize methacrylic acid, for 18 hours it was allowed to coated at room temperature in a buffer solution, in two-steps approaches. Firstly, for 5 hours we dissolved 0.23 g of chitosan in a methacrylic acid aqueous solution (0.5%, v/v) with magnetically stirring. Secondly, with stirring continuously and individually, 0.2 mmol of mix ammonium nitrate salt NH_4NO_3 and urea ($\text{CO}(\text{NH}_2)_2$) (1:1), and 0.4 mmol of Calcium phosphate ($\text{Ca}(\text{H}_2\text{PO}_4)_2 \cdot \text{H}_2\text{O}$), were added to the solution separately tile the solution get clear. Subsequently the polymerization was carried out at 45°C for 7 hours with magnetically stirring, this result in formation of a nanomaterial solution that was consequently exposed to pressure 1.5 psi for 72 hours, discontinuously 7 hours /day, and after that it was centrifuged at 1200 rpm for 35 minutes, and then cooled for two hrs. in an ice bath.

2.5.2. Synthesis of Zinc and Iron nanoparticles

Zinc nanomaterial (Figure 2 c) was synthesized by using an aqueous solution of zinc chloride (ZnCl_2). Solution of sodium hydroxide was slowly added drope by drope in a molar ratio of 1:2 with strongly mixing for 8 hours. The obtained precipitate was filtered and washed

thoroughly with deionized water, then stirred with water/toluene system by high speed stirrer, then rewashed with deionized water only for 2 hours, after that the precipitate was centrifuged for 40 minutes at 1000 rpm, and then allowed to cool in an ice bath for two hrs. The precipitate get dried at 100 °C by using an oven, subsequently exposed to pressure 1.5 psi for 72 hours discontinuously (7 hours / day) (**Daneshvar *et al.*, 2007**) and (**Abdel Wahab *et al.*,2020**). As for Nano iron (Figure 2 d) was synthesized with a little modified method from magnetite (Fe_3O_4) of Fe^{3+} and Fe^{2+} at a molar ratio of 3:2 by reduction–precipitation in aqueous ammonia (0.3 mol/L) and coated tetramethyl ammonium hydroxide under strong stirring for 2 hours by magnetically stirring, subsequently the precipitate was separated by using magnetic filtration with a permanent magnet, then it was washed with dis. water tile reached to a neutral pH, afterward centrifuged at 1400 rpm for 25 minutes according to **Qu *et al.* (1999)** and **Abdel Wahab and Sahar (2018)**.

The size and state of nano-particles were noticed straight by Transmission Electron Microscopy (TEM) utilizing an electron speed increase voltage of 60 kV. The TEM tests were ready by dropping a couple of drops of the arrangement on a carbon-covered copper framework, the work were finished in TEM labs (Histo. Unit) in Fac. of Agric. Cairo College Research Park (FA-CURP).




Figure (2): Showing Nano particles a) Nano-nitrogen, b) Nano-Phosphorus, c) Nano-Zinc, d) Nano-Iron.

2.6. Pot experiment

The current investigation was carried out at the greenhouse of Agricultural Research Center, Giza, Egypt within two summer growing seasons 2023 and 2024. In this experiment the Soybean cultivar was Giza 111, that obtained from Field Crops Research Institute, Agricultural Research Center, Giza, A.R. Egypt.

This experiment was designed to investigate the impact of four elements of nanofertilizers: nitrogen (NF), Phosphorus (PF), Iron (FeF) and Zinc (ZF), on morphology, yield components and chemical composition of soybean *Glycine max* (L.), plants were grown in sandy soil Table (1) obtained from Cairo-Alexandria Desert Road.

Table 1: The physical and chemical characteristics of sandy soil

Physical properties		Cations (meq / l)		Anion (meq / l)	
Sand (%)	90.5	Ca ⁺⁺	4.16	CO ₃ ⁼	0
Silt (%)	6.50	Mg ⁺⁺	1.52	HCO ₃ ⁻	0.54
Clay (%)	3.0	Na ⁺	3.24	Cl ⁻	3.31
Texture grade	Sandy	K ⁺	0.21	SO ₄ ⁼	5.28
pH (1:2.5)	7.8	Micronutrients (ppm)			
EC (dS/m)	1.37	Fe	2.4		
Total-N (%)	0.03	Zn	0.4		
Organic-C (%)	0.38	Cu	0.3		

2.6.1. Plant materials and experimental procedures

Seeds of soybean were sown on May 12, 2023, in plastic pots (40 cm in diameter), that filled with sandy soil after washed it three times with water and then mixed with compost Table (2) at a rate of 5 tons / fadden for all treatments.

Table 2. The physical and chemical analysis of the compost applied

Property	Value
Moisture content (%)	25
PH (1:5)	7.6
EC (1: 5 extract) dSm ⁻¹	3.2
Organic-C (%)	30.1
Organic matter (%)	51.7
Total-N (%)	1.4
Total-K (%)	1.27
C/N ratio	20:1
Total-P (%)	1.2
Fe (ppm)	1018
Zn (ppm)	279
Total content of Bacteria (cfu.g ⁻¹)	2.3 x 10 ⁷
Phosphate dissolving Bacteria (cfu.g ⁻¹)	2.6 x 10 ⁶
Weed seeds	0

The experiment design was coordinated in a randomized complete block design (RCBD) with triplicates. Each block had fifteen different nano-fertilizer treatments Table (3) in randomized

distribution. Each replicate had three pots; contains three holes for each pot, after the start of germination, thin out and leaved 2 plants in each hole.

Table 3: Treatments used in pot experiment

No.	Treatments	Concentration
1	T0 (Control)	(as a recommended)
2	T1 (NF)+rhizo.	100ppmN
3	T2 (NF & PF) +rhizo.	100ppmN +100ppmP
4	T3 (NF & ZnF) +rhizo.	(100ppmN)+(40ppmZn)
5	T4 (NF & FeF) +rhizo.	(100ppmN)+(40ppmFe)
6	T5 (NF & PF & ZnF) +rhizo.	(100ppmN)+(100ppmP)+(40ppmZn)
7	T6 (NF & PF & FeF) +rhizo.	(100ppmN)+(100ppmP)+(40ppmFe)
8	T7 (NF & PF & ZnF&FeF) +rhizo.	(100ppmN)+(100ppmP)+(40ppmZn)+(40ppmFe)
9	T8 (PF) +rhizo.	(100ppmP)
10	T9 (PF & ZnF) +rhizo.	(100ppmP)+(40ppmZn)
11	T10 (PF & FeF) +rhizo.	(100ppmP)+(40ppmFe)
12	T11 (PF & ZnF&FeF) +rhizo.	(100ppmP)+(40ppmZn)+(40ppmFe)
13	T12 (ZnF) +rhizo.	(40ppmZn)
14	T13 (ZnF&FeF) +rhizo.	(40ppmZn)+(40ppmFe)
15	T14 (FeF) +rhizo.	(40ppmFe)

NF: nitrogen as nanofertilizers, PF: Phosphorus as nanofertilizers, FeF: Iron as nanofertilizers and ZF: Zinc as nanofertilizers, rhizo.: rhizobium.

2.6.2. Recommended additions

For the all treatments phosphorus and potassium fertilizers were applied as a basal dressing. Each pot delivered the following basic treatments: 5.6 g/ pot (225 kg/ fadden), two weeks before sowing calcium super phosphate (15.5% P_2O_5) was added, then two weeks after sowing 1.5 g/ pot (60 kg/ fadden) potassium sulphate (48% K_2O) was added and repeated after one month later, mineral nitrogenous fertilizer (MN): Ammonium sulphate (20.6%N) was added at (20 units of nitrogen / fadden) It was added at the time of planting, and after 30 days, 20 units of nitrogen were added,

2.7. Statistical analysis

The obtained results represented average of two seasons was undergrown to proper analysis of variance as revealed by **Snedecor and Cochran (1980)**. At 5% the least significant differences (L.S.D.) were calculated for each tested character under various assigned treatments.

3. Results

3.1. Effect of the synthesized nanoparticles on rhizobium growth.

The viable total count of rhizobia showed different growth response to different concentrations of nanoparticles and also different range of colony size Table (4), rhizobia recorded high growth rate and large colony size with N, P NPs and Fe, Zn NPs at concentration 100 ppm, 40ppm respectively while N, P NPs showing low growth rate at concentration 200 and 150 ppm, respectively.

Table (4): Effect of the synthesized nanoparticles on *B. japonicum* growth.

Parameters	<i>B. japonicum</i> &N conc. (ppm)			<i>B. japonicum</i> &P conc. (ppm)			<i>B. japonicum</i> &Fe conc. (ppm)			<i>B. japonicum</i> &Zn conc. (ppm)			<i>B. japonicum</i>
	100	150	200	50	100	150	35	40	50	35	40	50	-
Colonies number	8-10	3-5	2	2-3	7-8	1-2	4-5	6-7	1-2	2-3	4-6	1-2	5-6
Colonies diameter mm	2-3	1-1.5	0.8	1.5-2	3-4	0.8-0.7	1.5-2	2-3	0.6-0.7	1-1.5	3-3.5	0.8-0.9	1.5-2
Growth rate	+++	++	+	+	+++	+	++	++	+	+	++	+	++

Conc.: concentration +++: strong growth

++: moderate growth

+: poor growth

The current results showed that rhizobia strain was thermotolerant, it grow well at 30°C as well as 37°C with 100ppm concentration of N and P NPs and 40ppm concentration of Fe and Zn NPs, it also able to grow below and above this degrees Table (5), moreover rhizobia strain it can grow at wide range of pH it recorded high growth rate under slightly acidic condition at concentration 100and 150ppm, 50and 100ppm, 35 and 40ppm for N, P, Fe and Zn, respectively Table (6).

Table (5): Effect of different temperature on *B. japonicum* growth in the presence of nanoparticles.

Measurement Temperature C°	N conc. (ppm)			P conc. (ppm)			Fe conc. (ppm)			Zn conc. (ppm)		
	100	150	200	50	100	150	35	40	50	35	40	50
15	+	+	-	+	+	-	+	+	-	+	+	-
25	+	+	-	+	+	-	+	+	-	+	+	-
30	+++	+	+	+	+++	+	+	++	+	+	++	+
35	++	+	-	+	++	-	+	++	-	+	++	-
40	+	+	-	+	+	-	+	+	-	+	+	-
45	+	+	-	-	+	-	-	-	-	-	-	-

Conc.: concentration +++: strong growth ++: moderate growth +: poor growth- : no growth

Table (6): Effect of different pH values on *B. japonicum* growth in the presence of nanoparticles.

Measurement pH	N conc. (ppm)			P conc. (ppm)			Fe conc. (ppm)			Zn conc. (ppm)		
	100	150	200	50	100	150	35	40	50	35	40	50
5.5	-	-	-	-	+	-	+	+	-	+	+	-
6	+	+	-	+	+	-	++	++	-	++	++	-
6.5	+++	+	+	++	+++	+	++	++	+	++	++	+
7	++	++	-	++	++	-	++	++	-	++	++	-
8	+	+	-	+	+	-	+	+	-	+	+	-
8.5	+	+	-	+	+	-	+	+	-	+	+	-
9	+	+	-	-	-	-	-	-	-	-	+	-

3.2. Plant growth promoting (PGP) traits of *B. japonicum*.

The results showed varying levels of PGP traits were reproduced by *B. japonicum* at different concentrations of NPs. The quantitative assaying of IAA in broth YM culture supplemented with (0.5%) L- Tryptophan, showed variation in the quantity of IAA produced by *B. japonicum*. The highest quantity of IAA (12.01 µg/ml) was recorded at 40 ppm of Zn NPs, followed by (11.26 µg/ml) for 100 ppm of N NPs, where produced low concentration of IAA 1.886 µg/ml at 50 ppm of Fe (Figure 3a).

As shown in (Figure 3b), the maximum amount of EPS production was 31.5 µg ml⁻¹ followed by 27.5 µg ml⁻¹ observed at 100 ppm of N and P NPs respectively, on the other hand the amount of EPS by *B. japonicum* individually was 20.8 µg ml⁻¹, on the other hand the lowest concentration of EPS obtained at 50 ppm of Fe 5.9 µg ml⁻¹.




Figure (3): Plant growth promoting (PGP) traits of *B. japonicum*, a) IAA Production b) exopolysaccharides (EPS) production at different concentrations of NPs.

Table (7) showing that the effect of various concentrations of NPs on nitrogenase activity and minerals solubilization by *B. japonicum*. The nitrogenase activity was more efficient at low concentration of N NP 6.49 ($\mu\text{mol C}_2\text{H}_4 \text{ h}^{-1}$), whereas *B. japonicum* only recorded 4.02 ($\mu\text{mol C}_2\text{H}_4 \text{ h}^{-1}$). The highest phosphate solubilization by *Bradyrhizobium japonicum* was 22.64ppm recorded at 100ppm P NPs while the rhizobium strain only recorded 15.02, furthermore 40ppm conc. of Fe and Zn recorded high phosphate solubilization rather than the other two concentrations 35 and 50ppm of them. Additionally, the high concentration of nano-Nitrogen resulted low P solubilization 5.77ppm. The efficiency of zinc solubilization by *B. japonicum* was found to vary relying on the concentration of nanomaterials, the highest zinc solubilization was 0.73ppm determined at 40ppm Zn NPs, while the lowest Zn solubilization was 0.08ppm at 150ppm P NPs. The *Bradyrhizobium* strain treated with 100 ppm of Fe NPs showed high Fe solubilization 0.93ppm compared to *Bradyrhizobium* only where it recorded 0.44ppm while 150ppm of P NPs observed the lowest Fe solubilization 0.11ppm.

Table (7): Showing that the effect of different concentrations of NPs on minerals solubilization by *B. japonicum*.

Treatments	Concentrations	Mean of Nitrogenase activity ($\mu\text{mol C}_2\text{H}_4 \text{ h}^{-1}$)	Mean of P conc.(ppm)	Mean of Zn conc.(ppm)	Mean of Fe conc.(ppm)
<i>B. japonicum</i> &N conc. (ppm)	100	6.49a	17.81b	0.55cd	0.79bc
	150	4.07e	13.92g	0.31g	0.42e
	200	1.86i	5.77m	0.1h	0.13h
<i>B. japonicum</i> &P conc. (ppm)	50	3.55f	14.29f	0.32g	0.23g
	100	5.94c	22.64a	0.51de	0.63d
	150	2.03i	10.07j	0.08h	0.11h
<i>B.japonicum</i> &Fe conc. (ppm)	35	2.75h	11.73h	0.49e	0.75c
	40	4.97d	15.88d	0.61b	0.93a
	50	1.51j	8.86k	0.1h	0.42e
<i>B.japonicum</i> & Zn conc. (ppm)	35	3.29g	10.92i	0.57bc	0.64d
	40	6.21b	16.19c	0.73a	0.81b
	50	1.20k	7.1l	0.28g	0.33f
<i>B. japonicum</i>	-	4.02e	15.02e	0.38f	0.44e
	LSD	0.194612	0.209294	0.041982	0.044005

3.3. Effect of co-application of N, P (100ppm) NPs and of Fe, Zn (40ppm) NPs and *B. japonicum* on soil's minerals concentration.

As shown in Table (8), the co-applying *B. japonicum* and NPs in sandy soil improved significantly the soil fertility, physical, chemical, and biological characteristics of soil. The data obviously show that the co-applying *B. japonicum* and NPs significantly enhanced the soil's minerals concentration (N, K, Fe and Zn) compared to the control treatment. N, K, Fe and Zn content in soil showed significant increase with T7 47.69, 187.60, 13.27 and 6.49ppm, respectively compared to T0 40.17, 180.65, 8.12 and 4.26ppm, respectively, where T1 showed high significant in P content 9.85ppm whereas T7 come in the second order 9.38ppm. The increase of N, P, K, Fe and Zn contents in soil associated with N fixation and P, Zn, Fe mobilization in the rhizosphere by *B. japonicum* may be supplied by NPs. The viability of minerals is maybe due to the exopolysaccharides produced by *B. japonicum* that improve the texture of soil, water and nutrients retention by sandy soil.

Table (8): Effect of foliar co-application of N, P Fe, Zn NPs and *B. japonicum* on soil's minerals concentration. (Results represented the average of two seasons).

Treatments	N ppm	P ppm	k ppm	Fe ppm	Zn ppm
T0	40.17b	6.42ba	180.65a	8.12b	4.26a
T1	32.30d	3.39c	168.38c	7.63b	2.39b
T2	37.79c	7.72a	170.58b	7.90b	2.77b
T3	38.29cb	5.33b	175.93b	7.95b	4.52a
T4	38.97cb	4.94b	177.14b	8.12b	3.13b
T5	40.78b	8.43a	179.38a	7.33b	4.79a
T6	42.86b	8.51a	181.61a	11.47a	4.33a
T7	47.69a	9.38a	187.60a	13.27a	6.49a
T8	26.20e	7.61a	166.23c	7.28b	2.40b
T9	27.47e	8.22a	168.56c	7.10b	3.53b
T10	26.42e	8.61a	169.86c	10.20a	3.65b
T11	39.90b	9.85a	160.74d	12.16a	6.24a
T12	25.20e	5.41b	165.83d	7.18b	4.62a
T13	27.71e	6.83ba	173.57b	10.43a	5.05a
T14	24.55e	5.44b	161.20d	11.85a	2.13b

Results represented the average of two seasons. The values of mean followed by different letters verify significantly by LSD at $p < 0.05$.

3.4. Nutritional status of the leaves

Table (9) displays the impact of Foliar spraying of both nanoparticles and rhizobium on the leaves chemical composition. With NPs and *B. japonicum* treatments, the treated soybean leaves recorded a significant increase in the mounts of all nutritional elements (N, P, K, Fe, Zn and Mn) of soybean leaves compared to control plants. The uppermost increase in nutritional status of the leaves N, K and Zn 6.26, 1.67 and 50.36%, respectively were observed in the plants exposed to T7 , where T1 was less significant in N and Zn 3.79 and 22.7%, respectively, additionally T12 resulted in low K percent 1.33% compared to control . Likewise T11 showed significant increase in P, Fe, and Mn 0.52,70.79 and 23.20 respectively.

Table (9): Effect of foliar co-application of N, P Fe, Zn NPs and *B. japonicum* on leaves's minerals concentration. (Results represented the average of two seasons).

Treatments	N %	P %	k %	Fe %	Zn %	Mn%
T0	4.06b	0.32c	1.34d	40.21d	25.06c	15.12c
T1	3.79c	0.32c	1.38d	41.24d	22.70d	18.07a

T2	3.89c	0.35c	1.48c	47.33c	24.72cd	18.13a
T3	4.69ab	0.37c	1.52b	46.25c	44.79b	20.16a
T4	4.67ab	0.44b	1.50b	52.69b	26.84c	19.18a
T5	4.59b	0.42b	1.55b	48.51c	46.98a	20.93a
T6	5.61a	0.45b	1.53b	53.10b	27.45c	21.33a
T7	6.26a	0.50a	1.67a	68.35a	50.36a	22.18a
T8	4.41b	0.42b	1.39d	40.08d	26.19c	18.07a
T9	5.16a	0.40b	1.43c	43.12d	28.74c	19.15a
T10	5.11a	0.48a	1.42c	55.28b	30.86c	17.67ab
T11	6.16a	0.52a	1.55b	70.19a	49.81a	23.20a
T12	4.65ab	0.42b	1.33d	41.64d	45.29a	16.92b
T13	6.08a	0.44b	1.37d	54.20b	48.29a	19.05a
T14	4.16b	0.40b	1.37d	52.36b	27.34c	17.36ab

Results represented the average of two seasons. The values of mean followed by different letters verify significantly by LSD at $p < 0.05$.

3.5. Chlorophyll content in leaves

Figure (4) shows the influence of foliar co-application of P&Zn&Fe NPs with *B. japonicum* on chlorophyll content in leaves of soybean plant. With respect to the other treatments, T11 treatment significantly increased chlorophyll content, achieving the highest values 9.54 mg/g FW followed by T7 8.49 mg/g FW, whereas the lowest values were realized with the corresponding plants exposed to T1 3.82 mg/g FW. With regard to the interaction effect of NPs, the most significant improvement in all measured chlorophyll content was observed when plants treated with P&Zn&Fe NPs with *B. japonicum*.




Figure (4): Effect of foliar co-application of NPs and *B. japonicum* on leaves chlorophyll content. (Results represented the average of two seasons). The significant differences among treatments was indicated by different letters ($P < 0.05$).

3.6. Effect of foliar co-application of NPs and *B. japonicum* on leaves Total Carbohydrates content and Leaf area

The foliar application of NPs and *B. japonicum* enhanced significantly the total carbohydrates of soybean leaves and leaf area (Figure 5, 6). The highest increases in total content of carbohydrates and leaf area of soybean's leaves were recorded for the co-application of NPs and *B. japonicum*, especially with (N, P, Zn, Fe) NPs and *B. japonicum* T7 followed by T11. Such increases were higher than the corresponding ones recorded for the application of other treatments. In this concern, T7 seemed to be the most efficient treatment recording significant increases in total carbohydrate content and leaf area by 20.67% and 122.74 cm², respectively higher than the non-amended control treatment, on the other hand T1 recorded significant decreases in both total carbohydrate content and leaf area by 15.54% and 115.41 cm².




Figure (5): Effect of foliar co-application of NPs and *B. japonicum* on leaves total carbohydrates. (Results represented the average of two seasons). The significant differences among treatments was indicated by different letters (P<0.05).




Figure (6): Effect of co application of NPs and *B. japonicum* on leaf area. (Results represented the average of two seasons). The significant differences among treatments was indicated by different letters (P<0.05).

3.7. Effect of foliar application of various nanoparticles (N&P&Zn&Fe) and *B. japonicum* on yield attributes of soybean in sandy soil.

Table (10) illustrates that the synergistic effect of *B. japonicum* and the nanoparticles (N& P& Zn& Fe) as foliar application have a positive impact on seed's macronutrients (N 6.33%, P 0.42%), micronutrients (Zn 46.87ppm, Fe 110.87ppm), however T10 (P& Fe NPs and *B. japonicum*) was significant increase in K 1.51%. The data illustrate that the highest positive influence on the investigated seed properties was achieved due to the foliar co-application of (N& P& Zn& Fe) nanoparticles and *B. japonicum*. This mean that th e companied NPs and *B. japonicum* enhanced the seed quality of soybean compared to control treatment.

Table (10): The Effect of various co-application nanomaterials (N&P&Zn&Fe) and *B. japonicum* on yield attributes of soybean in sandy soil.

Treatments	N%	P%	k%	Zn ppm	Fe ppm
T0	3.90b	0.31b	1.42b	40.84b	80.97c
T1	4.27ab	0.30b	1.42b	43.98ab	89.18b
T2	4.33ab	0.33b	1.45ab	43.99ab	93.50b
T3	5.16a	0.36ab	1.45ab	45.93a	102.24a
T4	5.12a	0.35b	1.45ab	44.54ab	101.61a
T5	5.09a	0.38ab	1.47a	44.40ab	101.25a
T6	5.10a	0.37ab	1.47a	44.34ab	100.46a
T7	6.33a	0.42a	1.37b	46.87a	110.87a
T8	3.87b	0.22c	1.41b	40.90b	81.79c
T9	4.92ab	0.36ab	1.44ab	44.21ab	97.39ab
T10	4.92ab	0.33b	1.51a	44.09ab	97.35ab
T11	6.24a	0.40a	1.47a	46.70a	108.98a
T12	5.15a	0.39a	1.40b	45.63a	102.71a
T13	6.15a	0.41a	1.41b	45.64a	106.12a
T14	3.91b	0.29b	1.41b	40.88b	82.47c

Results represented the average of two seasons. The values of mean followed by different letters verify significantly by LSD at $p < 0.05$.

The nutritional value of soybean (Seed protein and oil content) and the weight of 100 seed showed notable enhancement in response to the foliar spray of (N& P& Zn& Fe) NPs and *B. japonicum*. The plants treated with T7 significantly increase the oil percent 19.12%, the amount of protein in seed (32.43%) and also recorded the heaviest weight 100 seed (8.07gm) compared to T0 (control).Despite T7 recorded highly significant on seed quality but T11

(P&Zn&Fe and *B. japonicum*) was significant increase in oil percent 18.93%, the amount of protein in seed (38.97%) and also recorded increase in the weight 100 seed (10.27gm) compared to T0 (control) (Figure7). Our above results proved that *B. japonicum* has the ability to fix nitrogen, P Zn and Fe solubilization, therefore these results have positively reflect on the percent of N, P, Zn and Fe in seed.




Figure (7). yield attributes of soybean as affected by co application nanoparticles (N&P&Zn&Fe) and *B. japonicum*. The significant differences among treatments was indicated by different letters (P<0.05).

4. Discussion

Although, nanoparticles exert inhibitory activity against most microorganisms, but rhizobia able to grow strongly in the presence of N, P, Fe and Zn NPs at 100 and 40 ppm respectively compared to control (rhizobia only). This may be due to nanoparticles would increase the viability of nutrients for rhizobia and provide a conducive habitat for its

establishment or even acting as a source of nutrients for bacteria. According to **Dhas *et al.* (2014)**, bacteria probably accommodate to grow in the presence of nanomaterials or transfer them to another form to lower their toxicity during the lag phase of bacterial growth curve.

From the results it was concluded that nanoparticles induce rhizobium to tolerate a wide range of pH and temperature this explained by Niño *et al.* (2019) bacteria have ability to conjure numerous response mechanisms to adapt with nanoparticles in their growth habitats such as efflux pumps (membrane-bound transport proteins), electrostatic repulsion, altered morphology, formation of biofilm, extracellular matrices, transfer of genes, mutations, and metabolic responses.

On the other hand, nanoparticles have a positively effect on plant growth regulating characteristics, including production of growth hormones, siderophores and fixation of nitrogen by regulating the expression of genes *amoA* and *amoC2* that responsible for nitrification (**Amanda *et al.*, 2021**), this result is harmony with (**Dimkpa *et al.*, 2012b**) who found that the plant growth-promoting bacterium *Pseudomonas chlororaphis* showed increasing in IAA synthesis at 200 mg/L of CuO NPs, where the lowest concentration of IAA investigated at (1.89 µg/ml) for 50 ppm of Fe NPs. In this study, the obtained results were significantly lower than that recorded by **Seiduet *et al.* (2025)** where the amount of IAA produced by *Bradyrhizobium sp.* ranged from 54.97 ± 3.21 – 108 ± 12.10 µg/ml. The amount of EPS produced by *B. japonicum* contrast to **Ali and Hussein (2021)** who found that *Rhizobial* strains 3612, 1799 and 3629 showed EPS production 4.13, 3.62 and 3.54 g/L respectively, this due to the presence of NPs in growth culture at suitable concentration induced EPS production by *B. japonicum*. Our concerns were in line with the prior findings of **Khalid *et al.* (2020)**, who reported that different species of *rhizobium* are efficient in phosphorus solubilization.

These findings investigated that *B. japonicum* in the presences of NPs at certain concentration increase minerals solubilization, this support **Welch and Vandevivere's (1994)** hypothesis that naturally occurring polymers can affect mineral dissolution by acting as attachment structures to the mineral or rock surface. Moreover, the fresh microbial EPS increases the rate of feldspar dissolution, probably by forming complexes with framework ions in solution this confirmed by **Meena *et al.* (2014)**. **Alharbi *et al.* (2023)** reported that the co-application of nanoparticles and plant growth promoting rhizobacteria was generally found to increase the

content of mineral nutrient in rhizosphere rather than separately applied of nanoparticles or PGPR.

The foliar spray of NPs and *B. japonicum* resulted in increment in macro and micronutrient in soil and different plant parts like leaves and seed of soybean plant increase minerals solubilization and plant nutrient- uptake. This result is harmony (**De Souza *et al.*, 2021**) who demonstrated that the contents of Fe in soil and plant parts such as nodules, roots and shoots was increased by (Fe_3O_4 nanoparticules and *Rhizobium* sp. treatments) reveals that they enhance plants uptake of Fe from soil. Additionally (**Liu *et al.*, 2022; Verma *et al.*, 2023a**) reported that the content of soil nitrate was increased as a result of co-applying of nanomaterials and plant growth regulating rhizobacteria in comparison to a single dose of nitrogen. Moreover, NPs and PGPR boosted content of organic carbon, phosphorus, and nitrogen availability more than normal plants. Furthermore, NPs boosted rhizobial exopolysaccharides production, which plays a crucial role in enhancing structure of soil and retention of water, ultimately benefiting plant growth. EPS act as "glue," binding soil particles together, improving soil aggregation and reducing water and wind erosion. Interestingly, **Raliya *et al.* (2014)** added ZnO nanomaterials to culture medium of *Bacillus subtilis* strain JCT1 for enhancement EPS production that was improved aggregation in 33–83% in sandy soil, hence induce nutrient availability.

As the total minerals content increase in treated plants this may be regarded to accumulation of significant quantities of these minerals rather than control plants. This improvement in minerals content results from enhancement of nutrient uptake that promoted by introduced NPs and *Rhizobium*. The obtained results were close agreement with the previous results of **De Souza *et al.* (2021)** who found an increase in total nitrogen content in shoot (26.5–50.2%) and root (24.1–48.2%) of treated plants compared to control plants due to the effective combination between Fe_3O_4 NPs and *Rhizobium* inoculation. From this result, nanoparticles and *Rhizobium* sprayed on plant leaves increase in minerals in leaves more than the control this due to integration of *rhizobium* and NPs enhanced plant nutrient- uptake, this is in consistency with **Szopa *et al.* (2022)** who stated that the interaction between microbes and nanoparticle can boost nutrient uptake by plants encapsulated in nanoparticles, especially at detrimental conditions. Similarly, (**De Souza *et al.*, 2019**) investigated the synergistic action of NPs and *Rhizobium* lead to an increase in growth parameters of plants and also concentration of mineral content including Ca,

K, P, Fe, and Mn in all plant parts. These results were in similarity **Bashandy et al. (2024)** who found that the content of N, P, Zn, and K was high in seeds of faba bean plants due to triple application of microbial inoculation with *Rhizobium*, seaweed extract, and Zinc Oxide Nanoparticles.

The important role of *B. japonicum* in increase of nutrient solubility in addition to soil and foliar application of nanoparticles that enhance root absorption and pass on nutrient through stomata and allowing plants to receive essential nutrients, hence boosted the activity of the photosynthesis process, resulting in high chlorophyll content in leave. Our result is harmony with **Mahmoud and Sahar (2018) Ding et al. (2025)** revealed that the foliar application of combined of Fe and Zn-NPs showed high chlorophyll content (16.6–44.3 %) in wheat more than combination of (Zn), (Fe), and silicon NPs. Accordingly, it can be clarified that the usage of Zn and Fe-NPs in combination aggravated the level Fe and Zn in the chloroplast and therefore under drought conditions the photosynthetic pigments in wheat plants were enhanced. Additionally, **Sheta et al. (2024)** documented that the content of carbohydrate in cowpea leaves increased after Zn-NP and Fe-NP applications due to good growth of plant. On the other hand, **Rehman et al. (2023)** illustrated that the application of Zn-NP and Fe-NP increased, total phenol, proline, and carbohydrate content in wheat grown in saline soil.

The co-applying of *B. japonicum* and N&P&Zn&Fe showing significantly increase in leaf area in comparison to the other treatments. The obtained results could be explained as stated by **Ahmed et al. (2012) and Mahmoud et al. (2022)** who reported that Zinc (Zn) is responsible for elongation and enlargement of plant cells by stimulate tryptophan synthesise and its enzymes, that are responsible for biosynthesis of indole acetic acid beside that IAA produced by *B. japonicum* which exhibited more leaf growth.

On the other hand, co-application of NPs with *B. japonicum* boosted the increment seed's protein percent this due to nitrogen availability, both N taken by *B. japonicum* from the air, N provided by NPs and N available in the soil that influence amio acids formation and increase protein formation in seed this result is harmony with **Surtiningsih et al. (2009)** who reported that the availability of N from soil and air nitrogen fixation by *Rhizobium* bacteria is initially stacked on the stem and leaves and when the pods are formed, nitrogen is collected into pods, and about (30 - 90%) of N being absorbed into the seeds. Additionally, **Grangah et al. (2020)**

stated that nano-fertilizers especially Fe + Zn treatment improved the content of protein in bean seeds. Our results investigated that the co-application of nanoparticles and *B. japonicum* has positive effect on seed-oil content this may be due to Zn mobilized by *B. japonicum* and Zn supplied by soil and foliar application of NPs, which is effect on oil quantity as a result of activation of NADPH dependent dehydrogenase included in fat synthesis by application of zinc. These findings were also documented by those founded by **Raghuwanshi *et al.* (2017)**. Additionally, the obtained results are in agreement with **Yacoub *et al.* (2020)** who found a positive effect of foliar application of neutral rock P Nps+ ZnSO₄ or ZnO NPs recorded the maximum seed weight and straw yield, content of protein and protein (%) as well as content of oil and oil (%) of soybean. **Vaghar *et al.* (2021)** found that under drought conditions the lowest oil % at seed filling stage was (17.47%), while the highest oil % (23.73%) was recorded by foliar co-application of Zn and Mn nanoparticles in soybean plants.

The contribution of *B. japonicum* in N-fixation, which is responsible for amino acid formation and many other important biological compounds that have essential role in photosynthetic process and subsequently increase seed weight. This is in harmony with **Fituma *et al.* (2018)** who founded that the treated soybean plants with *Bradyrhizobium* recorded the highest 100-seeds weight (22 g), whereas the un-inoculated treatment (control) recorded the lowest 100-seeds weight (16.9 g).

5. Conclusions

Co-application of N & P & Zn & Fe NPs and *B. japonicum* showed positive effect on the growth and productivity of soybean in sand soil. The soil applications of *B. japonicum* enhanced minerals solubility in rhizosphere besides their production for exopolysaccharides that have a vital role in improve texture and properties of sandy soil. Additionally, both soil and foliar application of NPs and *B. japonicum* increased nutrient uptake and improved morphological and physiological traits of plant. Furthermore, the synergistic performance of NPs and rhizobium sp. increase chlorophyll content, carbohydrates percent and leaf area of soybean plant. The plant growth promoting traits of *B. japonicum* and NPs application provide highest quality and quantity of soybean seeds. From the results of these study, it was suggested that the soil addition and

foliar spray of N &P &Zn &Fe nanoparticles and *B. japonicum* could be recommended to improve soybean plants cope growth in sandy soil.

6. References

Abdel Wahab, M. M., Hassan, M. R., Sanaa, E. A. E., Hameed, A. and Emad, A. A. (2023). Application of Silicon, Zinc, and Zeolite Nanoparticles—A Tool to Enhance Drought Stress Tolerance in Coriander Plants for Better Growth Performance and Productivity. *Plants* 12: 2838. <https://doi.org/10.3390/plants12152838>

Abdel Wahab, M. M. , Sanaa, E. A. E. , Asmaa, B. E, Eman, Z. O. and Reham, M. E. (2022). Prospective Practice for Compound Stress Tolerance in Thyme Plants Using Nanoparticles and Biochar for Photosynthesis and Biochemical Ingredient Stability. *Agronomy*12: 1069. <https://doi.org/10.3390/agronomy12051069>

Abdel Wahab, M. M. and Sahar, S. T. (2018). Main Sulphur Content in Essential Oil of *Eruca Sativa* as Affected By Nano Iron and Nano Zinc Mixed with Organic Manure. *Agriculture Poľnohospodárstvo* 64 (2): 65–79. DOI: 10.2478/agri-2018-0007

Abdel Wahab, M. M., El-Attar, A.B., Abeer, A. M. (2017). Economic Evaluation of Nano and Organic Fertilizers as an Alternative Source to Chemical Fertilizers on *Carum Carvi* L. Plant Yield and Components. *Agriculture (Poľnohospodárstvo)* 63 (1): 33–49

Abdel Wahab, M. M., Emad, A. A., Suzy, M. A., Mohamed, B. I. E. and Shady, A. M. (2020). Synergetic Effects of Zinc, Boron, Silicon, and Zeolite Nanoparticles on Confer Tolerance in Potato Plants Subjected to Salinity. *Agronomy* 10: 19. doi:10.3390/agronomy10010019

Abdel Wahab M. Mahmoud and Sahar S. Taha (2018). Main Sulphur Content in Essential Oil of *Eruca Sativa* as Affected By Nano Iron and Nano Zinc Mixed with Organic Manure. *Agriculture (Poľnohospodárstvo)* 64 (2): 65–79. DOI: 10.2478/agri-2018-0007

Ahmed, H.A.H., Khalil, M.K., Abd-EiRahman, A.M., Nadia, A.M.(2012). Effect of zinc, tryptophan and indole acetic acid on growth, yield and chemical composition of Valencia orange trees. *J. Appl. Sci. Res* 8: 901–914

Ahemad, M. and Kibret, M.(2014). Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *J. King. Saud. Univ. Sci.* 26: 1–20

Alharbi, K., Hafez, E. M., Omara, A. E. D., Rashwan, E. and Alshaal, T. (2023). Zinc oxide nanoparticles and PGPR strengthen salinity tolerance and productivity of wheat irrigated with saline water in sodic-saline soil. *Plant Soil* 493: 475–495

Ali, A.A. and Hussein, M.H.A. (2021). Co-Inoculation Effect of Rhizobia with Potassium and Zinc Solubilizing Bacteria for Enhancing Some Growth Parameters of *Phaseolus vulgaris* Plants. *Middle East Journal of Agriculture Research* 10(4): 1431-1441

Amanda, C. P. M., Lucas, S. R., Emerson, R. C. and Paulo, T. L.(2021). The potential of nanomaterials associated with plant growth promoting bacteria in agriculture. *Biotech* 11:318

Aoda, M.I., Smucker, A.J.M., Majeed, S.S., Mohammed, H.A., Al-Sahaf, F.H. and Robertson, G.P. (2021). Novel Root Zone Soil Water Retention Improves Production with Half the Water in Arid Sands. *Agron J* 113 : 2398–2406

Ayman. A., Hassan A. Z. A., Rania A. El-bialy, Abdel Wahab M. M.(2020). Fate of Nano Soil Particles and Their Influence on Geotechnical and Physiochemical Properties of Sandy Soil. *American Journal of Nanomaterials* 8 (1): 48-57. DOI:10.12691/ajn-8-1-5

Bandemer, S. L. and Schaible.P.J. (1944). Determination of iron. *Ind Eng Chem* 16: 317-319

Bashandy, S.O., Abd El-Hafeez, A.M. and Sarhan, M.G.R.(2024). Influence of Zinc Oxide Nanoparticles (ZnONPs), Seaweed Extract and Microbial Inoculation with *Rhizobium* on Faba Bean Nodulation, Yield, and Quantity. *Journal of Soil Sciences and Agricultural Engineering* 15 (10):279 – 285

Brick, J. M., Bostock, R. M. and Silverstone, S. E. (1991). Rapid in situ assay for indole acetic acid production by bacteria immobilized on nitrocellulose membrane. *Appl Environ Microbiol.* 57: 535–538

Campos, E. V. R., Pereira, A., Aleksieienko, I., do Carmo, G. C., Gohari, G. and Santaella, C.(2023). Encapsulated plant growth regulators and associative microorganisms: nature-based solutions to mitigate the effects of climate change on plants. *Plant Sci* 331:111688

Castro, J., Calijuri, M., Ferreira, J., Assemany, P. and Ribeiro, V.(2020). Microalgae based biofertilizer: a life cycle approach. *Sci Total Environ* 724: 138138

Damery, J.T. and Alexander, M.(1969). Physiological differences between effective and ineffective strains of *Rhizobium*. *Soil Science* 108(3): 209-216

Daneshvar, N., Aber, S., Sayed, D., Khataee, M.S. and Rasoulifard, A.R.(2007). Preparation and investigation of photocatalytic properties of ZnO nanocrystals: Effect of operational parameters and kinetic study. *Int J Nucl. Quantum Eng* **1**: 24–29

Dawood, M. G. (2021). Effect of priming with proline on the performance of two wheat cultivars. *Agricultural Engineering International: CIGR Journal* **23**(3)

De Souza, A., Govea-Alcaide, E., Masunaga, S.H., Fajardo-Rosabal, L., Effenberger, F., Rossi, L.M. and Jardim, R.F.(2019). Impact of Fe₃O₄ nanoparticle on nutrient accumulation in common bean plants grown in soil. *Springer Nat Appl Sci* **1**: 308

De Souza, A.T., Ernesto, G. A., Ernesto, G. P., Sueli, H. M. F., Effenberger, Liane, M. R., Raúl, L. and Renato, F. J.(2021). Fe₃O₄ nanoparticles and *Rhizobium* inoculation enhance nodulation, nitrogen fixation and growth of common bean plants grown in soil. *Rhizosphere* **17** : 100275

Dhas, S.P., Shiny, P.J., Khan, S., Mukherjee, A. and Chandrasekaran, N.(2014). Toxic behavior of silver and zinc oxidenanoparticles on environmental microorganisms. *J Basic Microbiol* **54**: 916–927.

Dimkpa, C.O., Zeng, J. and McLean, J. E.(2012b). Production of indole-3-acetic acid via the indole-3-acetamide pathway in the plant beneficial bacterium *pseudomonas chlororaphis* O6 is inhibitedby ZnO nanoparticles but enhanced by CuO nanoparticles. *Appl Environ Microbiol* **78**:1404–1410

Ding,M., Chigui, X., Shaoxiang, L., Jia, L., Yunxin, K.,Cuiping, Z., Hongsheng, L., Yan, Y., Kun, L., Muzammal, R., Zahid, M., Guiqiang, F.,Sobhy, M. I, Chaowu, Z., Shah, F.and Gang, D.(2025). Effects of iron, zinc, and silicon nanoparticles on morpho-physiological growth, yield, and quality of wheat (*Triticumaestivum* L.) under drought stress.*Journal of Hazardous Materials Advances* **19**:100789

FAO (2021).The State of Food and Agriculture. Climate Change, Agriculture and Food Security; FAO: Rome, Italy

Fituma, T., Tana, T. and Alemneh, A. A.(2018). Response of soybean to *Bradyrhizobium japonicum* inoculation and phosphorus application under intercropping in the Central Rift Valley of Ethiopia. *South African Journal of Plant and Soil* **35**(1): 33-40

Francis, A. J., Dodge, C., Chendrayan, K. and Quinby, H.(1988). An-aerobic microbial dissolution of lead oxide and production of organicacids.US Patent No. 4758345

Grangah, F. M., Rashidi, V., Mirshekari, B., Khalilv,B. E. and Farahvash, F.(2020). Effects of nano-fertilizers on physiological and yield characteristics of pinto bean cultivars under water deficit stress. *J Plant Nutr* 43: 2898–2910

Jackson, M. L.(1973). Soil Chemical Analysis. Prentice Hall, New Delhi. Kamilova F, Kravchenko L V, Shaposhnikov A I. 2006, Organic acids, sugars, and L-tryptophan in exudates of vegetables growing on stone wool and their effects on activities of rhizosphere bacteria. *Molecular Plant-Microbe Interactions* 19: 250–256

Khalid, R., Zhang, X.X., Hayat, R. and Ahmed, M.(2020). Molecular characteristics of therhizobia isolated from Arachishypogaea grown under stress environment. *Sustainability* 12: 6259

Khalid, M., Liu, X., Ur Rahman, S., Rehman, A., Zhao, C. and Li, X.(2023). Responses of microbial communities in rhizocompartments of king grass to phytoremediation of cadmium-contaminated soil. *Sci Total Environ* 904: 167226. doi: [10.1016/j.scitotenv.2023.167226](https://doi.org/10.1016/j.scitotenv.2023.167226)

Kolapo, A., Temitope, O. O., Nolwazi, Z. K., Khalid, M. E., Hazem, S. K. and Olajide, J. F.(2025). Enhancing land nutrient through rhizobia biofertilization: modeling the joint effects of rhizobium inoculants and improved soybean varieties on soybean productivity in North Central, Nigeria. *Frontiers in Sustainable Food Systems* 9: 1509230

Liu, L., Nian, H.; and Lian, T(2022). Plants and rhizospheric environment: affected by zinc oxide nanoparticles (ZnO NPs). A review. *Plant Physiol Biochem* 185: 91–100

Ma, D., Yin, L., Ju, W., Li, X., Liu, X. and Deng, X.(2021). Meta-analysis of green manure effects on soil properties and crop yield in northern China. *Field Crop Res* 266: 108146

Mahmoud, N., Ahmed, M. A.,Tingting, W. , Shi, S. and Han, T.(2024). Soybean in Egypt: current situation, challenges, and futureperspectives. *Discover Sustainability* 5425

Mahmoud, A.W.M.; and Swaefy, H.M. (2020). Comparison between commercial and nano NPK in presence of nano zeolite on sage plant yield and its components under water stress. *Agriculture* 66: 24–39

Meena, V.S., Maurya, B.R. and Verma, J.P. (2014).Does a rhizospheric microorganism enhance K⁺ availability in agricultural soils? *Microbiol Res* 169: 337-347

Musei, S.K., Kuyah, S., Nyawira, S., Ng' ang' a, S.K., Karugu, W.N., Smucker, A. and Nkurunziza, L.(2024). Sandy soil reclamation technologies to improve crop productivity and soil health: a review. *Front. Soil Sci* 4: 1345895

Nett, R., Bender, K. and Peters, R.(2022). Production of the plant hormone gibberellin by rhizobia increases host legume nodule size. *ISME J* 16: 1809–1817

Ninama, J., Razauddin, Kushal, S., Sulochna, Babita, Y., Satya, N., Satapathy, Jogendra, K. and Singh, B.V.(2023). Effects and consequences of nano fertilizer application on plant growth and developments: a review. *Int J Environ Clim* 13: 2288–2298

Niño-Martínez, N., Salas Orozco, M.F., Martínez-Castañón, G.A., Torres Méndez, F. and Ruiz, F(2019). Molecular mechanisms of bacterial resistance to metal and metal oxide nanoparticles. *Int J Mol Sci* 20 : 2808

Qu, S., Yang, H., Ren, D., Kan, S., Zou, G., Li, D. and Li, M.J. (1999). Magnetite nanoparticles prepared by precipitation from partially reduced ferric chloride aqueous solutions. *Journal of colloid and interface science* 215(1): 190–192

Raliya, R., Tarafdar, J. C., Mahawar, H., Kumar, R., Gupta, P. and Mathur, T.(2014). ZnO nanoparticles induced exopolysaccharide production by *B. subtilis* strain JCT1 for arid soil applications. *Int J Biol Macromol* 65: 362–368

Raghuwanshi, N., Sharma, B.L., Ulkey,I. and Prajapati, S.(2017). Residual and cumulative effect of zinc on yield, quality of soybean (*Glycine max* L.) and various pools of zinc in a vertisol of Madhya Pradesh, CV. JS 97-52. *International Journal of Bio-resource and Stress Management* 8(3): 444-449

Rehman, M.Z., Anayatullah, S., Irfan, E., Hussain, S.M., Rizwan, M., Sohail, M.I., Jafir, M., Ahmad, T., Usman, M. and Alharby, F.(2023). Nanoparticles assisted regulation of oxidative stress and antioxidant enzyme system in plants under salt stress: A review. *Chemosphere* 314: 137649

Salama, H.S.A., Nawar, A.I., Khalil, H.E. and Shaalan, A.M(2021). Improvement of maize productivity and n use efficiency in a no-tillage irrigated farming system: effect of cropping sequence and fertilization management. *Plants* 40

- Seidu, O.A., Stephen, M. G., John, M. W. and Victoria, W. N.(2024).** Isolation, screening and *in-vitro* characterization of plant growth-promoting *Bradyrhizobium* isolates from the nodules of Bambara groundnut (*Vigna subterranean*) for potential use as bioinoculants. *Front Sustain Food Syst* 8:1506346
- Shang, H., Ma, C., Li, C., Zhao, J., Elmer, W., White, J.C. and Xing, B.(2021).** Copper oxide nanoparticle-embedded hydrogels enhance nutrient supply and growth of lettuce (*Lactuca sativa*) infected with *Fusarium oxysporum* f. sp. *lactucae*. *Environ Sci Technol* 55: 13432–13442
- Sheta, M.H., Ahmed, H. M. A., Mohammed, A. E., Hala, M. B., Nawal, A. O., Mohamed, A. M. A., Abdel-Nasser, A. E., Rasha, S. E. and Mahmoud, M. I. M.(2024).** Green Synthesis of Zinc and Iron Nanoparticles Using *Psidiumguajava* Leaf Extract Stimulates Cowpea Growth, Yield, and Tolerance to Saline Water Irrigation. *Horticultura*. 10: 915 1-20
- Singh, A., Rajput, V.D., Pandey, D., Sharma, R., Ghazaryan, K. and Minkina ,T.(2023).** Nano zinc-enabled strategies in crops for combatting zinc malnutrition in human health. *Front Biosci* 28: 158
- Snedecor, G.W. and Cochran, W.G.(1980).** Statistical methods.7th Ed., Iowa State Univ. Press, Towa, USA
- Soumare, A., Diedhiou, A., Thuita, M., Hafidi, M., Ouhdouch, Y. and Gopalakrishnan, S.(2020).** Exploiting biological nitrogen fixation: a route towards a sustainable agriculture. *Plan Theory* 9: 1011
- Surtiningsih, T., Farida, dan T. and Nurhariyati (2009).** Biofertilisasi Bakteri Rhizobium Pada Tanaman Kedelai. *Berk Penel Hayati* 15: 31–35
- Szopa, D., Mielczarek, M., Skrzypczak, D., Izydorczyk, G., Mikula, K. and Chojnacka, K.(2022).** Encapsulation efficiency and survival of plant growth-promoting microorganisms in an alginate-based matrix – a systematic review and protocol for a practical approach. *Ind Crop Prod* 181:114846
- Tafida, I., Nazifi, B. and Adam, A. S.(2022).** Socio-economic analysis of the adoption of selected improved soybean (*Glycine max*) varieties in Tofa local government area of Kano state, Nigeria. *FUDMA J Agric Agric Technol* 8 : 99–107
- Vaghar, M. S., Sayfzadeh, S., Zakerin, H. R., Kobraee, S. and Valadabady, A. R.(2021).** Effect of foliar application of iron, zinc and manganese nanochelate on some quantitative and qualitative characteristics of soybean (*Glycine Max* L.) under water deficit stress. *Environ Stresses Crop Sci* 14: 703–718

Vardumyan, H., Singh, A., Rajput, V.D., Minkina, T., Ragab, E. H. and Ghazaryan, K.(2024). Additive-mediated phytoextraction of copper-contaminated soils using *Medicago lupulina* L. Egypt J Soil Sci 64:599–618

Verma, K. K., Song, X., Verma, C. L., Huang, H., Singh, M. and Xu, L.(2023a). Mathematical modeling of climate and fluoride effects on sugarcane photosynthesis with silicon nanoparticles. Plant Physiol Biochem 204: 108089

Vincent, J.M.(1970). A Manual for the Practical Study of the Root Nodule Bacteria. I B P, Hand book, No. 15, Bluckwell Publishers, Oxford, UK

Welch, S.A. and Vandevivere, P.(1994). Effect of microbial and other naturally occurring polymers on mineral dissolution. Geomicrobiol J 12: 227-238

Wilcockson J. and Werner D. (1978). Nitrogenase Activity of *Rhizobium japonicum* Growing on Agar Surfaces in Relation to Slime Production, Growth and Survival. Journal of General Microbiology 108: 151-160

Yacoubi, N.M., Ismail, S.A., Raslan, M. and Khedr, M.H.(2020). The effect of using nanoparticles phosphorus and zinc on quality and quantity of soybean (*Glycine Max* l.). Plant Archives 20: 8863-8876

Zahra, Z., Habib, Z., Hyun, H. and Shahzad, H. M. A.(2022). Overview on recent developments in the design, application, and impacts of nanofertilizers in agriculture. Sustainability 14: 9397. <https://doi.org/10.3390/su14159397>