

Genome-wide analysis of the SWEET gene family in *Hemerocallis citrina* and functional characterization of HcSWEET4a in response to salt stress

Lihong Cao

Shanxi Agricultural University

Jinyao Wang

Shanxi Agricultural University

Lixuan Wang

Shanxi Agricultural University

Huili Liu

Shanxi Agricultural University

Wenjing Wu

Shanxi Agricultural University

Feifan Hou

Shanxi Agricultural University

Yuting Liu

Shanxi Agricultural University

Yang Gao

Shanxi Agricultural University

Xiaojing Cheng

Shanxi Agricultural University

Sen Li

saulisen@163.com

Shanxi Agricultural University

Guoming Xing

Shanxi Agricultural University

Research Article

Keywords: *Hemerocallis citrina*, SWEET, Genome-wide, Abiotic stressors, Functional characterization

Posted Date: May 24th, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-4408049/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Sugars will be eventually effluxed transporter (SWEET), one of the sugar transporters in plants, plays significant roles in numerous biological processes and responding to abiotic stressors. However, the characteristics and functions of the *SWEET* genes in *Hemerocallis citrina* remain unclear and poorly elucidated. In this study, the whole genome of *Hemerocallis citrina* was utilized to conduct bioinformatics analysis and a total of 19 *HcSWEET* genes were successfully identified. Analysis of the physicochemical properties indicated dominant differences among these *HcSWEETs*. A phylogenetic analysis revealed that *HcSWEET* proteins can be divided into 4 clades ranging from Clade I to IV, where proteins within the same clade exhibited shared conserved motifs and gene structures. Five to six exons were contained in the majority of *HcSWEET* genes, which were unevenly distributed across 11 chromosomes. The gene duplication analysis showed the presence of 4 gene pairs. Comparative syntenic maps revealed that the *HcSWEET* gene family might present more closed homology in monocotyledons than dicotyledons. Cis-acting element analysis of *HcSWEET* genes indicated key responsiveness to various hormones, light, and stresses. Additionally, transcriptome sequencing analysis suggested that most *HcSWEET* genes had a relatively higher expression in roots, and *HcSWEET4a* was significantly up-regulated under salt stress. Overexpression further verified the possibility that *HcSWEET4a* is involved in response to salt stress, which provides novel insights and facilitates in-depth studies of the functional analysis of *HcSWEETs* in resistance to abiotic stress.

Background

Growth and development processes are often severely affected when plants are exposed to various abiotic stresses such as drought and salinity due to their sessile characteristics [1, 2]. Sugars, as major sources of carbon skeletons for cell wall and energy metabolism, are crucial signals that regulate the expression of related genes under abiotic stress [3, 4]. Cell-to-cell transport of sugars is mediated by transport proteins and three transporters including MSTs (monosaccharide transporters), SUTs (sucrose transporters), and SWEET have been identified so far [5, 6]. Unlike two other transporters, the SWEETs can typically act as bidirectional transmembrane transporter for sugars to cross the biomembrane system along a pH-independent concentration gradient via MtN3/saliva structural domains [7]. Members of the *SWEET* gene family encode membrane proteins consisting of conserved transmembrane (TM) domains linked by a PQ-loop-repeat, which are conserved across eukaryotes and prokaryotes [8, 9]. However, Eukaryotic SWEET proteins generally contain seven transmembrane helices (TMHs), obviously distinguishable from semiSWEETs consisting of 3 TMHs in prokaryotes, which may be correlated to the evolution process of plants [10, 11].

As a transporter identified about sugars recently, the SWEETs are found to be present in all kingdoms of life [5, 9]. SWEET proteins play central roles in plant growth and development such as plant defense, long-distance transport of sucrose, vegetative and reproduction growth, senescence, and stress responses [12–14]. Currently, four SWEETs clades have been reported in plants, respectively Clade I, II, III, and IV, furthermore genes belonging to the same clade hold similar structures and functions [8, 15]. Considering the physiological relevance of the fertility of sucrose, the AtSWEET13 sucrose transporter contributes to male fertility restoration [16]. Vacuolar Sugar Carrier *AtSWEET16* overexpression lines promote stress tolerance under non-favorable conditions [17]. OsSWEET11 and OsSWEET15 participate in seed development in rice [18]. Clade III OsSWEETs, which are taken as interaction factors, mediate bacterial blight susceptibility by interacting with OsHMGB1 and OsHsp20L to breed broad-spectrum disease-resistant rice [19]. SISWEET7a and SISWEET14 negatively regulate the sugar transport and storage of fruits in tomatoes [20]. MdSWEET9b positively regulates sugar accumulation in apples, providing the foundation for

exploring the metabolism between hormone and sugar pathways [21]. GmSWEET15 in soybean mediates sucrose export from the endosperm to the early embryo seed development [22]. CsSWEET17 is activated to positively regulate cold tolerance by interacting with CsLHY in tea plants [23]. Pineapple SWEET10 is found to act as a potential glucose transporter, by manipulating whose activity fruit crop yield and quality may be improved [24]. SWEET11b transports both sugar and cytokinin to maintain normal development in barley grains [25].

The plant cell membrane is the first cell organelle to transfer chemical signals to the cytoplasm when receiving a stress signal under harsh environmental conditions. Interestingly, the acclimatization makes plants more tolerant to be defended against most stresses during exposure to abiotic stressors [26]. In the long term, environmental stresses like drought and salinity can activate stress-related response genes of plants to adjust to osmotic pressure [27–30]. Therefore, innovative solutions are inevitable to develop to mitigate soil salinity and drought stress. Sucrose transport, a key physiological process, is impacted by abiotic stresses [31]. Overexpression of *OsMST6* confers improved resilience to drought and salt stress in *Arabidopsis thaliana* [32]. Up-regulation of SUTs may be necessary to maintain phloem transport despite decreased sucrose available under salt and drought stresses in plants [33]. OsSUTs show significant importance in coping with drought and salt stress for rice plants [34]. The increase of soluble sugar contents and tolerances to both drought and salt stress are enhanced by elevating the production of trehalose in *OsTPSP* overexpression lines [35]. OsSWEET13 and OsSWEET15 transporters of Clade III pose critical importance of sucrose translocation and distribution in adaptive responses of rice plants to abiotic stresses [31].

Considering edible, medicinal, and ornamental value in horticultural perennial crops, *Hemerocallis citrina* Borani (night lily) is paid relatively more attention and widely cultivated worldwide. However, the lack of the night lily genome restrains the progression that molecular mechanisms and gene function analysis are explored. Therefore, as genomic information was released in 2021, new insights will be provided for a deeper understanding of the evolution, development, and biological functions of night lily [36]. As previous results reported, the *SWEET* gene family has become the research focus on responsiveness to abiotic stress. To investigate how night lily responds to drought and salt stress, we performed genome-wide identification of the *SWEET* members. In this study, members of the *HcSWEET* gene family were identified at the genomic level, followed by physicochemical properties, phylogenetic trees, conserved motifs, gene structures, chromosomal positions, cis-acting elements, and collinearity relationships were comprehensively analyzed using bioinformatics methods. Additionally, tissue-specific expression patterns of *HcSWEET* genes, as well as relative expression levels of genes under drought and salt treatments, were analyzed by transcriptome sequencing (RNA-seq). And that RNA-seq data from stress treatments were validated by RT-qPCR. The function of *HcSWEET4a* in stress response was further investigated by the *Agrobacterium*-mediated watermelon transformation method. The research can serve as a promising reference for offering strategies for improving tolerance to abiotic stress in night lily.

Results

Identification and characterization of the SWEET gene family in night lily

A total of 19 genes that encode HcSWEET proteins were identified after homology alignment and conservative domain verification. These genes were named *HcSWEET1* to *HcSWEET16* according to their homology with dicotyledons such as *Arabidopsis* AtSWEETs and monocotyledons such as rice OsSWEETs. *HcSWEET* gene characteristics of open reading frame (ORF) size, number of amino acid (AA/aa), protein molecular weight (MW), theoretical isoelectric point (pI), instability index (II), aliphatic index (AI), grand average of hydropathicity (GRAVY), number of predicted TMHs and conserved domain position are analyzed in Table 1. As a consequence, the deduced

number of amino acids in the night lily SWEET proteins was from 229 to 299, with the MW varying widely from 25.617 kDa to 32.938 kDa, and the PI index is between 5.49 to 9.66. All proteins ranged from 27.08 to 42.19 on the instability index, from 105.08 to 133.65 on the aliphatic index, and from 0.419 to 0.934 on GRAVY. The majority of predicted TMHs number was seven other than 6 TMHs of *HcSWEET5* and *HcSWEET13a/d*. Coupled with subcellular localization, the secondary structure including α -helix, extended strand, β -turn and random coil are summarized in Supplementary Table S1. Subcellular localization predictions suggested that the majority of *HcSWEET* genes located in the plasma membrane, are also found in the chloroplast, vacuole or cytoplasm (Supplementary Table S1). Considering the transmembrane characteristic, we predicted 3D structural models of 19 HcSWEET proteins using a homology modeling approach, shown in Supplementary Figure S1. The results demonstrated that various properties existed in different *HcSWEET* members.

Table 1
Characterization of the *SWEET* gene family in night lily

Gene name	ORF length (bp)	AA(aa)	MW(kDa)	pI	II	AI	GRAVY	THM	MtN3/saliva(PQ-loop repeat) domain position
<i>HcSWEET1a</i>	735	244	27190.28	9.26	40.09	112.58	0.666	7	7–95,132–214
<i>HcSWEET1b</i>	765	254	28508.65	9.08	27.08	105.08	0.504	7	7–95,132–214
<i>HcSWEET2</i>	690	229	25617.54	9.28	35.03	121.16	0.721	7	15–97,134–218
<i>HcSWEET3</i>	717	238	26521.72	9.60	42.19	117.52	0.684	7	7–98,132–217
<i>HcSWEET4a</i>	774	257	28446.10	9.13	30.78	125.1	0.846	7	11–94,134–218
<i>HcSWEET4b</i>	735	244	26987.61	8.95	34.37	133.65	0.95	7	10–98,133–217
<i>HcSWEET4c</i>	735	244	26992.56	8.89	37.04	130.82	0.934	7	11–98,133–217
<i>HcSWEET5</i>	765	254	28089.69	9.30	30.26	129.61	0.84	6	11–94,134–218
<i>HcSWEET6</i>	753	250	27597.16	9.39	35.56	134.04	0.903	7	10–95,134–218
<i>HcSWEET11</i>	798	265	29691.26	8.99	36.51	111.09	0.583	7	14–99,133–219
<i>HcSWEET13a</i>	747	248	27668.36	8.95	27.33	123.79	0.751	6	12–73,107–188
<i>HcSWEET13b</i>	867	288	32084.82	5.51	36.81	117.4	0.554	7	12–82,134–215
<i>HcSWEET13c</i>	846	281	31554.43	8.86	34.76	118.9	0.52	7	12–98,134–215
<i>HcSWEET13d</i>	747	248	28178.51	8.63	30.77	118.23	0.447	6	1–75,110–196
<i>HcSWEET14a</i>	837	278	31378.75	9.23	33.44	124.5	0.626	7	12–99,133–215
<i>HcSWEET14b</i>	858	285	32145.67	8.96	29.52	123.12	0.629	7	12–99,133–215
<i>HcSWEET14c</i>	858	285	32321.79	8.49	28.01	115.58	0.524	7	12–99,133–215
<i>HcSWEET15</i>	870	289	32114.98	5.49	40.11	124.15	0.754	7	12–98,134–215
<i>HcSWEET16</i>	900	299	32938.05	9.66	32.26	111.07	0.419	7	8–90,128–211

Phylogenetic analysis of HcSWEET proteins

To better understand the evolutionary interrelatedness among these HcSWEET members, we constructed a phylogenetic tree containing SWEETs from six species (*Hemerocallis citrina*, *Hemerocallis fulva*, *Oryza sativa*, *Zea mays*, *Arabidopsis thaliana*, *Vitis vinifera*) based on amino acid sequences using the TBtools software (Fig. 1). SWEET protein sequences of six species were listed in Supplementary Information 1. The results revealed that 19 HcSWEET members were classified into four major clades, namely Clade I, II, III, and IV respectively. Multiple HcSWEETs are tightly clustered with other SWEETs. Among these clades, the member number varied significantly. Clade III contained the most HcSWEET members (9), followed by Clade II (5), I (4), and IV (1), indicating a similar distribution pattern and gene function to these of other species.

Conserved motif composition, domain, and gene structure analysis of HcSWEET genes

According to the results from phylogenetic analysis, we divided the identified 19 HcSWEET proteins into 4 clades (Fig. 2A). Ten different conserved motifs were identified by analyzing amino acid sequences of HcSWEET proteins using the MEME online tool (Fig. 2B). The number and distribution of these motifs showed variations in different proteins. Each protein had 5–7 motifs, in which Motif 1, 3, and 4 were found in all proteins, moreover, different clades had distinct motifs. Motif analysis suggested that the *HcSWEET* gene family showed both conservation and differences, which might result from the diversity of protein functions within clades. Conserved domain analysis revealed that HcSWEET proteins possessed domains consisting of MtN3_slv and MtN3_slv superfamily (Fig. 2C), implying high conservation during evolution. In addition, to understand the structural compositions of *HcSWEET* genes, a structural map including the untranslated region (UTR) and coding sequence (CDS) was constructed based on the genome sequence (Fig. 2D). The number of CDSs ranged from 5 to 6. Coding sequences of 19 *HcSWEETs* were displayed in Supplementary Information 2. Gene structure analysis suggested that members within the same evolutionary clade exhibited similar gene structures.

Chromosomal localization, gene duplication, and collinearity analysis of HcSWEETs in *Hemerocallis citrina*, *Arabidopsis thaliana* and *Oryza sativa*

To investigate the chromosomal localization of *HcSWEET* genes, we analyzed the distribution of identified 19 genes on night lily chromosomes (Fig. 3). The distributions of these genes were determined via the night lily genome reported in 2021 [36]. The results showed that 19 genes were evenly dispersed across 9 of 11 chromosomes (LGs) with gene counts ranging from 1 to 4 in total. Notably, the largest number of genes were located in LG2, in contrast, LG6 and LG8 hosted only one gene and no gene located in LG7 and LG9.

To set forth whether gene duplication events forced the evolution of the *HcSWEET* gene family, an intra-species collinear analysis was performed (Fig. 4A). Among all gene pairs of the night lily genome, four pairs of duplicated segments were identified in the *HcSWEETs*, implying that segmental duplication events were likely to contribute to genetic diversity. Subsequently, we constructed inter-species syntenic maps of *Hemerocallis citrina* with other two representative species including monocotyledon *Oryza sativa* and dicotyledon *Arabidopsis thaliana* to identify orthologous genes through utilizing TBtools (Fig. 4B). It was observed that seven genes exhibited collinearity relationships with two genes in *Arabidopsis thaliana* and ten genes in *O. sativa*. These results suggest that *HcSWEET* genes likely share more similarities in function with *OsSWEET* genes in *Oryza sativa*.

Identification and distributions of cis-regulatory elements in HcSWEETs promoters

We specifically extracted 2000 base pairs upstream of the *HcSWEET* genes transcription start codon to explore their potential biological functions through the database named PlantCARE online. As shown in Fig. 5, a total of three types of main responsive elements were identified, hormone responses, plant growth and development, and

responses to abiotic stress respectively. It was found that 19 genes contained 6 to 26 light-responsive elements, indicating that the majority of *HcSWEET* genes may be regulated by light signaling. Moreover, all 19 genes contained hormone-related and development-related cis-acting elements. Hormone-related elements were referred to as MeJA, salicylic acid, abscisic acid, auxin, and gibberellin response elements. In contrast, development-related elements contained zein metabolism regulation, anaerobic induction, cell cycle regulation, seed-specific regulation, circadian control, flavonoid biosynthesis, endosperm, and meristem expression, anoxic specific inducibility and phytochrome down-regulation response elements. Additionally, *HcSWEET* genes other than *HcSWEET4c*, *HcSWEET13d*, and *HcSWEET14b* had cis-acting elements linked to abiotic stress, involving defense and stress, drought, and low-temperature responsive elements. *HcSWEET13d* also contained the wound-responsive element. Overall, various types of cis-elements were contained in the promoter regions of 19 *HcSWEETs*, indicating their possible involvement in diverse biological processes, environmental stresses, and regulatory pathways.

Expression patterns of HcWEET genes across different tissues and phenotype observations under drought and salt stress

To further analyze the possible roles of *HcSWEET* genes in abiotic stress, 19 genes were analyzed for relative expression profiles in different tissues and developmental stages (tender root, mature root, bud, tender leaf, mature leaf, tender scape, and mature scape) (Fig. 6A). As listed in Supplementary Table S3, RNA-seq results of tissue-specific expression revealed that most *HcSWEETs* in Clade II (*HcSWEET4a*, *HcSWEET4b*, *HcSWEET5*) and III (*HcSWEET13c*, *HcSWEET13d*, *HcSWEET14a*, *HcSWEET14b*, *HcSWEET14c*, *HcSWEET15*) exhibited the higher transcript levels in roots than other organs. Among them, *HcSWEET4a* and *HcSWEET13c* were simultaneously in a related highly expression in the scape. Except these genes, *HcSWEET2* in Clade I was expressed especially and abundantly in roots throughout the developmental process.

The transporters of the *SWEET* gene family in *Arabidopsis thaliana* have been reported to increase root proliferation by enhancing sucrose transport from shoot to root during abiotic stress [37]. Therefore, we carried out stress experiments to determine whether the case is suitable for *HcSWEET* genes in night lily. We treated 30 d seedling roots with drought stress for 0 h, 24 h, 48 h, 72 h, and 108 h, and identically salt stress for 0 h, 24 h, 48 h, 72 h, and 96 h. As was displayed in Fig. 6B and 6C, night lily plants primarily showed a slight symptom of loss of water when treated for 24 h, after treatment for 48 h, leaves presented severe yellow and rotten roots.

Expression pattern analysis of HcSWWTs under drought and salt stress

Based on the phenotypes observed in night lily plants treated by drought and salinity, roots were collected, followed by RNA-Seq analysis (Supplementary Table S4/5). Depicted in Fig. 7A, expressions of genes including *HcSWEET1b*, *HcSWEET4a*, *HcSWEET4b*, *HcSWEET5*, *HcSWEET13b*, *HcSWEET13c*, and *HcSWEET13d* were induced compared with the control group (0 h) under drought stress. On the contrary, *HcSWEET2*, *HcSWEET13a* and *HcSWEET14a* expressed in a down-regulation trend. Relative expression levels of two genes (*HcSWEET1a*, *HcSWEET14b*) first increased and then decreased, while *HcSWEET14c* exhibited a contrary expression pattern.

We further investigated expression levels of *HcSWEETs* under salt treatments (Fig. 7B). Five *HcSWEET* genes (*HcSWEET4a*, *HcSWEET5*, *HcSWEET11*, *HcSWEET13c*, *HcSWEET13d*) showed significantly elevated expression levels. By contrast, the expression of *HcSWEET2*, *HcSWEET6*, *HcSWEET13b*, and *HcSWEET14b* increased during different salt stresses. After salt treatments, there were 4 genes (*HcSWEET1b*, *HcSWEET14a*, *HcSWEET14c*, and *HcSWEET16*) initially up-regulating, subsequently decreasing again in expression levels. *HcSWEET4c*,

HcSWEET13a, *HcSWEET15* went down and then up. The various expression types showed that functional differences were likely to exist in *HcSWEET* genes in response to abiotic stresses.

To unveil the potential function of *HcSWEET* genes in responding to different stresses including drought and salinity, nine *HcSWEET* genes showing significant responses were selected and RNA-Seq results were further confirmed using RT-qPCR experiments (Fig. 8). *HcSWEET4a*, *HcSWEET4b*, and *HcSWEET5* showed an up-regulation in expressions under two stresses, notably, the expressions of *HcSWEET4a* and *HcSWEET5* significant increased after salt stresses, suggesting the potential possibilities of *HcSWEET4a* and *HcSWEET5* involved in salt stress. *HcSWEET2* exhibited a pattern of decreased expression compared with 0 h. Given whole wavelike patterns, the expressions of *HcSWEET1a*, *HcSWEET4a*, *HcSWEET4b*, *HcSWEET5*, *HcSWEET13c* had a sharp increase at the time point of 24–72 h, but decreased in 96–108 h throughout stresses. Overall, our analysis revealed that expression trends of 19 genes were consistent with those of the RNA-Seq and the majority of *HcSWEETs* presented a dynamic response pattern.

Subcellular localization of HcSWEET4a and HcSWEET5

HcSWEET4a and HcSWEET5 were predicted to function in the plasma membrane (Supplementary Table S1). To evaluate the validity of results, we performed co-expressed experiments in *N. benthamiana* leaf cells (Fig. 9). The fusion constructs pSuper:HcSWEET4a-GFP and pSuper:HcSWEET5-GFP were co-expressed with the plasma membrane marker (mCherry). Excluding empty vectors pSuper-GFP, GFP fluorescent signals were all exclusively observed in the plasma membrane. Results indicated that HcSWEET4a and HcSWEET5 belonged to plasma membrane-localized proteins.

Overexpression of HcSWEET4a in watermelon and response to salt stress

Previous experiments have implied that both *HcSWEET4a* and *HcSWEET5* exhibited significant up-regulation under salt stress. To further interpret the function of HcSWEETs in response to salt stress. We ectopically overexpressed alternative *HcSWEET4a* in watermelon through *Agrobacterium*-mediated transformation. Two transgenic lines (OE-1, OE-2) were obtained and expression levels were 3.67–4.18 times higher than the wild-type (WT) watermelon plant (Fig. 10B). Under NaCl treatment (150 mM) for 24 h, transgenic plants exhibited slightly curled leaf margins, while WT plants have been almost wilted (Fig. 10A). Phenotype observations further showed the up-regulation of *HcSWEET4a* might improve tolerance to salinity of plants.

Discussion

Hemerocallis citrina, belonging to Liliaceae *Hemerocallis*, is an important specialty crop widely cultivated around the world. With continuously expanded consumption patterns, night lily tends to be a model vegetable species for understanding deeply of perennial plant growth and development [38, 39]. Thereby, its tolerance to abiotic stress has become one of most key elements which influence production and quality. *SWEET* genes are plant-specific and taken as an essential class of sugar transporters in multiple plants [14, 40]. No studies were reported on the genome-wide identification of *HcSWEETs* or their function validation in perennial night lily previously. The present study investigated the response patterns of *HcSWEET* genes through genome-wide and expression analysis under drought and salt stress, aiming at promising new insights into enhancing stress resistances of night lily.

The study isolated 19 functional *HcSWEETs* successfully (Table 1), whose characteristics were similar to those reported in other plants especially *HfSWEETs* [41]. Secondary (Supplementary Table S1) and tertiary (Supplementary Figure S1) structure of HcSWEET proteins were predicted, revealing potential functions and

interactions between proteins [42]. HcSWEETs fell into four phylogenetic clades by bioinformatics analysis, which were uniform with HfSWEETs, AtSWEETs, VvSWEETs, and LcSWEETs [5, 41, 43–44]. There are not only similarities but also discrepancies in the members' number of the same clade, for instance, as was shown in (Fig. 1), nine members were found in Clade III contrast sharply to only one in AtSWEET and OsSWEET [45–48]. Phylogenetic analyses were supported by analyses of conserved motifs and gene structures. Conserved motif analysis indicated that most HcSWEETs harbored seven conserved transmembrane domains (TMDs), while genes within each clade also shared more unique motifs (Fig. 2A-B). Above conclusion is consistent with *Arabidopsis thaliana*, *Solanum lycopersicum*, and *Oryza sativa* [5, 49, 50]. Gene structure model demonstrated that *HcSWEETs* within each clade were similar in an exon-intron organization (Fig. 2D). For instance, each member of Clade I, III (except *HcSWEET13d*), and IV harbors 6 exons, whereas genes in Clade II contain 5 exons. Previous relative report in *Oryza sativa* indicated the loss-of-exon rate is larger than the gain-of-exon rate after undergoing segmental duplication [51]. Therefore it is proposed that all clades except Clades II might contain the original ancestor *HcSWEET* genes, and genes in Clade II might result from gene duplication and divergence.

HcSWEET genes were uniformly located in 9 chromosomes (Fig. 3). Intra-species collinear analysis identified four pairs of duplicated segments (Fig. 4A), and inter-species syntenic maps identified seven homologous genes in *Arabidopsis thaliana* but two in *Oryza sativa* (Fig. 4B). Therefore, we speculated that tandem or segmental duplication might result in species evolution by gene family expansion, consistent with other plant species reported [9, 48, 52]. The promoters of *HcSWEETs* mainly contain stress response elements, growth and development-related elements, hormone response elements (Fig. 5), suggesting that *HcSWEET* genes are indispensably in the process of various biological activities and coping with abiotic stresses of night lily. At the same time, some previous reports establish a theoretical foundation for further exploring gene functions of *HcSWEETs* [53–62].

SWEETs in the same or different species may have different cellular localizations [5, 17, 63]. AtSWEET4 in *Arabidopsis* is localized in the plasma membrane, mediating sugar transport to axial sinks as a hexose facilitator [64]. OsSWEET4 and ZmSWEET4c are located in the basal endosperm transfer layer membrane, transferring hexoses across BETL to sustain the development of large endosperm in cereal grains [65]. SWEET4 in grapevine as a glucose transporter, was also identified as a plasma membrane protein [44]. Sugar transporter gene *PbSWEET4* in pear, localizing in the plasma membrane, causes the reduction of sugar content and leads to early senescence of leaves [66]. In our present study, HcSWEET4a and HcSWEET5, falling in Clade II, were localized in the plasma membrane (Fig. 9). AtSWEET11 and close paralog AtSWEET12 localize to the plasma membrane of the phloem, suggesting the same localization also may exist in different clades [67].

At present, the SWEET family of transporters has been characterized as one of the key players in sugar translocation in higher plants [5, 6]. The major types of translocated sugar and their regulation in partition are necessary concerning plant development and stress responses [68, 69]. In divided four clades, genes in Clade I and II prefer hexose, Clade III sucrose, and Clade IV SWEETs likely fructose transporters in *Arabidopsis* [17, 67]. Expression patterns of genes are tightly correlated with functions, thus, tissue-specific analysis can be used to predict biological functions [70]. In this study, genes in the same clade tended to present the same expression pattern (Fig. 6). In Clade II, *HcSWEET4a*, *HcSWEET4b*, and *HcSWEET5*, highly expressing in roots except for *HcSWEET4c* and *HcSWEET6* in buds, might play central roles in mediating stress response as reported in *Arabidopsis* [64]. Our finding indicated that similar biological functions are performed more possibly in the same subfamily.

SWEETs in cabbage and *Medicago truncatula* have been reported in response to chilling stress [59, 71]. MaSWEETs are involved in abiotic/biotic stress responses in bananas [45]. *ZmSWEET* gene in maize responds to abiotic stress

[72]. The *SWEET* gene family in wheat response widely to abiotic stress [73]. Our RNA-seq data preliminary showed that expression levels of most *HcSWEETs* first increased after treatments for 24 h compared with 0 h, then changed towards different trends (Fig. 7), RT-qPCR assay further verified the expression levels of some differential expression genes (DEGs) in the *HcSWEET* gene family to ensure the reliability (Fig. 8). These experiments jointly suggested that most *HcSWEETs* might respond to drought and salt stress in different regulatory pathways with functional redundancy.

The members of the *SWEET* gene family may act as sugar transporters exposed under adverse environmental conditions [74, 75]. To elucidate precise roles of *HcSWEET4a* in night lily, ectopic expression of *HcSWEET4a* was conducted in watermelon. The results showed that up-regulated expression of *HcSWEET4a* in transgenic lines was distinct compared with WT plants (Fig. 10A). Phenotype observations indicated transgenic lines have greater resistance against salinity than WT plants. All data integrated in our present study, It is concluded that *HcSWEET4a* enhances salt tolerance by positively regulating sugar transport. However, the internal regulatory network remains poorly elaborated.

Conclusions

In summary, our present study first analyzed all extracted *SWEET* sequences in *Hemerocallis citrina*. Nineteen *HcSWEET* genes were selected as members comprehensively characterized, including basic properties, evolutionary relationships, conserved motifs, protein domains, gene structure analysis, chromosomal localization, gene duplication, and collinearity analysis. The phylogenetic tree classified the *HcSWEET* gene into four clades. Each clade had similar motif compositions and gene structures. Tissue-specific expression patterns of *HcSWEETs* and DEGs under drought and salt treatments suggested the function diversity and potential response mechanism to environments. Ectopic expression in watermelon further assured the response of *HcSWEET4a* to salt stress. Our research lays a foundation for casting light on the regulatory mechanism of *HcSWEET* genes in response to abiotic stress in detail.

Materials and methods

Identification and characterization of the *HcSWEET* genes in night lily

The whole genome sequences of night lily were downloaded from NCBI(https://www.ncbi.nlm.nih.gov/assembly/GCA_017893485.1_ASM1789348v1_genomic). To identify all members of *HcSWEET* genes, the protein sequence of *SWEETs* in *Hemerocallis citrina*, *Zea mays* and *Vitis vinifera* also were obtained from National Center for Biotechnology Information (NCBI), *Arabidopsis thaliana* from the Arabidopsis Information Resource (TAIR) [76], *Oryza sativa* from Ensembl Genome database [77], and *Hemerocallis fulva* from reported data previously [41], respectively. Functional genes with incomplete domains were supplemented and screened by searching conserved domains (CD-Search) [78]. Molecular weight (MV) and theoretical isoelectric points (PI) were detected using TBtools [79]. Transmembrane topology is predicted by Deep TMHMM Server v1.0.24. Subcellular localization analysis was preliminarily predicted via WoLF PSORT software online [80].

Secondary and tertiary structure predictions

The secondary structures of 19 *HcSWEET* proteins were predicted by the SOPMA tool online. The three-dimensional (3D) structures of these *HcSWEET* proteins were constructed using homology modeling. All 19 protein sequences

were aligned with all other proteins of the Protein Data Bank (PDB), and the optimum models were finally built with a standard of relatively high GMQE and seq identity values by Swiss-Model [81].

Phylogenetic tree construction, gene structure analysis, and prediction of conserved motifs and domains

Multiple Alignment Trimming in TBtools was used to carry out protein sequence alignments in night lily and other four plants. Based on the primary results of sequence alignment, trimAL Wrapper proceeded to have a shear to remove non-conserved regions as soon as possible. After shears were completed, a phylogenetic tree was constructed by One Step Build an ML Tree with 5000 bootstrap replicates. The visualization of gene structure was performed by Visualize Gene Structure of TBtools. MEME online and NCBI conserved domain database were respectively used to predict the conserved motifs and domains of the HcSWEETs.

Chromosomal location, collinearity, and cis-acting element (CREs) analysis

The genome annotation file in GFF format of *HcSWEETs* was obtained from the NCBI database. Both *Arabidopsis* and rice genomes were downloaded from the Ensembl Plants database. Intra-species and inter-species collinear analysis about night lily were constructed via MCScanX and TBtools software [79, 82]. A 2000 bp sequence in the promoter region was extracted utilizing TBtools from the genome. The online PlantCARE database was employed to predict the CREs of interest [83].

Plant materials and stress treatments

Night lily F1 hybrid population 116 (Dongzhuang Huanghua as female parent and Chonglihua as male parent) was used as treated plant material in this study, which was grown in the experimental base of Shanxi Agricultural University in Taigu under natural conditions and managed in the same regular agricultural practice. Florescence is concentrated in June to July and population 116 showed moderate resistance to abiotic stresses compared with the daylily cultivar 'Golden Doll'. Plants used for experimental treatments were from tissue cultivation. These seedlings through domestication were cultured in the same conditions for one month, after which drought and salt treatments were carried out. Drought and salt stress were respectively simulated using PEG6000 (10%) and NaCl (250 mM). Drought stresses were designed into 5 groups (0 h, 24 h, 48 h, 72 h, 96 h), and salt treatments were divided into 5 groups (0 h, 24 h, 48 h, 72 h, 108 h). Roots of treated plants were collected and immediately stored at -80°C for follow-up experiments.

Comparative transcriptome analysis

To analyze tissue-specific expression patterns, different tissues including tender root, mature root, bud, tender leaf, mature leaf, tender scape, and mature scape were collected. Total RNA of different tissues and treated roots was isolated using the TIANGEN RNAsimple Total RNA Kit, and RNA quality was evaluated by 1.5% (w/v) agarose gel. Samples with three biological replicates were selected and sent to BMK Biotechnology Company to conduct an RNA-seq assay. The average value of FPKM (reads per kilobase per million reads) was used to measure gene expression levels in the study. The heat map was drawn with $\log_2$ FPKM values to show different expression levels intuitively.

RT-qPCR analysis

Based on RNA-seq results, an RT-qPCR assay was performed for further verification of *HcSWEETs* expression levels with *HcACTIN* as the standard. Real-time quantitative PCR amplification was performed by PerfectStart® Green

SuperMix (TransGen, China). RT-qPCR procedure with triplicate was completed using a Real-Time PCR System (ABI QuantStudio 5, USA), and $\lg 2^{-\Delta\Delta C_t}$ values were taken as a measure of relative expression levels. SPSS 20 software was used to calculate statistical data and different letters represented significant differences between different treatments in accordance with Duncan's multiple test. All gene-specific primers are given in Supplementary Table S2.

Construction of HcSWEET4a transient expression vector and subcellular localization

The first-strand cDNA of night lily was synthesized using TransScript® One-Step gDNA Removal and cDNA Synthesis SuperMix (TransGen, China). Night lily WT cDNA was used as template to amplify the coding sequence without stop codon of *HcSWEET4a* using primers containing the XbaI/KpnI restriction sites. PCR product purified was sub-cloned into enzyme-digested pSuper1300-GFP vector by the homologous recombination method using Monclone Single Assembly Cloning Mix (Monad, China). The recombinant pSuper1300-HcSWEET4a-GFP was transformed into *Agrobacterium tumefaciens* strain GV3101. Tobacco leaf cells were transiently injected by the cultures. After 16–24 h, fluorescence images were captured under a fluorescence microscope (Leica STELLARIS 5, Germany). Relative primers are listed in Supplementary Table S2.

Overexpression vector construction and transformation of HcSWEET4 in watermelon

The PCR product with the KpnI and BamHI restriction sites was obtained by using binary pSuper1300-HcSWEET4a-GFP as template, then was cloned into 1305.4-GFP vector and sequenced. Constructed 1305.4-HcSWEET4a-GFP vector was transformed into watermelon YL using an optimized genetic transformation system [84]. Transgenic watermelon plants were screened by GFP fluorescence observation and PCR analysis. WT and positive plants were transplanted to the same growing conditions to cultivate. The drought and salt tolerances of transgenic plants were analyzed with WT watermelon plants as the negative control. Primers referred to in the experiment are given in Supplementary Table S2.

Declarations

Acknowledgments

We thank Prof. Zheng Li from College of Horticulture in Northwest A&F University for providing watermelon materials. We also thank intermediate laboratory technician Yi-Ping Wang from Experimental Teaching Center for offering technological assistance in subcellular localization.

Authors' contributions

S.L. and G.X. planned and designed the research. L.C., J.W., L.W., H.L., W.W., F.H., Y.L., Y.G. and X.C. performed experiments and analyzed the data. S.L., G.X., and L.C. wrote and revised the paper.

Funding

This study was supported by The first Shanxi postdoctoral innovation competition project, Shanxi provincial breeding joint research project (YZGG-10), Shanxi provincial key research and development project (202102140601009), Biological breeding engineering program of Shanxi Agricultural University (YZGC122).

Availability of data and materials

The data that support results are included in this manuscript and supplementary information files. Other relevant materials also are available from corresponding authors upon reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that there are no conflicts of interest.

References

1. Boyer JS. Plant productivity and environment. *Science*. 1982;218(4571):443–8.
2. EJD E, Norlyn JD, Rush DW, Kingsbury RW, Wrona AF. Saline culture of crops: a genetic approach. *Science*. 1980;210(4468):399–404.
3. Conde A, Neves A, Breia R, Pimenel D, Dinis LT, Bernardo S, Correia CM, Cunha A, Moutinho-Pereira J. Kaolin particle film application stimulates photoassimilate synthesis and modifies the primary metabolome of grape leaves. *J Plant Physiol*. 2018;1617(18):30026–9.
4. Conde A, Soares F, Breia R, Gerós H. Postharvest dehydration induces variable changes in the primary metabolism of grape berries. *Food Res Int*. 2018;105(03):261–70.
5. Chen LQ, Hou BH, Lalonde S, Takanaga H, Hartung ML, Qu XQ, Guo WJ, Kim JG, Underwood W, Chaudhuri B, et al. Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature*. 2010;468:527–32.
6. Broek VD, Gompel EV, Luttik MA, Pronk JT, Leeuwen CV. Mechanism of glucose and maltose transport in plasma-membrane vesicles from the yeast *Candida utilis*. *Biochem J*. 1997;321:487–95.
7. Zhang XH, Wang S, Ren Y, Gan CY, Li BB, Fan YYW, Zhao XQ, Yuan ZH. Identification, analysis and gene cloning of the *SWEET* gene family provide insights into sugar transport in Pomegranate (*Punica granatum*). *Int J Mol Sci*. 2022;23(5):2471.
8. Hamada M, Wada S, Kobayashi K, Satoh N. *Ci-Rga*, a gene encoding an MtN3/saliva family transmembrane protein, is essential for tissue differentiation during embryogenesis of the ascidian *Ciona intestinalis*. *Differentiation*. 2005;73(7):364–76.
9. Jia BL, Zhu XF, Pu ZJ, Duan YX, Hao LJ, Zhang J, Chen LQ, Jeon CO, Xuan YH. 2017. Integrative view of the diversity and evolution of SWEET and semiSWEET sugar transporters. *Front. Plant Sci*. 2017;8:2178.
10. Xuan YH, Hu YB, Chen LQ, Sosso D, Ducat DC, Hou BH, Frommer WB. Functional role of oligomerization for bacterial and plant SWEET sugar transporter family. *Proc Natl Acad Sci U S A*. 2013;110(39):E3685–94.
11. Mizuno H, Kasuga S, Kawahigashi H. The sorghum *SWEET* gene family: stem sucrose accumulation as revealed through transcriptome profiling. *Biotechnol Biofuels*. 2016;9:127.
12. Singh J, Das S, Jagadis Gupta K, Ranjan A, Foyer H, Kumar Thakur C. Physiological implications of SWEETs in plants and their potential applications in improving source-sink relationships for enhanced yield. *Plant Biotechnol J*. 2023;21(8):1528–41.

13. Singh Jeena G, Kumar S, Kumar Shukla R. Structure, evolution and diverse physiological roles of SWEET sugar transporters in plants. *Plant Mol Biol.* 2019;100(4–5):351–65.
14. Breia R, Conde A, Badim H, Margarida Fortes A, Geros H, Granell A. Plant SWEETs: from sugar transport to plant-pathogen interaction and more unexpected physiological roles. *Plant Physiol.* 2021;186(2):836–52.
15. Yuan M, Zhao JW, Huang RY, Li XH, Xiao JH, Wang SP. Rice MtN3/saliva/SWEET gene family: Evolution, expression profiling, and sugar transport. *J Integr Plant Biol.* 2014;56(6):559–70.
16. Isoda R, Palmai Z, Yoshinari A, Chen LQ, Tama F, Frommer B, Nakamura W. M. SWEET13 transport of sucrose, but not gibberellin, restores male fertility in *Arabidopsis sweet13;14*. *Proc Natl Acad Sci U S A.* 2022;119(42):e2207558119.
17. Klemens PAW, Patzke K, Deitmer J, Spinner L, Le Hir R, Bellini C, Bedu M, Chardon F, Krapp A, Ekkehard Neuhaus H. Overexpression of the vacuolar sugar carrier *AtSWEET16* modifies germination, growth, and stress tolerance in *Arabidopsis*. *Plant Physiol.* 2013;163(3):1338–52.
18. Yang J, Luo DP, Yang B, Frommer B, Eom W. SWEET11 and 15 as key players in seed filling in rice. *New Phytol.* 2018;218(2):604–15.
19. Wang X, Ju YH, Wu T, Kong LG, Yuan M, Liu HF, Chen XS, Chu ZH. The clade III subfamily of OsSWEETs directly suppresses rice immunity by interacting with OsHMGB1 and OsHsp20L. *Plant Biotechnol J.* 2024;14338:1–15.
20. Zhang XS, Feng CY, Wang MN, Li TL, Liu X, Jiang J. Plasma membrane-localized SISWEET7a and SISWEET14 regulate sugar transport and storage in tomato fruits. *Hortic Res.* 2021;8(1):186.
21. Zhang SH, Wang H, Wang T, Zhang J, Liu WJ, Fang HC, Zhang ZY, Peng FT, Chen XS, Wang N. Absciscic acid and regulation of the sugar transporter gene *MdSWEET9b* promote apple sugar accumulation. 2023;192(3):2081–2101.
22. Wang SD, Yokosho K, Guo R, Whelan J, Ruan YL, Ma JF, Shou HX. The Soybean Sugar Transporter GmSWEET15 Mediates Sucrose Export from Endosperm to Early Embryo. 2019;180(4):2133–41.
23. Wu YD, Di TM, Wu ZJ, Peng J, Wang J, Zhang KX, He MM, Li NN, Hao XY, Fang WP, Wang XC, Wang L. CsLHY positively regulates cold tolerance by activating *CsSWEET17* in tea plants. *Plant Physiol Biochem.* 2024;207:108341.
24. Fakher B, Arif Ashraf M, Wang LL, Wang XM, Zheng P, Aslam M, Qin Y. Pineapple SWEET10 is a glucose transporter. *Hortic Res.* 2023;10(10):uhad175.
25. Radchuk V, Belew ZM, Gundel A, Mayer S, Hilo A, Hensel G, Sharma R, Neumann K, Ortleb S, Wagner S et al. SWEET11b transports both sugar and cytokinin in developing barley grains. 2023;35(6):2186–207.
26. Nishad A, Kumar Nandi A. Recent advances in plant thermomemory. *Plant Cell Rep.* 2021;40(1):19–27.
27. Forni C, Duca D, Glick BR. Mechanisms of plant response to salt and drought stress and their alteration by rhizobacteria. *Plant Soil.* 2017;410:335–56.
28. Tiwari S, Lata C, Singh Chauhan P, Prasad V, Prasad M. 2017. A functional genomic perspective on drought signalling and its crosstalk with phytohormone-mediated signaling pathways in plants. *Curr Genom.* 2017;18:469–482.
29. Yang YQ, Guo Y. Elucidating the molecular mechanisms mediating plant salt-stress responses. *New Phytol.* 2018;217:523–39.
30. Muhammad M, Waheed A, Wahab A, Majeed M, Nazim M, Liu YH, Li L, Li WJ. Soil salinity and drought tolerance: An evaluation of plant growth, productivity, microbial diversity, and amelioration strategies. *Plant Stress.* 2024;11:100319.

31. Mathan J, Singh A, Ranjan A. Sucrose transport in response to drought and salt stress involves ABA-mediated induction of *OsSWEET13* and *OsSWEET15* in rice. *Physiol Plant*. 2021;171(4):620–37.
32. Monfared HH, Chew JK, Azizi P, Xue GP, Ee SF, Kadkhodaei S, Hedayati P, Ismail I, Zainal Z. Overexpression of a rice monosaccharide transporter gene (*OsMST6*) confers enhanced tolerance to drought and salinity stress in *Arabidopsis thaliana*. *Plant Mol Biol Rep*. 2020;38:151–64.
33. Kaur H, Manna M, Thakur T, Gautam V, Salvi P. Imperative role of sugar signaling and transport during drought stress responses in plants. *Physiol Plant*. 2021;171(4):833–48.
34. Ibraheem O, Dealtry G, Roux S, Bradley G. The effect of drought and salinity on the expressional levels of sucrose transporters in rice (*Oryza sativa* nipponbare) cultivar plants. *Plant Omics J*. 2011;4(2):68–74.
35. Redillas MC, Park SH, Lee JW, Kim YS, Jeong JS, Jung H, Jung H, Bang SW, Hahn TR, Kim JK. Accumulation of trehalose increases soluble sugar contents in rice plants conferring tolerance to drought and salt stress. *Plant Biotechnol Rep*. 2012;6:89–96.
36. Qing ZX, Liu JH, Yi XX, Liu XB, Hu GA, Lao J, He W, Yang ZH, Zou XY, Sun MS, et al. The chromosome-level *Hemerocallis citrina* Baroni genome provides new insights into the rutin biosynthesis and the lack of colchicine. *Hortic Res*. 2021;8(1):89.
37. Fatima U, Anjali A, Senthil-Kumar M. AtSWEET11 and AtSWEET12: the twin traders of sucrose. *Trends Plant Sci*. 2022;27(10):958–60.
38. Li S, Ji FF, Hou FF, Shi QQ, Xing GM, Chen H, Weng YQ, Kang XP. Morphological, palynological and molecular assessment of *Hemerocallis* core collection. *Sci Hortic*. 2021;285:110181.
39. Zuo GY, Li K, Guo YN, Niu XR, Yin LJ, Wu ZQ, Zhang XM, Cheng XJ, Yu J, Zheng SW, et al. Development and optimization of a rapid in vitro micropropagation system for the perennial vegetable night lily. *Hemerocallis citrina* Baroni Agron. 2024;14:244.
40. Yue WH, Cai KF, Xia X, Liu L, Wang JM. Genome-wide identification, expression pattern and genetic variation analysis of *SWEET* gene family in barley reveal the artificial selection of *HvSWEET1a* during domestication and improvement. *Front Plant Sci*. 2023;14:1137434.
41. Huang DM, Chen Y, Liu X, Bai NDA, Qin L. Genome-wide identification and expression analysis of the *SWEET* gene family in daylily (*Hemerocallis fulva*) and functional analysis of *HfSWEET17* in response to cold stress. *BMC plant Biol*. 2022;22(1):211.
42. Su H, Meng LJ, Qu ZC, Zhang W, Liu N, Cao PJ, Jin JJ. Genome-wide identification of the N6-methyladenosine regulatory genes reveals *NtFIP37B* increases drought resistance of tobacco (*Nicotiana tabacum* L). *BMC plant Biol*. 2024;24:134.
43. Xie HH, Wang D, Qin YQ, Ma AN, Fu JX, Qin YH, Hu GB, Zhao JT. Genome-wide identification and expression analysis of *SWEET* gene family in *Litchi chinensis* reveal the involvement of *LcSWEET2a/3b* in early seed development. *BMC Plant Biol*. 2019;19(1):499–512.
44. Chong J, Piron MC, Meyer S, Merdinoglu D, Bertsch C, Mestre P. The SWEET family of sugar transporters in grapevine: VvSWEET4 is involved in the interaction with *Botrytis cinerea*. *J Exp Bot*. 2014;65(22):6589–601.
45. Miao HX, Sun PG, Liu Q, Miao YL, Liu JH, Zhang KZ, Hu W, Zhang JB, Wang JY, Wang Z, et al. Genome-wide analyses of SWEET family proteins reveal involvement in fruit development and abiotic/biotic stress responses in banana. *Sci Rep*. 2017;7(1):3536.
46. Miao LM, Lv YX, Kong LJ, Chen QZ, Chen CQ, Li J, Zeng FH, Wang SY, Li JB, Huang L. Genome-wide identification, phylogeny, evolution, and expression patterns of MtN3/saliva/SWEET genes and functional analysis of BcNS in *Brassica rapa*. *BMC Genomics*. 2018;19(1):174.

47. Chandran D, Reinders A, Ward JM. Substrate specificity of the *Arabidopsis thaliana* sucrose transporter AtSUC2. *J Biol Chem*. 2003;278(45):44320–5.
48. Patil G, Valliyodan B, Deshmukh R, Prince S, Nicander B, Zhao MZ, Sonah H, Song L, Lin L, Chaudhary J, et al. Soybean (*Glycine max*) *SWEET* gene family: insights through comparative genomics, transcriptome profiling and whole genome re-sequence analysis. *BMC Genomics*. 2015;16:520.
49. Feng L, Frommer WB. Structure and function of SemiSWEET and SWEET sugar transporters. *Trends Biochem Sci*. 2015;40:480–6.
50. Yuan M, Wang S. Rice MtN3/saliva/SWEET family genes and their homologs in cellular organisms. *Mol Plant*. 2013;6:665–74.
51. Nuruzzaman M, Manimekalai R, Sharoni AM, Satoh K, Kondoh H, Ooka H, Kikuchi S. Genome-wide analysis of NAC transcription factor family in rice. *Gene*. 2010;465(1–2):30–44.
52. Gautam T, Saripalli G, Gahlaut V, Kumar A, Sharma PK, Balyan HS, Gupta PK. Further studies on sugar transporter (SWEET) genes in wheat (*Triticum aestivum* L). *Mol Biol Rep*. 2019;46:2327–53.
53. Lin IW, Sosso D, Chen LQ, Gase K, Kim SG, Kessler D, et al. Nectar secretion requires sucrose phosphate synthases and the sugar transporter SWEET9. *Nature*. 2014;508:546–9.
54. Guan YF, Huang XY, Zhu J, Gao JF, Zhang HX, Yang ZN. *Ruptured pollen grain1*, a member of the MtN3/saliva gene family, is crucial for exine pattern formation and cell integrity of microspores in *Arabidopsis*. *Plant Physiol*. 2008;147:852–63.
55. Sun MX, Huang XY, Yang J, Guan YF, Yang ZN. *Arabidopsis* RPG1 is important for primexine deposition and functions redundantly with RPG2 for plant fertility at the late reproductive stage. *Plant Reprod*. 2013;26:83–91.
56. Kanno Y, Oikawa T, Chiba Y, Ishimaru Y, Shimizu T, Sano N, Koshiba T, Kamiya Y, Ueda M, Seo M. AtSWEET13 and AtSWEET14 regulate gibberellin-mediated physiological processes. *Nat Commun*. 2016;7:13245.
57. Zhou Y, Liu L, Huang W, Yuan M, Zhou F, Li X, et al. Overexpression of *OsSWEET5* in rice causes growth retardation and precocious senescence. *PLoS ONE*. 2014;9(4):e94210.
58. Wang L, Yao LN, Hao XY, Li NN, Qian WJ, Yue C, Ding CQ, Zeng JM, Yang YJ, Wang XC. Tea plant SWEET transporters: expression profiling, sugar transport, and the involvement of CsSWEET16 in modifying cold tolerance in *Arabidopsis*. *Plant Mol Biol*. 2018;96:577–92.
59. Zhang W, Wang SY, Yu FW, Tang J, Shan X, Bao K, Yu L, Wang H, Fei ZJ, Li JB. Genome-wide characterization and expression profiling of SWEET genes in cabbage (*Brassica oleracea* var. *capitata* L.) reveal their roles in chilling and clubroot disease responses. *BMC Genomics*. 2019;20(1):93.
60. Sosso D, Luo D, Li QB, Sasse J, Yang J, Gendrot G, et al. Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport. *Nat Genet*. 2015;47:1489–93.
61. Guo C, Li H, Xia X, Liu X, Yang L. Functional and evolution characterization of SWEET sugar transporters in *Ananas comosus*. *Biochem Biophys Res Commun*. 2018;496:407–14.
62. Zhang Z, Zou LM, Ren C, Ren FR, Wang Y, Fan PG, Li SH, Liang ZC. *VvSWEET10* mediates sugar accumulation in grapes. *Genes*. 2019;10:255.
63. Chardon F, Bedu M, Calenge F, Klemens PAW, Spinner L, Clement G, Chietera G, Leran S, Ferrand M, Lacombeet B. Leaf fructose content is controlled by the vacuolar transporter SWEET17 in *Arabidopsis*. *Curr Biol*. 2013;23:697–702.
64. Liu XZ, Zhang Y, Yang C, Tian ZH, Li JX. AtSWEET4, a hexose facilitator, mediates sugar transport to axial sinks and affects plant development. *Sci Rep*. 2016;6:24563.

65. Sosso D, Luo DP, Li QB, Sasse J, Yang JL, Gendrot G, Suzuki M, Koch KE, McCarty DR, Chourey PS, et al. Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport. *Nat Genet.* 2015;47(12):1489–93.
66. Ni JP, Li JM, Zhu RX, Zhang MY, Qi KJ, Zhang SL, Wu J. Overexpression of sugar transporter gene *PbSWEET4* of pear causes sugar reduce and early senescence in leaves. *Gene.* 2020;743:144582.
67. Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, Fernie AR, Frommer WB. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science.* 2012;335(6065):207–11.
68. Wind J, Smeekens S, Hanson J. Sucrose: metabolite and signaling molecule. *Phytochemistry.* 2010;71:1610–4.
69. Xie HH, Wang D, Qin YG, Ma AN, Fu JX, Qin YH, Hu GB, Zhao JT. Genome-wide identification and expression analysis of *SWEET* gene family in *Litchi chinensis* reveal the involvement of *LcSWEET2a/3b* in early seed development. *BMC Plant Biol.* 2019;19(1):499.
70. Gao Y, Wang ZY, Kumar V, Xu XF, Yuan DP, Zhu XF, Li TY, Jia BL, Xuan YH. Genome-wide identification of the SWEET gene family in wheat. *Gene.* 2018;642:284–92.
71. Hu B, Wu H, Huang W, Song J, Zhou Y, Lin Y. *SWEET* gene family in *Medicago truncatula*: Genome-wide identification, expression and substrate specificity analysis. *Plants.* 2019;8(9):338.
72. Zhu JL, Zhou L, Li TF, Ruan YY, Zhang A, Dong XM, Zhu YS, Li C, Fan JJ. Genome-wide investigation and characterization of *SWEET* gene family with focus on their evolution and expression during hormone and abiotic stress response in maize. *Genes.* 2022;13(10):1682.
73. Qin JX, Jiang YJ, Lu YZ, Zhao P, Wu BJ, Li HX, Wang Y, Xu SB, Sun QX, Liu ZS. Genome-wide identification and transcriptome profiling reveal great expansion of *SWEET* gene family and their wide-spread responses to abiotic stress in wheat (*Triticum aestivum* L.). *J Integr Agric.* 2020;19(7):1704–20.
74. Valifard M, Le Hir R, Müller J, Scheuring D, Neuhaus HE, Pommerrenig B. Vacuolar fructose transporter SWEET17 is critical for root development and drought tolerance. *Plant Physiol.* 2021;187(4):2716–30.
75. Yao LN, Ding CQ, Hao XY, Zeng JM, Yang YJ, Wang XC, Wang L. CsSWEET1a and CsSWEET17 mediate growth and freezing tolerance by promoting sugar transport across the plasma membrane. *Plant Cell Physiol.* 2020;61(9):1669–82.
76. Rhee SY, Beavis W, Berardini TZ, Chen G, Dixon D, Doyle A, Garcia-Hernandez M, Huala E, Lander G, Montoya M, et al. The Arabidopsis Information Resource (TAIR): a model organism database providing a centralized, curated gateway to Arabidopsis biology, research materials and community. *Nucleic Acids Res.* 2003;31(1):2248.
77. Yates AD, Allen J, Amode RM, Azov AG, Barba M, Becerra A, Bhai J, Campbell LI, Carbajo MM, Chakiachvili M, et al. Ensembl genomes 2022: an expanding genome resource for non-vertebrates. *Nucleic Acids Res.* 2022;50(D1):D996–1003.
78. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI, Marchler GH, Song JS, et al. CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res.* 2020;48(D1):D265–8.
79. Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, Xia R. TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Mol Plant.* 2020;13(8):1194–202.
80. Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K. WoLF PSORT: protein localization predictor. *Nucleic Acids Res.* 2007;35(1):W585–7.
81. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer T, Rempfer C, Bordoli L, et al. SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res.* 2018;46(W1):W296–303.

82. Wang YP, Tang HB, Debarry JD, Tan X, Li JP, Wang XY, Lee TH, Jin HZ, Marler B, Guo H, et al. MCSscanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res.* 2012;40(7):e49.
83. Lescot M, Déhais P, Thijs G, Marchal K, Moreau Y, Peer YVD, Rouzé Pi, Rombautsets S, Plant CARE. a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res.* 2002;30:325–7.
84. Cao LH, Wei W, Shen JJ, Li Z, Xu ZM. Study on the optimization of transformation systems in watermelon. *Vegetable Res.* 2022;2:12.

Figures




Figure 1

Phylogenetic tree analysis and classification of the SWEETs in night lily and some other plants. Gene members of *Hemerocallis citrina* (Hc), *Hemerocallis fulva* (Hf), *Oryza sativa* (Os), *Zea mays* (Zm), *Arabidopsis thaliana* (At), *Vitis vinifera* (Vv) were classified into 4 clades. A phylogenetic tree was constructed via the TBtools software with 5000 bootstrap replicates.




Figure 2

Constructed phylogenetic tree, conserved motif, conserved domain, and gene structure analysis of SWEETs in *Hemerocallis citrina*. (A) Intraspecific evolutionary tree of 19 HcSWEET members. (B) The compositions and distribution of *HcSWEET*s conserved motifs. (C) The conserved functional domains of HcSWEETs. (D) Gene structures of *HcSWEET* genes.




Figure 3

Chromosome distributions of *SWEET* genes on night lily (LG1-11). All *HcSWEET* genes named are displayed on night lily LGs, with the LG number marked at the edge of each strip. The 0-500 Mb scale represents chromosome length. The lines inside the LGs represent gene density.




Figure 4

Collinearity of *HcSWEET* gene pairs. (A) Chromosome location and inter-chromosomal relationships of *SWEET*s in *Hemerocallis citrina*. The identified duplication events are marked by red lines. The LG number is labeled within different colored rectangles. (B) Collinearity analysis between the *SWEET*s of *Hemerocallis citrina* (*H. citrina*), *Arabidopsis thaliana* (*A. thaliana*), and *Oryza sativa* (*O. sativa*).



Figure 5
Distributions of cis-acting elements of the *HcSWEET* genes.




Figure 6
Expression dynamics of the *HcSWEET* genes in different tissues of night lily and phenotypes under different abiotic stresses. (A) Expression heatmap of *HcSWEET*s in the tender root, mature root, bud, tender leaf, mature leaf, tender scope, and mature scope. (B, C) The phenotypic observations of night lily under drought stress (B) and salt stress (C).




Figure 7

Expression heatmaps of *HcSWEETs* under drought (A) and salt (B) stresses based on RNA-seq data.




Figure 8

Expression patterns of 19 *HcSWEET* genes after drought and salt treatment. The different lowercase letters indicate significant differences by t-test ($p<0.05$).




Figure 9

Subcellular localization of *HcSWEET4a* and *HcSWEET5*. pSuper:GFP represents the negative control. The second panel represents a positive marker for the plasma membrane. The rightmost panel indicates the fusion of green fluorescence, red fluorescence, and bright field. Bar=25 µm.




Figure 10

Phenotype observations of *HcSWEET4a* overexpression lines under salt treatment. (A) Phenotype changes of transgenic lines (OE-1, OE-2) after treated for 0-24 h with NaCl (150 mM). Bar=17 mm. (B) The relative expression levels of *HcSWEET4a* in OE-1 and OE-2 lines. The significance of difference is marked by two asterisks ($p<0.05$).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFiguresandTables.docx](#)
- [SupplementaryInformation1.docx](#)
- [SupplementaryInformation2.doc](#)