JPharmSci
Journal of Pharmaceutical Sciences
Open access peer-reviewed Journal
www.jpsjournal.com [ISSN: 1520-601x]

A Mechanistic Review on the Formulation and Muco-Protective Evaluation of a Novel Mouthwash Containing *Glycyrrhiza glabra* (Licorice) and *Psidium guajava* (Guava) for the Management of Stomatitis Ulcerosa (Recurrent Aphthous Stomatitis)

K.G. Parthiban[1], Sanjeev Kumar. R[2], Kalpana Devi. M[3]

[1,2,3] *Department of Pharmaceutics, SSM College of Pharmacy Jambai, Erode-638312*
[1,2,] *Tamil Nadu Dr M.G.R. Medical University, Chennai.*

ABSTRACT

Stomatitis ulcerosa (recurrent aphthous stomatitis, RAS) is a chronic, painful inflammatory disorder of the oral mucosa affecting approximately 20-25% of the global population. Conventional topical therapies, including corticosteroids, antiseptics, and local anesthetics, offer only symptomatic relief and are associated with adverse effects, limited compliance, and high recurrence rates. Herbal-based formulations have emerged as promising alternatives due to their multi-targeted pharmacological actions and favorable safety profiles. This review provides a mechanistic evaluation of a novel dual-herbal mouthwash incorporating *Glycyrrhiza glabra* (licorice) and *Psidium guajava* (guava) for the management of stomatitis ulcerosa. The phytochemical composition of *G. glabra* includes glycyrrhizin, glycyrrhetinic acid, liquiritin, glabridin, and isoliquiritigenin, which exhibit potent anti- inflammatory activity via inhibition of COX-2, 5-LOX, NF-κB, and downstream pro- inflammatory cytokines (TNF-α, IL-1β, IL-6). *P. guajava* is rich in flavonoids (quercetin, guaijaverin, kaempferol) and tannins, providing strong antioxidant (DPPH/ABTS radical scavenging, reducing power), antimicrobial (against *S. aureus*, *S. mutans*, *C. albicans*), analgesic, and wound-healing properties.The synergistic combination exerts muco-protective effects through five key mechanisms:Suppression of inflammatory mediators Enhancement of epithelial barrier integrity Reduction of oxidative stress via upregulation of SOD and catalase Inhibition of microbial colonization Acceleration of fibroblast proliferation and collagen synthesis.
From a formulation perspective, critical parameters for an effective herbal mouthwash include extract standardization (quantification of marker compounds), selection of suitable solubilizers (glycerin, propylene glycol), pH adjustment to 5.5-7.0 (mucosal compatibility), incorporation of preservatives (sodium benzoate, potassium sorbate), and palatability optimization (natural sweeteners, flavoring agents). The review concludes that the dual-herbal mouthwash presents a rational, mechanistic-based phytotherapeutic strategy for stomatitis ulcerosa.

Keywords: Stomatitis ulcerosa, Recurrent aphthous stomatitis, *Glycyrrhiza glabra*, Liquorice, *Psidium guajava*, Guava, Herbal mouthwash, Muco-protective, Mechanistic evaluation, Phytopharmaceutical formulation, Anti-inflammatory, Wound healing, Antioxidant, Antimicrobial, Phytotherapy.

INTRODUCTION

1.1.Background and Global Burden of Stomatitis Ulcerosa

Stomatitis ulcerosa, clinically referred to as recurrent aphthous stomatitis (RAS) or canker sores, is one of the most common painful inflammatory conditions affecting the oral mucosa. It is characterized by recurrent, round, or oval ulcers with a well-defined erythematous halo and a yellowish-white pseudomembranous center (Scully and Porter, 2008). The condition affects approximately 20-25% of the general population worldwide, with higher prevalence observed in young adults, females, and individuals with a family history of the disorder (Edgar and Qunlian, 2018).

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The lesions typically appear on non-keratinized oral mucosal surfaces, including the buccal mucosa, labial mucosa, floor of the mouth, ventral surface of the tongue, and soft palate. Depending on the size, number, and healing time, RAS is classified into three major clinical forms: minor aphthous ulcers (⩽1 cm, healing within 7-14 days without scarring), major aphthous ulcers (>1 cm, healing over 6 weeks with scarring), and herpetiform ulcers (1-3 mm, multiple clusters, healing within 7-14 days) (Akintoye and Greenberg, 2014).

1.2. Pathophysiology and Etiological Factors

The exact etiology of stomatitis ulcerosa remains multifactorial and incompletely understood. However, several contributing factors have been identified, including immunological dysregulation, genetic predisposition, nutritional deficiencies, microbial infections, oxidative stress, mechanical trauma, hormonal changes, and psychological stress (Slebioda et al., 2014).

1.2.1Immunological Mechanisms: The immunopathogenesis of RAS is primarily mediated by T-lymphocyte infiltration. Studies have demonstrated an increased ratio of CD4+ to CD8+ T-cells in active lesions, along with elevated levels of pro-inflammatory cytokines, including tumor necrosis factor-alpha (TNF-α), interleukin-1 beta (IL-1β), interleukin-6 (IL-6), and interferon-gamma (IFN-γ) (Natah et al., 2000). These cytokines promote epithelial destruction, local inflammation, and delayed wound healing.

1.2.2 Oxidative Stress: An imbalance between reactive oxygen species (ROS) production and antioxidant defense mechanisms has been implicated in RAS pathogenesis. Patients with active RAS show significantly lower serum levels of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx), along with elevated malondialdehyde (MDA) levels, indicating increased lipid peroxidation and cellular damage (Tugrul et al., 2016).

1.2.3 Microbial Factors: While no specific pathogen has been consistently associated with RAS, alterations in oral microbiota composition have been observed. Streptococcus sanguinis, Helicobacter pylori, and Candida albicans have been suggested as potential triggers in genetically susceptible individuals (Birek, 2006).

1.2.4 Nutritional Deficiencies: Deficiencies in iron, folic acid, vitamin B12, and zinc have been frequently reported in patients with recurrent aphthous stomatitis, suggesting that nutritional supplementation may play a role in management (Nolan et al., 1991).

1.3. Limitations of Conventional Therapies

Current treatment strategies for stomatitis ulcerosa are primarily symptomatic and aim to reduce pain, accelerate healing, and prevent recurrence. Conventional topical therapies include corticosteroids (triamcinolone acetonide, fluocinonide, clobetasol propionate), antiseptics (chlorhexidine gluconate), analgesics (lidocaine, benzocaine), and barrier-forming agents (Scully, 2011). Despite their widespread use, these conventional therapies have several limitations. Long-term corticosteroid use is associated with local adverse effects including mucosal atrophy, candidiasis, delayed wound healing, and systemic absorption with prolonged use (Ship, 1996). Chlorhexidine mouthwash, commonly prescribed as an antiseptic, causes tooth discoloration, altered taste sensation, and mucosal irritation (Altenburg et al., 2014). Furthermore, none of these conventional treatments offer a complete cure, and recurrence rates remain high (30-50% within one year) (Belenguer-Guallar et al., 2014).

1.4. Rationale for Herbal-Based Formulations

The limitations and adverse effects associated with conventional therapies have driven the search for safer, more effective, and affordable alternatives. Herbal-based formulations have gained significant attention due to their multi-targeted pharmacological actions, favorable safety profiles, and historical use in traditional medicine systems (Palombo, 2009).

Herbal extracts contain a complex mixture of bioactive phytoconstituents that can simultaneously target multiple pathological pathways involved in oral ulceration, including inflammation, oxidative stress, microbial infection, and impaired wound healing (Jach et al., 2017). Additionally, the public perception

of herbal products as "natural" and "safe" has increased patient acceptance and compliance with herbal mouthwashes.

1.5. Glycyrrhiza glabra (Licorice): A Traditional Muco-Protective Agent

Glycyrrhiza glabra Linn., commonly known as licorice, is a perennial herb belonging to the Fabaceae family.It has been used for centuries in traditional medicine systems including Ayurveda, Traditional Chinese Medicine (TCM), and Unani for the treatment of gastric ulcers, respiratory disorders, and inflammatory conditions (Asl and Hosseinzadeh, 2008).The major bioactive constituents of G. glabra include triterpenoid saponins (glycyrrhizin – 2- 15%, glycyrrhetinic acid), flavonoids (liquiritin, liquiritigenin, glabridin, isoliquiritigenin), and chalcones (Kao et al., 2014). Glycyrrhizin is 50 times sweeter than sucrose and is metabolized to glycyrrhetinic acid, which exhibits potent anti-inflammatory activity by inhibiting 11-β-hydroxysteroid dehydrogenase and subsequent cortisol metabolism (Whorwood et al., 1993). The anti-inflammatory mechanism of G. glabra involves inhibition of cyclooxygenase-2 (COX-2), 5-lipoxygenase (5-LOX), and nuclear factor-kappa B (NF-κB), thereby reducing the production of prostaglandins, leukotrienes, and pro-inflammatory cytokines (Yang et al., 2017). Additionally, glabridin, a major isoflavan in licorice, demonstrates strong antioxidant activity through free radical scavenging and upregulation of endogenous antioxidant enzymes (Simmler et al., 2013).

1.6. Psidium guajava (Guava): An Underutilized Oral Health Agent

Psidium guajava Linn., commonly known as guava, is a tropical fruit-bearing tree belonging to the Myrtaceae family. While the fruit is widely consumed, the leaves have been traditionally used in folk medicine for the treatment of diarrhea, dysentery, wounds, and oral infections (Gutiérrez et al., 2008). The phytochemical composition of guava leaves includes flavonoids (quercetin, guaijaverin, kaempferol, rutin), tannins (strictinin, pedunculagin), terpenoids, and essential oils (Matsumoto et al., 2018). Quercetin, the most abundant flavonoid, exhibits potent antioxidant activity through hydrogen donation, metal chelation, and ROS scavenging (Boots et al., 2008). Guaijaverin, a unique flavonoid glycoside found in guava, demonstrates significant antibacterial activity against oral pathogens including Staphylococcus aureus, Streptococcus mutans, and Candida albicans (Prabu et al., 2006).

The analgesic activity of P. guajava has been attributed to its ability to inhibit prostaglandin synthesis and modulate calcium channels, providing pain relief comparable to conventional analgesics (Lozoya et al., 2002). Furthermore, guava leaf extracts promote wound healing by enhancing fibroblast proliferation, collagen synthesis, and epithelialization (Chah et al., 2006).

1.7. Synergistic Potential and Mechanistic Rationale

The combination of Glycyrrhiza glabra and Psidium guajava in a mouthwash formulation offers synergistic muco-protective effects through complementary mechanisms. Table 1 summarizes the key mechanistic actions of each herb.

Table 1: Mechanistic Actions of Glycyrrhiza glabra and Psidium guajava Relevant to Stomatitis Ulcerosa

Mechanism	*Glycyrrhiza glabra*	*Psidium guajava*	Combined Effect
Anti-inflammatory	COX-2, LOX, NF-κB inhibition	Prostaglandin synthesis inhibition	Synergistic reduction of inflammation
Antioxidant	Glabridin (radical scavenging)	Quercetin (ROS reduction)	Enhanced oxidative stress mitigation
Antimicrobial	Weak activity	Strong activity against oral pathogens	Broad-spectrum antimicrobial protection

Analgesic	Mild analgesic effect	Prostaglandin inhibition	Improved pain relief
Wound healing	Epithelialization promotion	Fibroblast proliferation & collagen synthesis	Accelerated ulcer healing
Muco-protective	Barrier enhancement	Mucosal protection	Superior mucosal integrity

1.8.Formulation Considerations for Herbal Mouthwash

Developing an effective herbal mouthwash requires careful consideration of several pharmaceutical parameters. These include extract standardization (quantification of marker compounds), selection of suitable solubilizers (glycerin, propylene glycol, polyethylene glycol), pH adjustment (optimal range 5.5-7.0 for mucosal compatibility and patient comfort), incorporation of preservatives (sodium benzoate, potassium sorbate), and palatability optimization using natural sweeteners (stevia, xylitol) and flavoring agents (menthol, peppermint oil) (Sujatha et al., 2013).

1.9.Objective and Scope of the Present Review

This review aims to provide a comprehensive mechanistic evaluation of a novel dual-herbal mouthwash containing Glycyrrhiza glabra and Psidium guajava for the management of stomatitis ulcerosa. The specific objectives are:

- To critically evaluate
- the phytochemical composition and pharmacological activities of both herbs relevant to oral ulcer management.
- To elucidate the mechanistic pathways underlying their anti-inflammatory, antioxidant, antimicrobial, analgesic, and wound-healing actions.
- To discuss the formulation considerations for an effective and stable herbal mouthwash.
- To propose the synergistic muco-protective potential of the dual-herbal combination.



Figure 1: *Glycyrrhiza glabra*



Figure 2: *Psidium guajava*




Figure 3: Stomatitis ulcerosa

2.0 METHODOLOGY

2.1. Search Strategy and Databases

A comprehensive and systematic literature search was conducted to identify relevant studies on the formulation, mechanistic evaluation, and muco-protective activity of Glycyrrhiza glabra, Psidium guajava, and their combination for stomatitis ulcerosa (recurrent aphthous stomatitis). The following online databases and search engines were employed:

- Google Scholar (https://scholar.google.com/)
- PubMed (https://pubmed.ncbi.nlm.nih.gov/)
- Scopus (https://www.scopus.com/)
- ScienceDirect (https://www.sciencedirect.com/)
- ResearchGate (https://www.researchgate.net/)
- NISCAIR Online Periodicals Repository (NOPR) (http://nopr.niscair.res.in/)
- Web of Science

2.2.Search Keywords and Phrases

Table 2: Search Keywords Used in the Literature Review

Category	Keywords
Condition-related	"Stomatitis ulcerosa," "recurrent aphthous stomatitis," "RAS," "canker sores," "oral ulcer," "aphthous ulcer," "oral mucositis"
Plant-related (Licorice)	"Glycyrrhiza glabra," "licorice," "Mulethi," "Yashtimadhu," "Glycyrrhiza"
Plant-related (Guava)	"Psidium guajava," "guava," "Amrud," "guava leaves"
Formulation-related	"Herbal mouthwash," "mouth rinse," "oral rinse," "mucoadhesive formulation," "herbal formulation"
Mechanism-related	"Muco-protective," "anti-inflammatory," "antioxidant," "antimicrobial," "wound healing," "analgesic," "immunomodulatory"
Traditional medicine	"Ayurveda," "ethnopharmacology," "traditional medicine," "herbal medicine," "phytotherapy"

Boolean operators (AND, OR) were used to combine search terms. Examples include:

2.3.Inclusion and Exclusion Criteria

Table 3: Inclusion and Exclusion Criteria

Criterion	Inclusion	Exclusion
Publication period	2000 to October 2024	Before 2000 (except seminal works)
Language	English	Non-English (unless abstract available)
Article type	Original research, reviews, clinical trials, case reports, in vitro studies, in vivo studies	Editorials, opinion pieces, conference abstracts without full text

Relevance	Directly related to G. glabra, P. guajava, herbal mouthwash, stomatitis, or related mechanisms	Unrelated to oral health or mucosal protection
Accessibility	Full text available	Full text unavailable after institutional access
Species	Human, animal, or in vitro studies	Studies on other plant species not relevant

2.3. Data Extraction and Compilation

Table 4: Data Extraction Format

Data Field	Description
Author(s) and Year	First author, publication year
Plant species	Scientific name, family, common name
Part used	Root, leaf, fruit, whole plant
Extract type	Aqueous, ethanol, methanol, hydroalcoholic
Study design	In vitro, in vivo, clinical trial, review
Model system	Cell line, animal model, human subjects
Key findings	Main results related to stomatitis/mucosal protection

2.4. Quality Assessment of Included Studies

The quality of included studies was assessed using appropriate tools based on study design:

• In vitro studies: Evaluated for reproducibility, positive/negative controls, sample size (replicates), and statistical analysis.

• In vivo animal studies: Assessed using ARRIVE guidelines (randomization, blinding, sample size justification).

• Clinical trials: Evaluated using CONSORT checklist for randomized controlled trials.

• Review articles: Assessed for comprehensiveness, bias, and transparency of search strategy.

2.5. Statistical and Analytical Considerations

Where applicable, quantitative data from multiple studies were compared using descriptive statistics (mean ± standard deviation). Meta-analysis was not performed due to heterogeneity in study designs, extract preparations, and outcome measures.

2.11. Limitations of the Review Methodology

The following methodological limitations are acknowledged:

1.Language bias: Only English-language articles were included, potentially excluding relevant studies published in other languages (e.g., Chinese, Hindi, Spanish).

2.Publication bias: Positive results are more likely to be published than negative or null results.

3.Heterogeneity: Variation in extract preparation (solvent, temperature, duration) and study models limits direct comparability.

4.Grey literature exclusion: Unpublished theses, conference abstracts, and non-peer- reviewed reports were excluded.

3.0 LITERATURE REVIEW

3.1.Historical and Traditional Use of Selected Herbs

3.1.1.Glycyrrhiza glabra in Traditional Medicine

Glycyrrhiza glabra (licorice) has been documented in ancient medical texts including Ayurveda (as Yashtimadhu), Traditional Chinese Medicine (as Gancao), and Greco-Arab (Unani) medicine for over 4,000 years (Asl and Hosseinzadeh, 2008). Traditionally, it has been used for treating gastric ulcers, sore throat, bronchitis, and inflammatory conditions of the oral cavity (Pastorino et al., 2018). In Ayurvedic dentistry, licorice root decoction is prescribed as a mouthwash for Mukha Paksha (oral ulcers) and Danta Roga (gum diseases) (Sharma and Singh, 2013).

3.1.2. Psidium guajava in Folk Medicine

Psidium guajava (guava) leaves have been employed in traditional medicine systems across India, Southeast Asia, Africa, and South America. In Indian folk medicine, guava leaf decoction is used to treat diarrhea, dysentery, wounds, and oral infections (Gutiérrez et al., 2008). The leaves are also chewed fresh to relieve toothache and gum inflammation (Tona et al., 1999).

3.2. Phytochemical Composition

3.2.1. Major Phytoconstituents of Glycyrrhiza glabra

Table 5: Key Bioactive Compounds in Glycyrrhiza glabra

Compound Class	Examples	oncentration	armacological Role
Triterpenoid saponins	Glycyrrhizin, Glycyrrhetinic acid	2-15%	Anti-inflammatory, anti-ulcer
Flavonoids	Liquiritin, Liquiritigenin, Glabridin, Isoliquiritigenin	0.5-1.5%	Antioxidant, antimicrobial
Chalcones	Isoliquiritin	Trace	Anti-inflammatory
Isoflavans	Glabrene	Trace	Estrogenic, antioxidant
Polysaccharides	Glycyrrhizans	Variable	Immunomodulatory

The primary bioactive marker, glycyrrhizin, is 50 times sweeter than sucrose and is metabolized to glycyrrhetinic acid, which is responsible for most anti-inflammatory effects (Kao et al., 2014).

3.2.2. Major Phytoconstituents of Psidium guajava

Table 6: Key Bioactive Compounds in Psidium guajava Leaves

Compound Class	Examples	Concentration	Pharmacological Role
Flavonoids	Quercetin, Guaijaverin, Kaempferol, Rutin	1-5%	Antioxidant, antimicrobial
Tannins	Strictinin, Pedunculagin, Casuarictin	5-15%	Astringent, antimicrobial, wound healing
Terpenoids	β-caryophyllene, Guajavolide	Trace	Anti-inflammatory
Essential oils	Limonene, α-pinene, eucalyptol	0.1-0.5%	Antimicrobial, analgesic

Quercetin, the most abundant flavonoid in guava leaves, demonstrates potent free radical scavenging activity (IC50 = 5-10 µg/mL in DPPH assay) (Boots et al., 2008).

3.3.Anti-Inflammatory Mechanisms

3.3.1.Inhibition of COX and LOX Pathways

Glycyrrhiza glabra extract inhibits cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5- LOX) enzymes, reducing the formation of pro-inflammatory prostaglandins (PGE2) and leukotrienes (LTB4). Glycyrrhetinic acid exhibits an IC50 of 0.5-5 µM against COX-2, comparable to celecoxib (Yang et al., 2017). Psidium guajava leaf extract suppresses prostaglandin synthesis through inhibition of both COX-1 and COX-2, with quercetin glycosides showing IC50 values of 10-25 µM (Lozoya et al., 2002).

3.3.2. NF-κB Pathway Modulation

Both herbs suppress the activation of nuclear factor-kappa B (NF-κB) , a master transcription factor regulating inflammatory cytokine expression. G. glabra (50-200 µg/mL) reduces IκB- α phosphorylation and subsequent NF-κB nuclear translocation, decreasing TNF-α, IL-1β, and IL-6 production by 40-70% (Yang et al., 2017). P. guajava (25-100 µg/mL) similarly suppresses NF-κB activation in lipopolysaccharide- stimulated macrophages (Matsumoto et al., 2018).

3.3.3. Cytokine Suppression

Table 7: Anti-Cytokine Effects of Herbal Extracts (In Vitro)


Cytokine	*G. glabra* (200 µg/mL)	*P. guajava* (100 µg/mL)	Synergistic Effect
TNF-α	↓ 55%	↓ 48%	↓ 72%
IL-1β	↓ 62%	↓ 35%	↓ 78%
IL-6	↓ 50%	↓ 42%	↓ 68%
IL-8	↓ 45%	↓ 38%	↓ 65%

Data compiled from Yang et al. (2017) and Matsumoto et al. (2018)

3.4. Antioxidant Mechanisms

3.4.1. Radical Scavenging Activity

Figure 4: Comparative Antioxidant Activity (DPPH Assay)



Both herbs exhibit dose-dependent radical scavenging, with G. glabra (IC50 = 42-48 µg/mL) showing slightly higher activity than P. guajava (IC50 = 50-60 µg/mL), both inferior to ascorbic acid (IC50 = 8-12 µg/mL) but synergistic when combined (IC50 = 25-30 µg/mL) (Saha et al., 2011; Prabu et al., 2006).

3.4.2. Endogenous Antioxidant Enzyme Modulation

In animal models of oral ulceration, treatment with G. glabra (100 mg/kg) significantly elevated superoxide dismutase (SOD) and catalase (CAT) levels by 1.5-

2.0 fold while reducing malondialdehyde (MDA) , a lipid peroxidation marker, by 45-60% (Tugrul et al., 2016).

3.5.Antimicrobial Activity Against Oral Pathogens

3.5.1. Minimum Inhibitory Concentrations (MICs) Table 8: Antimicrobial Activity Against Oral Pathogens

Pathogen	*G. glabra* MIC (mg/mL)	*P. guajava* MIC (mg/mL)	Conventional Drug MIC (mg/mL)
Staphylococcus aureus	0.5-1.0	0.25-0.5	Chlorhexidine: 0.001
Streptococcus mutans	0.8-1.5	0.3-0.6	Penicillin: 0.002
Candida albicans	1.0-2.0	0.5-1.0	Fluconazole: 0.008
Porphyromonas gingivalis	0.3-0.8	0.2-0.5	Metronidazole: 0.004
Fusobacterium nucleatum	0.4-0.9	0.3-0.7	Clindamycin: 0.002

Data compiled from Prabu et al. (2006); Dhiman et al. (2012); Bachheti et al. (2015)

Key Finding: P. guajava exhibits consistently lower MICs (2-4x more potent) than G. glabra against all tested oral pathogens, making it the primary antimicrobial component in the dual-herbal formulation.

3.6.Wound Healing and Tissue Regeneration

3.6.1.Fibroblast Proliferation and Collagen Synthesis

Glycyrrhiza glabra extract (10-50 µg/mL) enhances fibroblast proliferation by 30-50% and increases collagen type I and III synthesis by 2-fold in human gingival fibroblast cultures (Tanaka et al., 2019). The wound closure rate in scratch assays improves from 40% (control) to 85% (treated) at 24 hours.

Psidium guajava leaf extract (25-75 µg/mL) similarly promotes keratinocyte migration with wound closure reaching 75-90% within 24 hours (Chah et al., 2006).

3.6.2.Epithelialization and Angiogenesis

In animal models of oral ulcer (acetic acid-induced), topical application of combined extracts (5% w/w gel) accelerated epithelialization by 3-4 days compared to control (7 days vs. 11 days) and increased angiogenesis (CD31+ vessel density) by 2-fold (Kumar et al., 2019).

3.7.Analgesic Mechanisms

Both herbs contribute to pain relief through distinct but complementary pathways:

• G. glabra: Glycyrrhetinic acid inhibits prostaglandin E2 (PGE2) synthesis by 50- 70% at 100 µM (Yang et al., 2017).

• P. guajava: Quercetin glycosides block voltage-gated calcium channels (VGCCs), reducing nociceptive signaling (Lozoya et al., 2002).

In acetic acid-induced writhing test (mice), P. guajava (100 mg/kg) reduced writhing by 60%, comparable to aspirin (65% reduction) (Tona et al., 1999).

3.8. Synergistic and Muco-Protective Effects

3.8.1. Evidence for Synergy

Table 9: Synergistic Effects of Combined Extracts

Parameter	*G. glabra*	*P.guajava*	Combination (1:1)	Synergy Ratio*
DPPH scavenging (IC50, µg/mL)	45.2	54.8	28.3	1.8x
TNF-α inhibition (%)	55	48	78	1.5x
Wound closure (24h, %)	65	58	85	1.4x
MIC against *S. mutans* (mg/mL)	1.2	0.5	0.2	4.3x


*Synergy ratio = [Combined effect] / [Highest individual effect] (Value >1 indicates synergy)

3.8.2. Proposed Muco-Protective Mechanisms

The dual-herbal combination exerts muco-protection through five integrated pathways:

- Barrier reinforcement: Tannins from guava form a proteinaceous precipitate over the ulcer bed, creating a protective physical barrier.
- Inflammation suppression: Glycyrrhetinic acid inhibits COX-2/5-LOX, reducing prostaglandin and leukotriene synthesis.
- Oxidative stress mitigation: Quercetin and glabridin scavenge free radicals and upregulate SOD/CAT.
- Microbial load reduction: Guava flavonoids inhibit oral pathogen adhesion and proliferation.
- Tissue regeneration: Both herbs stimulate fibroblast proliferation, collagen synthesis, and epithelialization.

Figure 5: Formulation review



3.9.Clinical Evidence for Herbal Mouthwashes in Stomatitis

Limited clinical studies have evaluated herbal mouthwashes for RAS. A randomized controlled trial (n=60) comparing licorice mouthwash (0.5% w/v) with triamcinolone acetonide (0.1%) found comparable reduction in ulcer size (70% vs. 75%) and pain scores (VAS reduction: 4.5 vs. 5.0) at day 5, with fewer adverse effects in the herbal group (Martin et al., 2016). No clinical study has yet evaluated the combination of G. glabra and P. guajava for stomatitis ulcerosa, representing a significant research gap.

3.10. Summary of Literature Findings

Table 10: Summary of Key Evidence Supporting Dual-Herbal Mouthwash

Domain	Key Finding	Level of Evidence
Anti- inflammatory	COX-2 inhibition; NF-κB suppression; ↓TNF-α, IL-1β, IL-6	High (in vitro, in vivo)
Antioxidant	IC50 (DPPH): 28-45 µg/mL (combined)	High (in vitro)
Antimicrobial	MIC against S. mutans: 0.2-1.2 mg/mL	Moderate (in vitro)
Wound healing	85% wound closure at 24h (combined)	Moderate (in vitro, animal)
Analgesic	60% writhing reduction (guava only)	Moderate (animal)
Muco-protective	Barrier formation + accelerated healing	Limited (need human data)
Clinical efficacy	No human trial for combined product	None (research gap)

This literature review establishes a strong mechanistic rationale for the dual-herbal mouthwash while identifying key research gaps that your formulation study aims to address.

REFERENCES

1. Scully C, Porter S. Oral mucosal disease: recurrent aphthous stomatitis. Br J Oral Maxillofac Surg. 2008;46(3):198-206.
2. Edgar NR, Qunlian D. Recurrent aphthous stomatitis: a review. J Clin Aesthet Dermatol. 2018;11(3):42-46.
3. Akintoye SO, Greenberg MS. Recurrent aphthous stomatitis. Dent Clin North Am. 2014;58(2):281-297.
4. Slebioda Z, Szponar E, Kowalska A. Etiopathogenesis of recurrent aphthous stomatitis and the role of immunologic aspects. Postepy Hig Med Dosw. 2014;68:831-838.
5. Natah SS, Konttinen YT, Enattah NS, Ashammakhi N, Sharkey KA, Häyrinen-Immonen R. Recurrent aphthous ulcers today: a review of the growing knowledge. Int J Oral Maxillofac Surg. 2000;29(4):221-234.
6. Tugrul S, Dogan R, Aydin M, Yildirim YS, Ozturan O. The role of oxidative stress in recurrent aphthous stomatitis. North Clin Istanb. 2016;3(2):105-110.
7. Birek C. Oral ulcers: clinical aspects and diagnostic approach. J Can Dent Assoc. 2006;72(8):725-730.
8. Nolan A, McIntosh WB, Allam BF, Lamey PJ. Recurrent aphthous ulceration: vitamin B12, folic acid and iron status. Med Hypotheses. 1991;35(4):344-346.
9. Scully C. Clinical practice: aphthous ulceration. N Engl J Med. 2011;365(8):735- 741.
10. Ship II. Epidemiologic aspects of recurrent aphthous ulcerations. Oral Surg Oral Med Oral Pathol. 1996;81(4):422-427.
11. Altenburg A, Abdel-Naser MB, Seeber C, Abdallah M, Zouboulis CC. Practical aspects of management of recurrent aphthous stomatitis. J Dtsch Dermatol Ges. 2014;12(9):773-781.
12. Belenguer-Guallar I, Jiménez-Soriano Y, Claramunt-Lozano A. Treatment of recurrent aphthous stomatitis. J Clin Exp Dent. 2014;6(2):e168-e173.

13. Palombo EA. Traditional medicinal plant extracts and natural products with activity against oral bacteria. J Ethnopharmacol. 2009;122(2):205-218.
14. Jach G, Ginalski D, Masłyk M, Kubiński K. Herbal preparations in the treatment of oral diseases. Herba Pol. 2017;63(2):48-58.
15. Asl MN, Hosseinzadeh H. Review of pharmacological effects of Glycyrrhiza glabra and its bioactive compounds. Phytother Res. 2008;22(6):709-724.
16. Kao TC, Shyu MH, Yen GC. Neuroprotective effects of glycyrrhizic acid and its derivatives. Neurochem Res. 2014;39(9):1805-1815.
17. Whorwood CB, Sheppard MC, Stewart PM. Licorice inhibits 11β-hydroxysteroid dehydrogenase messenger ribonucleic acid levels. J Clin Endocrinol Metab. 1993;77(4):1049-1054.
18. Yang R, Yuan BC, Ma YS, Zhou S, Liu Y. The anti-inflammatory activity of licorice. Nat Prod Res. 2017;31(14):1563-1573.
19. Simmler C, Pauli GF, Chen SN. Phytochemistry and biological properties of glabridin. Fitoterapia. 2013;86:52-61.
20. Gutiérrez RM, Mitchell S, Solis RV. Psidium guajava: a review of its traditional uses, phytochemistry and pharmacology. J Ethnopharmacol. 2008;117(1):1-27.
21. Matsumoto Y, Kageyama C, Tsukuda S, Ito Y. Flavonoids from Psidium guajava leaves and their biological activities. Nat Prod Commun. 2018;13(6):709- 712.
22. Boots AW, Haenen GR, Bast A. Health effects of quercetin: from antioxidant to nutraceutical. Eur J Pharmacol. 2008;585(2-3):325-337.
23. Prabu GR, Gnanamani A, Sadulla S. Guaijaverin – a plant flavonoid as potential antiplaque agent against Streptococcus mutans. J Appl Microbiol. 2006;101(2):487-495.
24. Lozoya X, Meckes M, Abou-Zaid M, Tortoriello J, Nozzolillo C, Arnason JT. Quercetin glycosides in Psidium guajava and their analgesic activity. Arch Med Res. 2002;33(3):241-245.
25. Chah KF, Eze CA, Emuelosi CE, Esimone CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. J Ethnopharmacol. 2006;104(1-2):164-167.
26. Sujatha B, Krithi K, Anjali M, Narayana Swamy VB. Formulation and evaluation of polyherbal mouthwash. J Pharm Sci Res. 2013;5(9):176-179.
27. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. BMJ. 2009;339:b2535.
28. Chase MW, Christenhusz MJM, Fay MF, et al. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Bot J Linn Soc. 2016;181(1):1-20.
29. The Plant List. Version 1.1. Royal Botanic Gardens, Kew and Missouri Botanical Garden; 2013. Available from: http://www.theplantlist.org/
30. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. PLoS Biol. 2010;8(6):e1000412.
31. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. BMJ. 2010;340:c332.
32. Asl MN, Hosseinzadeh H. Review of pharmacological effects of Glycyrrhiza glabra and its bioactive compounds. Phytother Res. 2008;22(6):709-724.
33. Pastorino G, Cornara L, Soares S, Rodrigues F, Oliveira MB. Liquorice (Glycyrrhiza glabra): A phytochemical and pharmacological review. Phytother Res. 2018;32(12):2323-2339.
34. Sharma P, Singh V. Role of Yashtimadhu (Glycyrrhiza glabra) in oral health: A review. Int J Ayurveda Res. 2013;4(1):45-49.
35. Gutiérrez RM, Mitchell S, Solis RV. Psidium guajava: A review of its traditional uses, phytochemistry and pharmacology. J Ethnopharmacol. 2008;117(1):1-27.
36. Tona L, Kambu K, Ngimbi N, Cimanga K, Vlietinck AJ. Antiamoebic and phytochemical screening of some Congolese medicinal plants. J Ethnopharmacol. 1999;66(2):205-211.
37. Kao TC, Shyu MH, Yen GC. Neuroprotective effects of glycyrrhizic acid and its derivatives. Neurochem Res. 2014;39(9):1805-1815.
38. Boots AW, Haenen GR, Bast A. Health effects of quercetin: from antioxidant to nutraceutical. Eur J Pharmacol. 2008;585(2-3):325-337.

39. Yang R, Yuan BC, Ma YS, Zhou S, Liu Y. The anti-inflammatory activity of licorice: A review. Nat Prod Res. 2017;31(14):1563-1573.
40. Lozoya X, Meckes M, Abou-Zaid M, Tortoriello J, Nozzolillo C, Arnason JT. Quercetin glycosides in Psidium guajava and their analgesic activity. Arch Med Res. 2002;33(3):241-245.
41. Matsumoto Y, Kageyama C, Tsukuda S, Ito Y. Flavonoids from Psidium guajava leaves and their biological activities. Nat Prod Commun. 2018;13(6):709- 712.
42. Saha S, Verma RJ, Singh A, Chopra K. Antioxidant activity of Glycyrrhiza glabra extract. Asian J Pharm Clin Res. 2011;4(2):97-100.
43. Prabu GR, Gnanamani A, Sadulla S. Guaijaverin – a plant flavonoid as potential antiplaque agent against Streptococcus mutans. J Appl Microbiol. 2006;101(2):487-495.
44. Tugrul S, Dogan R, Aydin M, Yildirim YS, Ozturan O. The role of oxidative stress in recurrent aphthous stomatitis. North Clin Istanb. 2016;3(2):105-110.
45. Dhiman A, Nanda A, Ahmad S. Antimicrobial activity of Psidium guajava leaves against oral pathogens. J Pharm Res. 2012;5(8):4215-4218.
46. Bachheti RK, Sharma S, Rana V. Antimicrobial activity of Glycyrrhiza glabra against oral microflora. Int J Pharm Sci Res. 2015;6(4):1528-1532.
47. Tanaka A, Shibata H, Ishii N, Kataoka T. Wound healing activity of Glycyrrhiza glabra in human gingival fibroblasts. J Oral Sci. 2019;61(2):278-284.
48. Chah KF, Eze CA, Emuelosi CE, Esimone CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. J Ethnopharmacol. 2006;104(1-2):164-167.
49. Kumar S, Alam MS, Alam N, Ahmad S. Wound healing activity of Glycyrrhiza glabra and Psidium guajava combination gel in rat model. J Herbal Med. 2019;15:100-105.
50. Martin MD, Sherman J, van der Ven P, Burgess J. A clinical trial of a licorice mouthwash for recurrent aphthous stomatitis. Oral Surg Oral Med Oral Pathol Oral Radiol. 2016;122(2):156-163.

HOW TO CITE: K.G. Parthiban, Sanjeev Kumar. R, Kalpana Devi. M, A Mechanistic Review on the Formulation and Muco-Protective Evaluation of a Novel Mouthwash Containing Glycyrrhiza glabra (Licorice) and Psidium guajava (Guava) for the Management of Stomatitis Ulcerosa (Recurrent Aphthous Stomatitis), J. Pharm. Sci., 2026, 2 (6), 42-54. https://doi.org/10.5281/zenodo.20516527