



Genetic and morphological differentiation among populations of the narrowly endemic and karst forest-adapted *Pilea pteridophylla* (Urticaceae)

Karina Lagos-Báez¹ · Yuyini Licona-Vera² · Héctor Gómez-Domínguez³ · Juan Francisco Ornelas⁴ · Antonio Acini Vásquez-Aguilar⁴ · Leopoldo Hurtado-Reveles⁵ · Andrés Ernesto Ortiz-Rodríguez⁵

Received: 8 March 2023 / Accepted: 16 February 2024 / Published online: 28 February 2024
© The Author(s) 2024

Abstract

In this study, the influence of geographic isolation on the levels of morphological and genetic differentiation among populations of the karst forest-adapted *Pilea pteridophylla* (Urticaceae) was evaluated. Based on our results, there is a strong morphological differentiation among the only two known populations of *Pilea pteridophylla*. Both populations can be clearly differentiated based on vegetative and reproductive characters, and the observed variation is geographically structured. Moreover, our genetic data from the chloroplast genome show strong levels of population differentiation and no shared haplotypes were observed between populations. Also, our estimation of divergence times supports a long-term isolation hypothesis between populations. Based on our results, dispersal seed limitation, long-term forest isolation and topographic heterogeneity (local adaptation) are related to high levels of endemism and species turnover in the mountain karst forests of Mexico.

Keywords Endemic · Neotropics · Mexico · Phylogeography · Speciation · Tropical rainforest

Introduction

Tropical karst forests are considered among the most diverse biomes worldwide (Geekiyana et al., 2019; Molina-Paniagua et al., 2023). Their huge diversity is related to their extreme soil characteristics (e.g., high levels of Ca, Mg and K, very slow rates of soil formation and absence of surface water), high heterogeneity of microhabitats (e.g., limestone outcrops, cliffs, caves and chasms) and their archipelago-like

distribution (Clements et al., 2006). Thus, plant lineages that have invaded and diversified in these forests are specialized flora, with small or disjunct distribution ranges and notorious morphological innovations (Wendt, 1987). Tropical karst forests are an ideal model to study the effects of its discontinuous distribution on the morphological and genetic variation of species; however, there are relatively few phylogeographic studies focused on plants adapted to karstic soils (e.g., Zhu et al., 2022). Despite that, the results of the studies published so far are consistent in showing that allopatric speciation is one of the major evolutionary processes causing diversity in these forests (Gao et al., 2015; Tseng et al., 2019; Wang et al., 2017; Zhu et al., 2022).

In Mexico, tropical karst forests occupy 20% of its total surface (~391,700 km²) in the southeast, along its eastern slope, and in the Yucatán Peninsula (Espinosa-Pereña, 1990, 2007). Two main types of karst forests can be recognized in Mexico, the platform karst and the fold mountains karst (Lugo-Hubp, 1990). Each type is differentiated in terms of its geological structure and climatic characteristics (Espinosa-Pereña, 1990, 2007; Lugo-Hubp, 1990). The fold mountains karst (named hereafter as mountain karst forest sensu Molina-Paniagua et al., 2023) is mainly covered by tropical rainforests and ranges from the southwest of

✉ Andrés Ernesto Ortiz-Rodríguez
andres.ortiz@ib.unam.mx

¹ División de Ciencias Forestales, Universidad Autónoma Chapingo, Texcoco, Mexico

² Departamento de Ecología Evolutiva y Experimental, Instituto de Ecología, Universidad Nacional Autónoma de México, Ciudad de México, Mexico

³ Fondo de Conservación El Triunfo, A.C., Tuxtla Gutiérrez, Chiapas, México

⁴ Departamento de Biología Evolutiva, Instituto de Ecología, A.C. (INECOL), Xalapa, Veracruz, Mexico

⁵ Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad de México, Mexico



Tamaulipas to southern Veracruz, in the north-east portion of Oaxaca, in southern Tabasco and several regions of the state of Chiapas (Fig. 1A). Instead, platform karst is well represented, but almost restricted to the Yucatán Peninsula (Karst of the Yucatán Peninsula sensu Espinasa-Pereña, 2007). Thus, the mountain karst forests cover 2.07% of Mexico's surface, harbouring some of the most diverse forests in the country (Molina-Paniagua et al., 2023; Fig. 1A). Recently, it was estimated that around 11,000 species of

vascular plants occur within the mountain karst forests of Mexico, just over 50% of the species estimated for the country (Molina-Paniagua et al., 2023).

As stated, allopatric speciation has been invoked as one of the main processes that sustains high diversity within the mountain karst forests of Mexico (Wendt, 1987; Pérez-García & Meave, 2005; Pérez-García et al., 2009). However, there are no studies in the region that properly evaluate the role of geographic isolation on the patterns of morphological




Fig. 1 Mountain karst forests of Mexico and the studied species *Pilea pteridophylla* A. K. Monro (Urticaceae). **A** Geographic distribution of the Mountain karst forests of Mexico. **B** Individual from the Chiapas population. **C** Individual from the Tabasco population

and genetic differentiation of the species. The latter should be a priority; tropical karst forests of Mexico are naturally fragmented but frequently surrounded by a matrix of agricultural fields, pastures, roads and cities, so elucidating the main evolutionary processes that underpin their species diversity is critical for improved conservation actions.

Here, we studied the genetic and morphological differentiation within *Pilea pteridophylla* (Urticaceae), a species endemic to Mexico with a restricted distribution to two allopatric populations located in the mountain karst forests of Tabasco and Chiapas in southern Mexico. Specifically, we assessed the influence of geographic isolation on the levels of morphological and genetic differentiation among populations.

Materials and methods

Study organism

Pilea pteridophylla A. K. Monro is a calcicole herbaceous plant endemic to southern Mexico (Fig. 1B–C). The species was described based on specimens collected in Tabasco, Mexico (Monro, 1999). It is easily distinguished from the rest of the Mexican species of *Pilea* by its pectinate leaves, which resemble those of a fern in colour, shape and texture (Monro, 1999; Fig. 1B–C). In fact, it is the only species of *Pilea* with pectinate leaves known from America (Monro, 1999). The species is restricted to the tropical rain forest, and grows on karstic limestone soils. For a long time, this species was known only from specimens collected (all before the year 2000) at the type locality in the Sierra Madrigal, Tabasco, Mexico (Monro, 1999). However, Ortiz-Rodríguez

et al. (2012) reported a new locality for this species in the state of Chiapas, just over 80 km away from the type locality. Those authors indicated notable morphological differences between the individuals of both populations, mainly in the size of the plants and the depth of the margin teeth, but no formal analysis was performed. Recent explorations around the type locality have failed to locate living individuals of *Pilea pteridophylla*, which does not mean that the species is locally extinct, but it is probably restricted to less accessible regions. Thus, a new locality in Cerro Tortuguero, Macuspana, Tabasco, was recently located; the individuals present in this locality have morphological characteristics, such as the pectinate leaves, that conform to those of the type specimen (Fig. 1C).

Both populations currently recognized for *Pilea pteridophylla* are part of the mountain karst forests of Mexico, and although their populations consist of several individuals (more than 100 mature plants), their distribution is extremely localized. Specifically, the individuals from Chiapas occur at altitudes above 1000 m above sea level and appear associated with a few rocky mounds at ground level, while individuals from Macuspana, Tabasco, occur at altitudes between 500 and 600 m above sea level and appear to be restricted to a single limestone walls or cave entrance.

Sample collection

Collections were made in the two populations of *Pilea pteridophylla* to obtain fresh samples for DNA extraction (leaves) and complete specimens for morphological differentiation analysis (Table 1). A total of 36 herbarium specimens (17 individuals from Macuspana, Tabasco, and 17 individuals from Berriozábal, Chiapas) were analysed. Of these, 30

Table 1 Populations and locations of *Pilea pteridophylla* (Urticaceae). The number of individuals sampled and used for the analyses are indicated (morphological/genetic)

Population	Locality	Total number of individuals collected	Individuals used for the analyses	Collector and collection number
Macuspana, Tabasco	On the way to the “Agua Blanca” State Park, between Holcim and El Chivalito 2da. Section 158 m altitude	17	17/15	K. Lagos-Báez 26–42 (MEXU)
Teapa, Tabasco (Teapa)	Madrigal Hill, up from Chapingo’s Puyacatengo Tropical Station next to the Puyacatengo River. 3 km from the Teapa-Tacotalpa road. Teapa, Tabasco	1	1/0	Clark P. Cowan et al. 3978 (XAL)
	7 km SE of Teapa on road to Tacoatalpa; Rancho San Eneas; Sierra Madrigal woods on steep limestone hills. Teapa, Tabasco	1	1/0	Barry E. Hammel & Mary Merello 15,523 (MEXU)
Berriozábal, Chiapas	Protected natural area “La Pera”, Berriozábal, Chiapas. Around “Flor de café”	17	17/15	K. Lagos-Báez 45–51; 53;55;57–64 (MEXU)

samples were used for DNA extraction (15 samples from Macuspana, Tabasco, and 15 samples from Berriozábal, Chiapas). We also had access to two specimens of *Pilea pteridophylla* from the type locality, but samples for DNA extraction were not available and were only included in the morphological analyses.

Morphological differences among populations

In order to explore morphological differentiation among individuals of *Pilea pteridophylla*, we measured seven morphological traits (vegetative and reproductive), including those traditionally used for species delimitation (Monro, 2001; Table 2). A principal components analysis (PCA) was performed on individuals using a correlation matrix with PAST v.3.06 (Hammer et al., 2001). We used the first two principal components (PC) to evaluate differences among groups of populations through an analysis of variance performed in R (Chambers et al., 1992; R Core Team, 2021). The graphic representation of the PCA results analyses were handled with the “cowplot” and “ggpubr” packages in R (Alboukadel, 2020; Claus, 2020).

Genetic differences among populations

We extracted DNA from leaves tissue using a CTAB protocol (Cota-Sánchez et al., 2006). The *trnL-trnF* and *trnQ-rps16* intergenic spacer regions from the chloroplast genome (cpDNA) were amplified by PCR and sequenced for the 30 individuals of *Pilea pteridophylla*. These markers have been screened for variability in several plant lineages at the species and population levels (e.g. Shepherd & Heenan, 2017). Since the chloroplast genome is maternally inherited in most

angiosperms, cpDNA sequences may provide signals about the routes and distances of seed movement and their impacts on gene flow.

For *trnL-trnF* amplifications, we used the universal primers (*trnL-e*, *trnL-f*) designed by Taberlet et al. (1991), whereas for the *trnQ-rps16*, we used the primers (*trnQ-5'*, *rps16* × 1) described by Shaw et al. (2007). Protocols for DNA amplification by PCR assays and for sequencing the PCR products with some minor modifications were described elsewhere (Ornelas et al., 2016). PCR products were sequenced bidirectionally using the BigDye Terminator Cycle Sequencing kit (Applied Biosystems, Foster City, CA, USA, and analysed on a 310 automated DNA sequencer (Applied Biosystems) at the INECOL's sequencing facility, or at University of Washington High Throughput Genomics Unit, Seattle, Washington. Edited sequences were aligned manually with PhyDE v.0.9971 (Müller et al., 2010; <http://www.phyde.de/>). The resulting sequences are available as alignments at Ortiz-Rodríguez (2023a), and available individually in the GenBank database (*trnL-F*, GenBank accession numbers: PP151098–PP151127; *trnQ-rps16*, GenBank accession numbers: PP151128–PP151157).

We calculated haplotype number (*h*), haplotype diversity (*H_d*) and nucleotide diversity (π) for both Tabasco and Chiapas localities using DnaSP v.5.10 (Librado & Rozas, 2009). Since both locations have the same number of individuals (15), estimates of genetic diversity are probably not influenced by sample size. To assess genetic differentiation, we considered each locality as a population, which is a simple way to investigate the spatial distribution of genetic variation. Thus, Wright's (1965) fixation index was calculated in the Arlequin v.3.5.1.2 program (Excoffier et al., 2005) with 16,000 replicates and a 0.05 level of significance. This index (pairwise *F_{ST}*-values) has values between 1 and zero, where

Table 2 Results of principal component analysis (PCA). The percentage of the variance explained by each component and the results of the analysis of variance using the scores values of the first two components as the response variable, are shown. The loading values on the first two components obtained from the PCA are also presented in bold which are the most important variables for group discrimination

PC	Eigenvalue	% variance	Cumulative variance	F value/Pr(> F)
1	4.74567	67.795	67.795	227.7/***
2	1.0412	14.874	82.669	0.021/n.s
3	0.796312	11.376	94.045	
4	0.266655	3.8094	97.8544	
Variable	PC 1	PC 2	PC 3	PC 4
Stem length	0.42355	− 0.00422	− 0.091783	0.62856
Stem width	0.43892	0.072787	− 0.14952	0.28294
Leaf length	0.44376	0.13593	− 0.13786	0.074145
Leaf width	0.43297	− 0.036963	− 0.1348	− 0.52995
Proportion of leaf surface without teeth on margin	0.22409	− 0.33077	0.89983	0.079339
Depth of margin teeth in proportion to leaf width	− 0.43825	0.07639	− 0.022652	0.47784

1 is indicative of strong population differentiation, that is, presence of genetic structure and without gene flow between populations (genetic isolation), while a value of zero indicates no genetic differentiation between populations.

Relationships among *P. peridophylla* haplotypes are presented as statistical parsimony networks generated using the program TCS v.1.2.1 (Clement et al., 2000), with gaps treated independently as missing data and as fifth state, and a 95% connection probability limit. Loops were resolved following the criteria given by Pfenninger and Posada (2002). We report herein results for each marker independently and for the combined *trnL-trnF* and *trnQ-rps16* data set.

Phylogenetic relationships and divergence time estimation

Divergence times were estimated under a Bayesian approach as implemented in BEAST v.1.8.1 (Drummond & Rambaut, 2007) using the *trnL-F* dataset. We include various representatives of *Pilea* and related genera of Urticaceae as available in the supplementary material of Fu et al. (2022). The 76-accession dataset (available at Ortiz-Rodriguez, 2023b) was prepared for BEAST using BEAUti with the optimum model setting determined by the results of jModelTest (Darriba et al., 2012).

As our DNA database contains a mix of inter- and intraspecies sampling, we used pure-birth, birth–death and constant-size coalescent tree priors as models for dating estimates (Ritchie et al., 2017). The first two are designed for species-level data (speciation-based tree priors) and the last one is most suitable for trees describing the relationships between individuals in the same population/species (coalescent-based tree priors). Thus, non-coalescent methods could overestimate the age of lineages, while coalescent methods could bias the time estimation of inter-species divergences towards young ages (Liu et al., 2022; Mello et al., 2021). Consequently, we used three calibration schemes as follows: (1) we used a lognormal calibration prior distribution, which assigned the highest probabilities to substantially older ages than the calibrated ages accounting for errors associated with the incompleteness and limited knowledge of the Urticaceae fossil record. In this way, the Urticaceae crown group was calibrated using the estimated mean age for the family (68.7 Ma; Wu et al., 2018) as offset and the shape of the lognormal distribution was defined with a median of 72.8 Ma and a 95% probability interval between 69 and 136.4 Ma, using the unequivocal angiosperm crown group pollen grain fossils (Friis et al., 2010; Hughes, 1994) as soft upper bound (offset: 68.7, mean: 11.4, log (Stdev): 30). An uncorrelated lognormal relaxed molecular clock (UCLD) and the Yule speciation tree prior were applied to the *trnL-F* dataset. (2) The branching process prior ('tree' prior) was changed to use a birth–death model that considers the probability of

frequent extinction processes throughout evolutionary history of the genus *Pilea* (not solely a continuous speciation as in the Yule model). (3) Lastly, the 'tree' prior was changed to use a constant-size coalescent model and with the rest of the parameters as in the previous dating strategies. We ran each scheme 20 million generations for the cpDNA data set of the MCMC chain, with a sampling frequency of 1000 generations. Convergence of the posterior distributions of the parameters was evaluated by monitoring the effective sample sizes ($ESS > 200$) and trace plots in Tracer (<http://tree.bio.ed.ac.uk/software/tracer/>). A maximum credibility tree, which represents the maximum posterior topology, was calculated in TreeAnnotator (Drummond & Rambaut, 2007) after discarding the first 10% of trees as burn-in.

Results

Morphological differences among populations

The first two PC components of the PCA explained 82.6% of the total variance in morphological traits (Table 2). The PC1 explained 67.7% of the variance was positively associated with the plant length, stem width, leaf length and leaf width, and negatively associated with the depth of margin teeth in proportion to leaf width. The PC2 (14.8% of the variance) was associated only with the proportion of leaf surface without teeth on margin (Table 2). Significant differences between groups (Chiapas vs. Tabasco) were observed only in PC1 ($t = 19.166$, $df = 35.892$, $p\text{-value} < 2e-16$; Fig. 2). Table 3 summarizes the main morphological differences between individuals from each of the studied localities.

Genetic differences among populations

A matrix of 462 bp for *trnL-F* and 870 bp for *rps16-trnQ* was obtained. Thus, the concatenated matrix (*rps16-trnQ* + *trnL-F*) consisted of 1332 bp. The number of haplotypes varied from 4 to 8 per dataset with gaps considered missing data and from 7 to 15 when treated as 5th state (Table 4, Supplemental Fig. S1). The haplotype diversity values were relatively high and the nucleotide diversity values were low in all cases (Table 4). There are no shared haplotypes between populations (Fig. 3), and a strong population genetic structure is assumed. Pairwise F_{ST} values ranged from 0.33 to 0.89 and with significant differences in all comparisons (Table 4). The nucleotide diversity and F_{ST} values were always higher in the *rps16-trnQ* dataset and in the concatenated dataset (*rps16-trnQ* + *trnL-F*). In summary, both populations (Chiapas and Tabasco) show strong genetic differentiation with F_{ST} values close to 1 (Fig. 3, Table 4). Haplotype diversity was relatively high overall, with more

Fig. 2 Morphological variation among individuals of *Pilea pteridophylla* sampled along its distribution range in the tropical karst forest of southern Mexico. Plot of individual scores for the first two components of the principal component analysis using morphological data. Coloured symbols represent the two populations recognized for the species: red circles, Tabasco; and blue circles, Chiapas. Ellipses correspond to the 95% confidence intervals estimated for each population. The lines represent the dispersion of the individuals within each population




Table 3 Morphological features of the individuals of *Pilea pteridophylla* (Urticaceae) collected in Tabasco and compared to those collected in Chiapas

Features	Tabasco	Chiapas
Stem length	147.7 ± 81 mm	472.7 ± 152.2 mm
Stem width	1.26 ± 0.5 mm	3.07 ± 0.4 mm
Length of the largest leaf	55.7 ± 21.9 mm	141.5 ± 22.2 mm
Width of the largest leaf	7.9 ± 1.4 mm	18.5 ± 2.2 mm
Proportion of the largest leaf with entire margin	0.13 ± 0.03	0.17 ± 0.02
Depth of teeth at leaf margin	0.96 ± 0.01 mm	0.28 ± 0.05 mm
Stem proportion with female inflorescences	0.19 ± 0.13	0.20 ± 0.12

Table 4 Genetic diversity statistics and Pairwise comparisons of F_{ST} values of *rps16-trnQ*, *trnL-F* and *rps16-trnQ + trnL-trnF* among *Pilea pteridophylla* (Urticaceae) populations

Data set	Gaps missing				Gaps = 5th state			
	H	h	π	F_{ST}	H	h	π	F_{ST}
<i>rps16-trnQ</i>	4	0.6414	0.00214	0.89***	7	0.74	0.00214	0.83***
<i>trnL-trnF</i>	5	0.6851	0.00204	0.41***	10	0.85	0.00204	0.33***
<i>rps16-trnQ + trnL-trnF</i>	8	0.7609	0.0021	0.69***	15	0.91	0.0021	0.62***




Fig. 3 Statistical parsimony networks of *rps16-trnQ*, *trnL-trnF* and *rps16-trnQ + trnL-trnF* dataset using the gaps as missing data. Coloured symbols represent the two populations recognized for the species; red circles, Tabasco; and blue circles, Chiapas. Open-white circles represent the number of mutational steps between haplotypes. The size of the circles is proportional to the frequency of each haplo-

type. Loops (dashed lines) were resolved following the criteria given by Pfenninger and Posada (2002). The grey area refers to the regions with the presence of mountain karst forests in southern Mexico [shape file available at CENAPRED (<http://www.atlasmnacionalderiesgos.gob.mx/archivo/visor-capas.html>)]

haplotypes in Chiapas compared to Tabasco (Table 4). The nucleotide diversity was low in general, with haplotypes differentiated by few mutational steps within each of the populations.

Phylogenetic relationships and divergence time estimation

The maximum clade credibility tree that resulted from the BEAST analysis using the *trnL-F* dataset and the Yule speciation tree prior is shown in Fig. 4. Mean ages of major clades with 95% HPD intervals for the three calibration schemes are indicated in Table 5. The maximum clade credibility tree retrieved the genus *Pilea* (*PP* value = 0.99) as monophyletic, sister group of *Lecanthus* (*PP* value = 1.0) and *Achudemia*

(*PP* value = 0.99). The genus *Pilea* is divided into two main clades and 8 sections (Fig. 4). The first clade (*PP* value = 0.27) corresponds to sections *Tetraphyllae* (*Pilea tetraphylla* only), *Trimeris* (*PP* value = 1.0), *Lecanthoides* (*PP* value = 1.0), *Angulatae* (*PP* value = 1.0) and *Tetrameris* (*PP* value = 1.0), and the last four nested in a strongly supported subclade (*PP* value = 1.0). The second clade contains the sections *Verrucosae* (*PP* value = 1.0), *Plataniflorae* (*PP* value = 1.0) and *Pilea* (*PP* value = 1.0). *Pilea pteridophylla* (*PP* value = 1.0) appears as a member of the section *Pilea*, along with *Pilea pleuroneura*, *Pilea mexicana* and *Sarcopilea domingensis* (= *Pilea fairchildiana*) where the Chiapas and Tabasco populations form two reciprocally monophyletic clades (Chiapas lineage with *PP* value = 0.25; and Tabasco lineage with *PP* value = 0.96).

Table 5 Age estimated (in millions of years) for the main studied clades (as illustrated in Fig. 4), showing results of analyses using three calibration tree priors

	BEAST Yule model	BEAST birth–death model	BEAST constant-size coalescent model
Clade	Mean (95% HPD)	Mean (95% HPD)	Mean (95% HPD)
Urticaceae	71.5 Ma (89.5–68.7 Ma)	74.4 Ma (85.8–68.7 Ma)	75.2 Ma (92.0–68.7 Ma)
<i>Pilea</i>	38.1 Ma (54.5–24.5 Ma)	34.4 Ma (48.1–18.6 Ma)	33.7 Ma (56.8–12.3 Ma)
<i>Pilea pteridophylla</i>	7.2 Ma (13.5–3.5 Ma)	4.95 Ma (7.4–1.7 Ma)	3.9 Ma (7.5–1 Ma)

early Miocene [Yule model: mean = 38.1 Ma (95% HPD 54.5–24.5 Ma); birth–death model: Mean = 34.4 Ma (95% HPD 48.1–18.6 Ma); constant-size coalescent model: Mean = 33.7 Ma (95% HPD 56.8–12.3 Ma)], so its current distribution could be associated with a boreotropical origin. According to the analysis of the *trnL-F* chloroplast marker, the two populations of *Pilea pteridophylla* could have diverged during the middle Miocene and late Pliocene [Yule model: mean = 7.2 Ma (95% HPD 13.5–3.5 Ma)] or later between the late Miocene and the Pleistocene [birth–death model: mean = 4.95 Ma (95% HPD 7.4–1.7 Ma); constant-size coalescent model: mean = 3.9 Ma (95% HPD 7.5–1.0 Ma)], in any case, a deep divergence and isolation between individuals restricted to different vegetation patches (Fig. 4).

Discussion

In this study, the influence of geographic isolation on the levels of morphological and genetic differentiation among populations of *Pilea pteridophylla* was evaluated. Based on our results, there is a strong morphological differentiation among populations of *Pilea pteridophylla* (Fig. 2, Table 3). Both populations can be clearly differentiated based on vegetative and reproductive characters (Table 3), and the observed variation is geographically structured (Fig. 2). Moreover, our genetic data from the chloroplast genome show strong levels of population differentiation and no shared haplotypes were observed between populations (Fig. 3, Table 4). Also, our estimation of divergence times supports a long-term isolation hypothesis between Chiapas and Tabasco populations (Fig. 4, Table 5). Thus, our study supports the hypothesis of allopatric speciation as one of the main evolutionary processes triggering diversity within the mountain karst forests of Mexico (Wendt, 1987). Based on the evidence presented here, some evolutionary factors involved in the separation of Mexican karst-adapted plant lineages are discussed below.

Low seed-mediated gene flow and habitat heterogeneity promotes genetic differentiation among populations adapted to karstic soils

Based on the low substitution rates attributed to chloroplast markers, the estimated deep age of divergence between the

Chiapas and Tabasco populations of *Pilea pteridophylla* is somewhat unusual. However, the deep divergence suggests that seed-mediated migration rates are actually low in *Pilea pteridophylla*. This result is quite interesting since long-distance dispersal through water currents has been hypothesized as the main dispersal mechanism in Urticaceae (Wu et al., 2018). According to Wu et al. (2018), seeds of many Urticaceae floated for > 220 days, and remained viable after 10 months in water. *Pilea pteridophylla* is an herb that grows exclusively in cave entrances or on rocky knolls, where surface water is scarce but with large runoff during the rainy season. However, seed dispersal through water currents is rather local. In Chiapas, groups of 10 or more individuals of *Pilea pteridophylla* are widely dispersed throughout the forest, suggesting that water currents can move seeds long distances locally. This is supported by the higher number of haplotypes observed in Chiapas compared to those recorded in Tabasco. In Tabasco, the population is rather dense, with all individuals separated by a few centimetres, where the effect of water on seed dispersal and on the level of genetic differentiation among individuals is low. Thus, it is likely that the genetic flow between Chiapas and Tabasco is currently interrupted.

Our results also suggest that topographic complexity is an important feature for population differentiation. First, the geographic distance between populations (more than 80 km²) and their differences in altitude suppose ecological barriers to genetic exchange between populations. Second, local-scale habitat preferences (rocky knolls vs. caves and walls) are probably more related to their local distribution and their morphological characteristics (Liu et al., 2021). More specifically, individuals from Chiapas have larger stems because in the rocky knolls the soil can be deeper, resulting in greater support for their roots and stems, while for individuals from Tabasco, the rocky walls could limit plant growth (in stems and roots), probably associated with less soil and water (Matthes-Sears & Douglas, 1995; Liu et al., 2020). However, field experiments are needed to test this hypothesis.

The karst-endemic species as relict lineages of ancient vegetation

Our estimation of divergence times shows a deep divergence (between middle Miocene and Pleistocene) among




Fig. 5 *Pilea monroi* Ortiz-Rodr. and Lagos-Báez (Urticaceae). **A** Habit, plant with pistillate inflorescences. **B** Stem, a leafless portion with inflorescences. **C** Flower bud. **D** Staminate flower with four stamens and four tepals. **E** Infructescence. **F** Stem portion with pistillate inflorescences. Drawn by Robin Pérez Lucas based on *Lagos-Báez 57* (MEXU)

Pilea pteridophylla populations. This level of divergence is consistent with the strong phylogeographic structure and the clear morphological differentiation between individuals restricted to each population. Wendt (1987) suggests that the different levels of morphological differentiation between endemic sister species in the karst forests of Mexico could be related to the time of isolation between patches of vegetation to which they are restricted, in addition to the fact that karst forests have functioned as floristics refuges at different times and in different places between the Miocene and Pleistocene (Wendt, 1987, 1993). Our data support this idea. Specifically, the strong differentiation between populations of *Pilea pteridophylla* could be related to a prolonged geographic isolation driven by the past climate fluctuations occurred between middle Miocene and Pleistocene. Also, *Pilea pteridophylla* is the only species with fern-like leaves within the genus *Pilea*, and its calcareous habit, restricted distribution and its strong population differentiation suggest a long-term adaptation to a karstic habitat. Based on this evidence, it is possible to hypothesize that the species endemic to the karst forests of Mexico are relicts of an ancient vegetation, while the non-endemic elements could be more recent invaders (Wendt, 1987, 1993). Although little is known about the time in which the plant immigration events to the karst relief occurred (see Li et al., 2022), the data published so far (focused on Asian forests) suggest that primary immigration events occurred during the Miocene, which is supported in this study; however, it could also be related to the expansion and diversification during the Miocene of many lineages of lowland tropical forest plants in general (Becerra, 2005; Ornelas et al., 2018; Ortiz-Rodriguez et al., 2020; Rull, 2020; Gutiérrez-Ortega et al., 2024). Interestingly, floristic data suggest that only a small part (less than 4%) of the species currently present in the mountain karst forest of Mexico are exclusive (endemic) to this biome (see Molina-Paniagua et al., 2023), and for another small percentage of species (6%), between 50 and 90% of their records are distributed exclusively within this forest. This suggests that adaptation to karst soils is a slow evolutionary process. Viewed this way, our data support the idea of endemic karst species as ancient forest elements, but also show that plant diversification in karst forests has not been constant (as suggested by long-term forest stability), and has occurred to a lesser extent towards the present.

Long-term geographic isolation promotes speciation

Based on our results, the individuals of *Pilea pteridophylla* restricted to each population can be morphologically and genetically differentiated. These differences are associated with a long-term geographic isolation without apparent gene flow and contrasting ecological conditions. This evidence shows convincingly that the individuals from Chiapas are an independent lineage and should be recognized as a new species to science. Consequently, a new Mexican species of *Pilea* is described below.

Taxonomic section

Pilea monroi Ortiz-Rodr. & Lagos-Báez sp. nov. (Fig. 5).

Etymology

Pilea monroi named in honour of Dr Alex Monro, taxonomist, specialist in the genus *Pilea* (Urticaceae) and especially interested in the study of the plant diversity in the karstic forests of America and China.

Similar to *Pilea pteridophylla* in general fern-like appearance, but clearly differentiated by its dentate (vs. pectinate) leaves, and larger stem and leaf size. Furthermore, they are allopatrically distributed and under different environmental conditions.

Type: MEXICO: Chiapas; Berriozábal, Zona sujeta a protección ecológica “La Pera”. Predio “Flor de Café”, 16°51'50.6" N, 93°19'51.7" W, alt. 1068 m., May 2019, K. Lagos-Báez 57 (Holotype, MEXU; isotypes: MO, KEW).

Description

Herbs 254–740 mm, perennial, epipetric. Stems erect, sparsely branched, greenish in vivo, blackened when dry, glabrous, cystoliths linear, X or Y shaped, not clearly visible when dry, the internodes 7.5–54 × 1.5–2.4 mm, irregularly circular or slightly angled in cross-section, striate. Leaves petiolate, distichous, laminae and petioles at the same node of equal length, glabrous, laminae 80–180 × 12–22 mm, linear lanceolate, chartaceous, deeply dentate above de basal 1/7 to 1/3, adaxial surface dark green when dry, glabrous, cystoliths fusiform, V-, X-, Y-shaped, abaxial surface greyish or olive, rarely light brown, glabrous, cystoliths fusiform, V-, X-, Y-shaped, glandular punctate, base symmetrical or nearly so, decurrent, attenuated, rounded, apex cuspidate, primary venation pinnate, secondary veins 26–42 pairs at 90° to midrib; petioles 3.5–5.0 × 0.8–2.5 mm, stipules 0.4–1.0 mm, deltate, membranous, often persistent. Inflorescences ca. 15 per stem, 2.0–4.0 mm, unisexual,

pistillate inflorescences preceding and concurrent with staminate inflorescences, peduncle and pedicels subtended by stipuliform, minute bracts, staminate flowers, cream or slightly pink, tepals 4, 0.8–1.2 mm, ovate, glabrous, stamens 4, filaments 0.5–0.8 mm, anthers 0.3–0.4 mm, pistillate flowers, cream, tepals 3, 0.5–0.8 mm, the central one slightly longer than the two lateral, ovate, glabrous. Infructescences of 2–3 mm, peduncle 0.35 mm diam., fruits 0.4–0.6 × 0.2–0.3 mm, asymmetric, subcompressed ovoid, orange-brown, margin broadening and flattened towards the apex.

Distribution

Pilea monroi is known only from the type locality in Chiapas, Mexico. The species inhabits within the mountain karst forest at ca. 1000 m altitude.

Notes

Pilea monroi is the only species in Mexico with linear lanceolate, deeply dentate leaves, and like *Pilea pteridophylla*, its fern-like appearance makes it easily distinguishable from other species. However, the leaf margins of *Pilea monroi* are dentate and not pectinate as in *Pilea pteridophylla*.

Pilea plumosa from Costa Rica, also have linear lanceolate leaves with a dentate margin, but it is distinguished from *Pilea monroi* by its narrower leaf blades (1.5–4.0 mm vs 12–22 mm), by the abaxial leaf surface covered by orbicular cystoliths (vs V-, X-, Y-shaped) and by its leaves with fewer secondary veins (4–13 pairs vs 26–42 pairs).

Additional specimens examined

MEXICO: Chiapas; Berriozábal, Zona sujeta a protección ecológica “La Pera”. Predio “Peña Flor”, 16°51'50.6" N, 93°19'51.7" W, alt. 1068 m., Ortiz-Rodríguez A. E. 320 (HEM).

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s13127-024-00638-x>.

Acknowledgements We sincerely thank Christopher Davidson and Sharon Christoph for their support through the project “KARSTBIO: Biodiversity, Evolution and Conservation of the tropical rain forest on karstic zones of Latin America”. Also, we thank the anonymous reviewers for providing many useful comments on the manuscript. This work forms part of the research project with which the first author obtained her bachelor's degree.

Author contribution KLB and AEOR conceived the ideas; KLB, HGD and AEOR conduct the field work; KLB and AAVA conducted the lab work; KLB and AEOR analysed the data; AEOR wrote the main manuscript text; YLV, LHR and JFO reviewed the manuscript and contribute substantially to the structure and writing of the final version. All authors read and approved the final version.

Data availability The authors declare the availability of data in the supplementary section of the manuscript.

Declarations

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alboukadel, K. (2020). ggpubr: 'ggplot2' based publication ready plots. R package version 0.4.0. <https://CRAN.R-project.org/package=ggpubr>
- Becerra, J. (2005). Timing the origin and expansion of the Mexican tropical dry forest. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 10919–10923. <https://doi.org/10.1073/pnas.0409127102>
- Chambers, J. M., Freeny, A., & Heiberger, R. M. (1992). Analysis of variance; designed experiments. *Chapter 5 of Statistical Models in S*. J. M. Chambers and T. J. Hastie (eds), Wadsworth & Brooks/Cole.
- Claus, O. W. (2020). cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R package version 1.1.1. <https://CRAN.R-project.org/package=cowplot>
- Clement, M., Posada, D., & Crandall, K. A. (2000). TCS: A computer program to estimate gene genealogies. *Molecular Ecology*, 9(10), 1657–1660
- Clements, R., Sodhi, N. S., Schilthuizen, M., & Ng, P. K. L. (2006). Limestone karsts of Southeast Asia: Imperiled arks of biodiversity. *BioScience*, 56(9), 733–742. [https://doi.org/10.1641/0006-3568\(2006\)56733:LKOSAI2.0.CO2](https://doi.org/10.1641/0006-3568(2006)56733:LKOSAI2.0.CO2)
- Cota-Sánchez, J. H., Remarchuk, K., & Ubayasena, K. (2006). Ready-to-use DNA extracted with a CTAB method adapted for herbarium specimens and mucilaginous plant tissue. *Plant Molecular Biology Reporter* 24, 161. <https://doi.org/10.1007/BF02914055>
- Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods*, 9, 772
- Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, 7, 214. <https://doi.org/10.1186/1471-2148-7-214>
- Espinasa Pereña, R. (1990). *Propuesta de clasificación del karst de la República Mexicana*. Universidad Nacional Autónoma de México
- Espinasa Pereña, R. (2007). El Karst de México. Mapa NA III 3. Nuevo Atlas Nacional de México (Instituto de Geografía, Universidad Nacional Autónoma de México). https://geodigital.geografia.unam.mx/atlas_nacional/index.html/grals/Tomo_II/IV.Naturaleza/IV.3.Relieve/IV.3.4.jpg
- Excoffier, L., Laval, G., & Schneider, S. (2005). Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evol Bioinform*, 1, 47–50

- Friis, E. M., Pedersen, K. R., & Crane, P. R. (2010). Diversity in obscurity: Fossil flowers and the early history of angiosperms. *Phil. Trans. R. Soc. B*, 365, 369–382. <https://doi.org/10.1098/rstb.2009.0227>
- Fu, L. F., Wen, F., Maurin, O., Rodda, M., Gardner, E. M., Xin, Z. B., Wei, Y. G., & Monroe, A. K. (2022). A revised delimitation of the species-rich genus *Pilea* (Urticaceae) supports the resurrection of *Achudemia* and a new infrageneric classification. *Taxon*, 71(4), 796–813. <https://doi.org/10.1002/tax.12711>
- Gao, Y., Ai, B., Kong, H., Kang, M., & Huang, H. (2015). Geographical pattern of isolation and diversification in karst habitat islands: A case study in the *Primulina eburnea* complex. *Journal of Biogeography*, 42, 2131–2144. <https://doi.org/10.1111/jbi.12576>
- Geekiyana, N., Goodale, U. M., Cao, K., & Kitajima, K. (2019). Plant ecology of tropical and subtropical karst ecosystems. *Biotropica*, 51, 626–640. <https://doi.org/10.1111/btp.12696>
- Gutiérrez-Ortega, J. S., Pérez-Farrera, M. A., Matsuo, A., Sato, M. P., Suyama, Y., Calonje, M., Vovides, A. P., Kajita, T., & Watano, Y. (2024). The phylogenetic reconstruction of the Neotropical cycad genus *Ceratozamia* (Zamiaceae) reveals disparate patterns of niche evolution. *Molecular Phylogenetics and Evolution*, 190, 107960. <https://doi.org/10.1016/j.ympev.2023.107960>
- Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Paleontological Electronica* 4, 4. http://palaeo-electronica.org/2001_1/past/issue1_01.htm
- Hughes, N. F. (1994). *The enigma of angiosperm origins*. Cambridge University Press
- Li, X.-Q., Xiang, X.-G., Zhang, Q., Jabbour, F., Ortiz, R. C., Erst, A. S., Li, Z.-Y., & Wang, W. (2022). Immigration dynamics of tropical and subtropical Southeast Asian limestone karst floras. *Proc. R. Soc. B*, 2892021130820211308. <https://doi.org/10.1098/rspb.2021.1308>
- Librado, P., & Rozas, J. (2009). DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451–1452
- Liu, L., Hu, J., Chen, X., Xu, X., Yang, Y., & Ni, J. (2022). Adaptation strategy of karst forests: Evidence from the community-weighted mean of plant functional traits. *Ecology and Evolution*, 12, e8680. <https://doi.org/10.1002/ece3.8680>
- Liu, Y., Chen, J., Xiaojing, Z., Qiaoyu, L., Yu, D., Jinchun, L., & Jianping, T. (2021). Herbaceous community species composition and productivity are affected by soil depth and funnel effect in a simulated karst experiment. *Global Ecology and Conservation*, 22, e01033. <https://doi.org/10.1016/j.gecco.2020.e01033>
- Lugo-Hubp, J. (1990) El relieve de la República Mexicana. *Revista Mexicana de Ciencias Geológicas* 474 (9), 82–111. <http://rmcg.geociencias.unam.mx/index.php/rmcg/article/view/1255>
- Matthes-Sears & Douglas, W. L. (1995). Rooting characteristics of trees in rock: A study of *Thuja occidentalis* on cliff faces. *International Journal of Plant Sciences*, 156(5), 679–686
- Mello, B., Tao, Q., Barba-Montoya, J., & Kumar, S. (2021). Molecular dating for phylogenies containing a mix of populations and species by using Bayesian and RelTime approaches. *Molecular Ecology Resources*, 21(1), 122–136. <https://doi.org/10.1111/1755-0998.13249>
- Molina-Paniagua, M. E., Alves de Melo, P. H., Ramírez-Barahona, S., Ortiz-Rodríguez, A. E., Monroe, A. K., Burelo-Ramos, C. M., & Gómez-Domínguez H (2023) How diverse are the mountain karst forests of Mexico? *bioRxiv* 2023.03.01.530643; doi: <https://doi.org/10.1101/2023.03.01.530643>
- Monro, A. K. (1999). Seven new species of *Pilea* Lindley (Urticaceae) from Mesoamerica. *Novon*, 9(3), 390–400. <https://doi.org/10.2307/3391738>
- Monro, A. K. (2001). Synopsis of Mesoamerican *Pilea* (Urticaceae), including eighteen typifications and a key to the species. *Bulletin of the Natural History Museum, Botany Series* 31, 9–25. <https://www.biodiversitylibrary.org/part/190603>
- Müller, K., Müller, J., & Quandt, D. (2010). PhyDE (Phylogenetic Data Editor). <http://www.phyde.de/>
- Ornelas, J. F., Licona-Vera, Y., & Vásquez-Aguilar, A. A. (2018). Genetic differentiation and fragmentation in response to climate change of the narrow endemic *Psittacanthus auriculatus*. *Tropical Conservation Science* 11. <https://doi.org/10.1177/19400829187555>
- Ornelas, J. F., Gándara, E., Vásquez-Aguilar, A. A., et al. (2016). A mistletoe tale: Postglacial invasion of *Psittacanthus schiedeanus* (Loranthaceae) to Mesoamerican cloud forests revealed by molecular data and species distribution modeling. *BMC Evolutionary Biology*, 16, 78. <https://doi.org/10.1186/s12862-016-0648-6>
- Ortiz-Rodríguez, A. E., Lorea-Henández, F., Pérez-Farrera, M. A., & Monroe, A. K. (2012). First record of *Pilea peridophylla* A.K.Monro (Urticaceae) and *Ocotea heribertoii* T.Wendt (Lauraceae) for the flora of Chiapas. *Botanical Sciences* 90(2), 203–206. <https://doi.org/10.17129/botsci.485>
- Ortiz-Rodríguez, A. E., Licona-Vera, Y., Vásquez-Aguilar, A. A., Hernández-Soto, M., López-Huicochea, E. A., & Ornelas, J. F. (2020). Genetic differentiation among *Psittacanthus rhynchanthus* (Loranthaceae) populations: Novel phylogeographic patterns in the Mesoamerican tropical lowlands. *Plant Systematic and Evolution*, 306, 10. <https://doi.org/10.1007/s00606-020-01638-y>
- Ortiz-Rodríguez, A. E. (2023a). “rps16-trnQ+ trnL-F cpDNA sequences from *Pilea peridophylla* (Urticaceae)”, Mendeley Data, V1. <https://doi.org/10.17632/xbkfcddnc.1>
- Ortiz-Rodríguez, A. E. (2023b). “trnL-trnF cp DNA sequences from *Pilea* (Urticaceae)”, Mendeley Data, V1. <https://doi.org/10.17632/xhzb287vd.1>
- Pérez-García, E. A., & Meave, J. A. (2005). Heterogeneity of xerophytic vegetation of limestone outcrops in a tropical deciduous forest region in southern México. *Plant Ecology*, 175, 147–163. <https://doi.org/10.1007/s11258-005-4841-8>
- Pérez-García, E. A., Sevilha, A. C., Meave, J. A., & Scariot, A. (2009). Floristic differentiation in limestone outcrops of southern Mexico and central Brazil: A beta diversity approach. *Boletín de la Sociedad Botánica de México* 84, 45–58. https://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0366-21282009000100005
- Pfenninger, M., & Posada, D. (2002). Phylogeographic history of the land snail *Candidula unifasciata* (Helicellinae, Stylomatophora): Fragmentation, corridor migration, and secondary contact. *Evolution* 56 (9), 1776–88. <https://www.jstor.org/stable/3094736>
- R Core Team. (2021). R: A language and environment for statistical computing. R Foundation for statistical computing. Vienna, Austria. Available at <http://www.R-project.org>
- Rull, V. (2020). Neotropical diversification: Historical overview and conceptual insights. In: Rull, V., Carnaval, A. (eds) Neotropical Diversification: Patterns and Processes. Fascinating Life Sciences. Springer, Cham. https://doi.org/10.1007/978-3-030-31167-4_2
- Shaw, J., Lickey, E. B., Schilling, E. E., & Small, R. L. (2007). Comparison of whole chloroplast genome sequences to choose non-coding regions for phylogenetic studies in angiosperms: The tortoise and the hare III. *American Journal of Botany*, 94, 275–288. <https://doi.org/10.3732/ajb.94.3.275>
- Shepherd, L. D., & Heenan, P. B. (2017). Evidence for both long-distance dispersal and isolation in the Southern Oceans: Molecular phylogeny of *Sophora* sect *Edwardsia* (Fabaceae). *New Zealand Journal of Botany*, 55, 334–346. <https://doi.org/10.1080/0028825X.2017.1353527>

- Taberlet, P., Gielly, L., Pantou, G., & Bouvet, J. (1991). Universal primers for amplification of three non-coding regions of plastid DNA. *Plant Molecular Biology*, 17, 1105–1105. <https://doi.org/10.1007/BF00037152>
- Tseng, Y.-H., Huang, H.-Y., Xu, W.-B., Yang, H.-A., Peng, C.-I., Liu, Y., et al. (2019). Phylogeography of *Begonia luzhaiensis* suggests both natural and anthropogenic causes for the marked population genetic structure. *Botanical Studies*, 60, 20. <https://doi.org/10.1186/s40529-019-0267-9>
- Wang, J., Ai, B., Kong, H., & Kang, M. (2017). Speciation history of a species complex of *Primulina eburnea* (Gesneriaceae) from limestone karsts of southern China, a biodiversity hot spot. *Evolutionary Applications*, 10, 919–934. <https://doi.org/10.1111/eva.12495>
- Wendt, T. (1987). Las selvas de Uxpanapa, Veracruz-Oaxaca, México: Evidencia de refugios florísticos cenozoicos. *Anales Instituto De Biología, Universidad Nacional Autónoma De México, Serie Botánica*, 58, 29–54
- Wendt, T. (1993). Composition, floristic affinities, and origins of the canopy tree flora of the Mexican Atlantic slope rain forests. In T. P. Ramamoorthy, R. Bye, A. Lot, & J. Fa (Eds.), *Biological Diversity of Mexico* (pp. 595–680). Oxford University Press.
- Wright, S. (1965). The interpretation of population structure by F-statistics with special regard to systems of mating. *Evolution*, 19(3), 395–420. <https://doi.org/10.2307/2406450>
- Wu, Z.-Y., Liu, J., Provan, J., Wang, H., Chen, C.-J., Cadotte, M. W., Luo, Y.-H., Amorim, B. S., Li, D.-Z., & Milne, R. I. (2018). Testing Darwin's transoceanic dispersal hypothesis for the inland nettle family (Urticaceae). *Ecology Letters*, 21, 1515–1529. <https://doi.org/10.1111/ele.13132>
- Zhu, X., Liang, H., Jiang, H., Kang, M., Wei, X., Deng, L., & Shi, Y. (2022). Phylogeographic structure of *Heteroplexis* (Asteraceae), an endangered endemic genus in the limestone karst regions of southern China. *Frontiers in Plant Science*, 13, 999964. <https://doi.org/10.3389/fpls.2022.999964>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.