

A Review On The Role Of Herbal Medicine In Modulating Hypertension-Related Complications: Mechanistic Insights

Bhoomika Swarnkar, Abhishek Nand, Lokprabha Hirwani, Harkesh Dadsena, Pushpendra Kumar, Chhavi Rahangdale, Yashika Israni, Helina Tandon, Umakant Sahu, Vishal Jain*, Narendra Kumar

University Institute Of Pharmacy, Pt. Ravishankar Shukla University, Raipur, Chhattisgarh- 492010

ABSTRACT

One of the most common chronic conditions in the world, hypertension significantly increases the risk of kidney, cardiovascular, and neurological problems. Despite the wide availability of synthetic antihypertensive medications, adverse effects, high prices, and poor adherence—particularly in areas with limited resources—often restrict long-term control. Due to these restrictions, there is now more interest in herbal remedies, many of which have long been essential components of traditional Chinese medicine, Ayurveda, and Unani. Vascular reconstruction, dysfunction of endothelial cells, oxidative stress, and abnormalities in signalling molecules like nitric oxide & hydrogen sulphide are some of the underlying causes of hypertension that are examined in this review. It also emphasizes how particular medicinal plants alter these pathways. Herbs with a variety of mechanisms of action, such as calcium channel inhibition, renin-angiotensin-aldosterone system regulation, nitric oxide enhancement, and antioxidant effects, such as *Terminalia arjuna*, *Crataegus monogyna*, *Ginkgo biloba*, *Withania somnifera*, *Boerhaavia diffusa*, *Tribulus terrestris*, *Allium sativum*, and *Camellia sinensis*, have antihypertensive activity. Their phytoconstituents, including withanolides, ginsenosides, allicin, and punarnavine, have demonstrated promise in reducing the risk of problems such as erectile dysfunction, nephropathy, atherosclerosis, stroke, and left ventricular hypertrophy. By combining pharmacological facts and molecular insights, this review emphasizes the importance of herbal drugs as long-term and effective adjuncts or alternatives for managing hypertension and its associated difficulties.

Keywords: Renin-angiotensin-aldosterone system (RAAS), Phytoconstituents, Hypertension (HTN), Endothelial dysfunction, Vascular smooth muscle cell (VSMC).

INTRODUCTION

Hypertension, often known as high blood pressure, is a major public health issue around the world. By 2025, it is expected that more than 1.5 billion persons worldwide—or almost one in every three people aged 30 to 79—will have hypertension⁶. Despite advances in diagnosis and care, less than 20% of these people successfully control their blood pressure⁷. The issue is particularly grave in India, where around 24% of adult men and 21% of adults are affected¹. Urban populations have prevalence rates that range from 25% to 30%, while rural areas have a bit lower estimate between 10% and 14%². Even more concerning is that only about 37% of persons with hypertension obtain a diagnosis, 30% receive

treatment, and less than 15% maintain ongoing blood pressure management³.


This growing burden, along with limited treatment accessibility and low long-term adherence, has rekindled interest in both traditional and alternative therapies, particularly those involving medicinal plants. Herbal therapies have been used for millennia in traditional practices such as Ayurveda, Traditional Chinese Medicine (TCM), and Unani, where botanicals are used to lower blood pressure, relieve stress, and improve ⁶vascular health. Recent pharmacological investigations have verified the blood pressure-lowering properties of several plants, include *Rauwolfia serpentina*, *Allium sativum* (garlic), and *Hibiscus sabdariffa*. These kinds of plants are high in bioactive substances—such as

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

alkaloids, flavonoids, polyphenols, and saponins, which give antihypertensive benefits via mechanisms such as ACE inhibition, vasodilation, antioxidant effects, and increased urine output⁴.

Given the prevalence of hypertension worldwide and the disadvantages of synthetic medications—such as side effects, increased costs, and the possibility of polypharmacy—herbal treatments for hypertension are gaining recognized as viable and sustainable supplementary solutions^{5,6,7}. This review evaluates the existing evidence on herbal antihypertensive therapies, stressing their pharmacological effects, clinical effectiveness, safety profiles, and potential role in the holistic management of hypertension⁸.

Even though there are different therapeutic options to manage hypertension and cholesterol using regular drugs, blood pressure control remains unsatisfactory, particularly in low and middle-income countries⁹. This may be due to delays in intensifying treatment regimens like lifestyle changes and traditional medications, as well as a lack of awareness regarding the significance of adhering to prescribed treatments^{10,11}. Worldwide, it has been discovered that 80% of individuals are open to the idea of using herbs for treating health issues, while the exclusive use of traditional herbal medicine among those with hypertension stood at 38.6%, with 47.5% using it in conjunction with antihypertensive medications^{12,13}.



GRAPHICAL ABSTRACT

2. METHODS

2.1 Search Strategy and Method for Identification of Studies

A literature search of published papers, periodicals, magazines, monographs, dissertations, and theses was part of this project. This time range was extended in certain cases because of incomplete information using the search terms "hypertensive," "hypotensive," "anti-hypertensive," and "blood pressure," along with their corresponding translations, the researchers identified medicinal plants that can alter blood pressure for this

study through literature and other resources found in the Professor Eurico Back Library at UNESC. Research was also conducted using Medline, PubMed, Science Direct, academic Google, and Scientific Electronic Library Online to survey plant/drug interactions. The terms "antihypertensive drugs" (antihypertensives), "high blood pressure" (hypertension), and "drug interactions" were linked to the scientific names of plants that were found to alter blood pressure. The Gray search was conducted using Google Scholar as an extra platform, and substantial reference mining was done from the chosen papers.

This was done to make sure that this systematic review did not omit any pertinent publications.

2.2 Pathophysiology of hypertension

The pathophysiological processes linked to the development of hypertension include increased vascular resistance, which is primarily identified by reduced vascular diameter due to increased arterial remodelling and vascular contraction¹⁴. Various elements play a role in the pathophysiology of hypertension (HTN), such as heightened activity of the renin-angiotensin-aldosterone system (RAAS), increased sympathetic nervous system stimulation, vasopressin effects, disruption of G protein-coupled receptor signalling, inflammatory responses, various functions of T-cells, and the range of vasoactive peptides released by other endothelial and smooth muscle cells¹⁵.

Molecular Pathogenesis of Hypertension

Hypertension is defined by abnormalities in the arteries throughout the vascular system, impacting major arteries like the aorta, smaller resistance vessels (150–400µm), and the microcirculation, which includes arterioles and capillaries. Heightened arterial reactivity (responsiveness and strength) results from a lack of proper regulation in the presence of endothelial nitric oxide synthase (e-NOS) and pro-oxidant enzymes, along with heightened basal and stimulated calcium levels resulting from excessive transmembrane calcium permeability via calcium channels, and/or the simultaneous occurrence of vascular smooth muscle cell (VSMC) hyperplasia and hypertrophy (vascular remodelling), can result in increased vasoconstriction. These pathological processes contribute to a greater ratio of the thickness of the vessel wall compared to the size of the arterial lumen¹⁶. The heightened ratio significantly contributes to the onset of hypertension.




Figure. 1 Pathophysiology of Hypertension

2.2.1. Vascular Smooth Muscle Cell (VSMC) Proliferation

Vascular smooth muscle cells (VSMCs) play a role in the development of hypertension, and their growth leads to heightened peripheral resistance by reducing the diameter of arteries¹⁷. To gain an understanding of

these complex changes, it is necessary to investigate the modifying variables that encourage or inhibit VSMC formation in the therapy of hypertension. Growth factors stimulate the cell to enter the cell cycle until the G1 phase, the first checkpoint. VSMC changes are a primary contributor to the development

of hypertension. Normally, these cells are contractile and relaxed, which helps to regulate blood flow and vascular tone. However, in hypertension conditions, VSMCs become more active and lose their contractile properties. This shift, known as phenotypic flipping, allows them to proliferate, migrate, and produce more structural proteins, resulting in larger and more rigid blood vessels¹⁹.

Another signal, platelet-derived growth factor (PDGF), functions similarly by activating pathways that lead to enhanced VSMC proliferation and remodeling¹⁹. High blood pressure exerts mechanical

pressures on VSMCs as well. Mechanosensitive molecules detect these stresses and activate pathways such as YAP/TAZ and RhoA/ROCK, promoting cell growth and shape alterations²⁰. These modifications help to restrict and harden the arteries. Inflammation and oxidative stress exacerbate the situation. Increased amounts of reactive oxygen species (ROS) can activate pathways such as EGFR/AKT/ERK, resulting in additional cell growth. In addition, inflammatory molecules like as TLR4 and NF-KB are activated, promoting inflammation and driving VSMCs into an active, synthetic state²¹.




Figure. 2 Receptor-Mediated RAAS Pathways Contributing to Hypertension

A new study published in 2025 revealed the role of a protein known as ZFP36. This protein reduces the quantity of RGS2, which normally inhibits cell proliferation. When RGS2 is reduced, signals that induce contraction and cell growth become more powerful, exacerbating hypertension. In animal models, inhibiting ZFP36 in VSMCs helped lower blood pressure and enhance vascular shape²². Various signals cause this behaviour. One of the most significant is angiotensin II (Ang II), a chemical that causes high blood pressure. It activates the AT1R receptor on VSMCs, which stimulates intracellular pathways such as ERK and elevates calcium levels in the cells, boosting their development and migration²³.

Several potential therapeutics are being investigated to prevent or treat VSMC overgrowth. One of these is Resolving D1, a molecule that reduces inflammation and inhibits damaging signalling pathways. Research indicates that it can inhibit Ang II-induced vascular remodelling and lower blood pressure in animals²³.

2.2.2. Endothelial Cells

Endothelial cells form a single layer that lines the interior surface of blood arteries and are critical for vascular function. These cells control processes like blood flow, vascular tone, inflammation, blood coagulation, and barrier permeability. In healthy settings, endothelial cells release chemicals like nitric oxide (NO), which helps relax blood arteries and prevent clot formation or severe inflammation²⁴.

Endothelial cells frequently fail in the presence of hypertension and cardiovascular disease. This dysfunction can be caused by high levels of angiotensin II (Ang II), oxidative stress, and disrupted blood flow. Ang II not only increases blood pressure but also disturbs the protective role of endothelial cells by increasing the formation of reactive oxygen species (ROS), lowering nitric oxide bioavailability, and harming the inner lining of blood vessels²⁵.

According to recent research, Ang II can cause endothelial cells to die through a process known as ferroptosis. This process is initiated by lipid

peroxidation and involves receptors such as CD36, which absorb oxidized lipids and decrease the protective barrier created by these cells. As a result, the vascular wall is more susceptible to inflammation and injury²⁶.

Oxidative stress is a primary cause of endothelial damage. When ROS levels rise due to mitochondrial

malfunction or NADPH oxidase activation, they activate inflammatory pathways such as NF-KB, JAK/STAT, and MAPK. These signalling pathways promote the production of adhesion molecules like ICAM-1 and VCAM-1, which attract immune cells to the vascular wall and increase inflammation and remodeling²⁷.




Figure. 3 Oxidative stress and role of NOS on blood vascular system

2.2.3. Repertoire of Signalling Molecules

The following signalling molecules become essential to the pathophysiology of hypertension assuming a homeostatic imbalance²⁸. Fortunately, various plant and herbal extracts, along with their individual metabolites, can influence signalling cascades related to the physiology of the cardiovascular system (refer to Section Herbs and Spices Most Commonly Used for Treatment of Hypertension, below). These herbs not only protect blood vessels but could also potentially reverse the changes associated with hypertension. This is especially true if changes in hypertensive patients are dealt with before they reach a decompensated state.

A. Reactive Oxygen Species

In a healthy state, pro-oxidant activity is counterbalanced by antioxidative agents. Nevertheless, when this balance is disrupted, the disorganized environment gives rise to pathological conditions like hypertension, atherosclerosis, and various other vascular complications²⁹. ROS, such as superoxide anions ($O_2^{\bullet -}$) and hydroxyl ions ($OH^{\bullet -}$), play a critical role in hypertension pathophysiology

by causing oxidative stress³⁰. Reactive oxygen species (ROS) are produced by nicotinamide-adenine dinucleotide phosphate (NADPH) oxidase and other enzymatic activities, particularly those associated with the electron transport chain in mitochondria³¹. ROS are also liable for the oxidation of low-density lipoprotein (LDL), which causes inflammation and increases VSMC proliferation. Inflammation and increased proliferation both play important roles in causing plaque formation, which may contribute to high blood pressure. Interestingly, in animals, blood pressure drops when ROS generation is reduced with antioxidant treatment³².

B. Nitric Oxide

Nitric oxide (NO), a key chemical produced by endothelial cells, has a direct effect on vascular smooth muscle cells (VSMCs) and is required for the control of vascular tone, blood pressure, and pathologic remodelling. Under normal settings, NO is mainly generated by endothelial nitric oxide synthesis (e-NOS) and spreads into vascular smooth muscle cells (VSMCs), which activates soluble guanylate cyclase (sGC). This activation increases the quantity of cyclic guanosine monophosphate (cGMP), which

stimulates protein kinase G (PKG). PKG reduces cellular calcium levels, smooth muscle elasticity, and decreases VSMC proliferation and migration³³.

Beyond the typical cGMP-dependent mechanisms, NO has cGMP-independent actions. One important method is S-nitrosation, in which NO covalently changes proteins such as RhoA and Raf-1, inhibiting their signaling capabilities. These proteins are implicated in the ERK/MAPK pathway, which typically promotes VSMC development and migration. Inhibiting this mechanism causes cell cycle arrest and decreased proliferation³⁴.

Recent research has demonstrated that NO can control VSMC growth by modulating ubiquitin-proteasome pathways. Specifically, NO promotes the degradation of UbcH10, an E2 enzyme required for cell cycle progression. Downregulating UbcH10 causes arrest during the G1/S phase transition and lessens neointimal thickening after vascular damage³⁵. Stimulating NADPH oxidase produces superoxide anion ($O_2^{\bullet-}$), which combines with NO to form peroxynitrite, a potent oxidant. This peroxynitrite then induces oxidative breakdown of the e-NOS cofactor tetrahydrobiopterin (BH4), yielding the inactive dihydrobiopterin (BH2). This produces more superoxide anion, which eventually lowers BH4 levels. As a result, a shift in balance from NO to superoxide production occurs, which is known as the uncoupling of NOS³⁶.

C. Hydrogen Sulphide

In VSMCs and perhaps endothelial cells, cystathionine γ -lyase (CSE) converts L-cysteine to produce hydrogen sulphide (H_2S)³⁷. H_2S has a vasorelaxant impact on VSMCs via increasing intracellular cGMP levels and activating ATP-dependent potassium channels^{37,38}.

Hydrogen sulphide (H_2S) is recognized as an important gas transmitter for maintaining vascular health. In VSMCs, H_2S regulates physiological activities such relaxation, proliferation, migration, and oxidative stress resistance. The circulatory system manufactures it naturally through the enzyme cystathionine γ -lyase (CSE). H_2S relaxes vascular smooth muscle by activating ATP-sensitive potassium (K_{ATP}) channels, resulting in membrane hyperpolarization and decreased calcium influx. This

reduces intracellular Ca^{2+} and promotes vasodilation³⁹. H_2S prevents vascular calcification by activating the KEAP1-NRF2-NQO1 pathway, a key antioxidant mechanism. This reduces oxidative damage and inhibits signals that promote calcification in VSMCs⁴⁰. H_2S 's actions make it an attractive target for therapies to prevent vascular disorders such as hypertension, atherosclerosis, and restenosis.

3. Herbal Approaches to Addressing Hypertension-Related Issues: An In-Depth Physiological Analysis

1. Left Ventricular Hypertrophy (LVH) and Heart Failure: -

Pathophysiology of Disease - Chronic high blood pressure increases afterload, which is the pressure the left ventricle must overcome to pump blood. As a result, myocardial cells expand, causing left ventricular hypertrophy (LVH) to maintain cardiac output. Nonetheless, this favorable hypertrophy can turn maladaptive over time, resulting in increased stiffness of the myocardium, fibrosis, poor relaxation (diastolic dysfunction), and eventually heart failure, whether with preserved or reduced ejection fraction⁴¹.

At the cellular level, elevated blood pressure activates hypertrophic signalling pathways such MAPKs and alters calcium control, resulting in contractile dysfunction and poor energy efficiency⁴².

Herbal Solutions and Mechanisms: -

A. *Terminalia arjuna* - The deciduous tree *Terminalia arjuna*, also referred to as Arjuna, is indigenous to the Indian subcontinent and has long been used as a cardiac tonic in Ayurvedic medicine. Its effectiveness in treating cardiovascular disorders, especially left ventricular hypertrophy (LVH) and heart failure (HF), which are frequently brought on by chronic hypertension, is supported by recent pharmacological and clinical research.

Pathophysiology of *Terminalia arjuna* - The cardi tonic properties of *Terminalia arjuna* are well established. According to clinical research, taking 400 mg of Oxyjun® (standardized Arjuna extract) daily enhanced cardiac output by 6.3%, decreased tiredness by 22.5%, and improved left ventricular ejection

fraction⁴³. Mechanistically, its triterpenoids and flavonoids modulate intracellular calcium cycling, enhance myocardial contractility, and reduce oxidative stress-induced apoptosis in cardiac cells⁴².

The heart uses left ventricular hypertrophy to make up for elevated systemic vascular resistance in persistent hypertension. After maintaining cardiac output for a while, this adaptive mechanism turns maladaptive because of fibrosis, higher myocardial oxygen demand, and poorer diastolic relaxation. Eventually, these alterations lead to cardiac failure with a preserved or decreased ejection fraction⁴³.

Key Pathways Involved include –

- Activation of MAPK and NF-KB, which results in the production of inflammatory and fibrotic genes.
- Oxidative stress and malfunctioning mitochondria.
- Diastolic filling and contractility are hampered by improper calcium management⁴³.

Phytoconstituents like Triterpenoids (arjunic acid, arjunolic acid), Flavonoids (arjunone, arjunolone), Glycosides, tannins, and minerals (especially CoQ10-like activity) constituents contribute to Arjuna's antioxidant, anti-inflammatory, and cardiogenic properties. Terminalia arjuna modulates calcium ion channels and sarcoplasmic reticulum function, improving myocyte relaxation and contraction cycles, thus supporting both systolic and diastolic function. By downregulating caspase-3 and maintaining mitochondrial membrane potential, it prevents myocardial cell death. This is important for sustaining cardiac energy metabolism, particularly during ischemia stress⁴⁴.

B. *Crataegus monogyna* - Native to Europe and Asia, hawthorn (*Crataegus monogyna*) is a medicinal shrub that has long been used in both traditional and contemporary herbal therapy to treat circulatory disorders⁴⁵. It is highly relevant for the treatment of heart failure (HF) and left ventricular hypertrophy (LVH), two common complications of long-term hypertension, because its berries, leaves, and flowers contain a wide variety of bioactive compounds that have

cardiotonic, anti-hypertrophic, and vasodilatory effects^{46,47}.

Pathophysiology of *Crataegus monogyna* - To maintain cardiac output, chronic hypertension raises afterload over time, which induces adaptive myocardial hypertrophy. This process eventually turns pathological because of:

- Fibrosis of the heart.
- Stiffness of the left ventricle increased.
- Calcium signalling impairment.
- Modified expression of some genes in cardiomyocytes⁴⁸.

These changes disrupt diastolic function, promote electrical remodelling, and increase the risk of systolic heart failure.

2. Stroke (Ischemic and Haemorrhagic): -

Pathophysiology of Disease - Chronic hypertension harms small blood vessels in the brain, leading to arteriosclerosis, lipohyalinosis, and the development of microaneurysms. This vascular condition increases the risk of haemorrhagic stroke because of blood vessel due to vessel blockage. Furthermore, hypertension weakens cerebral autoregulation, affects the integrity of the blood-brain barrier (BBB), elevates levels of inflammatory cytokines, and triggers neuronal apoptosis^{49,50}.

Herbal Solutions and Mechanisms: -

- A. *Ginkgo biloba*** - In the search for complementary or alternative therapies, *Ginkgo biloba*, a traditional medicinal plant native to China, has received interest for its neuroprotective properties. The standardized extract, notably EGb 761, contains flavonoids (such as quercetin and kaempferol) and terpenoids (such as ginkgolides and bilobalide) with antioxidant, anti-inflammatory, antiplatelet, and vasodilatory properties that are useful in stroke therapy. Extracts, notably ginkgo diterpene lactone meglumine (GDLM), have been shown to help with stroke rehabilitation. In a randomized research involving 3,448 individuals with ischemic stroke, there was a 6.8% increase in the rate of positive recovery (as defined by the modified Rankin Scale

scores of 0-1) compared to those receiving a placebo⁴⁹.

Pathophysiology of *Ginkgo biloba* - GDLM enhances cerebral blood flow, inhibits platelet activating factor reducing thrombosis, protects endothelial cells, and promotes neuronal resistance to hypoxia through antioxidant mechanisms⁵⁰. Ischemic stroke is caused by arterial obstruction, which leads to diminished cerebral blood flow, ATP depletion, glutamate-induced excitotoxicity, oxidative stress, and neuronal death. Hemorrhagic stroke occurs when blood vessels break, resulting in hematoma formation, oxidative damage, and inflammation. Ischemic stroke is caused by arterial obstruction, which leads to diminished cerebral blood flow, ATP depletion, glutamate-induced excitotoxicity, oxidative stress, and neuronal death. Hemorrhagic stroke occurs when blood vessels break, resulting in hematoma formation, oxidative damage, and inflammation.⁵¹ *Ginkgo biloba*, particularly its extract EGb 761, provides neuroprotection via antioxidant, anti-inflammatory, and vasodilating properties. Its flavonoids and terpenoids absorb free radicals, enhance mitochondrial activity, and increase nitric oxide-mediated vasodilation⁵². In hemorrhagic stroke, *Ginkgo* decreases edema and maintains the blood-brain barrier, limiting neuronal damage⁵³.

B. *Withania somnifera* - *Withania somnifera* (Ashwagandha), an important plant in Ayurvedic medicine, is recognized for its adaptogenic, anti-inflammatory, neuroprotective, and antioxidant effects. The root extract contains bioactive substances such as withanolides, sitoindosides, and alkaloids, which enhance its medicinal potential. Recent experimental and clinical research have highlighted its neurorestorative efficacy in central nervous system illnesses, including stroke⁵⁴.

Pathophysiology of *Withania somnifera* - In ischemic stroke, the cessation of cerebral blood flow causes oxidative stress, excitotoxicity, and activation of the inflammatory cascade. Ashwagandha mitigates these effects by reducing lipid peroxidation, inhibiting pro-inflammatory cytokines (e.g., TNF- α , IL-6), and enhancing endogenous antioxidant defences (e.g., superoxide dismutase, catalase)⁵⁵.

In haemorrhagic stroke, where bleeding into brain tissue causes edema and neuronal injury,

Ashwagandha reduces neuroinflammation and supports the integrity of the blood-brain barrier. Its ability to upregulate brain-derived neurotrophic factor (BDNF) promotes neurogenesis and neuronal recovery post-stroke⁵⁶.

3. Chronic Kidney Disease (Hypertensive Nephropathy) –

Pathophysiology of Disease - Renal arteriole constriction and thickening brought on by hypertension results in ischemia and damage to the glomeruli and tubules. Proteinuria, a decline in glomerular filtration rate (GFR), and a gradual loss of nephrons lead to chronic kidney disease (CKD)⁵⁷.

Herbal Solutions and Mechanisms: -

A. *Boerhaavia diffusa* - Punarnava, also known as *Boerhaavia diffusa*, is a traditional Ayurvedic herb that is widely used for its nephroprotective, diuretic, anti-inflammatory, and antioxidant qualities. Important bioactive substances found in the plant, including flavonoids, lignans, boeravinones, and punarnavine, have been shown to have protective effects on renal tissues⁵⁸. Long-term uncontrolled hypertension causes glomerular sclerosis and tubulointerstitial fibrosis, which in turn causes hypertensive nephropathy, a progressive form of chronic kidney disease (CKD). Punarnava has shown promise in decreasing these side effects while maintaining renal function.

Pathophysiology of *Boerhaavia diffusa* - Glomerular ischemia, arteriolar constriction, glomerulosclerosis, and progressive nephron loss because of persistently elevated intraglomerular pressure are the hallmarks of hypertensive nephropathy. The processes behind these changes include oxidative stress, fibrosis, inflammation, and activation of the renin-angiotensin-aldosterone system (RAAS)⁵⁹.

Boerhaavia diffusa has nephroprotective properties by:

- Reducing oxidative damage by increasing the levels of endogenous antioxidant enzymes, such as superoxide dismutase and catalase.

- Inhibiting pro-inflammatory cytokines including TNF- α and IL-6 to modify inflammatory pathways.
- Displaying diuretic properties that alleviate renal stress and reduce systemic blood pressure.
- Reducing fibrotic signaling (such as TGF- β 1), which prevents the formation of extracellular matrix in renal tissues.

B. *Tribulus terrestris* - lowers intraglomerular pressure and promotes natriuresis⁶⁰ by inhibiting RAAS and lowering aldosterone levels. In Ayurvedic and traditional medical systems, *Tribulus terrestris*, also referred to as Gokshura, is a well-known medicinal plant that has nephroprotective, antihypertensive, diuretic, and anti-inflammatory qualities. Glycosides⁶¹, alkaloids, flavonoids, and saponins—particularly protodioscin—are the main bioactive components. *Tribulus terrestris* has been investigated for its potential to treat renal problems, notably those brought on by chronic hypertension, and has long been used as a renal tonic.

Pathophysiology of *Tribulus terrestris* - Consequently, the strain on the renal vasculature is reduced. By increasing the quantities of endogenous enzymes such as glutathione peroxidase and superoxide dismutase, it also strengthens antioxidant defences and combats oxidative damage. Additionally, it helps prevent immune-related tissue damage through its anti-inflammatory properties, which are mediated by the reduction of cytokines including TNF- α and IL-6. Crucially, new research indicates that *T. terrestris* may block pathways linked to fibrosis, including the TGF- β 1/Smad axis, which is essential for the development of chronic renal scarring.^{62,63}

4. Atherosclerosis and Vascular Dysfunction: -

Pathophysiology of Disease - The endothelium deteriorates more quickly due to high blood pressure, which causes LDL to oxidize and plaque to form. Because of the increased arterial stiffness and clot formation, this increases the risk of peripheral artery disease and heart attacks. High blood pressure-

induced endothelial damage leads to LDL oxidation and the formation of atherosclerotic plaques, increasing the risk of cardiovascular events⁴². Chronic arterial disease known as atherosclerosis is brought on by endothelial dysfunction, which is frequently brought on by smoking, diabetes, hypertension, or hyperlipidaemia⁶⁴. Endothelial damage promotes the migration of low-density lipoprotein (LDL) into the intima, where it is oxidatively modified to become oxidized LDL (oxLDL), which sets off an inflammatory cascade⁶⁵. When monocytes enter the artery wall, mature into macrophages, and generate foam cells, early fatty streaks are created. Fibrous plaque development is facilitated by the proliferation of smooth muscle cells and the deposition of extracellular matrix. TNF- α and IL-1 β are examples of inflammatory mediators that increase matrix metalloproteinase activity, weakening plaques and raising the chance of rupture⁶⁶. Reduced nitric oxide (NO) bioavailability, oxidative stress, and decreased vasodilation all contribute to vascular dysfunction, which in turn encourages thrombosis and vascular stiffness. Acute ischemia episodes and thrombus development are brought on by plaque rupture, which reveals thrombogenic material⁶⁵.

Herbal Solutions and Mechanisms: -

A. *Allium sativum* (garlic): - Garlic, or *Allium sativum*, is a popular medical herb known for its heart-healthy properties. Garlic is rich in bioactive sulfur-containing compounds that have anti-inflammatory, anti-thrombotic, antioxidant, and lipid-lowering properties, including allicin, ajoene, and S-allyl cysteine⁶⁷. Its medical benefits are particularly notable in cardiovascular disease, where it helps to lower risk factors such as hyperlipidemia, hypertension, and oxidative stress. It has traditionally been utilized in both a culinary and therapeutic environment⁶⁸.

Pathophysiology of *Allium sativum* - Plaques in atherosclerosis form because of endothelial dysfunction and lipid accumulation. Oxidized low-density lipoprotein (oxLDL) causes an inflammatory response that draws monocytes, which then change into foam cells, resulting in the creation of fatty streaks and the progression of plaques⁶⁹. Garlic counteracts this process by blocking HMG-CoA reductase, lowering LDL cholesterol production, and

decreasing LDL oxidation via antioxidant activity⁷⁰. It increases nitric oxide (NO) generation, improves endothelial function and vasodilation, and lowers oxidative stress. Reducing cytokines like IL-6 and TNF- α can reduce plaque destabilization and vascular damage^{71,72}. *Allium sativum* regulates lipid metabolism, oxidative stress, and vascular inflammation via various pathways, thereby addressing the fundamental pathophysiological causes of atherosclerosis and vascular dysfunction.

B. *Camellia sinensis* (Green Tea) - Green tea, referred to as *Camellia sinensis*, is a rich source of polyphenolic compounds, especially catechins like epigallocatechin-3-gallate (EGCG), which have powerful antioxidant, anti-inflammatory, and lipid-modifying actions⁷³. Green tea is linked to lower cardiovascular morbidity and mortality, according to epidemiological studies, because it improves lipid profiles, endothelial function, and reduces oxidative stress. Catechins, particularly EGCG, improve arterial flexibility and prevent plaque formation by decreasing LDL oxidation and increasing endothelial NO synthase (eNOS) activity^{42,74}. Green tea's cardiovascular advantages originate from its capacity to decrease LDL oxidation, increase vascular responsiveness, and regulate important signalling pathways involved in vascular health.

Pathophysiology of *Camellia sinensis* - Endothelial damage triggers atherosclerosis, which is then followed by the entry of LDL cholesterol into the intima, oxidative modification, and the recruitment of monocytes, which develop into foam cells and aid in the creation of plaque⁷⁵. Vascular dysfunction is sustained by oxidative stress and inflammation, which lowers the bioavailability of nitric oxide. By scavenging reactive oxygen species, preventing LDL oxidation, and increasing endothelial nitric oxide synthase (eNOS) activity, green tea catechins—EGCG in particular—oppose these processes and improve vasodilation⁷⁶. Additionally, catechins reduce vascular inflammation and stabilize plaques by downregulating inflammatory mediators such NF- κ B, TNF- α , and interleukins⁷⁷.

5. Erectile Dysfunction (ED) –

Pathophysiology of Disease - Hypertension-induced arterial stiffness and endothelial dysfunction impair

blood flow to the penile tissue. Nitric oxide (NO) deficiency inhibits the smooth muscle relaxation necessary for erection. Reduced nitric oxide availability and increased vascular stiffness, which hinder blood flow to the erectile tissue, are the causes of erectile dysfunction^{42,78}. Furthermore, cGMP is broken down more quickly by elevated phosphodiesterase type 5 (PDE5) activity, which restricts smooth muscle relaxation. Hormonal abnormalities like low testosterone can lower libido and erectile function, while chronic inflammation and vascular stiffness further impair penile hemodynamics^{79,80}.

Herbal Solutions and Mechanisms: -

A. *Panax ginseng* - *Panax ginseng*, sometimes referred to as Korean or Asian ginseng, is a herb used extensively in East Asian traditional medicine to improve sexual health, energy, and general physiological robustness. Ginsenosides, its active components, have neuroendocrine-modulating, vasodilatory, and antioxidant properties, making it a promising treatment for erectile dysfunction (ED)⁸¹. According to preclinical and clinical research, *P. ginseng* promotes nitric oxide (NO) generation, enhances endothelial function, and increases libido, all of which lead to better erectile performance⁸².




Pathophysiology of *Panax ginseng* - Endothelial dysfunction, reduced NO bioavailability, oxidative damage, hormonal dysregulation, and neurovascular impairment all contribute to poor penile hemodynamics⁸³. Ginsenosides have been shown in animal studies to improve erectile function by influencing central dopaminergic pathways and increasing erectile responses. Ginsenosides also increase the activity of endothelial nitric oxide synthase (eNOS) and cyclic guanosine monophosphate (cGMP), which relaxes smooth muscles in the corpus cavernosum and improves penile blood flow. Additionally, ginsenosides prevent oxidative damage to the vascular endothelium, which preserves erectile capacity⁸⁴.





B. *Mucuna pruriens* - *Mucuna pruriens*, often known as velvet bean, is a traditional Ayurvedic herb used to improve male sexual performance. Its seeds are rich in L-DOPA, the natural precursor to dopamine, and contain antioxidant

constituents such as flavonoids and alkaloids. Emerging research suggests these components contribute to improved erectile function by acting on central and peripheral neuroendocrine pathways⁸⁵.

Pathophysiology of *Mucuna pruriens* - Erectile function depends critically on nitric oxide (NO)-

mediated vasodilation in penile tissue. *M. pruriens* enhances dopamine levels in the brain, activating the hypothalamic–pituitary–gonadal axis, leading to increased gonadotropins and testosterone release, which in turn supports NO synthesis and endothelial function. It also mitigates oxidative stress in the corpus cavernosum via antioxidant actions, protecting NO availability and penile tissue structure^{85,86}.

S . N .	Botanic al Name (Common Name)	Mechanism (s) of Action (Target receptor)	Part Used	Antihypertensive Phytoconstituents (Major Active)	Chemical Structure	References
1 .	<i>Terminalia arjuna</i> (Arjuna)	PPAR α receptor activation Ca ²⁺ -handling proteins Endothelial nitric oxide (NO)	Stem bark	Arjunolic acid , arjunic acid, arjunosides (glycosides), tannins, ellagic acid, gallic acid, flavonoids and oligomeric proanthocyanidins		87, 88, 89
2 .	<i>Crataegus monogyna</i> (Hawthorn)	Na ⁺ /K ⁺ -ATPase inhibition Endothelial NO synthase (eNOS) activation TGF- β 1/Smad signaling inhibition	Leaves/flowers, and berries	Vitexin , Hyperoside, rutin, quercetin, procyanidins.		90, 91, 92, 93, 94
3 .	<i>Ginkgo biloba</i> (Ginkgo)	Endothelial NO synthase (eNOS) activation Platelet-activating factor (PAF) receptor antagonism NF- κ B pathway inhibition	Leaves	Ginkgolide B Flavonoids (quercetin, kaempferol), ginkgolides, bilobalide		95, 96, 97, 98

4	<i>Withania somnifera</i> (Ashwagandha)	NMDA receptor modulation Nrf2/ARE pathway activation NF-κB pathway inhibition	Roots, leaves	Withanolides Withaferin A, withanolide A sitoindosides		99, 100, 101
5	<i>Boerhavia diffusa</i> (Punarnava)	TGF-β1/Smad signaling inhibition Renin–Angiotensin–Aldosterone System (RAAS) modulation	Roots, leaves	Punarnavine, boeravinones Flavonoids alkaloids		102, 103, 104, 105
6	<i>Tribulus terrestris</i> (Puncture Vine)	TGF-β1/Smad signaling inhibition Renin–Angiotensin–Aldosterone System (RAAS) modulation Nrf2/ARE pathway activation	Fruits, roots	Protodioscin Saponins (dioscin), flavonoids Diosgenin, Polyphenols, alkaloids		106, 107, 108, 109
7	<i>Allium sativum</i> (garlic)	HMG-CoA reductase inhibition NF-κB inhibition eNOS activation → ↑ NO	Bulb	Allicin, S-allyl cysteine, ajoene, diallyl disulfide, flavonoids, diallyl trisulfide, Organosulfur compounds		110, 111, 112, 113




8	<i>Camellia sinensis</i> (Green Tea)	HMG-CoA reductase inhibition NF-κB pathway inhibition eNOS activation → ↑ NO	Leaves	Epigallocatechin gallate (EGCG), catechins, theaflavins, flavonoids		114, 115, 116, 117
9	<i>Panax ginseng</i> (Ginseng)	eNOS activation → ↑ NO release Nitrogenic nerve signaling modulation Nrf2/ARE pathway activation, ↓ ROS	Root	Ginsenosides Rg1 (Rb1, Rg3, Re, Rd), polysaccharides, flavonoids, Total ginsenosides		118, 119, 120,
10	<i>Mucuna pruriens</i> (Velvet Bean)	Dopamine D2 receptor activation Hypothalamic–pituitary–gonadal (HPG) axis activation Nrf2/ARE pathway activation	Seeds	L-DOPA (Levodopa), alkaloids, flavonoids, saponins, phenolic compounds, antioxidants (quercetin, catechins)		121, 122, 123

Table No. 1 Herbal Approaches for Managing Hypertension with Their Mechanisms of Action.

4. Herbal medicines used for the treatment of hypertension

Side effects are common with several antihypertensive medications used to treat hypertension. To treat it, scientific research suggests a variety of lifestyle changes as well as the application of appropriate medicinal plants.⁴¹ Certain herbs and spices contain secondary metabolites that have antihypertensive effects. Many herbal remedies help

manage and lower hypertension by providing antioxidant, anti-inflammatory, and anti-apoptotic benefits, activating the e-NOS-NO signalling pathway, reducing endothelial permeability, and promoting angiogenesis.⁴²

The list of medicinal plants that have been found to be useful in managing and treating hypertension is provided in Table 1.

S.N .	Botanical Name	Common Name	Mechanism(s) of Action	Part Used	Antihypertensive Phytoconstituents (Major Active)	Reference
1.	<i>Allium sativum</i>	Garlic	Vasodilation, ACE inhibition, antioxidant	Bulb	Allicin, S-allyl cysteine, diallyl disulfide	124
2.	<i>Hibiscus sabdariffa</i>	Rosella	Vasodilation via Anthocyanins, ACE inhibition, diuretic	Calyx	Anthocyanins (delphinidin-3-sambubioside), protocatechuic acid	125
3.	<i>Crataegus oxyacanthine</i>	Hawthorn	Ca ²⁺ Channel blockage, Antioxidant, Vasodilation	Berry	Flavonoids (vitexin, rutin), oligomeric procyanidins	126
4.	<i>Nigella sativa</i>	Black Cumin	↓SNS stimulation, antioxidants, Ca ²⁺ blockage	Seed	Thymoquinone, α-hederin, nigellidine	127
5.	<i>Rauwolfia serpentina</i>	Sarpa Gandha	Sympatholytic via reserpine, catecholamine depletion	Root	Reserpine, ajmaline, serpentine (indole alkaloids)	128
6.	<i>Boerhavia diffusa</i>	Punarnava	Diuretic, Ca ²⁺ channel blockade, antioxidant	Root	Punarnavine, boeravinones (rotenoids)	129
7.	<i>Moringa oleifera</i>	Drumstick	Vasodilation, ACE inhibition, antioxidant	Leaf	Quercetin, chlorogenic acid, isothiocyanates	130
8.	<i>Coptis chinensis</i>	Goldthread	ACE inhibition, ↓Ang II, ↑e-NOS	Rhizome	Berberine, palmatine, coptisine	131
9.	<i>Panax ginseng</i>	Ginseng	↑NO production, vasorelaxation, anti-inflammatory	Root	Ginsenosides (Rb1, Rg1, Re)	132

10.	<i>Apium graveolens</i>	Celery	Ca ²⁺ channel inhibition, diuretic	Seed	3-n-butylphthalide, apigenin, luteolin	133
11.	<i>Salvia Miltiorrhiza</i>	Red sage	K ⁺ channel activation, ACE inhibition, antioxidant	Root	Tanshinones, salvianolic acid B, rosmarinic acid	134
12.	<i>Peganum harmala</i>	Esfand	NO modulation, vasodilation, monoamine oxidase inhibition	Seed	Harmine, harmaline (β -carboline alkaloids)	135
13.	<i>Peperomia pellucida</i>	Shiny Bush	ACE inhibition, antioxidant, diuretic	Whole Plant	Pellucidin A, quercetin, tannins	136
14.	<i>Syzygium polyanthum</i>	Indonesian Bay-leaf	Vasorelaxation via phenolics, NO-dependent pathways	Leaf	Phenolic acids, flavonoids (quercetin, rutin)	137
15.	<i>Bidens Pilosa</i>	Blackjack	Diuretic, NO pathway enhancement, Ca ²⁺ channel inhibition	Leaf	Polyacetylenes, rutin, caffeic acid	138
16.	<i>Andrographis paniculata</i>	Kalmegh	ACE inhibition, Ca ²⁺ & β -blockade, $\uparrow$ NO	Leaf/Aerial Parts	Andrographolide, neoandrographolide	139
17.	<i>Terminalia arjuna</i>	Arjuna	Vasorelaxation, antioxidant, cardiogenic	Bark	Arjunolic acid, arjunosides, tannins	140
18.	<i>Camellia sinensis</i>	Green Tea	Antioxidant, vasorelaxant (polyphenols), ACE inhibition	Leaf	EGCG (epigallocatechin gallate), catechins, theaflavins	141

19.	<i>Ocimum sanctum</i>	Holy basil	Diuretic, antioxidant, vasodilatory flavonoids	Leaf	Eugenol, rosmarinic acid, ursolic acid	142
20.	<i>Glycyrrhiza glabra</i>	Licorice	Mineralocorticoid effect, though use limited due to ↑BP risk	Root	Glycyrrhizin, liquiritigenin (note: ↑BP risk)	143
21.	<i>Cinnamomum zeylanicum</i>	Cinnamon	Insulin sensitization, vasodilation, antioxidant	Bark	Cinnamaldehyde, procyanidins	144
22.	<i>Withania Somnifera</i>	Ashwagandha	↓Cortisol, antioxidant, ↑NO	Root	Withanolides, sitoindosides	145
23.	<i>Valeriana officinalis</i>	Valerian	CNS depressant, ↓SNS, sedative	Root	Valerenic acid, flavonoids	146
24.	<i>Crocus sativus</i>	Saffron	Ca ²⁺ channel blocker, antioxidant	Stigma	Crocin, safranal, crocetin	147
25.	<i>Phyllanthus amarus</i>	Bhumi amla	Antioxidant, ACE inhibition	Whole Plant	Phyllanthin, hypophyllanthin, ellagic acid	148
26.	<i>Justicia gendarussa</i>	Willow-leaved justicia	Vasodilation, Ca ²⁺ blockade	Leaf	Gendarusin A, flavonoids	149
27.	<i>Tribulus terrestris</i>	Gokshura	Diuretic, vasorelaxant saponins	Fruit	Protodioscin, saponins	150
28.	<i>Eclipta alba</i>	Bhringraj	Vasodilator, NO activation	Whole plant	Wedelolactone, ecliptine	151

29.	<i>Vitex negundo</i>	Nirgundi	Vasodilator, anti-inflammatory	Leaf	Flavonoids (luteolin, casticin), iridoids	152
30.	<i>Barleria prionitis</i>	Vajradanti	Diuretic, vasorelaxant	Leaf	Barlerin, iridoid glycosides	153
31.	<i>Azadirachta indica</i>	Neem	ACE inhibition, antioxidant, vasodilation	Leaf	Nimbin, nimbolide, quercetin	154
32.	<i>Piper betle</i>	Betel leaf	Vasorelaxation, calcium antagonism	Leaf	Hydroxychavicol, eugenol	155
33.	<i>Leea indica</i>	Bandicot berry	Ca ²⁺ channel blocker, anti-inflammatory	Leaf	Flavonoids, phenolic acids	156
34.	<i>Centella asiatica</i>	Gotu Kola	NO-mediated vasodilation, diuretic	Leaf	Asiaticoside, madecassoside	157
35.	<i>Sida cordifolia</i>	Bala	Adrenergic modulation, mild hypotensive	Leaf	Ephedrine, vasicinol (low doses hypotensive)	158
36.	<i>Aegle marmelos</i>	Bael	Antioxidant, vasodilation via polyphenols	Leaf	Marmelosin, aegeline, flavonoids	159
37.	<i>Cissampelos pareira</i>	velvetleaf	ACE inhibition, diuretic, antioxidant	Root	Cissampeline, hayatin (alkaloids)	160
38.	<i>Solanum nigrum</i>	Black nightshade	Diuretic, β -blocking potential	Leaf	Solamargine, solanine, flavonoids	161

39.	<i>Cassia occidentalis</i>	Coffee senna	Ca ²⁺ channel blockade, diuretic	Leaf	Emodin, chrysophanol, anthraquinones	162
40.	<i>Mentha piperita</i>	Peppermint	Vasodilation, NO modulation	Leaf	Menthol, rosmarinic acid	163
41.	<i>Tamarindus indica</i>	Tamarind	ACE inhibition, diuretic, antioxidant	Fruit pulp	Procyanidins, tartaric acid, catechins	164
42.	<i>Rosmarinus officinalis</i>	Rosemary	Vasodilation, ACE inhibition, Ca ²⁺ blockade	Leaf	Rosmarinic acid, carnolic acid, ursolic acid	165
43.	<i>Mangifera indica</i>	Mango	Polyphenol-mediated vasodilation	Leaf/fruit	Mangiferin, gallic acid, quercetin	166
44.	<i>Curcuma longa</i>	Turmeric	Antioxidant, NO modulation, anti-inflammatory	Rhizome	Curcumin, demethoxycurcumin	167
45.	<i>Avena sativa</i>	Oats	Diuretic, endothelial function improvement	Grain	Avenanthramides, β -glucan	168
46.	<i>Olea europaea</i>	Olive leaf	ACE inhibition, vasodilation, antioxidant	Leaf	Oleuropein, hydroxytyrosol, verbascoside	169
47.	<i>Orthosiphon stamineus</i> (syn. <i>O. aristatus</i>)	Java tea	Diuretic, RAAS modulation, antioxidant	Aerial	Rosmarinic acid, sinensetin, eupatorin	170
48.	<i>Apocynum venetum</i>	Luobum	β -blocker-like, Ca ²⁺ antagonism,	Leaf	Flavonoids (hyperoside), quercetin glycosides	170

			anxiolytic (↓SNS)			
49	<i>Eucommia ulmoides</i>	Du-zhong	ACE inhibition, NO↑, vasodilation	Bark/Leaf	Geniposidic acid, pinosresinol diglucoside	171
50	<i>Zingiber officinale</i>	Ginger	Ca ²⁺ channel blockade, ACE↓, vasodilation	Rhizome	6-Gingerol, shogaols	170
51	<i>Allium cepa</i>	Onion	ACE↓, vasodilation, antioxidant	Bulb	Quercetin, sulfur compounds	170
52	<i>Elettaria cardamomum</i>	Green Cardamom	Diuretic, vasodilation, antioxidant	Seed	1,8-Cineole, terpinyl acetate, flavonoids	172, 173
53	<i>Beta vulgaris</i>	Beetroot	Dietary nitrate → NO↑ vasodilation	Root/juice	Nitrate, betalains	174, 175
54	<i>Vitis vinifera (seed)</i>	Grape seed	Endothelial function, ACE↓, antioxidant	Seed Extract	Procyanidins (OPCs), catechins	176, 177
55	<i>Linum usitatissimum</i>	Flaxseed	NO↑, anti-inflammatory, RAAS modulation	Seed	α-Linolenic acid, lignans (SDG)	178, 179
56	<i>Theobroma cacao</i>	Cocoa	NO↑ (endothelial), ACE↓	Bean	Flavanols (epicatechin, catechin)	180, 181
57	<i>Punica granatum</i>	Pomegranate	ACE↓, antioxidant, endothelial protection	Juice/Peel	Punicalagins, ellagic acid	182, 183

58	<i>Vaccinium myrtillus</i> / <i>spp.</i>	Bilberry /Blueberry	NO↑, antioxidant	Berry	Anthocyanins (delphinidin/cyanidin glycosides)	170
59	<i>Glycine max</i>	Soybean	Endothelial/estrogenic, RAAS modulation	Seed	Isoflavones (genistein, daidzein)	170
60	<i>Sesamum indicum</i>	Sesame	Antioxidant, NO/endothelial effects	Seed/Oil	Sesamin, sesamolin, lignans	170
61	<i>Coriandrum sativum</i>	Coriander	Ca ²⁺ blockade, diuretic	Seed/leaf	Linalool, flavonoids	170
62	<i>Cuminum cyminum</i>	Cumin	Vasodilation, antioxidant	Seed	Cuminaldehyde, terpenes	170
63	<i>Trigonella foenum-graecum</i>	Fenugreek	Insulin sensitization, diuretic, NO↑	Seed	Saponins (diosgenin), 4-hydroxyisoleucine	170
64	<i>Passiflora incarnata</i>	Passionflower	Anxiolytic (↓SNS), vasodilation	Aerial Parts	Flavonoids (vitexin), harmala alkaloids (trace)	170
65	<i>Achillea millefolium</i>	Yarrow	Vasodilation, diuretic	Aerial Parts	Flavonoids, azulenes	170
66	<i>Tilia cordata/platyphyllos</i>	Linden	Vasodilation, mild diuretic, anxiolytic	Flower/bract	Flavonoids (tiliroside), volatile oils	170
67	<i>Leonurus cardiaca</i>	Motherwort	Negative chronotropy, vasodilation	Aerial parts	Leonurine, stachydrine	170
68	<i>Urtica dioica</i>	Stinging nettle	Diuretic, RAAS modulation	Leaf/root	Flavonoids, lignans, sterols	170
69	<i>Cynara scolymus</i>	Artichoke leaf	Endothelial protection, diuretic	Leaf	Cynarin, chlorogenic acid	170

70	<i>Petroselinum crispum</i>	Parsley	Diuretic, vasodilation	Leaf/seed	Apiol, myristicin, flavonoids	170
71	<i>Berberis vulgaris</i>	Barberry	ACE↓, vasodilation	Root/Bark	Berberine, berbamine	170
72	<i>Berberis aristata</i>	Tree Turmeric	ACE↓, vasodilation	Root/Bark	Berberine, palmatine	170
73	<i>Salvia officinalis</i>	Sage	ACE↓, antioxidant	Leaf	Rosmarinic acid, carnosol	170
74	<i>Melissa officinalis</i>	Lemon Balm	Anxiolytic, vasodilation	Leaf	Rosmarinic acid, flavonoids	170
75	<i>Morus alba</i>	White Mulberry	ACE↓, vasodilation	Leaf	DNJ, quercetin, chlorogenic acid	170
76	<i>Nelumbo nucifera</i>	Lotus leaf	Diuretic, lipids↓, vasodilation	Leaf/embryo	Nuciferine, flavonoids	170
77	<i>Stephania tetrandra</i>	Fangji	Ca ²⁺ channel blockade	Root	Tetrandrine, fangchinoline	170
78	<i>Panax notoginseng</i>	Notoginseng	Endothelial NO↑, anti-inflammatory	Root	Notoginsenosides R1, Rg1, Rb1	184
79	<i>Vernonia amygdalina</i>	Bitter Leaf	Vasodilation, antioxidant	leaf	Vernodalin, flavonoids	170
80	<i>Uncaria tomentosa</i>	Cat's claw (Peru)	Vasodilation, anti-inflammatory	Bark	Oxindole alkaloids	170
81	<i>Hibiscus rosa-sinensis</i>	Chinese hibiscus	Vasodilation, diuretic	Flower/leaf	Anthocyanins, flavonoids	170

82	<i>Houttuynia cordata</i>	Fish Mint	Diuretic, endothelial effects	Aerial parts	Houttuynoside, quercitrin	170
83	<i>Bacopa monnieri</i>	Brahmi	Anxiolytic (↓SNS), NO↑	Aerial parts	Bacosides	170
84	<i>Tinospora cordifolia</i>	Guduchi	Endothelial/anti-inflammatory	Stem	Tinosporaside, berberine-like alkaloids	170
85	<i>Persea americana</i>	Avocado	Vasodilation, antioxidant	Leaf/seed	Flavonoids, phenolics	170
86	<i>Satureja khuzestanica/hortensis</i>	Savory	ACE↓, antioxidant	Aerial parts	Carvacrol, thymol, rosmarinic acid	170
87	<i>Carica papaya</i>	Papaya	Ca ²⁺ blockade (preclinical), antioxidant	Leaf/Seed	Benzyl isothiocyanate, flavonoids	170
88	<i>Eleutherococcus senticosus</i> (syn. <i>Acanthopanax senticosus</i>)	Siberian ginseng	Endothelial/NO modulation, adaptogenic (↓SNS)	Root	Eleutherosides	170
89	<i>Nardostachys jatamansi</i>	Spikenard	Sedative (↓SNS), vasodilation	Rhizome	Jatamansone (valeranone)	170
90	<i>Cydonia oblonga</i>	Quince	Vasodilation, antioxidant	Leaf/95fruit	Quercetin, chlorogenic acid	170
91	<i>Angelica sinensis</i>	Dong-quai	Vasodilation, anti-platelet	Root	Ferulic acid, ligustilide	184
92	<i>Plantago major/asiatica</i>	Plantain	Diuretic, anti-inflammatory	Leaf/Seed	Aucubin, acteoside	170

93	<i>Alisma orientale</i>	Ze-xie	Diuretic, RAAS modulation	Rhizome	Alisol A/B (triterpenes)	184
94	<i>Aspalathus linearis</i>	Rooibos	ACE↓, antioxidant	Leaf	Aspalathin, nothofagin	170
95	<i>Scutellaria baicalensis</i>	Chinese skullcap	ACE↓, vasodilation	Root	Baicalin, baicalein	184
96	<i>Portulaca oleracea</i>	Purslane	NO↑, K ⁺ channel activation, diuretic	Aerial parts	Omega-3s, flavonoids	170
97	<i>Ammi visnaga</i>	Khella	Ca ²⁺ channel blockade, vasodilation	Fruit	Khellin, visnagin	170
98	<i>Gynostemma pentaphyllum</i>	Jiaogulan	AMPK/NO/endothelial effects	Leaf	Gypenosides (saponins)	185
99	<i>Nigella arvensis</i>	Wild black cumin	Antioxidant, vasodilation, Ca ²⁺ channel modulation	Seed	Thymoquinone derivatives, alkaloids	186
100	<i>Erythrina variegata</i>	Indian coral tree	Ca ²⁺ channel blockade, CNS depressant (↓SNS activity)	Baak/Leaf	Isoflavonoids (erycristagallin), alkaloids	187

Table No. 2 Other Effective medicinal plants on Hypertension:

Abbreviations:

ACE – Angiotensin-Converting Enzyme

Ang II – Angiotensin II

AT1R – Angiotensin II Type 1 Receptor

BBB – Blood–Brain Barrier

CKD – Chronic Kidney Disease

CNS – Central Nervous System

CoQ10 – Coenzyme Q10

CSE – Cystathionine γ -Lyase

ED – Erectile Dysfunction

EGCG – Epigallocatechin-3-Gallate

GDLM – Ginkgo Diterpene Lactone Meglumine

GFR – Glomerular Filtration Rate

HPG Axis – Hypothalamic–Pituitary–Gonadal Axis

H₂S – Hydrogen Sulphide

IL – Interleukin

LDL – Low-Density Lipoprotein

LVH – Left Ventricular Hypertrophy

MAPK – Mitogen-Activated Protein Kinase

NO – Nitric Oxide

NOS – Nitric Oxide Synthase

PAF – Platelet-Activating Factor

PDE5 – Phosphodiesterase Type 5

PDGF – Platelet-Derived Growth Factor

PKG – Protein Kinase G

RAAS – Renin–Angiotensin–Aldosterone System

ROS – Reactive Oxygen Species

SNS – Sympathetic Nervous System

TGF-β1 – Transforming Growth Factor Beta 1

TNF-α – Tumour Necrosis Factor Alpha

VSMC – Vascular Smooth Muscle Cell

CONCLUSION

One of the most difficult worldwide health problems is still hypertension, which has a major role in neurological, renal, and cardiovascular problems. While traditional antihypertensive medications continue to be the mainstay of treatment, their drawbacks in terms of adverse effects, cost, and patient compliance underscore the need for safer and more long-lasting substitutes. The renin-angiotensin-aldosterone system modulation, calcium channel blockade, nitric oxide enhancement, and antioxidant defence are some of the mechanisms by which herbal medicines, enhanced with a variety of phytoconstituents like flavonoids, alkaloids, saponins, and polyphenols, offer promising antihypertensive effects.

Terminalia arjuna, *Crataegus monogyna*, *Ginkgo biloba*, *Withania somnifera*, *Allium sativum*, and *Camellia sinensis* are important medicinal plants that

show promise in managing hypertension and reducing its secondary complications, like cardiac hypertrophy, renal impairment, cerebrovascular injury, vascular dysfunction, and erectile disorders.

To fully realize the benefits of these herbal compounds, future research should focus on thorough clinical validation, standardized formulations, and long-term safety monitoring. By acting through a variety of pathways, their diverse phytoconstituents provide a comprehensive therapeutic strategy. By combining these ancient therapies with modern biomedical research, they can be used as supportive or complementary interventions.

REFERENCES

1. Dai N, Zhao FF, Fang M, Pu FL, Kong LY, Liu JP. *Gynostemma pentaphyllum* for dyslipidemia: A systematic review of randomized controlled trials. *Front. Pharmacol.* 2022; 13. doi:10.3389/fphar.2022.917521.
2. Xiong XJ, Yang X chen, Liu W, Duan L, Wang PQ, You H *et al.* Therapeutic efficacy and safety of traditional Chinese medicine classic herbal formula Longdanxiegan decoction for hypertension: A systematic review and meta-analysis. *Front. Pharmacol.* 2018; 9. doi:10.3389/fphar.2018.00466.
3. Bahari H, Omidian K, Goudarzi K, Rafiei H, Asbaghi O, Hosseini Kolbadi KS *et al.* The effects of pomegranate consumption on blood pressure in adults: A systematic review and meta-analysis. *Phytotherapy Research.* 2024; 38. doi:10.1002/ptr.8170.
4. McGuire J, Utset-Ward TJ, Reed DR, Lynch CC. Re-calculating! Navigating through the osteosarcoma treatment roadblock. *Pharmacol. Res.* 2017; 117. doi:10.1016/j.phrs.2016.12.004.
5. Rendina D, D'Elia L, Abate V, Rebellato A, Buondonno I, Succoio M *et al.* Vitamin D Status, Cardiovascular Risk Profile, and miRNA-21 Levels in Hypertensive Patients: Results of the HYPODD Study. *Nutrients* 2022; 14. doi:10.3390/nu14132683.
6. Ried K, Fakler P, Stocks NP. Effect of cocoa on blood pressure. *Cochrane Database of Systematic Reviews.* 2017; 2017. doi:10.1002/14651858.CD008893.pub3.

7. Yuzefpolskiy Y, Baumann FM, Penny LA, Studzinski GP, Kalia V, Sarkar S. Vitamin D receptor signals regulate effector and memory CD8 T cell responses to infections in mice. *Journal of Nutrition* 2014; 144. doi:10.3945/jn.114.202895.
8. Alemu W, Argaw D, Adimasu M, Ewunie TM. Predictors of mortality among severe acute malnourished children. A multi-center prospective follow-up study. *Clin Nutr ESPEN* 2023; 53. doi:10.1016/j.clnesp.2022.12.014.
9. Lipert A, Szadkowska I, Matusiak-Wieczorek E, Kochan E. The Effect of Herbal Supplements on Blood Pressure: Systematic Review and Meta-Analysis. *Antioxidants*. 2022; 11. doi:10.3390/antiox11081419.
10. Javed S, Mangla B, Ahsan W. From propolis to nanopropolis: An exemplary journey and a paradigm shift of a resinous substance produced by bees. *Phytotherapy Research*. 2022; 36. doi:10.1002/ptr.7435.
11. Benjamim CJR, Porto AA, Valenti VE, Sobrinho AC da S, Garner DM, Gualano B *et al*. Nitrate Derived From Beetroot Juice Lowers Blood Pressure in Patients With Arterial Hypertension: A Systematic Review and Meta-Analysis. *Front. Nutr.* 2022; 9. doi:10.3389/fnut.2022.823039.
12. Tovoli F, Faggiano C, Raiteri A, Giamperoli A, Catenaro T, Sansone V *et al*. Risk of Drop-Out from Follow-Up Evaluations for Celiac Disease: Is It Similar for All Patients? *Nutrients* 2022; 14. doi:10.3390/nu14061223.
13. Carpio-Arias TV, Solís Manzano AM, Sandoval V, Vinueza-Veloz AF, Rodríguez Betancourt A, Betancourt Ortiz SL *et al*. Relationship between perceived stress and emotional eating. A cross sectional study. *Clin Nutr ESPEN* 2022; 49. doi:10.1016/j.clnesp.2022.03.030.
14. Kazemi S, Yaghooblou F, Siassi F, Rahimi Froushani A, Ghavipour M, Koohdani F *et al*. Cardamom supplementation improves inflammatory and oxidative stress biomarkers in hyperlipidemic, overweight, and obese pre-diabetic women: a randomized double-blind clinical trial. *J Sci Food Agric* 2017; 97. doi:10.1002/jsfa.8414.
15. Xu X, Tian W, Duan W, Pan C, Huang M, Wang Q *et al*. Quanduzhong capsules for the treatment of grade 1 hypertension patients with low-to-moderate risk: A multicenter, randomized, double-blind, placebo-controlled clinical trial. *Front Pharmacol* 2023; 13. doi:10.3389/fphar.2022.1014410.
16. Anywar G, Kakudidi E, Byamukama R, Mukonzo J, Schubert A, Oryem-Origa H. Indigenous traditional knowledge of medicinal plants used by herbalists in treating opportunistic infections among people living with HIV/AIDS in Uganda. *J Ethnopharmacol* 2020; 246. doi:10.1016/j.jep.2019.112205.
17. Susalit E, Agus N, Effendi I, Tjandrawinata RR, Nofiarny D, Perrinjaquet-Moccetti T *et al*. Olive (*Olea europaea*) leaf extract effective in patients with stage-1 hypertension: Comparison with Captopril. *Phytomedicine* 2011; 18. doi:10.1016/j.phymed.2010.08.016.
18. Liska DAJ, Dioum EH, Chu Y, Mah E. Narrative Review on the Effects of Oat and Sprouted Oat Components on Blood Pressure. *Nutrients*. 2022; 14. doi:10.3390/nu14224772.
19. Anwar MA, Al Disi SS, Eid AH. Anti-hypertensive herbs and their mechanisms of action: Part II. *Front Pharmacol* 2016; 6. doi:10.3389/fphar.2016.00050.
20. Baldé ES, Megalizzi V, Traoré MS, Cos P, Maes L, Decaestecker C *et al*. In vitro antiprotozoal, antimicrobial and antitumor activity of Pavetta crassipes K. Schum leaf extracts. *J Ethnopharmacol* 2010; 130. doi:10.1016/j.jep.2010.05.042.
21. Deiana M, Incani A, Rosa A, Corona G, Atzeri A, Loru D *et al*. Protective effect of hydroxytyrosol and its metabolite homovanillic alcohol on H2O2 induced lipid peroxidation in renal tubular epithelial cells. *Food and Chemical Toxicology* 2008; 46. doi:10.1016/j.fct.2008.05.037.
22. Seo MS, Hong SW, Yeon SH, Kim YM, Um KA, Kim JH *et al*. Magnolia officinalis attenuates free fatty acid-induced lipogenesis via AMPK phosphorylation in hepatocytes. *J Ethnopharmacol* 2014; 157. doi:10.1016/j.jep.2014.09.031.
23. McKay DL, Blumberg JB. A review of the bioactivity and potential health benefits of peppermint tea (*Mentha piperita* L.). *Phytotherapy Research*. 2006; 20. doi:10.1002/ptr.1936.

24. Malik K, Ahmad M, Bussmann RW, Tariq A, Ullah R, Alqahtani AS *et al.* Ethnobotany of anti-hypertensive plants used in Northern Pakistan. *Front Pharmacol* 2018; 9. doi:10.3389/fphar.2018.00789.
25. Aloufi BH, Atwan MA, Alshammari AM. Treatment of Hypertension by Using Natural Herbs and their Mechanism of Action. *J Biochem Technol* 2022; 13. doi:10.51847/wx7mN3flrC.
26. Prangthip P, Panbangred W, Reamtong O. Potential antihypertensive activity of novel peptides from green basil leaves. *BMC Complement Med Ther* 2023; 23. doi:10.1186/s12906-023-04098-2.
27. Verma T, Sinha M, Bansal N, Yadav SR, Shah K, Chauhan NS. Plants Used as Antihypertensive. *Nat. Prod. Bioprospect.* 2021; 11. doi:10.1007/s13659-020-00281-x.
28. Fatehi-Hassanabad Z, Jafarzadeh M, Tarhini A, Fatehi M. The antihypertensive and vasodilator effects of aqueous extract from *Berberis vulgaris* fruit on hypertensive rats. *Phytotherapy Research* 2005; 19. doi:10.1002/ptr.1661.
29. Son TG, Gong EJ, Bae MJ, Kim SD, Heo K, Moon C *et al.* Protective effect of genistein on radiation-induced intestinal injury in tumor bearing mice. *BMC Complement Altern Med* 2013; 13. doi:10.1186/1472-6882-13-103.
30. Singh D, Siew YY, Chong TI, Yew HC, Ho SSW, Lim CSES *et al.* Identification of Phytoconstituents in *Leea indica* (Burm. F.) Merr. Leaves by high performance liquid chromatography micro time-of-flight mass spectrometry. *Molecules* 2019; 24. doi:10.3390/molecules24040714.
31. Sowemimo AA, Adio O, Fageyinbo S. Erratum to 'Anticonvulsant activity of the methanolic extract of *Justicia extensa* T. Anders' (Journal of Ethnopharmacology (2011) 138, 3, (697-699)). *J. Ethnopharmacol.* 2012; 142. doi:10.1016/j.jep.2012.04.036.
32. Subapriya R, Nagini S. Medicinal properties of neem leaves: A review. *Curr. Med. Chem. Anticancer Agents.* 2005; 5. doi:10.2174/1568011053174828.
33. Hsu CD, Hsieh LH, Chen YL, Lin IC, Chen YR, Chen CC *et al.* Complementary effects of pine bark extract supplementation on inattention, impulsivity, and antioxidative status in children with attention-deficit hyperactivity disorder: A double-blinded randomized placebo-controlled cross-over study. *Phytotherapy Research* 2021; 35. doi:10.1002/ptr.7036.
34. Xinke Z, Yingdong L, Mingxia F, Kai L, Kaibing C, Yuqing L *et al.* Chinese herbal medicine for the treatment of primary hypertension: A methodology overview of systematic reviews. *Syst Rev* 2016; 5. doi:10.1186/s13643-016-0353-y.
35. Ingle N, Ojha N, Kumar A. Clinical study to evaluate the Brinhaniya effect of Vidarikandadi Yog to enhance the sport performance in children. *J Ayurveda Integr Med* 2013; 4. doi:10.4103/0975-9476.118709.
36. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K *et al.* Global disparities of hypertension prevalence and control. *Circulation* 2016; 134. doi:10.1161/CIRCULATIONAHA.115.018912.
37. Zareie A, Sahebkar A, Khorvash F, Bagherniya M, Hasanzadeh A, Askari G. Effect of cinnamon on migraine attacks and inflammatory markers: A randomized double-blind placebo-controlled trial. *Phytotherapy Research* 2020; 34. doi:10.1002/ptr.6721.
38. Park H, Hwang YH, Choi JG, Ma JY. In vitro and in vivo evaluation of systemic and genetic toxicity of Citrus unshiu peel. *J Ethnopharmacol* 2018; 215. doi:10.1016/j.jep.2017.12.029.
39. Vlachojannis C, Al-Ahmad A, Hellwig E, Chrubasik S. Listerine® Products: An Update on the Efficacy and Safety. *Phytotherapy Research* 2016; 30. doi:10.1002/ptr.5555.
40. Eun MK, Hye JK, Kim S, Woo HC, Yeon HC, Shi YR *et al.* Modulation of macrophage functions by compounds isolated from *Zingiber officinale*. *Planta Med* 2009; 75. doi:10.1055/s-0028-1088347.
41. Saleem R, Rani R, Ahmed M, Sadaf F, Ahmad SI, ul Zafar N *et al.* Effect of cream containing *Melia azedarach* flowers on skin diseases in children. *Phytomedicine* 2008; 15. doi:10.1016/j.phymed.2008.02.002.
42. Yates T, Davies MJ, Gorely T, Talbot D, Bull F, Sattar N *et al.* The effect of increased ambulatory activity on markers of chronic low-grade inflammation: Evidence from the PREPARE programme randomized controlled trial. *Diabetic*

- Medicine* 2010; 27. doi:10.1111/j.1464-5491.2010.03091.x.
43. Isbrucker RA, Burdock GA. Risk and safety assessment on the consumption of Licorice root (*Glycyrrhiza* sp.), its extract and powder as a food ingredient, with emphasis on the pharmacology and toxicology of glycyrrhizin. *Regulatory Toxicology and Pharmacology* 2006; 46. doi:10.1016/j.yrtph.2006.06.002.
 44. Borhanuddin M. Study of antihypertensive effects of ocimum sanctum. *Bangladesh Journal of Medical Science* 2016; 15. doi:10.3329/bjms.v15i3.30189.
 45. Kuriyama S, Shimazu T, Ohmori K, Kikuchi N, Nakaya N, Nishino Y *et al.* Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: The Ohsaki study. *JAMA* 2006; 296. doi:10.1001/jama.296.10.1255.
 46. Dwivedi S. Terminalia arjuna Wight & Arn.-A useful drug for cardiovascular disorders. *J. Ethnopharmacol.* 2007; 114. doi:10.1016/j.jep.2007.08.003.
 47. Stohs SJ, Miller H, Kaats GR. A review of the efficacy and safety of banaba (*Lagerstroemia speciosa* L.) and corosolic acid. *Phytotherapy Research* 2012; 26. doi:10.1002/ptr.3664.
 48. Salehi B, Quispe C, Sharifi-Rad J, Giri L, Suyal R, Jugran AK *et al.* Antioxidant potential of family Cucurbitaceae with special emphasis on Cucurbita genus: A key to alleviate oxidative stress-mediated disorders. *Phytotherapy Research*. 2021; 35. doi:10.1002/ptr.7045.
 49. Hu N, Wang C, Wang B, Wang L, Huang J, Wang J *et al.* Qianghuo Shengshi decoction exerts anti-inflammatory and analgesic via MAPKs/CREB signaling pathway. *J Ethnopharmacol* 2022; 284. doi:10.1016/j.jep.2021.114776.
 50. Kolla MC, Laya A, Bayang JP, Koubala BB. Effect of different drying methods and storage conditions on physical, nutritional, bioactive compounds and antioxidant properties of doum (*Hyphaene thebaica*) fruits. *Heliyon* 2021; 7. doi:10.1016/j.heliyon.2021.e06678.
 51. Zhao T, Hu S, Ma P, Che D, Liu R, Zhang Y *et al.* Neohesperidin suppresses IgE-mediated anaphylactic reactions and mast cell activation via Lyn-PLC-Ca²⁺ pathway. *Phytotherapy Research* 2019; 33. doi:10.1002/ptr.6385.
 52. Zhou L, Zuo Z, Chow MSS. Danshen: An overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J. Clin. Pharmacol.* 2005; 45. doi:10.1177/0091270005282630.
 53. Xie T, Li K, Gong X, Jiang R, Huang W, Chen X *et al.* Paeoniflorin protects against liver ischemia/reperfusion injury in mice via inhibiting HMGB1-TLR4 signaling pathway. *Phytotherapy Research* 2018; 32. doi:10.1002/ptr.6161.
 54. Salehi B, Konovalov DA, Fru P, Kapewangolo P, Peron G, Ksenija MS *et al.* Areca catechu—From farm to food and biomedical applications. *Phytotherapy Research*. 2020; 34. doi:10.1002/ptr.6665.
 55. Epstein M, Calhoun DA. Aldosterone blockers (mineralocorticoid receptor antagonism) and potassium-sparing diuretics. *J. Clin. Hypertens*. 2011; 13. doi:10.1111/j.1751-7176.2011.00511.x.
 56. Yu R, Zhou Y, Shi S, Wang X, Huang S, Ren Y. Icariside II induces ferroptosis in renal cell carcinoma cells by regulating the miR-324-3p/GPX4 axis. *Phytomedicine* 2022; 102. doi:10.1016/j.phymed.2022.154182.
 57. Oi-Kano Y, Iwasaki Y, Nakamura T, Watanabe T, Goto T, Kawada T *et al.* Oleuropein aglycone enhances UCP1 expression in brown adipose tissue in high-fat-diet-induced obese rats by activating β -adrenergic signaling. *Journal of Nutritional Biochemistry* 2017; 40. doi:10.1016/j.jnutbio.2016.11.009.
 58. Maurya R, Boini T, Misro L, Radhakrishnan T, Singh R. Review on Pharmacognostic, Phytochemical and Pharmacological Characteristics of Medicinal Plants which can be used to Design a Novel Ayurvedic Formulation to Treat Hypertension in Multidimensional Approaches. *Curr Hypertens Rev* 2023; 19. doi:10.2174/1573402119666230221084859.
 59. Kumar S, Kumari D, Singh B. Genus Rauvolfia: A review of its ethnopharmacology, phytochemistry, quality control/quality assurance, pharmacological activities and clinical evidence. *J. Ethnopharmacol.* 2022; 295. doi:10.1016/j.jep.2022.115327.
 60. Kavyani Z, Musazadeh V, Safaei E, Mohammadi Asmaroud M, Khashakichafi F, Ahrabi SS *et al.* Antihypertensive effects of *Nigella sativa*

- supplementation: An updated systematic review and meta-analysis of randomized controlled trials. *Phytotherapy Research*. 2023; 37. doi:10.1002/ptr.7891.
61. Liu B, Wang D, Luo E, Hou J, Qiao Y, Yan G *et al*. Role of TG2-Mediated SERCA2 Serotonylation on Hypoxic Pulmonary Vein Remodeling. *Front Pharmacol* 2020; 10. doi:10.3389/fphar.2019.01611.
 62. Heydarian A, Kashani AHF, Masoodi M, Aryaeian N, Vafa M, Tahvilian N *et al*. Effects of saffron supplementation on serum inflammatory markers and quality of life in patients with ulcerative colitis: A double blind randomized controlled clinical trial. *J Herb Med* 2022; 36. doi:10.1016/j.hermed.2022.100593.
 63. Yan S, Liu S, Qu J, Li X, Hu J, Zhang L *et al*. A Lard and Soybean Oil Mixture Alleviates Low-Fat–High-Carbohydrate Diet-Induced Nonalcoholic Fatty Liver Disease in Mice. *Nutrients* 2022; 14. doi:10.3390/nu14030560.
 64. Alarcón-Alonso J, Zamilpa A, Aguilar FA, Herrera-Ruiz M, Tortoriello J, Jimenez-Ferrer E. Pharmacological characterization of the diuretic effect of Hibiscus sabdariffa Linn (Malvaceae) extract. *J Ethnopharmacol* 2012; 139. doi:10.1016/j.jep.2011.12.005.
 65. Metwally M, Ong KJ, Ledger WL, Li TC. Does high body mass index increase the risk of miscarriage after spontaneous and assisted conception? A meta-analysis of the evidence. *Fertil Steril* 2008; 90. doi:10.1016/j.fertnstert.2007.07.1290.
 66. Alimoradi N, Firouzabadi N, Fatehi R. Metformin and insulin-resistant related diseases: Emphasis on the role of microRNAs. *Biomedicine and Pharmacotherapy*. 2021; 139. doi:10.1016/j.biopha.2021.111662.
 67. Jang DJ, Lee MS, Shin BC, Lee YC, Ernst E. Red ginseng for treating erectile dysfunction: A systematic review. *Br. J. Clin. Pharmacol*. 2008; 66. doi:10.1111/j.1365-2125.2008.03236.x.
 68. Dalaklioglu S, Ozbey G. The potent relaxant effect of resveratrol in rat corpus cavernosum and its underlying mechanisms. *Int J Impot Res* 2013; 25. doi:10.1038/ijir.2013.6.
 69. Zamani M, Kelishadi MR, Ashtary-Larky D, Amirani N, Goudarzi K, Torki IA *et al*. The effects of green tea supplementation on cardiovascular risk factors: A systematic review and meta-analysis. *Front. Nutr.* 2023; 9. doi:10.3389/fnut.2022.1084455.
 70. Zenk JL, Frestedt JL, Kuskowski MA. HUM5007, a novel combination of thermogenic compounds, and 3-acetyl-7-oxo-dehydroepiandrosterone: each increases the resting metabolic rate of overweight adults. *Journal of Nutritional Biochemistry* 2007; 18. doi:10.1016/j.jnutbio.2006.11.008.
 71. Romijn AR, Rucklidge JJ. Systematic review of evidence to support the theory of psychobiotics. *Nutr Rev* 2015; 73. doi:10.1093/nutrit/nuv025.
 72. Saeed M, Naveed M, Arif M, Kakar MU, Manzoor R, Abd El-Hack ME *et al*. Green tea (*Camellia sinensis*) and L-theanine: Medicinal values and beneficial applications in humans—A comprehensive review. *Biomedicine and Pharmacotherapy*. 2017; 95. doi:10.1016/j.biopha.2017.09.024.
 73. Ogunmola OJ, Olaifa AO, Oladapo OO, Babatunde OA. Prevalence of cardiovascular risk factors among adults without obvious cardiovascular disease in a rural community in Ekiti State, Southwest Nigeria. *BMC Cardiovasc Disord* 2013; 13. doi:10.1186/1471-2261-13-89.
 74. Huang R, Chen H. Comparison of Water-Assisted Decontamination Systems of Pulsed Light and Ultraviolet for Salmonella Inactivation on Blueberry, Tomato, and Lettuce. *J Food Sci* 2019; 84. doi:10.1111/1750-3841.14510.
 75. Iciek M, Kwiecień I, Włodek L. Biological properties of garlic and garlic-derived organosulfur compounds. *Environ. Mol. Mutagen*. 2009; 50. doi:10.1002/em.20474.
 76. Gonciarz W, Krupa A, Chmiela M. Proregenerative activity of IL-33 in gastric tissue cells undergoing Helicobacter pylori-induced apoptosis. *Int J Mol Sci* 2020; 21. doi:10.3390/ijms21051801.
 77. Lozada-Lechuga J, Villarreal ML, Fliniaux MA, Bensaddek L, Mesnard F, Gutiérrez M del C *et al*. Isolation of jacaranone, a sedative constituent extracted from the flowers of the Mexican tree Ternstroemia pringlei. *J Ethnopharmacol* 2010; 127. doi:10.1016/j.jep.2009.11.020.
 78. Peluffo GD, Diamant MJ, Klein S. The dual role of inflammation in cancer. *Medicina (B Aires)* 2007; 67. doi:10.1016/j.phymed.2007.07.009.

79. Neychev VK, Mitev VI. The aphrodisiac herb *Tribulus terrestris* does not influence the androgen production in young men. *J Ethnopharmacol* 2005; 101. doi:10.1016/j.jep.2005.05.017.
80. Do J, Choi S, Choi J, Hyun JS. Effects and mechanism of action of a *Tribulus terrestris* extract on penile erection. *Korean J Urol* 2013; 54. doi:10.4111/kju.2013.54.3.183.
81. Megala J, Geetha A. Antiulcerogenic activity of hydroalcoholic fruit extract of *Pithecellobium dulce* in different experimental ulcer models in rats. *J Ethnopharmacol* 2012; 142. doi:10.1016/j.jep.2012.05.011.
82. Tangjang S, Namsa ND, Aran C, Litin A. An ethnobotanical survey of medicinal plants in the Eastern Himalayan zone of Arunachal Pradesh, India. *J Ethnopharmacol* 2011; 134. doi:10.1016/j.jep.2010.11.053.
83. Akhter F, Sultan Alvi S, Ahmad P, Iqbal D, Alshehri BM, Khan MS. Therapeutic efficacy of *Boerhaavia diffusa* (Linn.) root methanolic extract in attenuating streptozotocin-induced diabetes, diabetes-linked hyperlipidemia and oxidative-stress in rats. *Biomedical Research and Therapy* 2019; 6. doi:10.15419/bmrat.v6i7.556.
84. Gupta RK, Kesari AN, Murthy PS, Chandra R, Tandon V, Watal G. Hypoglycemic and antidiabetic effect of ethanolic extract of leaves of *Annona squamosa* L. in experimental animals. *J Ethnopharmacol* 2005; 99. doi:10.1016/j.jep.2005.01.048.
85. Janecki A, Conrad A, Engels I, Frank U, Kolodziej H. Evaluation of an aqueous-ethanolic extract from *Pelargonium sidoides* (EPs® 7630) for its activity against group A-streptococci adhesion to human HEp-2 epithelial cells. *J Ethnopharmacol* 2011; 133. doi:10.1016/j.jep.2010.09.018.
86. Kuboyama T, Tohda C, Komatsu K. Neuritic regeneration and synaptic reconstruction induced by withanolide A. *Br J Pharmacol* 2005; 144. doi:10.1038/sj.bjp.0706122.
87. Oh BS, Shin EA, Jung JH, Jung DB, Kim B, Shim BS *et al.* Apoptotic effect of galbanic acid via activation of caspases and inhibition of Mcl-1 in H460 non-small lung carcinoma cells. *Phytotherapy Research* 2015; 29. doi:10.1002/ptr.5320.
88. Mansukhani MP, Bellolio MF, Kolla BP, Enduri S, Somers VK, Stead LG. Worse outcome after stroke in patients with obstructive sleep apnea: An observational cohort study. *Journal of Stroke and Cerebrovascular Diseases* 2011; 20. doi:10.1016/j.jstrokecerebrovasdis.2010.02.011.
89. Weisleder N, Ma JJ. Ca²⁺ sparks as a plastic signal for skeletal muscle health, aging, and dystrophy. *Acta Pharmacol. Sin.* 2006; 27. doi:10.1111/j.1745-7254.2006.00384.x.
90. Kwon KJ, Lee EJ, Cho KS, Cho DH, Shin CY, Han SH. Ginkgo biloba extract (Egb761) attenuates zinc-induced tau phosphorylation at Ser262 by regulating GSK3 β activity in rat primary cortical neurons. *Food Funct* 2015; 6. doi:10.1039/c5fo00219b.
91. Mitsui T, Taniguchi N, Kawasaki N, Kagami Y, Arita J. Fetal exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin induces expression of the chemokine genes Cxcl4 and Cxcl7 in the perinatal mouse brain. *Journal of Applied Toxicology* 2011; 31. doi:10.1002/jat.1612.
92. Tadić VM, Dobrić S, Marković GM, Dordević SM, Arsić IA, Menković NR *et al.* Anti-inflammatory, gastroprotective, free-radical-scavenging, and antimicrobial activities of hawthorn berries ethanol extract. *J Agric Food Chem* 2008; 56. doi:10.1021/jf801668c.
93. Astudillo L, Payoux P, Game X, Sailler L, Arné JL, Arlet-Suau E. Bilateral testicular and epididymal involvement in sarcoidosis [5]. *American Journal of Medicine.* 2004; 116. doi:10.1016/j.amjmed.2003.12.034.
94. Koch E, Malek FA. Standardized extracts from hawthorn leaves and flowers in the treatment of cardiovascular disorders preclinical and clinical studies. *Planta Med.* 2011; 77. doi:10.1055/s-0030-1270849.
95. Holubarsch CJF, Colucci WS, Meinertz T, Gaus W, Tendra M. The efficacy and safety of Crataegus extract WS® 1442 in patients with heart failure: The SPICE trial. *Eur J Heart Fail* 2008; 10. doi:10.1016/j.ejheart.2008.10.004.
96. Parveen A, Babbar R, Agarwal S, Kotwani A, Fahim M. Mechanistic clues in the cardioprotective effect of terminalia arjuna bark extract in isoproterenol-induced chronic heart failure in rats. *Cardiovasc Toxicol* 2011; 11. doi:10.1007/s12012-010-9099-2.

97. Kumar G, Saleem N, Kumar S, Maulik SK, Ahmad S, Sharma M *et al.* Transcriptomic validation of the protective effects of aqueous bark extract of *Terminalia arjuna* (Roxb.) on isoproterenol-induced cardiac hypertrophy in rats. *Front Pharmacol* 2019; 10. doi:10.3389/fphar.2019.01443.
98. Bansal T, Chatterjee E, Singh J, Ray A, Kundu B, Thankamani V *et al.* Arjunolic acid, a peroxisome proliferator-activated receptor agonist, regresses cardiac fibrosis by inhibiting non-canonical TGF-signaling. *Journal of Biological Chemistry* 2017; 292. doi:10.1074/jbc.M117.788299.
99. Suresh S, Prakash S. Effect of *Mucuna pruriens* (Linn.) on oxidative stress-induced structural alteration of Corpus cavernosum in streptozotocin-induced diabetic rat. *Journal of Sexual Medicine* 2011; 8. doi:10.1111/j.1743-6109.2011.02221.x.
100. Su MT, Chen CH, Kuo PH, Hsu CC, Lee IW, Pan HA *et al.* Polymorphisms of estrogen-related genes jointly confer susceptibility to human spermatogenic defect. *Fertil Steril* 2010; 93. doi:10.1016/j.fertnstert.2008.09.030.
101. Zhao F, Wang L, Liu K. In vitro anti-inflammatory effects of arctigenin, a lignan from *Arctium lappa* L., through inhibition on iNOS pathway. *J Ethnopharmacol* 2009; 122. doi:10.1016/j.jep.2009.01.038.
102. Tanrikut C, Feldman AS, Altemus M, Paduch DA, Schlegel PN. Adverse effect of paroxetine on sperm. *Fertil Steril* 2010; 94. doi:10.1016/j.fertnstert.2009.04.039.
103. Shores M. Testosterone treatment and cardiovascular events in prescription database studies. *Asian J Androl* 2018; 20. doi:10.4103/aja.aja_25_17.
104. Burnett AL, Nehra A, Breau RH, Culkin DJ, Faraday MM, Hakim LS *et al.* Erectile Dysfunction: AUA Guideline. *Journal of Urology* 2018; 200. doi:10.1016/j.juro.2018.05.004.
105. Eşkazan AE. Evolving treatment strategies in CML – moving from early and deep molecular responses to TKI discontinuation and treatment-free remission: is there a need for longer-term trial outcomes? *Br. J. Clin. Pharmacol.* 2018; 84. doi:10.1111/bcp.13637.
106. Leung KW, Wong AS. Pharmacology of ginsenosides: a literature review. *Chin. Med.* 2010; 5. doi:10.1186/1749-8546-5-20.
107. Moreno SA, Morgentaler A. Testosterone deficiency and Peyronie's disease: Pilot data suggesting a significant relationship. *Journal of Sexual Medicine* 2009; 6. doi:10.1111/j.1743-6109.2009.01250.x.
108. Vlachopoulos C, Jackson G, Stefanadis C, Montorsi P. Erectile dysfunction in the cardiovascular patient. *Eur. Heart J.* 2013; 34. doi:10.1093/eurheartj/ehs112.
109. Bezan A, Masic E, Krieger D, Stojakovic T, Pummer K, Zigeuner R *et al.* The preoperative AST/ALT (De Ritis) ratio represents a poor prognostic factor in a cohort of patients with nonmetastatic renal cell carcinoma. *Journal of Urology* 2015; 194. doi:10.1016/j.juro.2015.01.083.
110. Singh BN, Shankar S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): Mechanisms, perspectives and clinical applications. *Biochem. Pharmacol.* 2011; 82. doi:10.1016/j.bcp.2011.07.093.
111. Anandh Babu P, Liu D. Green Tea Catechins and Cardiovascular Health: An Update. *Curr Med Chem* 2008; 15. doi:10.2174/092986708785132979.
112. Hodgson JM, Croft KD. Tea flavonoids and cardiovascular health. *Mol. Aspects Med.* 2010; 31. doi:10.1016/j.mam.2010.09.004.
113. Sobenin IA, Pryanishnikov V V., Kunnova LM, Rabinovich YA, Martirosyan DM, Orekhov AN. The effects of time-released garlic powder tablets on multifunctional cardiovascular risk in patients with coronary artery disease. *Lipids Health Dis* 2010; 9. doi:10.1186/1476-511X-9-119.
114. Perini W, Kunst AE, Snijder MB, Peters RJG, van Valkengoed IGM. Ethnic differences in metabolic cardiovascular risk among normal weight individuals: Implications for cardiovascular risk screening. The HELIUS study. *Nutrition, Metabolism and Cardiovascular Diseases* 2019; 29. doi:10.1016/j.numecd.2018.09.004.
115. Rahman K, Lowe GM. Garlic and cardiovascular disease: A critical review. In: *Journal of Nutrition.* 2006 doi:10.1093/jn/136.3.736s.

116. Bawadi HA, Al-Kuran O, Al-Bastoni LAA, Tayyem RF, Jaradat AN, Tuuri G *et al.* Gestational nutrition improves outcomes of vaginal deliveries in Jordan: an epidemiologic screening. 2010; 30. doi:10.1016/j.nutres.2010.01.005.
117. Jacobs ET, Foote JA, Kohler LN, Skiba MB, Thomson CA. Re-examination of dairy as a single commodity in US dietary guidance. *Nutr Rev* 2020; 78. doi:10.1093/nutrit/nuz093.
118. Banerjee SK, Maulik SK. Effect of garlic on cardiovascular disorders: a review. *Nutr J* 2002; 1. doi:10.1186/1475-2891-1-4.
119. Garfinkel-Castro A, Kim K, Hamidi S, Ewing R. Obesity and the built environment at different urban scales: Examining the literature. *Nutr Rev* 2017; 75. doi:10.1093/nutrit/nuw037.
120. Panella S, Marcocci ME, Celestino I, Valente S, Zwergel C, Li Puma DD *et al.* MC1568 inhibits HDAC6/8 activity and influenza A virus replication in lung epithelial cells: Role of Hsp90 acetylation. *Future Med Chem* 2016; 8. doi:10.4155/fmc-2016-0073.
121. Shindou H, Hishikawa D, Harayama T, Yuki K, Shimizu T. Recent progress on acyl CoA: Lysophospholipid acyltransferase research. *J. Lipid Res.* 2009; 50. doi:10.1194/jlr.R800035-JLR200.
122. Libby P, Ridker PM, Hansson GK. Progress and challenges in translating the biology of atherosclerosis. *Nature.* 2011; 473. doi:10.1038/nature10146.
123. Gimbrone MA, García-Cardena G. Endothelial Cell Dysfunction and the Pathobiology of Atherosclerosis. *Circ Res* 2016; 118. doi:10.1161/CIRCRESAHA.115.306301.
124. Chhatre S, Nesari T, Somani G, Kanchan D, Sathaye S. Phytopharmacological overview of Tribulus terrestris. *Pharmacogn. Rev.* 2014; 8. doi:10.4103/0973-7847.125530.
125. Thanasakdech S, Tewtrakul S. Wound healing gel containing compound 2 α -acetoxysandaracopimaradien-1 α -ol from Kaempferia marginata rhizomes. *J Herb Med* 2021; 28. doi:10.1016/j.hermed.2021.100437.
126. Liu C, Zhen D, Du H, Gong G, Wu Y, Ma Q *et al.* Synergistic anti-inflammatory effects of peimine, peiminine, and forsythoside a combination on LPS-induced acute lung injury by inhibition of the IL-17-NF- κ B/MAPK pathway activation. *J Ethnopharmacol* 2022; 295. doi:10.1016/j.jep.2022.115343.
127. Abdelaziz DH, Boraii S, Cheema E, Elnaem MH, Omar T, Abdelraouf A *et al.* The intraperitoneal ondansetron for postoperative pain management following laparoscopic cholecystectomy: A proof-of-concept, double-blind, placebo-controlled trial. *Biomedicine and Pharmacotherapy* 2021; 140. doi:10.1016/j.biopha.2021.111725.
128. Gebissa E. Khat in the Horn of Africa: Historical perspectives and current trends. *J Ethnopharmacol* 2010; 132. doi:10.1016/j.jep.2010.01.063.
129. Samirana PO, Murti YB, Jenie RI, Setyowati EP. Marine Sponge-Derived Fungi: Fermentation and Cytotoxic Activity. *J Appl Pharm Sci* 2021; 11. doi:10.7324/JAPS.2021.110103.
130. MacDonald-Wallis C, Lawlor DA, Fraser A, May M, Nelson SM, Tilling K. Blood pressure change in normotensive, gestational hypertensive, preeclamptic, and essential hypertensive pregnancies. *Hypertension* 2012; 59. doi:10.1161/HYPERTENSIONAHA.111.187039.
131. Catelan TBS, Santos Radaí JA, Leitão MM, Branquinho LS, Vasconcelos PC de P, Heredia-Vieira SC *et al.* Evaluation of the toxicity and anti-inflammatory activities of the infusion of leaves of Campomanesia guazumifolia (Cambess.) O. Berg. *J Ethnopharmacol* 2018; 226. doi:10.1016/j.jep.2018.08.015.
132. Raghavan A, Shah ZA. Withania somnifera Improves Ischemic Stroke Outcomes by Attenuating PARP1-AIF-Mediated Caspase-Independent Apoptosis. *Mol Neurobiol* 2015; 52. doi:10.1007/s12035-014-8907-2.
133. Choudhary D, Bhattacharyya S, Bose S. Efficacy and Safety of Ashwagandha (Withania somnifera (L.) Dunal) Root Extract in Improving Memory and Cognitive Functions. *J Diet Suppl* 2017; 14. doi:10.1080/19390211.2017.1284970.
134. Xie Q, Li H, Lu D, Yuan J, Ma R, Li J *et al.* Neuroprotective Effect for Cerebral Ischemia by Natural Products: A Review. *Front. Pharmacol.* 2021; 12. doi:10.3389/fphar.2021.607412.
135. Núñez-Iglesias MJ, Novio S, García C, Pérez-Muñuzuri E, Soengas P, Cartea E *et al.*

- Glucosinolate-degradation products as co-adjuvant therapy on prostate cancer in vitro. *Int J Mol Sci* 2019; 20. doi:10.3390/ijms20204977.
136. Feigin VL, Stark BA, Johnson CO, Roth GA, Bisignano C, Abady GG *et al.* Global, regional, and national burden of stroke and its risk factors, 1990-2019: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet Neurol* 2021; 20. doi:10.1016/S1474-4422(21)00252-0.
137. Tian X, Xu Q, Xia X, Zhang Y, Meng X, Wang A. Efficacy of ginkgo diterpene lactone meglumine on cognitive function in patients with acute ischemic stroke: a predefined exploratory analysis of a multicenter, double-blind, randomized controlled trial. *J Neurol* 2024; 271. doi:10.1007/s00415-024-12272-w.
138. Linhart C, Tukana I, Lin S, Taylor R, Morrell S, Vatucaawaqa P *et al.* Continued increases in hypertension over three decades in Fiji, and the influence of obesity. *J Hypertens* 2016; 34. doi:10.1097/HJH.0000000000000807.
139. Liu C, Huang Y. Chinese herbal medicine on cardiovascular diseases and the mechanisms of action. *Front. Pharmacol.* 2016; 7. doi:10.3389/fphar.2016.00469.
140. Parveen A, Babbar R, Agarwal S, Kotwani A, Fahim M. Terminalia arjuna Enhances Baroreflex Sensitivity and Myocardial Function in Isoproterenol-Induced Chronic Heart Failure Rats. *J Cardiovasc Pharmacol Ther* 2012; 17. doi:10.1177/1074248411416816.
141. Maulik SK, Talwar KK. Therapeutic potential of Terminalia arjuna in cardiovascular disorders. *American Journal of Cardiovascular Drugs*. 2012; 12. doi:10.2165/11598990-000000000-00000.
142. Ramesh P, Palaniappan A. Terminalia arjuna, a Cardioprotective Herbal Medicine—Relevancy in the Modern Era of Pharmaceuticals and Green Nanomedicine—A Review. *Pharmaceuticals*. 2023; 16. doi:10.3390/ph16010126.
143. Dwivedi S, Chopra D. Revisiting terminalia arjuna-an ancient cardiovascular drug. *J Tradit Complement Med* 2014; 4. doi:10.4103/2225-4110.139103.
144. Srivastava S, Girandola R, Kokate AS. Effect of Oxyjun™ on Adipose Tissue Inflammation - A Randomized, Placebo-Controlled Clinical Study. *International Journal of Physical Education, Fitness and Sports* 2020. doi:10.34256/ijpefs2045.
145. Vondrisk TM, Wang Y. A new (heat) shocking player in cardiac hypertrophy. *Circ. Res.* 2008; 103. doi:10.1161/CIRCRESAHA.108.189118.
146. Achit H, Guillemin F, Karam G, Ladrière M, Baumann C, Frimat L *et al.* Cost-effectiveness of four living-donor nephrectomy techniques from a hospital perspective. *Nephrology Dialysis Transplantation*. 2020; 35. doi:10.1093/ndt/gfz143.
147. Wilson JL, Bouillaud F, Almeida AS, Vieira HL, Ouidja MO, Dubois-Randé JL *et al.* Carbon monoxide reverses the metabolic adaptation of microglia cells to an inflammatory stimulus. *Free Radic Biol Med* 2017; 104. doi:10.1016/j.freeradbiomed.2017.01.022.
148. Sun Y, Huang Y, Zhang R, Chen Q, Chen J, Zong Y *et al.* Hydrogen sulfide upregulates KATP channel expression in vascular smooth muscle cells of spontaneously hypertensive rats. *J Mol Med* 2015; 93. doi:10.1007/s00109-014-1227-1.
149. Bucci M, Papapetropoulos A, Vellecco V, Zhou Z, Zaid A, Giannogonas P *et al.* cGMP-Dependent Protein Kinase Contributes to Hydrogen Sulfide-Stimulated Vasorelaxation. *PLoS One* 2012; 7. doi:10.1371/journal.pone.0053319.
150. Pan LL, Liu XH, Gong QH, Yang HB, Zhu YZ. Role of cystathionine γ -Lyase/hydrogen sulfide pathway in cardiovascular disease: A novel therapeutic strategy? *Antioxid. Redox Signal.* 2012; 17. doi:10.1089/ars.2011.4349.
151. Michel T, Vanhoutte PM. Cellular signaling and NO production. *Pflugers Arch.* 2010; 459. doi:10.1007/s00424-009-0765-9.
152. Agorrody G, Peclat TR, Peluso G, Gonano LA, Santos L, van Schooten W *et al.* Benefits in cardiac function by CD38 suppression: Improvement in NAD⁺ levels, exercise capacity, heart rate variability and protection against catecholamine-induced ventricular arrhythmias. *J Mol Cell Cardiol* 2022; 166. doi:10.1016/j.yjmcc.2022.01.008.
153. Almeida J, Oliveira LA, Benini R, Crestani CC. Role of hippocampal nitrenergic neurotransmission in behavioral and cardiovascular dysfunctions

- evoked by chronic social stress. *Nitric Oxide* 2020; 94. doi:10.1016/j.niox.2019.11.004.
154. Qu J, Tan S, Xie X, Wu W, Zhu H, Li H *et al.* Dendrobium Officinale Polysaccharide Attenuates Insulin Resistance and Abnormal Lipid Metabolism in Obese Mice. *Front Pharmacol* 2021; 12. doi:10.3389/fphar.2021.659626.
155. Kissel CK, Anderson TJ. Understanding Coronary Microvascular Disease—New Insights for a Confusing and Underdiagnosed Entity. *Canadian Journal of Cardiology*. 2020; 36. doi:10.1016/j.cjca.2020.01.018.
156. Song P, Zou MH. Regulation of NAD(P)H oxidases by AMPK in cardiovascular systems. *Free Radic. Biol. Med.* 2012; 52. doi:10.1016/j.freeradbiomed.2012.01.025.
157. Montezano AC, Dulak-Lis M, Tsiropoulou S, Harvey A, Briones AM, Touyz RM. Oxidative stress and human hypertension: Vascular mechanisms, biomarkers, and novel therapies. *Canadian Journal of Cardiology*. 2015; 31. doi:10.1016/j.cjca.2015.02.008.
158. Montezano AC, Touyz RM. Reactive oxygen species, vascular noxs, and hypertension: Focus on translational and clinical research. *Antioxid. Redox Signal.* 2014; 20. doi:10.1089/ars.2013.5302.
159. Ong SLH, Whitworth JA. How do glucocorticoids cause hypertension: Role of nitric oxide deficiency, oxidative stress, and eicosanoids. *Endocrinol. Metab. Clin. North Am.* 2011; 40. doi:10.1016/j.ecl.2011.01.010.
160. Higashi Y. Roles of Oxidative Stress and Inflammation in Vascular Endothelial Dysfunction-Related Disease. *Antioxidants*. 2022; 11. doi:10.3390/antiox11101958.
161. Li S, Zhou C, Zhu Y, Chao Z, Sheng Z, Zhang Y *et al.* Ferrostatin-1 alleviates angiotensin II (Ang II)- induced inflammation and ferroptosis in astrocytes. *Int Immunopharmacol* 2021; 90. doi:10.1016/j.intimp.2020.107179.
162. Nakamaru R, Nakagami H, Hayashi H, Sun J, Tenma A, Yamamoto K *et al.* A novel angiotensin II peptide vaccine without an adjuvant inmice. *J Hypertens* 2021; 39. doi:10.1097/HJH.0000000000002597.
163. Gallo G, Savoia C. New Insights into Endothelial Dysfunction in Cardiometabolic Diseases: Potential Mechanisms and Clinical Implications. *Int. J. Mol. Sci.* 2024; 25. doi:10.3390/ijms25052973.
164. Du L, Zhang Y, Li X, Liu C, Li Z, Zhou J *et al.* Blood pressure excursion on admission and intravenous thrombolysis in ischemic stroke. *J Hypertens* 2023; 41. doi:10.1097/HJH.0000000000003456.
165. Liu S, Jiang X, Lu H, Xing M, Qiao Y, Zhang C *et al.* HuR (Human Antigen R) Regulates the Contraction of Vascular Smooth Muscle and Maintains Blood Pressure. *Arterioscler Thromb Vasc Biol* 2020; 40. doi:10.1161/ATVBAHA.119.313897.
166. Liu C, Li N, Li F, Deng W, Dai G, Tang Y *et al.* CircHIPK2 facilitates phenotypic switching of vascular smooth muscle cells in hypertension. *J Hum Hypertens* 2023; 37. doi:10.1038/s41371-023-00834-w.
167. Liu S, Lin Z. Vascular Smooth Muscle Cells Mechanosensitive Regulators and Vascular Remodeling. *J. Vasc. Res.* 2022; 59. doi:10.1159/000519845.
168. MacK CP. Signaling mechanisms that regulate smooth muscle cell differentiation. *Arterioscler. Thromb. Vasc. Biol.* 2011; 31. doi:10.1161/ATVBAHA.110.221135.
169. Marx SO, Totary-Jain H, Marks AR. Vascular smooth muscle cell proliferation in restenosis. *Circ Cardiovasc Interv* 2011; 4. doi:10.1161/CIRCINTERVENTIONS.110.957332.
170. Lacolley P, Regnault V, Nicoletti A, Li Z, Michel JB. The vascular smooth muscle cell in arterial pathology: A cell that can take on multiple roles. *Cardiovasc. Res.* 2012; 95. doi:10.1093/cvr/cvs135.
171. Folkow B. ‘Structural factor’ in primary and secondary hypertension. *Hypertension*. 1990; 16. doi:10.1161/01.hyp.16.1.89.
172. Rawat P, Singh PK, Kumar V. Anti-hypertensive medicinal plants and their mode of action. *J. Herb. Med.* 2016; 6. doi:10.1016/j.hermed.2016.06.001.
173. Kumari M, Krishnamurthy PT, Pindiprolu SKSS, Sola P. DR-5 and DLL-4 mAb functionalized SLNs of gamma-secretase inhibitors- An approach for TNBC treatment.

- Adv. Pharm. Bull. 2021; 11. doi:10.34172/APB.2021.070.
174. Shayganni E, Bahmani M, Asgary S, Rafieian-Kopaei M. Inflammaging and cardiovascular disease: Management by medicinal plants. *Phytomedicine* 2016; 23. doi:10.1016/j.phymed.2015.11.004.
175. Liwa AC, Smart LR, Frumkin A, Epstein HAB, Fitzgerald DW, Peck RN. Traditional herbal medicine use among hypertensive patients in sub-Saharan Africa: A systematic review. *Curr. Hypertens. Rep.* 2014; 16. doi:10.1007/s11906-014-0437-9.
176. Tabassum N, Ahmad F. Role of natural herbs in the treatment of hypertension. *Pharmacogn. Rev.* 2011; 5. doi:10.4103/0973-7847.79097.
177. Mackenzie IS, MacDonald TM. Identifying poor adherence to antihypertensive medications in patients with resistant hypertension. *Br. J. Clin. Pharmacol.* 2019; 85. doi:10.1111/bcp.13806.
178. Aabidha P, Cherian A, Paul E, Helan J. Maternal and fetal outcome in pre-eclampsia in a secondary care hospital in South India. *J Family Med Prim Care* 2015; 4. doi:10.4103/2249-4863.154669.
179. Schutte AE, Jafar TH, Poulter NR, Damasceno A, Khan NA, Nilsson PM *et al.* Addressing global disparities in blood pressure control: perspectives of the International Society of Hypertension. *Cardiovasc. Res.* 2023; 119. doi:10.1093/cvr/cvac130.
180. Zhou B, Carrillo-Larco RM, Danaei G, Riley LM, Paciorek CJ, Stevens GA *et al.* Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *The Lancet* 2021; 398. doi:10.1016/S0140-6736(21)01330-1.
181. Al Disi SS, Anwar MA, Eid AH. Anti-hypertensive herbs and their mechanisms of action: Part I. *Front. Pharmacol.* 2016; 6. doi:10.3389/fphar.2015.00323.
182. Kamyab R, Namdar H, Torbati M, Ghojzadeh M, Araj-Khodaei M, Fazljou SMB. Medicinal plants in the treatment of hypertension: A review. *Adv. Pharm. Bull.* 2021; 11. doi:10.34172/APB.2021.090.
183. Bhadane BS, Patil MP, Maheshwari VL, Patil RH. Ethnopharmacology, phytochemistry, and biotechnological advances of family Apocynaceae: A review. *Phytotherapy Research.* 2018; 32. doi:10.1002/ptr.6066.
184. Cuspidi C, Tadic M, Sala C, Gherbesi E, Grassi G, Mancia G. Extreme dipping: Is the cardiovascular risk increased? An unsolved issue. *J. Hypertens.* 2019; 37. doi:10.1097/HJH.0000000000002099.
185. Gupta R, Xavier D. Hypertension: The most important non communicable disease risk factor in India. *Indian Heart J.* 2018; 70. doi:10.1016/j.ihj.2018.02.003.
186. Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E *et al.* Hypertension in India: A systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *J. Hypertens.* 2014; 32. doi:10.1097/HJH.000000000000146.

HOW TO CITE: Bhoomika Swarnkar, Abhishek Nand, Lokprabha Hirwani, Harkesh Dadsena, Pushpendra Kumar, Chhavi Rahangdale, Yashika Israni, Helina Tandon, Umakant Sahu, Vishal Jain*, Narendra Kumar, A Review On The Role Of Herbal Medicine In Modulating Hypertension-Related Complications: Mechanistic Insights, Int. J. Sci. R. Tech., 2026, 3 (6), 457-490. <https://doi.org/10.5281/zenodo.20567000>

s