

ORIGINAL PAPER

Open Access



Comparative evaluation of anticancer efficacy and molecular mechanism of silver nanoparticles synthesized from methanolic and ethanolic extracts of *Citrus medica*

Ejaj Ahmad¹, Mohammad Sajid², Alina Athar^{1,3}, Arun Kumar¹, Nimisha Nimisha¹, Abhay Kumar Sharma¹, Mohammad Z. Ahmed⁴, Ali Saeed Alqahtani⁴, Mausumi Bharadwaj² and Sundeep Singh Saluja^{1,5*}

Abstract

The development of biosynthesized and cost-efficient nanomedicine is a promising strategy in cancer therapeutics. Herein, we have synthesized highly potent silver nanoparticles (AgNPs) using methanolic and ethanolic extracts of *Citrus medica* fruit, unraveling the significance of different solvents. Fourier transform infrared spectroscopy confirmed the occurrence of secondary metabolites on the synthesized AgNPs' surface. The Zetasizer analysis showed no significant particle-size differences between both solvents-based-synthesized AgNPs; however, zeta potential showed significant differences in the particle surface charge. The high-resolution transmission electron microscopy confirmed a predominantly spherical morphology of AgNPs. Notably, AgNPs synthesized from ethanolic extracts exhibited smaller sizes (5–7 nm) and a more uniform distribution compared to their methanolic counterparts. The X-ray diffraction demonstrated a polycrystalline nature with a face-centered cubic structure. Functional assays showed a significant increase in the anticancer efficacy of ethanolic extract-derived AgNPs against the breast cancer cells (MCF-7 and MDA-MB-231), accompanied by extensive DNA fragmentation and enhanced induction of apoptosis. Mechanistically, both methanolic and ethanolic AgNPs induce upregulation of pro-apoptotic gene expression (e.g., *TP53*, *BAX*, and *CYC3*) and down-regulation of anti-apoptotic gene (e.g., *BCL2*). The flow cytometry analysis showed a substantial increase of the ethanolic extract-based AgNPs in the late-stage apoptotic cells, which was further confirmed by morphological changes in treated MCF-7 cells. This study reveals the pivotal influence of the selection of solvent on the synthesis and anticancer potency of AgNPs, significantly impacting their ability to induce apoptosis in MCF-7 cells. Our findings introduce a novel strategy for optimizing green-synthesized AgNPs-based therapies, offering promising avenues for advancing nanoparticle-driven cancer treatments and potentially transforming future therapeutic approaches in oncology.

Highlights

- To analyze the role of organic solvent-based extraction of phytochemicals and their influence on the physiochemical properties of the synthesized AgNPs.
- Assessment of the anticancer efficacy of methanolic or ethanolic extracts of *Citrus medica* fruit-based synthesized AgNPs on MCF-7 and MDA-MB-231 cells.

*Correspondence:

Sundeep Singh Saluja
sundeepsaluja@yahoo.co.in

Full list of author information is available at the end of the article




© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

- The synthesized AgNPs induce apoptosis through an increased expression of pro-apoptotic genes (*BAX*, *p53*, and *CYCS*) and reduced expression of the anti-apoptotic gene (*BCL2*).
- AgNPs synthesized through ethanolic extract more effectively induce apoptosis in MCF-7 cells.

Keywords *Citrus medica*, Green synthesis, Silver nanoparticle, Anticancer, MCF-7

Graphical Abstract

Schematic representation of AgNPs synthesized from methanolic or ethanolic extract of *Citrus medica* fruit and evaluations of their anticancer efficacy against MCF-7 cell lines.



Introduction

Breast cancer (BC) falls among the cancers that are associated with a high mortality rate globally and causes 6.9% of all cancer-related deaths (Sung et al. 2021; Bray et al. 2024). The early stage of BC diagnosis would enable a complete cure of the disease. However, due to the lack of visible symptoms at an early stage, the disease is mainly diagnosed at a late stage, leading to a high death rate among cancer patients. The treatment modalities primarily include traditional approaches like surgery, chemotherapy, and radiotherapy. Chemotherapy kills most cancer cells; however, long-term exposure to chemotherapeutics promotes the origin of cancer stem cells (CSC), leading to the development of new cancer cells. These cells create a huge huddle in BC treatment owing to the development of resistance to chemotherapeutic agents, which plays a pivotal role in propagation, recurrence, and metastasis (Nahvi et al. 2022). Thus, strategies need to be adopted to minimize exposure to CSC to anticancer drugs to prevent the development of drug-resistant cells.

Nanomedicine has immense potential for BC treatment by minimizing the long-term exposure of the anti-cancer drugs to the cells by encapsulating them into nanocarriers that release a limited quantity of drugs in a sustained fashion. Thus, numerous innovative approaches have been designed that led to the development of new nanocarriers or ligands decorated nanocarriers in which anticancer drugs were encapsulated, and their efficacy has been analyzed against BC cell lines and xenograft tumors in animal models (Tagde et al. 2022; Rahman et al. 2023). Moreover, some commercial nanomedicines, such as Doxil and Abraxane, were developed against BC. These nanomedicines are now used as adjuvant therapy against BC and show favorable clinical outcomes (Wu et al. 2017; Jiang et al. 2022). However, these nanomedicines are expensive and cause some form of toxicity in healthy cells.

The development of nontoxic and economical nanomedicine hunt led to the exploration of novel metallic nanoparticle potential that includes the use of metals like Au, Ag, and Pt. These metallic nanoparticles have shown

utility in various fields and exhibit efficacy against cancer cells. The high efficacy of metallic nanoparticles against cancer cells is due to their small size distribution, shape, long-term activity, and capability to exhibit optical and magnetic properties (Khan et al. 2017). Metallic nanoparticles are readily taken up by the cells owing to their high density, which could help in the management of cancer (Odularu, 2018; Sharma et al. 2018). Among the metallic nanoparticles, the attributes of silver metal-based nanoparticles exhibited exceptional characteristics including exceptional physiochemical and biological properties, which make them unlike their macroscale counterparts (Sharma et al. 2009).

Moreover, silver is a nontoxic metal/element that shows high electrical and thermal conductivity as well as stability at ambient temperature. Also, it displays a wide array of absorption of visible and far IR regions, surface-enhanced Raman scattering, chemical stability, and non-linear optical behaviors (Zhang et al. 2016; Chouhan, 2018; Jain et al. 2021). These properties of silver metal made it more valuable than other metallic nanoparticles. However, silver nanoparticles prepared by physical and chemical methods may contain harsh and unfriendly environmental compounds and solvents that cause toxicity (Kharissova et al. 2019). Thus, an alternative nanoparticle synthesis method was envisioned, which led to the development of the green synthesis of metallic nanoparticles. The approach overcame some problems associated with physical and chemical methods of metallic nanoparticle toxicity.

The green synthesis of silver nanoparticles uses various parts of plants, algae, and microorganisms, which contain phytochemicals/secondary metabolites and compounds containing functional groups such as aldehyde, alcohol, and ketone, which cause the reduction of Ag^+ to Ag^0 (Zuhrotun et al. 2023). The chain propagation led to the synthesis of AgNPs. Further, these secondary metabolites cause the stabilization and capping of AgNPs. The prepared AgNPs are economical and eco-friendly. The extraction of compounds from the plants or their parts depends on the nature of the solvents owing to the physiochemical properties of the phytochemicals present in the plant (Bitwell et al. 2023). AgNPs were synthesized by extracting the phytoconstituents of plants and their parts in various organic solvents, and their anticancer/biological properties were evaluated (Das et al. 2013; Chitsazi et al. 2016; Heidari et al. 2018; Antropova et al. 2021). The synthesized AgNPs showed effective anticancer properties against some cancer cell lines (Das et al. 2013; Heidari et al. 2018). Kis et al. synthesized AgNPs from ethanolic extract of *Populi gemmae* extract and observed a significant decrease in the growth of MCF-7 and A549 cancer cell lines (Kis et al. 2022). *Citrus medica*

(citron) is a big and rough-surfaced lemon-like fruit belonging to the family Rutaceae (Davidson, 2006). The extensive research focus on its antipathogenic capabilities undoubtedly stems from its remarkable abundance of phytochemicals known for their potent antimicrobial effects (Benedetto et al. 2023).

Herein, we have prepared AgNPs from methanolic and ethanolic extracts of *C. medica* fruit separately and characterized them through various biophysical techniques. The anticancer efficacy of these formulations was checked on the breast cancer cells (MCF-7) to comprehend the influence of solvent in the extraction of secondary metabolites from the fruit and their influence in the synthesis of AgNPs. Further, the molecular mechanisms of cancer cell growth inhibition of these AgNPs-treated cells were evaluated by DNA fragmentation analysis and quantification of proapoptotic, anti-apoptotic, proto-oncogene of the cell cycle, mitochondria, and tumor suppressor gene by real-time PCR.

Material and methods

Material

The *C. medica* (citron) fruit was procured from the local market of Lucknow, Uttar Pradesh, India. Reagents, such as sulforhodamine B (SRB), 2,2-diphenyl-1-picrylhydrazyl (DPPH), silver nitrate, and potassium bromide, were procured from Sigma Aldrich (Sigma-Aldrich Chemicals Pvt. Ltd., Bangalore, India). Penicillin, gentamycin, streptomycin, DMEM/F-12, trypsin–EDTA (0.5%), FBS, and Verso cDNA synthesis kit were obtained from Thermo Scientific (Waltham, MA, USA). Methanol and ethanol were purchased from Qualigens Fine Chemicals, Maharashtra, India. Milli-Q and distilled water were prepared in the laboratory, using the Millipore purification system. A breast cancer cell line (MCF-7) was obtained from ATCC, whereas MDA-MB-231 was procured from the NCCS, Pune, India, and other chemicals used that are not mentioned were also of analytical grade.

Methods

Silver nitrate (AgNO_3) solution preparation

Silver nitrate solution (10 mM) was prepared in Milli-Q water under a magnetic stirrer for 2 h, and the solution was further filtered with Whatman paper no. 1 to remove any impurities. The prepared solution was kept in an amber-colored reagent bottle to avoid photodecomposition of AgNO_3 .

Methanolic and ethanolic *C. medica* fruit extract preparation

The *C. medica* fruit was cleaned thoroughly in running water followed by rinsing with Milli-Q water. Then, it was dried in a hot air oven under controlled 45–50 °C for 48 h. Next, the fruit was chopped into small pieces using a

surgical blade and further dried at the same temperature for 24 h. Subsequently, the dried fruit was made into fine powder by mortar and pestle. Then, 50 mL of methanol and ethanol (70%) was added into 5 g (10% w/v) of each dried powder in a separate Erlenmeyer flask and kept overnight (in the dark) at room temperature. Next, the prepared mixtures (10% w/v) were sonicated using a sonicator (Branson CPX5800H, Richmond, VA, USA) for 1 h and passed through a Whatman filter paper (Grade 1). The filtrates were stored at 4 °C and used (within 2 weeks) to synthesize AgNPs.

Biosynthesis and isolation of AgNPs from methanolic and ethanolic extract

The solubility of biochemical compounds depends on the polarity of the solvent. Thus, we have extracted the secondary metabolites/phytochemicals of *C. medica* fruit in two different organic solvents, viz., methanol and ethanol (70% v/v) that were used to synthesize AgNPs.

For methanolic extract-based green synthesis of silver nanoparticles (Met-AgNPs), we have prepared (10% w/v) methanolic extract of *C. medica* fruit. The influence of different parameters such as concentration of AgNO₃ solution, quantity of extract, temperature, pH, rotation (rpm), and incubation time was carefully evaluated to prepare small size AgNPs. After varying each parameter, the UV–Vis spectrum of the reaction mixture was analyzed to ascertain whether nanoparticles were synthesized or not. Once we got the designated peaks of the reaction mixture, then subsequently, the size of the particles was analyzed using Zetasizer. By varying these parameters, we successfully synthesized the smallest size of AgNPs when 10 mM of 30 mL AgNO₃ and 1 mL of extract (30:1) were allowed to continuously rotate at 500 rpm at 70 °C for 2 h on a magnetic stirrer (DAIHAN Scientific, South Korea). A gradual time-dependent change of color from a colorless solution to pale yellow and then light brown was observed. However, a distinct brown color appeared after 1 h. This optical color change advocates that biomolecules/secondary metabolites present in the *C. medica* fruit extract reduced the silver ions into AgNPs.

Similarly, the smallest size of AgNPs was prepared from the ethanolic extract (Eth-AgNPs) by varying the parameters mentioned above. We could prepare the smallest size of Eth-AgNPs from the same ratios of AgNO₃ and ethanolic *Citron* fruit extract, temperature, and rotation. However, the Eth-AgNPs are formed at a lower AgNO₃ solution (1 mM) and require a longer duration, viz., 2 h. A gradual color change of the solution (from light brown to dark brown) was observed within 2 h time which suggests AgNP synthesis owing to the reduction of Ag⁺ → Ag⁰ by secondary metabolites of the fruit extract. The

biosynthesized AgNPs were separated by centrifugation for 10 min at 13,500 ×g by a centrifuge (Eppendorf, Hamburg, Germany). The AgNPs pellet was washed three times with MQ water. Then, the isolated AgNPs were dried by placing them into a vacuum oven at 45 °C, 0.8 bar (DAIHAN Scientific, South Korea), for 2 h and frozen at –80 °C until further applications.

Biophysical characterization of synthesized AgNPs

For the characterization (size, shape, and charge) of prepared AgNPs, various techniques were used including UV–Vis spectroscopy, FTIR, Zetasizer, XRD, and TEM imaging. Initially, the UV–Vis spectra of the Met-AgNPs and Eth-AgNPs were acquired by adding the 200 µL of the formulations in microtiter plate wells and scanned from 300 to 700 nm range with 1 nm resolution through a multimode reader (Varioskan LUX, Thermo Scientific, USA). Next, the FTIR spectra of the Met-AgNPs and Eth-AgNPs were scanned within a wavelength 4000–500 cm^{–1} range with 1 cm^{–1} spectral resolution (Bruker, Optik GmbH, Karlsruhe, Germany). Further, the Met-AgNP's size (average), polydispersity index (PDI), and zeta potential were determined (at 25 °C temperature) by the Malvern Zetasizer Pro-blue instrument (Malvern Analytical Ltd., Malvern UK). To comprehend the phase and crystalline structure of the Met-AgNPs and Eth-AgNPs, an XRD pattern was noted using a Rigaku D/Max-IIIC diffractometer applying CuKα monochromatic radiation and AXIS Supra by a monochromatic source (Al anode Kα: 1486.6 eV) between 2θ of 10 to 80°. TEM images of either Met-AgNPs or Eth-AgNPs AgNPs were acquired to analyze their actual sizes and morphologies. Before imaging, the samples were sonicated (5 min) in a sonicator (Branson, 5800). The respective AgNPs (100 µL) were added to Milli-Q water (900 µL), and samples (5 µL) were gently positioned on a carbon-coated copper grid, and samples were dried in the air using a lamp heat. The TEM evaluation of both AgNPs was accomplished at an accelerating voltage of 200 kV (FEI Tecnai G² 20 S-TWIN).

Analysis of antioxidant activity

The antioxidant action of *C. medica* fruit extracts (methanolic/ethanolic) and their respective extract-based AgNPs (Met-AgNPs and Eth-AgNPs) was determined by the DPPH assay (Bhakya et al. 2016). Briefly, 100 µL of different concentrations of extracts and synthesized AgNPs prior dissolved in an 80% ethanol solution was added into 96 well plates (in triplicate). Varying concentrations of ascorbic acid (100 µL) were used as a positive control. Afterward, 0.2 mM DPPH reagent was prepared in ethanol (80%) and added to each well. The plate was incubated for 30 min (no light exposure) at room

temperature, and the plate was read at 517 nm by a multimode reader (Varioskan LUX, Thermo Scientific). The free radical scavenging activity was determined by the below-mentioned equation.

$$\text{DPPH} - \text{free radical scavenging activity} = \frac{\text{Abs (control)} - \text{Abs (sample)}}{\text{Abs (control)}} \times 100$$

Evaluation of anticancer analysis

Analysis of antiproliferative activity against breast cancer cells To measure the cytotoxic activity of Met-AgNPs and Eth-AgNPs, the SRB assay was accomplished as defined earlier by Vichai et al. (2006) with few changes (Vichai and Kirtikara, 2006). The cells (MCF-7/MDA-MB-231) were seeded (10^4 cells/well) in culture plates (96 wells) and incubated in a CO₂ incubator for 24 and 48 h at 37 °C. The DMEM medium was replaced with fresh medium (100 µL/well), and different concentrations of either Met-AgNPs or Eth-AgNPs ranging from 3.125 to 100 µg/mL prepared in DMEM medium were supplemented to the corresponding wells in triplicates. After incubation (24 and 48 h), the cells were washed with PBS and fixed using trichloroacetic acid (10%) for 1 h at 4 °C. Then, the plates were washed four times with sterile distilled water and dried. A total of 100 µL of SRB solution (0.057%) was added for 30 min at room temperature. The plates were immediately rinsed four times with acetic acid (1%) and kept for drying at room temperature. Finally, 10 mM Tris base buffer (pH 10.5) was added (200 µL/well) to the wells, and absorbance at 564 nm was taken by a microplate reader (SPECTROstar Nano, BMG Labtech, Ortenberg, Germany).

Analysis of DNA fragmentation

MCF-7 cancer cells were grown on 6-well plates and incubated with the IC₅₀ of AgNPs for 24 h. Later, the cells were harvested after washing with PBS and collected in the microcentrifuge tube by centrifuging at $4000 \times g$ for 5 min. The lysis buffer (600 µL/tube) comprising Tris-HCl (10 mM, pH-7.4), NaCl (400 mM), EDTA (100 mM), and SDS (6%) was added to the tubes. Then, RNase solution (10 µL/tube) treatment was performed to remove any RNA contaminant. Further, the solution was incubated at 37 °C for 5 min, and proteins were precipitated by adding 200 µL of 6 M NaCl solution. Then, the solution was centrifuged ($13000 \times g$, 10 min) at ambient temperature. The supernatant was collected in the microcentrifuge tube and mixed with isopropanol (600 µL/tube). The mixture was placed on ice for 15 min. The same steps were repeated by adding isopropanol and incubated for 20 min. Then, the DNA (precipitate) was carefully cleaned with 70% ethanol (600 µL) and dried

at room temperature. The purified DNA was dissolved in 200 µL of buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). Finally, agarose gel electrophoresis was performed to observe isolated DNA.

RNA extraction and cDNA construction

RNA was isolated from untreated cells (control) and Met-AgNPs- or Eth-AgNPs-treated MCF-7 using the TRIzol method. Briefly, the cell pellet was washed two times with chilled PBS and homogenized in TRIzol reagent followed by the addition of chloroform (200 µL) and then incubated at RT for 15–20 min. The tubes were centrifuged (12,000 rpm, 4 °C, 15 min), and the upper aqueous phase was collected into another microcentrifuge tube. The isopropanol (1 mL) was added to the tubes and incubated for 2 h at –20 °C followed by centrifugation (12,000 rpm, 4 °C, 15 min). The precipitate (RNA) was washed two times with 70% ethanol. Then, nuclease-free water was added to the pellet and kept at –80 °C for further processing. The Verso cDNA synthesis kit was used for cDNA synthesis from the isolated RNA. A master mix reaction (20 µL/tube) was prepared using reaction buffer (5×), dNTPs mix (5 mM each), anchored Oligo dT primer (0.5 µg/µL), random hexamer (0.4 µg/µL), Verso enzyme mix, and template RNA (1 ng). To verify the formation of the cDNA strand, it was subsequently amplified using glyceraldehyde 3-phosphate dehydrogenase (GAPDH) housekeeping gene primer pairs. The thermal cycler conditions include initial denaturation (95 °C, 5 min), 35 cycles of denaturation (95 °C, 30 s), annealing (60 °C, 35 s), and extension (72 °C, 35 s) followed by a final extension step (72 °C, 7 min). The amplified products were resolved on agarose gel (2%), and synthesized cDNA was stored at –20 °C for further use.

Quantitative PCR for mRNA expression analysis The expressions of apoptotic genes (*TP53*, *CYCS*, *BCL2L1*, *BAX*, and *CCND1*) and caspases (*CASP9*, *CASP8*, and *CASP3*) were assessed by using the specific primer pairs (Table 1). A housekeeping gene, *GAPDH*, was used as an internal control. The total reaction volume was made 10 µL by adding 1× SYBR green dye, 10 pM each of forward and reverse primer, and 100 ng/µL cDNA. Thermal cycler conditions comprise a denaturation step (95 °C, 5 min) followed by annealing and extension steps at 53 °C for 30 s for 40 cycles. The PCR reaction for gene-specific primers and the internal control (*GAPDH*) was carried out in triplicate. To minimize the rate of error, a non-template control was also included in each reaction. Finally, the

Table 1 Primer sequence of genes along with their annealing temperature and amplicon size

Genes	Primer sequence		Annealing temp (°C)	Amplicon size (bp)
	Forward (5′–3′)	Reverse (3′–5′)		
TP53	AAGTCTGTGACTTGACGTAATCC	GTCATGTGCTGTGACTGCTTGTA	57	152
CYCS	TGGGTGATGTTGAGAAAGGC	ATTGGCGGCTGTGTAAGAG	55	158
BCL2L1	GGTTGACTTTCTCTCTAC	CGATTGAGTCCCTTCTG	54	103
BAX	GCTGCAGAGGATGATTG	CCTTGAGCACCAGTTTG	53	146
CCND1	CGAAGTGGAACCATCC	CTCCTTCTGCACACATT	54	130
GAPDH	AGCGAGATCCCTCCAAA	CTTGAGGCTCTTGTCATACT	60	200
CAPS9	GGCTCTTCCTTTGTTTCATCTCC	TCACCAATCTCCAGAACCA	58	210
CAPS8	GATGTATTCCAGAGACTCCAG	GGTAGGTAATCAGCAATCCA	58	110
CASP3	ATGGAAGCGAATCAATGGAC	AAACATCACGCATCAATTC	57	240

analysis of the melting curve was performed to verify template specificity, and the relative fold change was estimated by the $2^{-\Delta\Delta C_t}$ formula.

Analysis of MCF-7 cells apoptosis by flow cytometry

MCF-7 cells that were present in the logarithmic growth phase were seeded (1×10^6 cells/well) in a 6-well plate and exposed to varying concentrations of Eth-AgNPs and Met-AgNPs (5, 10, and 25 $\mu\text{g/mL}$). The DMSO (0.01%) treatment was given to control cells as a vehicle control. After 24 h of incubation, apoptotic cells induced by Eth-AgNPs and Met-AgNPs were identified with the help of an Annexin V/PI staining kit (BioLegend, USA). In brief, the cells were collected, washed, and mixed with a binding buffer (100 μL) provided with the kit. Then, a mixture (1:1) of Annexin-FITC and PI was added to the tubes. After 30 min, binding buffer (400 μL) was supplemented to the labelled cells. Subsequently, cells were acquired in a Flow Cytometer (Cytomics FC 500; Beckman Coulter, Brea, CA, USA).

Analysis of apoptotic cells morphology

Morphological changes in the MCF-7 cell following treatment with either Met-AgNPs or Eth-AgNPs were

(37 °C, 24 h). Then, the cells were washed with PBS, and images of each well were captured by inverted fluorescence microscopy.

Analysis RBC hemolysis

The RBC hemolysis test was performed to comprehend the toxicity of the synthesized Met-AgNPs and Eth-AgNPs. The RBC hemolysis test was performed by following the published procedure (Bansal et al. 2015). Briefly, human blood was centrifuged at 2500 rpm for 10 min, and plasma and buffy coat were carefully removed. The RBC cells were washed twice with phosphate-buffered saline. One-hundred microliters of 4% RBC cells was taken in microfuge tubes. Then, 100 μL of varying concentrations of Met-AgNPs or Eth-AgNPs, such as 200, 100, 50, and 25 $\mu\text{g/mL}$, was added in respective tubes in triplicate, while 100 μL of 1% (v/v) Triton solution and PBS was taken in separate tubes as a positive and negative control, respectively. The reaction tubes were incubated at 37 °C for 1 h. Once the incubation was over, tubes were centrifuged at 2000 rpm for 10 min. Then, 100 μL of supernatants from the tubes was carefully taken and placed in the respective 96 wells in the ELISA plate. The absorbance of the supernatant was measured at 540 nm. The % lysis of RBC was calculated using the following formulas as below:

$$\% \text{ RBC hemolysis} = \frac{\text{Absorbance of sample} - \text{Absorbance of negative control}}{\text{Absorbance of positive control} - \text{Absorbance of negative control}}$$

directly observed using an inverted fluorescence microscope (Optica, Italy). The cells were placed (0.1×10^6 cells/well) into tissue culture plates (6 wells) and allowed to grow overnight in a CO_2 incubator at 37 °C. The next day, the media was discarded, and different concentrations of treatment for either Met-AgNPs or Eth-AgNPs were given, and the plates were placed in a CO_2 incubator

Statistical analysis

Analysis of variance (ANOVA) test was practiced to determine the statistical differences between a pair of data sets. The $p\text{-value} < 0.05$ was treated as statistically significant between a pair of data sets.

Result and discussion

The extraction of phytochemicals/secondary metabolites from the plants/parts of the plant depends on the polarity of solvents and their physiochemical properties (Nawaz et al. 2020). Both methanol and ethanol are polar organic compounds; however, methanol is more polar than ethanol due to the absence of $-CH_2-$ electron pumping groups, which reduce the partial negative charge on oxygen molecules. This decreases the partial charge difference between oxygen and hydrogen in the hydroxyl group of ethanol. Moreover, methanol is among the prominent solvents used to extract phytochemicals from herbs and spices. Ethanol 70% (v/v) contains 30% water molecules, which increases its polarity, leading to an increase in the extraction of more polar compounds/hydrophilic secondary metabolites.

Plants contain various secondary metabolites or phytochemicals like carbohydrates, amino acids, proteins, flavonoids, alkaloids, tannins, polyphenols, and saponins (Roy et al. 2019). The differences in solvent polarity, including the addition of 30% water to ethanol (70% v/v), led to varying levels of phytochemical extraction. This, in turn, resulted in differences in the compounds and their quantities, affecting the reduction of Ag^+ to Ag^0 and the subsequent stabilization and capping during the synthesis of AgNPs. As a result, AgNPs synthesized from different extracts contain distinct surface functional groups, leading to variations in their antioxidant properties and anticancer efficacy against MCF-7 cells. In contrast to traditional chemical synthesis techniques, this method avoids the usage of toxic compounds and high temperatures, providing a secure and more justifiable alternative. Thus, the current study provides pragmatic evidence that the AgNPs synthesized employing 70% ethanol extract exhibit better anticancer activity against MCF-7 cells as compared to the methanol extract-based AgNPs.

Silver nanoparticles characterization

UV-Vis spectroscopy analysis

The mixing of methanolic extract (1 mL) with 10 mM $AgNO_3$ solution (30 mL) causes a gradual color change from pale yellow to light brown, which is the best primary noticeable index of the formation of AgNPs. Similarly, previous studies observed a color change during the synthesis of AgNPs (Dehghanizade et al. 2018; Masum et al. 2019), which confirmed the completion of the reaction between the phytochemicals present in the methanolic extract and $AgNO_3$ solution. Further, the absorption spectrum of the formed AgNPs (light brown color) displayed maximum absorption between 300 and 400 nm, suggesting the reduction of $Ag^+ \rightarrow Ag^0$ in the presence of phytochemicals from *C. medica* fruit extract and the synthesis of AgNPs (Fig. 1A).

Similarly, the absorption spectrum of the synthesized Eth-AgNPs was analyzed and observed absorbance maxima between 320 and 380 nm (Fig. 1B). The high absorbance between 320 and 380 nm was due to the surface plasmon phenomenon of the electron cloud present on the surface of AgNPs as evident from the previous observations of ethanolic extract-based synthesis of AgNPs (Mahmoud et al. 2020; Samuggam et al. 2021).

Analysis of AgNPs by FTIR

The Ag ions reduction by the phytochemicals present in the plant extract is attributed to the synthesis of AgNPs; subsequently, these compounds can cause stabilization and capping of AgNPs. The functional groups of these phytochemicals present on the surface of AgNPs were identified by the FTIR technique. The FTIR spectrum of the Met-AgNPs exhibited major absorption peak shifts at 3468.51, 2927.3, 2854.5, 2090.89, and 1749,1634.89 cm^{-1} . Moreover, new absorption peaks appeared at lower wave numbers, such as 1385, 1304, 1250, 1096, 1022, 680, and




Fig. 1 UV-visible spectrum of the AgNPs synthesized from *C. medica* fruit extract. **A** Met-AgNPs. **B** Eth-AgNPs




Fig. 2 FTIR spectrum of biosynthesized AgNPs. **A** Methanolic extract and Met-AgNPs. **B** Ethanolic extract and Eth-AgNPs

456 cm^{-1} (Fig. 2A). The peak shifts, as well as the appearance of new peaks, indicate the reducing, stabilizing, and capping actions of the phytochemicals of *C. medica*. A shift in the peak from 3369.32 cm^{-1} to 3468.2 cm^{-1} was due to either O–H or N–H stretching of the phenols available in the plant extract (Bilal et al., 2019). The occurrence of C–H stretching belonging to the aliphatic functional group/methylene group is designated by the peak at 2927.3 cm^{-1} . This peak, 2927.3 cm^{-1} , also indicates the existence of triterpenoid saponins (Qais et al. 2019). The peak at 2854.5 cm^{-1} demonstrated the presence of alkenes' asymmetric and symmetric stretches. The absorption peak at 1634.89 cm^{-1} matches the C=C group of aromatic compounds (Raghunandan et al. 2010). The peak at 1385 cm^{-1} signifies the incidence of germinal methyl groups. The peak at 1022 cm^{-1} simplifies the C–O stretching of the ether group (Dehghanizade et al. 2018). On the other hand, the peak at 680 cm^{-1} was owing to the presence of the =CH group of the aromatic compounds.

Likewise, Met-AgNPs, the major absorption peaks of the ethanolic extract, were observed at 3468.78, 2922, 2852.3, 2087, and 1639.04 cm^{-1} (Fig. 2B). The biosynthesized Eth-AgNPs showed a very minor change in the shift of the major spectrum peaks. However, at the lower wave number, the spectrum showed new absorption peaks at 1463.6, 1377.3, 1341, 1320.5, 1120.5, 1100, and 523.25 cm^{-1} . Of these peaks, some of the absorption peaks like 1120.5, 1100, and 523.25 cm^{-1} were absent in the Met-AgNPs spectrum. The result can be concluded that most of the secondary metabolites were common in both extracts; however, some metabolites are different in both extracts.

Size and zeta potential analysis

The mean size and PDI of the Met-AgNPs were observed to be 154.8 nm and 0.206, respectively (Fig. 3A). The small PDI value suggests that the size distribution of the particle was narrowed down, suggesting




Fig. 3 AgNPs size analysis using DLS. Methanolic extract-based biosynthesized AgNPs. **A** Mean size, **B** ζ -potential, and **E** size vs. percentage of Met-AgNPs. Ethanolic extract-based biosynthesized AgNPs, **C** mean size, **D** ζ -potential, and **F** size vs. percentage of synthesized Eth-AgNPs

uniformity of the particles. Similarly, the mean size and PDI of the Eth-AgNPs were measured to be 163 nm and 0.37, respectively (Fig. 3C). The observed ζ -potential of the methanolic and ethanolic extract-based synthesized AgNPs were -23.6 mV and -16.75 mV, respectively, advocating the stability of these synthesized AgNPs (Fig. 3B & D). The negatively charged functional groups of the various phytochemicals attached to the nanoparticles' surface ascertained the colloidal capability of the prepared AgNPs (Ahn et al. 2019). Further, a graph between particle size and their percent distribution to better comprehend the size distribution was shown (Fig. 3E & F).

The comparative analysis of both Met-AgNPs and Eth-AgNPs suggests that there is no significant difference in the mean size of AgNPs synthesized from either methanolic extract or 70% ethanolic extract; however, a significant difference in the PDI was observed. The high PDI of Eth-AgNPs might be due to the inference of ethanol present in the reaction mixture, which was not completely evaporated owing to the synthesis being performed at 70°C , while the boiling point of ethanol is $\sim 78^\circ\text{C}$. On the contrary, methanol has a lower boiling point ($\sim 64^\circ\text{C}$). Thus, during the synthesis at 70°C , most of the methanol was evaporated; hence, it did not interfere with size measurement. Earlier, Esfanddarani et al. synthesized AgNPs from methanolic extract and 70% ethanolic extract of *Malva sylvestris* and observed that ethanolic extract-based AgNPs showed smaller particle size distribution as compared to methanolic extract-based AgNPs (Mahmoodi Esfanddarani et al. 2018).

The DLS technique measures the hydrodynamic size of the AgNPs which include an additional hydration layer around the nanoparticles. Hence, the size of the nanoparticles appears larger in comparison to the actual size as determined by the TEM imaging (Zia et al. 2017). Additionally, the hydrodynamic size of AgNPs can also

be influenced by the adsorption of phytochemicals on the synthesized AgNPs surface. The small variation in the AgNP hydrodynamic size could be due to the attachment of varying amounts of the phytochemicals on the particle's surface (Vanlalveni et al. 2021).

XRD analysis of AgNPs

The crystalline property of the prepared AgNPs was examined by the X-ray diffraction (XRD) analysis. The XRD spectrum of the Met-AgNPs synthesized from the methanolic extract showed prominent six 2θ peaks at 27.8 , 38.6 , 46.3 , 54.8 , 64.9 , and 77.1° (Fig. 4A). Besides these peaks, some other minor additional peaks were evident, which were due to noise or organic solvent contaminants. These Bragg's diffraction angles correspond to the lattice planes [122], [111], [200], [120], [220], and [311], respectively (Fig. 4A). Ojha et al. have synthesized AgNPs from *Ricinus communis* var. Carmencita leaf extract, and XRD diffraction was performed to ascertain the polycrystalline nature of the synthesized AgNPs. They observed diffraction angles at 27.81° , 32.19° , 38.16° , 44.43° , 46.23° , 54.93° , 64.65° , and 77.61° . These diffraction angles correspond to Bragg's diffraction planes [210], [122], [111], [200], [231], [142], [241], [220], and [311], which suggest a face-centered cubic structure (Ojha et al. 2017). Thus, the lattice planes of our experimental data suggest that the synthesized Met-AgNPs are polycrystalline in nature, with a face-centered cubic structure.

In contrast, XRD of the Eth-AgNPs showed five 2θ peaks at 27.8° , 32.2° , 46.3° , 57.4° , and 76.6° (Fig. 4B), which correspond to Bragg's lattice planes [210], [122], [231], [241], and [311], respectively. Meng et al. have synthesized AgNPs from *Ficus altissima* Blume leaf extract and have characterized its polycrystalline nature by performing X-ray diffraction analysis. They observed diffraction angle 2θ values at 27.81° , 32.16° , 38.12° , 44.3° , 46.21° , 54.83° , 57.39° , 64.42° , and 77.45° , corresponding to




Fig. 4 XRD spectrum of the biosynthesized AgNPs. **A** Met-AgNPs. **B** Eth-AgNPs

Bragg's diffraction planes [210], [122], [111], [200], [231], [142], [241], [220], and [311] (Meng, 2015). Similarly, Hemlata et al. have observed XRD peaks at 32.18°, 38.04°, 46.13°, 54.63°, and 77.08° of the AgNPs prepared from *R. communis* var. Carmencita leaf extract, which correspond to the [111], [200], [120], [202], and [311] Bragg's diffraction planes, respectively. Such planes are in agreement with the face-centered cubic structure and polycrystalline nature of the prepared Eth-AgNPs (Hemlata et al. 2020). Thus, the XRD diffraction angle of Eth-AgNPs confirmed the polycrystalline nature with a face-centered cubic structure.

TEM nalysis of AgNPs

HR-TEM pictures were captured at different magnifications for the size and morphology of the AgNPs (Fig. 5). Met-AgNPs TEM images suggest that the majorly formed particles were spherical with dimensions of 15–20 nm; however, some particles portray cylindrical morphology (Fig. 5A, B, C, D, E, F). Many authors synthesized AgNPs from methanolic extracts of different plants and observed similar sizes and morphologies (Singh and Navneet, 2021; Bawazeer et al. 2021). On the other hand, the TEM images of the Eth-AgNPs formed were relatively smaller in size, i.e., ranging from 5 to 7 nm (Fig. 5G, H, I, J, K, L). The ethanolic extract-based AgNPs showed mostly spherical morphology, while a few of them were also observed as cylindrical. Esfanddarani et al. have shown that AgNPs prepared from 70% ethanolic extract showed smaller size and a narrow polydispersity index compared to methanolic extract-based AgNPs (Mahmoodi Esfanddarani et al. 2018). At very high magnification (650,000×), the fringes were also visible on the particle's

surface of both Met-AgNPs and Eth-AgNPs, suggesting a uniform distribution of phytochemical constituents on AgNPs (Fig. 5D).

SAED image analysis of AgNPs SAED images give crucial information regarding the polycrystalline nature of the synthesized AgNPs (Rani et al., 2020). The result of the SAED image of both Met-AgNPs and Eth-AgNPs showed five bright electron diffraction rings, as seen in Fig. 5F and K, respectively. The presence of a bright ring suggests the pattern of crystal of Ag atom arrangement in the AgNPs. Our obtained data on crystal planes aligns with previous findings (Soshnikova et al. 2018; Yap et al. 2020). The Ag ions reduction in the presence of extracts (methanolic/ethanolic) of *C. medica* fruit led to the corresponding AgNPs synthesis, suggesting the polycrystalline nature of the formed AgNPs, which correlates well with the obtained XRD data. Following our data, an earlier study observed a similar face-centered cubic structure of the prepared AgNPs with five electron diffraction rings, which correspond to the lattice arrangement of [111], [200], [220], [311], and [222] (Kanniah et al. 2023).

Analysis of antioxidant efficacy of AgNPs

The antioxidant action of the extracts (methanolic/ethanolic) and the respective extracts-based AgNPs (Met-AgNPs and Eth-AgNPs) was detected by the DPPH assay. Met-AgNPs exhibited slightly higher scavenging capacity than the Eth-AgNPs (Fig. 6). The methanolic extract alone and Met-AgNPs demonstrated a dose-dependent increase in scavenging activity (Fig. 6A). Similarly, the ethanolic extract alone and its corresponding AgNPs showed comparable free radical scavenging efficacy, also




Fig. 5 TEM images of Met-AgNPs (A, B, C, D, E, F) with varying magnifications (A, B, C). Dark-field image (D), HR-TEM image with fringes (E), and SAED image of AgNPs (F). TEM images of Eth-AgNPs (G, H, I, J, K, L) with varying magnifications (G, H, I). Dark-field image (J). HR-TEM image with fringes (K) and SAED image of AgNPs (L)




Fig. 6 Antioxidant activity of the biosynthesized AgNPs analyzed by DPPH assay. **A** Methanolic extract and Met-AgNPs. **B** Ethanolic extract and Eth-AgNPs

displaying a dose-dependent scavenging effect (Fig. 6B). Moreover, the scavenging efficacy of the methanolic extract and methanolic extract-based AgNPs is significantly higher than that of the ethanolic extract and ethanolic extract-based AgNPs. The IC_{50} values of the free radical inhibition of the Met-AgNPs and Eth-AgNPs were observed to be 424 µg/mL and 382.45 µg/mL, respectively. This finding suggests that the methanolic extract and the ethanolic extract and their respective AgNPs could not show strong antioxidant activity.

Analysis of anticancer activity of the AgNPs

Treatment of cancer is still a challenging task for clinicians, especially in the advanced stages. Chemotherapy in the form of monotherapy or combination therapy received some success and prolonged patients' lives to some extent. However, anticancer drugs also cause severe toxicities to healthy cells, leading to deleterious side effects (Ahmad et al. 2021). This limits the patient's survival from a few months to a few years, depending on age and health. To overcome the toxicity, various strategies have been adopted, and the encapsulation of chemotherapeutics into nanocarriers is a promising approach. However, the approach is complex, expensive, and still causes some form of toxicity to the healthy cells (Ahmad et al. 2021).

The green synthesis of AgNPs was developed, and its potential is being extensively explored by many research groups in the treatment of various cancers. The approach has many attributes that enable it to be a potential candidate for the treatment of cancer (Ratan et al. 2020). Thus, the anticancer efficacy of the developed Met-AgNPs and Eth-AgNPs was analyzed at 24 h and 48 h of incubation on two different human breast cancer cells including MCF-7 and MDA-MB-231 cancer cells. Both formulations, Met-AgNPs and Eth-AgNPs, showed cytotoxic effects in a dose-dependent manner (Fig. 7). At a 50 µg/mL dose of

Met-AgNPs, nearly 45% growth inhibition was observed at 24 h of incubation, which was significantly enhanced to 75% when the incubation was extended to 48 h, suggesting the efficacy of Met-AgNPs against MCF-7 cells (Fig. 7A). As the dose further increased to 100 µg/mL, nearly 72% of the cell death was observed at 24 h, which increased to 94% with the extension of incubation time to 48 h (Fig. 7A). Similarly, Met-AgNPs treatment to MDA-MB-231 cells displayed a significant ($p < 0.0001$) reduction in the growth of cancer cells compared to the untreated group (control) with an IC_{50} value of 37.10 µg/mL at 24 h and 38.91 µg/mL for 48 h incubation (Fig. 7C). The result indicates that the Met-AgNPs were noticeably effective against both breast cancer cells. Muthukrishnan et al. have shown that methanolic extract-based AgNPs inhibit the growth of Dalton's lymphoma ascites tumor (Muthukrishnan et al. 2018). Additionally, our previous study showed that AgNPs synthesized from the methanolic extract of *Alpinia galanga* exhibit high efficacy against cervical cancer (Ahmad et al. 2024).

To understand whether AgNPs synthesized from extracts in different organic solvents induce varying levels of anticancer activity, and if so, to determine the extent of these differences, we also evaluated the anticancer activity of the AgNPs prepared from ethanolic extract against MCF-7 and MDA-MB-231 cancer cells. The result of the Eth-AgNPs also showed a dose-dependent cell growth inhibition in both cell lines (Fig. 7B, D). At 6.25 and 12.5 µg/mL doses, nearly 25% and 53% of the MCF-7 cancer cell growth inhibition was observed, respectively. Further increase of dosage to 25 µg/mL inhibited the growth of almost 87% of cancer cells after 24 h of incubation, which further enhanced to 90% with the increased incubation time (48 h). However, further increase of AgNP doses showed no significant inhibition both at 24 and 48 h of incubation time. The IC_{50} value of the Eth-AgNPs was 11.58 µg/mL, suggesting that Eth-AgNPs are more




Fig. 7 The anticancer activity of synthesized AgNPs was monitored against the MCF-7 and MDA-MB-231 breast cancer cells. The cells were treated with different concentrations (1.56–100 µg/ml) of synthesized AgNPs, and cytotoxicity was determined by the SRB method after 24 and 48 h of incubation. **A** Met-AgNPs effect on MCF-7 cells. **B** Eth-AgNPs effect on MCF-7 cells. **C** Met-AgNPs effect on MDA-MB-231 cells. **D** Eth-AgNPs effect on MDA-MB-231 cells

effective (than Met-AgNPs) in inhibiting the growth of MCF-7 cancer cells. Equivalently, Eth-AgNPs treatment showed a significant ($p < 0.0001$) decrease in the MDA-MB-231 cells viability as compared to the untreated group (control) with an IC_{50} value of 29.28 µg/mL and 10.15 µg/mL for 24 h and 48 h incubation, respectively (Fig. 7D). The high efficacy of Eth-AgNPs could be attributed to the additional extraction of hydrophilic secondary metabolites in 70% ethanol, which contains 30% water. Kis et al. synthesized AgNPs from the ethanolic extract of *Populi gammae* and observed significant, dose-dependent growth inhibition in both MCF-7 and lung carcinoma epithelial cell line A549 (Kis et al. 2022). The AgNPs synthesized by aqueous extracts of *Lasiurus scindicus* (L-AgNPs) and *Panicum turgidum* (P-AgNPs) seeds showed the cytotoxic effect on MDA-MB-231 cells, and IC_{50} values were 141.6 µg/mL and

77.59 µg/mL (Alburae et al. 2024). The synthesized AgNPs showed a better cytotoxic effect on MCF-7 cells compared to MDA-MB-231; therefore, further experiments were performed on MCF-7 cells.

Analysis of formulation toxicity by RBC hemolysis test

To analyze the toxicity of the synthesized Met-AgNPs and Eth-AgNPs, RBC hemolysis was performed. The results suggest that up to 50 µg/mL of either Met-AgNPs or Eth-AgNPs caused <1% of RBC hemolysis (Fig. 8). The concentration of AgNPs further increased to 100 µg/mL; the lysis was slightly increased in both extract-based AgNPs. However, in further increase of AgNPs concentration to 200 µg/mL, the RBC lysis is significantly increased in both AgNPs, especially more prominently increased in Eth-AgNPs. Importantly, our in-house formulations, Met-AgNPs and Eth-AgNPs, showed high toxicity to the




Fig. 8 RBC hemolysis (%) activity of the AgNPs treated for 1 h of methanolic and ethanolic extract-based AgNPs

cancer cells (MCF-7) at 100 µg/mL. However, this concentration causes little hemolysis to the RBC cells (< 2%). It was reported that < 5% RBC hemolysis of the formulation is safe for in vivo study.

Analysis of the molecular mechanism

DNA fragment analysis

DNA fragmentation is one of the peculiar features of apoptosis (Alharbi and Alsubhi 2022). DNA fragment analysis revealed a thinner band in samples treated with Met-AgNPs or Eth-AgNPs compared to the control, indicating smaller DNA fragments. *This was particularly evident in the smear in Eth-AgNP-treated cells. However, no clear smear in Met-AgNPs-treated samples (lane 3) was evident owing to relatively less fragmentation.* This suggests that Eth-AgNPs were more effective in breaking intact DNA, corroborating the MTT assay of Eth-AgNPs

treated MCF-7 cells. A similar observation has been reported in A549 cells treated with AgNPs prepared from *Azadirachta indica* (Alharbi and Alsubhi 2022) (Fig. 9).

Analysis of the molecular mechanism of apoptosis

To unravel the molecular mechanism behind the induction of apoptosis of the MCF-7 cells treated with either Met-AgNPs or Eth-AgNPs, the expression of some of the genes involved in the intrinsic pathways, viz., CCND1 (cyclin D1) and CYCS (cytochrome C) (Fig. 10A). The earlier gene implies the progression of the cell cycle's G1 phase to the S phase, while the later gene is a small heme-regulating protein of the mitochondria. Also, the extrinsic pathways, viz., proapoptotic TP53, BAX, and antiapoptotic BCL2L1 were analyzed by using their respective primers listed in Table 1. Earlier studies suggest that the AgNPs promote tumor cell death primarily through apoptosis, which is mediated by the activation




Fig. 9 DNA fragmentation study of the MCF cells treated with L1-Control, L2-Eth-AgNPs, and L3-Met-AgNPs (A). The image of RNA extracted from the MCF-7 cells treated with AgNPs on 1.2% agarose gel, L1-Control, L2-Eth-AgNPs, and L3-Met-AgNPs (B)




Fig. 10 **A** Relative mRNA expression of genes of MCF-7 cancer cell lines treated with either Met-AgNPs or Eth-AgNPs. **A** *TP53* gene expression, p -value between untreated vs. Meth-AgNPs < 0.05 . **B** *CCND1* gene expression, p -value untreated vs. Met-AgNPs < 0.05 ; untreated vs. Eth-AgNPs < 0.05 . **C** *CYCS* gene expression, p -value untreated vs. Met-AgNPs < 0.05 , untreated vs. Eth-AgNPs < 0.05 . **D** and **E** *BAX* gene expression, p -value between untreated vs. Met-AgNPs < 0.05 and untreated vs. Eth-AgNPs < 0.05 . **E** *BCL2L1* gene expression, p -values untreated vs. Met-AgNPs < 0.05 , untreated vs. Eth-AgNPs < 0.05 . **B** Relative mRNA expression of genes *CASP9* (**A**), *CASP8* (**B**), and *CASP3* (**C**) of the MCF-7 cells treated with either Met-AgNPs or Eth-AgNPs. The p -values are statistically not significant between groups

of the Bcl-2 protein family, encompassing both proapoptotic and antiapoptotic proteins that regulate cellular sensitivity at the mitochondrial level. Similarly, the key members of the Bcl-2 family play a crucial role in how tumor cells respond to the various AgNPs formulations by which they are treated. For instance, Bax, a proapoptotic protein target, p53, can trigger apoptosis in cancer cells, while BCL2L1, an antiapoptotic protein, can inhibit both autophagy and apoptosis, contributing to cancer cell survival (Bao et al. 2022).

The mRNA expression of the proapoptotic *BAX* gene either treated with Met-AgNPs or Eth-AgNPs displayed significantly increased expression as compared to control ($p < 0.05$). However, the antiapoptotic *BCL2L1* gene mRNA expression of the MCF-7 cells either treated with Met-AgNPs or Eth-AgNPs showed downregulation as compared to the control. *TP53*, another tumor suppressor gene, is essential for regulating cell cycle arrest,

senescence, and apoptosis, helping to eliminate severely damaged cells. It directly impacts the intrinsic apoptosis pathway by controlling mitochondrial outer membrane permeabilization. It is activated through specific post-translational modifications and majorly determines the cell's fate such as whether to repair DNA, halt the cell cycle, undergo apoptosis, or enter senescence (Yuan et al. 2018). Similarly, our results suggest that the Met-AgNPs showed relatively higher expression of the tumor suppressor *TP53* gene as compared to Eth-AgNPs -treated MCF-7 cancer cells. *Cyclin D1*, an antiapoptotic gene as well as an important regulator of the cell cycle, is involved in promoting cancer by leading to uncontrolled cell growth. In normal cells, the expression of *Cyclin D1* is controlled, but, in cancer cells, there is often an increase in the activity of *Cyclin D1* due to different mechanisms (Montalto and De Amicis, 2020). Similarly, our results also showed overexpression of mRNA of the *CCND1*

gene treated with either Met-AgNPs or Eth-AgNPs as compared to control which aligns with the previously established literature (Bao et al. 2022).

Another crucial gene in the process of apoptosis is cytochrome c. It plays a pivotal role in the functioning of mitochondria and is essential for the intrinsic pathway of apoptosis. Cancer cells often impede the release of cytochrome c from the mitochondria, allowing them to evade programmed cell death (PCD). This ability to avoid apoptosis supports the growth and survival of tumors (Pessoa, 2022). In the present study, it was found that the *CYCS* gene was overexpressed when treated with either Met-AgNPs or Eth-AgNPs when compared with the control. However, it was observed that the expression of the *CCND1* gene in both Met-AgNPs and Eth-AgNPs was not statistically significant. Thus, the above findings suggest that the synthesized Met-AgNPs or Eth-AgNPs induce the upregulation of proapoptotic *TP53* and *BAX* genes, an extrinsic pathway, which promotes PCD of MCF-7 cancer cells.

Yuan et al. have treated human cervical cancer with AgNPs synthesized from sinigrin, a reducing and stabilizing agent, and observed upregulation of *TP53*, *CYCS*, *BAX*, and *BAK1* genes, suggesting that the synthesized AgNPs induced apoptosis, leading to higher inhibition of cancer cell growth (Yuan et al. 2018). Our result suggests that the synthesized Met-AgNPs or Eth-AgNPs, besides inducing intrinsic pathways, can also activate extrinsic pathways to promote apoptosis in MCF-7 cancer cell lines. *BAX* and *CYCS* gene expression were statistically significant in AgNP-treated groups, while *TP53* and *BCL2L1* gene expression showed no significant difference. Our data does not distinctly differentiate gene expression levels between the AgNP-treated groups. However, it indicates that the synthesized AgNPs were highly effective in inducing both extrinsic and intrinsic apoptosis pathways, significantly killing MCF-7 cancer cells, as confirmed by the above cytotoxic assay (Fig. 7). To strengthen the findings of apoptotic and anti-apoptotic gene expressions, we evaluated the *CASP3*, *CASP8*, and *CASP9* mRNA expression in MCF-7 cells (Fig. 10B). We observed that cells treated with Met-AgNPs or Eth-AgNPs showed comparatively higher expression of *CASP3* (executor caspase) compared to *CASP8* and *CASP9*. Our results suggest that the slightly higher expression of *CASP3* in Eth-AgNPs-treated cells is supported by higher expression of *TP53*, *CYCS*, and *BAX*, which are important downstream molecules that initiate the mitochondrial apoptotic pathway (Ullah et al. 2020). Thus, the findings correlate that the activated molecules induce apoptosis, leading to cell death. The observed increased mRNA expression of *CASP3* implies that Eth-AgNPs significantly activate the transcriptional

machinery involved in apoptosis. Moreover, *CASP8* expression in Eth-AgNPs-treated cells is notably higher. Thus, the findings suggest that Eth-AgNPs induces apoptosis through the intrinsic mitochondrial pathway, which is described by increased expression of *P53*, *BAX*, *CYCS*, and downstream molecule *CASP3* (Baharara et al., 2015). In succinct, the AgNPs synthesized from ethanolic extract promote apoptosis in MCF-7 cells, whereas Met-AgNPs had less effect on these apoptotic genes.

Flow cytometry analysis of apoptosis

The process of apoptosis plays a pivotal role in the homeostasis of cells. Many morphological and biochemical changes enable us to characterize this phenomenon. The biosynthesized Met-AgNPs and Eth-AgNPs induce apoptosis in MCF-7 cell death in a concentration-dependent manner. Thus, Annexin V/PI was utilized to comprehend the MCF-7 cell death mechanism, viz., apoptosis or non-apoptosis. The result suggests that Met-AgNPs-treated cells did not show a dose-dependent increase of cells in the late stage of apoptosis; however, an increased number of cells were observed in the initial phase of apoptosis from 5 to 25 µg/mL of treated dose. On the other hand, at similar doses, Eth-AgNPs showed a consistent increase in the number of cells in the early stage of apoptosis; however, at 25 µg/mL of dose, a significant increase of the late stage of MCF-7 cell apoptosis (8.14%) (Fig. 11). This result is in concordance with the cell proliferation assay.

Analysis of morphological change The morphological changes of the MCF-7 cells treated with different doses, either Met-AgNPs or Eth-AgNPs, were analyzed by inverted fluorescence microscopy. The treatment with 5, 10, and 25 µg/mL doses of Met-AgNPs to MCF-7 cells showed a dose-dependent morphological change, highlighted with red circles (Fig. 12). On the other hand, MCF-7 cells treated with Eth-AgNPs showed notable changes in morphology in a concentration-dependent manner. At 5 µg/mL, very few cells showed morphological changes, while at 10 µg/mL, a remarkable number of cells exhibit morphological changes, and only a few cells are visible in a normal morphology. However, at 25 µg/mL of dose, almost all cells showed morphological changes. The result is in concordance with the cell proliferation assay.

Conclusion

The organic extracts of nutraceutical *C. medica* fruits, viz., methanolic and ethanolic-based AgNPs, were synthesized to evaluate the influence of different organic solvent extracts on the physiochemical characteristics of the particles and their anticancer efficacy against breast cancer cells. The functional metabolites of the *C. medica* fruit




Fig. 11 Flow cytometry analysis of apoptosis induced by different concentrations of Met-AgNPs or Eth-AgNPs. **A** Unstrained cells, **B** control, **C** 5 $\mu\text{g}/\text{mL}$, **D** 10 $\mu\text{g}/\text{mL}$, **E** 25 $\mu\text{g}/\text{mL}$ Met-AgNPs, **E** 5 $\mu\text{g}/\text{mL}$, **F** 10 $\mu\text{g}/\text{mL}$, and **G** 25 $\mu\text{g}/\text{mL}$ Eth-AgNPs




Fig. 12 Evaluation of morphological changes of MCF-7 cells (10 $\times$) after treatment with different concentrations of AgNPs. **A** Untreated MCF-7 cells. **B** 5 $\mu\text{g}/\text{mL}$ Eth-AgNPs. **C** 10 $\mu\text{g}/\text{mL}$ Eth-AgNPs. **D** 25 $\mu\text{g}/\text{mL}$ Eth-AgNPs. **E** 5 $\mu\text{g}/\text{mL}$ Met-AgNPs. **F** 10 $\mu\text{g}/\text{mL}$ Met-AgNPs. **G** 25 $\mu\text{g}/\text{mL}$ Met-AgNPs

were involved in the formation of both AgNPs and have a polycrystalline nature. In addition, the size of Eth-AgNPs was smaller than that of Met-AgNPs. The Eth-AgNPs showed high efficacy in inducing apoptosis in MCF-7 cancer cell lines compared to Met-AgNPs. Eth-AgNPs showed extensive DNA fragmentation as compared to Met-AgNPs-treated cells. The proapoptotic genes (*TP53* and *BAX*) showed upregulation, while the antiapoptotic gene (*BCL1L2*) showed downregulation, suggesting that either Eth-AgNPs or Met-AgNPs activate the extrinsic pathway to induce apoptosis in the MCF-7 treated cancer cells. Moreover, these AgNPs also induce upregulation of mRNA expression of *CCND1* and *CYCS* genes. Thus, besides inducing extrinsic pathways, these AgNPs can activate the intrinsic pathway of apoptosis. Further, Eth-AgNPs treatment induces extensive apoptosis in MCF-7 cells, which was further corroborated by morphological analysis. Therefore, the synthesized AgNPs were effective in killing MCF-7 cancer cells. In succinct terms, ethanolic extract-based AgNPs are more effective in inducing apoptosis, which led to high efficacy in inhibiting/killing MCF-7 breast cancer cells. Thus, we can conclude that preparing AgNPs from the organic extract of either methanol or ethanol is necessary to evaluate their anticancer efficacy. The addition of water to the ethanol would prove to generate effective AgNPs to kill/inhibit a higher number of MCF-7 cells compared to methanolic extract-based AgNPs. Future studies are warranted to decipher the role of effective plant metabolite extraction methods in generating more efficient anticancer AgNP-based therapeutics for tumors.

Limitation of the study

The plant extract containing phytocompounds is a better reducing and stabilizing agent for the synthesis of AgNPs compared to the chemically synthesized AgNPs (Nie et al., 2023). However, several factors including pH, temperature, choice of reductant, capping agents, and zeta potential determine the stability of AgNPs. Therefore, a comprehensive stability assessment is required for the synthesized AgNPs (Met-AgNPs and Eth-AgNPs). Further, the toxicity of AgNPs was attributed to their size, concentration, shape, surface charge, and surface chemistry. Therefore, the synthesized AgNPs (Met-AgNPs and Eth-AgNPs) need to be inspected in vivo conditions for their toxicity in biological environments.

Abbreviations

AgNPs	Silver nanoparticles
Met-AgNPs	Methanolic extract-based AgNPs
Eth-AgNPs	Ethanolic extract-based AgNPs
FTIR	Fourier transform infrared
HR-TEM	High-resolution transmission electron microscopy
SAED	Selected area electron diffraction
XRD	X-ray diffraction

MCF-7	Michigan Cancer Foundation
SRB	Sulforhodamine B
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
DMSO	Dimethyl sulfoxide
PDI	Polydispersity index
CYCS	Cytochrome C
CCND1	Cyclin D1
PCD	Programmed cell death

Acknowledgements

Govind Ballabh Pant Institute of Postgraduate Medical Education and Research, New Delhi, India, is admired for the facilities provided for this study. This work was supported by the Department of Biotechnology, Ministry of Science and Technology, Government of India, with grant no. BT/INF/22/SP33063/2019.

Authors' contributions

Conceptualization, EA and SSS; methodology, EA, MS, AA, AK, and MZA; formal analysis and investigation, EA, AA, MS, AK, AKS, MZA, and N; original draft, EA, AA, and MS; revise the draft critically, EA, MS, MB, and ASA; and funding acquisition, resources, and supervision, SSS. All authors read and approved the final manuscript.

Funding

We are grateful to the Department of Biotechnology, Ministry of Science and Technology, India, for providing financial support under the Grant ID: BT/INF/22/SP33063/2019.

Data availability

Not applicable.

Declarations

Competing interests

The authors declare that they have no competing interests.

Author details

¹Central Molecular Laboratory, Govind Ballabh Pant Institute of Postgraduate Medical Education and Research (GIPMER), New Delhi, India. ²Molecular Biology Group, Division of Molecular Genetics & Biochemistry, ICMR-National Institute of Cancer Prevention & Research, Noida, India. ³Genome Biology Lab, Department of Biosciences, Jamia Millia Islamia, New Delhi, India. ⁴Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia. ⁵Department of GI Surgery, Govind Ballabh Pant Institute of Postgraduate Medical Education and Research (GIPMER), New Delhi, India.

Received: 20 January 2025 Accepted: 3 May 2025

Published online: 15 October 2025

References

- Ahmad E, Ali A, Fatima MT et al (2021) Ligand decorated biodegradable nanomedicine in the treatment of cancer. *Pharmacol Res* 167:105544. <https://doi.org/10.1016/j.phrs.2021.105544>
- Ahmad E, Athar A, Nimisha, et al (2024) Harnessing nature's potential: Alpinia galanga methanolic extract mediated green synthesis of silver nanoparticle, characterization and evaluation of anti-neoplastic activity. *Bioprocess Biosyst Eng* 47:1183–1196. <https://doi.org/10.1007/s00449-024-02993-7>
- Ahn E-Y, Jin H, Park Y (2019) Assessing the antioxidant, cytotoxic, apoptotic and wound healing properties of silver nanoparticles green-synthesized by plant extracts. *Mater Sci Eng C Mater Biol Appl* 101:204–216. <https://doi.org/10.1016/j.msec.2019.03.095>
- Alburae N, Alshamrani R, Mohammed AE (2024) Bioactive silver nanoparticles fabricated using *Lasiurus scindicus* and *Panicum turgidum* seed extracts: anticancer and antibacterial efficiency. *Sci Rep* 14:4162. <https://doi.org/10.1038/s41598-024-54449-3>
- Alharbi NS, Alsubhi NS (2022) Green synthesis and anticancer activity of silver nanoparticles prepared using fruit extract of *Azadirachta indica*. *Journal*

- of Radiation Research and Applied Sciences 15:335–345. <https://doi.org/10.1016/j.jrras.2022.08.009>
- Antropova IG, Revina AA, Oo PM et al (2021) Synthesis of silver nanoparticles using reactive water-ethanol extracts from *Murraya paniculata*. *ACS Omega* 6:8313–8321. <https://doi.org/10.1021/acsomega.1c00019>
- Baharara J, Namvar F, Ramezani T, Mousavi M (2015) Mohamad R (2015) Silver nanoparticles biosynthesized using *Achillea biebersteinii* flower extract: apoptosis induction in MCF-7 cells via caspase activation and regulation of Bax and Bcl-2 gene expression. *Molecules* 20(2):2693–2706
- Bansal R, Pathak R, Jha D et al (2015) Enhanced antimicrobial activity of amine-phosphonium (N-P) hybrid polymers against gram-negative and gram-positive bacteria. *Int J Polym Mater Polym Biomater* 64:84–89. <https://doi.org/10.1080/00914037.2014.886246>
- Bao J, Jiang Z, Ding W et al (2022) Silver nanoparticles induce mitochondria-dependent apoptosis and late non-canonical autophagy in HT-29 colon cancer cells. *Nanotechnol Rev* 11:1911–1926. <https://doi.org/10.1515/ntrev-2022-0114>
- Bawazeer S, Rauf A, Shah SUA et al (2021) Green synthesis of silver nanoparticles using *Tropaeolum majus*: phytochemical screening and antibacterial studies. *Green Processing and Synthesis* 10:85–94. <https://doi.org/10.1515/gps-2021-0003>
- Benedetto N, Carlucci V, Faraone I et al (2023) An insight into *Citrus medica* Linn.: a systematic review on phytochemical profile and biological activities. *Plants* 12(12):2267. <https://doi.org/10.3390/plants12122267>
- Bhakya S, Muthukrishnan S, Sukumaran M, Muthukumar M (2016) Biogenic synthesis of silver nanoparticles and their antioxidant and antibacterial activity. *Appl Nanosci* 6:755–766. <https://doi.org/10.1007/s13204-015-0473-z>
- Bilal M, Khan S, Ali J et al (2019) Biosynthesized silver supported catalysts for disinfection of *Escherichia coli* and organic pollutant from drinking water. *J Mol Liq* 281:295–306. <https://doi.org/10.1016/j.molliq.2019.02.087>
- Bitwell C, Indra SS, Luke C, Kakoma MK (2023) A review of modern and conventional extraction techniques and their applications for extracting phytochemicals from plants. *Scientific African* 19:e01585. <https://doi.org/10.1016/j.sciaf.2023.e01585>
- Bray F, Laversanne M, Sung H et al (2024) Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 74:229–263. <https://doi.org/10.3322/caac.21834>
- Chitsazi MR, Korbekandi H, Asghari G et al (2016) Synthesis of silver nanoparticles using methanol and dichloromethane extracts of *Pulicaria gnaphalodes* (Vent.) Boiss. aerial parts. *Artif Cells Nanomed Biotechnol* 44:328–333. <https://doi.org/10.3109/21691401.2014.949726>
- Chouhan N (2018) Silver nanoparticles: synthesis, characterization and applications. In: Maaz K (ed) *Silver Nanoparticles - Fabrication, Characterization and Applications*. InTech1.
- Das S, Das J, Samadder A et al (2013) Biosynthesized silver nanoparticles by ethanolic extracts of *Phytolacca decandra*, *Gelsemium sempervirens*, *Hydrastis canadensis* and *Thuja occidentalis* induce differential cytotoxicity through G2/M arrest in A375 cells. *Colloids Surf B Biointerfaces* 101:325–336. <https://doi.org/10.1016/j.colsurfb.2012.07.008>
- Davidson A (2006) *The Oxford companion to food*. Oxford University Press
- Dehghanizadeh S, Arasteh J, Mirzaie A (2018) Green synthesis of silver nanoparticles using *Anthemis atropata* extract: characterization and in vitro biological activities. *Artif Cells Nanomed Biotechnol* 46:160–168. <https://doi.org/10.1080/21691401.2017.1304402>
- Heidari Z, Salehzadeh A, Sadat Shandiz SA, Tajdoust S (2018) Anti-cancer and anti-oxidant properties of ethanolic leaf extract of *Thymus vulgaris* and its bio-functionalized silver nanoparticles. *3 Biotech* 8(3):177. <https://doi.org/10.1007/s13205-018-1199-x>
- Hemlata MPR, Singh AP, Tejvath KK (2020) Biosynthesis of silver nanoparticles using *Cucumis prophetarum* aqueous leaf extract and their antibacterial and antiproliferative activity against cancer cell lines. *ACS Omega* 5:5520–5528. <https://doi.org/10.1021/acsomega.0c00155>
- Jain N, Jain P, Rajput D, Patil UK (2021) Green synthesized plant-based silver nanoparticles: therapeutic prospective for anticancer and antiviral activity. *Micro and Nano Syst Lett* 9:5. <https://doi.org/10.1186/s40486-021-00131-6>
- Jiang Y, Jiang Z, Wang M, Ma L (2022) Current understandings and clinical translation of nanomedicines for breast cancer therapy. *Adv Drug Deliv Rev* 180:114034. <https://doi.org/10.1016/j.addr.2021.114034>
- Kanniah P, Balakrishnan S, Subramanian ER et al (2023) Preliminary investigation on the impact of engineered PVP-capped and uncapped silver nanoparticles on *Eudrilus eugeniae*, a terrestrial ecosystem model. *Environ Sci Pollut Res Int* 30:25239–25255. <https://doi.org/10.1007/s11356-022-21898-0>
- Khan ZUH, Khan A, Chen Y et al (2017) Biomedical applications of green synthesized noble metal nanoparticles. *J Photochem Photobiol B, Biol* 173:150–164. <https://doi.org/10.1016/j.jphotobiol.2017.05.034>
- Kharisova OV, Kharisov BI, Oliva González CM et al (2019) Greener synthesis of chemical compounds and materials. *R Soc Open Sci* 6:191378. <https://doi.org/10.1098/rsos.191378>
- Kis B, Moacă E-A, Tudoran LB et al (2022) Green synthesis of silver nanoparticles using *Populi gemmae* extract: preparation, physicochemical characterization, antimicrobial potential and in vitro antiproliferative assessment. *Materials (Basel)* 15(14):5006. <https://doi.org/10.3390/ma15145006>
- Mahmoodi Esfanddarani H, Abbasi Kajani A, Bordbar A-K (2018) Green synthesis of silver nanoparticles using flower extract of *Malva sylvestris* and investigation of their antibacterial activity. *IET Nanobiotechnol* 12:412–416. <https://doi.org/10.1049/iet-nbt.2017.0166>
- Mahmoud A, Mohamed B, Ibraheem I et al (2020) Characterization of *Ulva fasciata* ethanolic extract-mediated biosynthesized silver nanoparticles and evaluation of their nephroprotective effects in Doxorubicin-injected Wistar rats | Semantic Scholar. *Adv Animal Vet Sci* 8(s2):98–111
- Masum MMI, Siddiqua MM, Ali KA et al (2019) Biogenic synthesis of silver nanoparticles using *Phyllanthus emblica* fruit extract and its inhibitory action against the pathogen *Acidovorax oryzae* strain RS-2 of rice bacterial brown stripe. *Front Microbiol* 10:820. <https://doi.org/10.3389/fmicb.2019.00820>
- Meng Y (2015) A sustainable approach to fabricating Ag nanoparticles/pVA hybrid nanofiber and its catalytic activity. *Nanomaterials (Basel)* 5:1124–1135. <https://doi.org/10.3390/nano5021124>
- Montalto FI, De Amicis F (2020) Cyclin D1 in Cancer: A Molecular Connection for Cell Cycle Control, Adhesion and Invasion in Tumor and Stroma. *Cells* 9(12):2648. <https://doi.org/10.3390/cells9122648>
- Muthukrishnan S, Vellingiri B, Murugesan G (2018) Anticancer effects of silver nanoparticles encapsulated by *Gloriosa superba* (L.) leaf extracts in DLA tumor cells. *Future Journal of Pharmaceutical Sciences* 4:206–214. <https://doi.org/10.1016/j.fjps.2018.06.001>
- Nahvi I, Belkahl S, Biswas S, Chakraborty S (2022) A review on nanocarrier mediated treatment and management of triple negative breast cancer: a Saudi Arabian scenario. *Front Oncol* 12:953865. <https://doi.org/10.3389/fonc.2022.953865>
- Nawaz H, Shad MA, Rehman N et al (2020) Effect of solvent polarity on extraction yield and antioxidant properties of phytochemicals from bean (*Phaseolus vulgaris*) seeds. *Braz J Pharm Sci* 56:e17129. <https://doi.org/10.1590/s2175-97902019000417129>
- Nie P, Zhao Y, Xu H (2023) Synthesis, applications, toxicity, and toxicity mechanisms of silver nanoparticles: a review. *Ecotoxicol Environ Saf* 15(253):114636. <https://doi.org/10.1016/j.ecoenv.2023.114636>
- Odularu AT (2018) Metal nanoparticles: thermal decomposition, biomedical applications to cancer treatment, and future perspectives. *Bioinorg Chem Appl* 2018:9354708. <https://doi.org/10.1155/2018/9354708>
- Ojha S, Sett A, Bora U (2017) Green synthesis of silver nanoparticles by *Ricinus communis* var. *Carmencita* leaf extract and its antibacterial study. *Adv Nat Sci: Nanosci Nanotechnol* 8:035009. <https://doi.org/10.1088/2043-6254/aa724b>
- Pessoa J (2022) Cytochrome c in cancer therapy and prognosis. *Biosci Rep* 42:BSR20222171. <https://doi.org/10.1042/BSR20222171>
- Qais FA, Shafiq A, Khan HM et al (2019) Antibacterial effect of silver nanoparticles synthesized using *Murraya koenigii* (L.) against multidrug-resistant pathogens. *Bioinorg Chem Appl* 2019:4649506. <https://doi.org/10.1155/2019/4649506>
- Ragunandan D, Bedre MD, Basavaraja S et al (2010) Rapid biosynthesis of irregular shaped gold nanoparticles from macerated aqueous extracellular dried clove buds (*Syzygium aromaticum*) solution. *Colloids Surf B Biointerfaces* 79:235–240. <https://doi.org/10.1016/j.colsurfb.2010.04.003>
- Rahman M, Afzal O, Ullah SNMN et al (2023) Nanomedicine-based drug-targeting in breast cancer: pharmacokinetics, clinical progress, and challenges. *ACS Omega* 8:48625–48649. <https://doi.org/10.1021/acsomega.3c07345>

- Rani P, Kumar V, Singh PP et al (2020) Highly stable AgNPs prepared via a novel green approach for catalytic and photocatalytic removal of biological and non-biological pollutants. *Environ Int* 2020;143:105924. <https://doi.org/10.1016/j.envint.2020.105924>
- Ratan ZA, Haidere MF, Nurunnabi M et al (2020) Green chemistry synthesis of silver nanoparticles and their potential anticancer effects. *Cancers (Basel)* 12(4):855. <https://doi.org/10.3390/cancers12040855>
- Roy A, Bulut O, Some S et al (2019) Green synthesis of silver nanoparticles: biomolecule-nanoparticle organizations targeting antimicrobial activity. *RSC Adv* 9:2673–2702. <https://doi.org/10.1039/C8RA08982E>
- Samuggam S, Chinni SV, Mutusamy P et al (2021) Green synthesis and characterization of silver nanoparticles using Spondias mombin extract and their antimicrobial activity against biofilm-producing bacteria. *Molecules* 26(9):2681. <https://doi.org/10.3390/molecules26092681>
- Sharma VK, Yngard RA, Lin Y (2009) Silver nanoparticles: green synthesis and their antimicrobial activities. *Adv Colloid Interface Sci* 145:83–96. <https://doi.org/10.1016/j.cis.2008.09.002>
- Sharma A, Goyal AK, Rath G (2018) Recent advances in metal nanoparticles in cancer therapy. *J Drug Target* 26:617–632. <https://doi.org/10.1080/1061186X.2017.1400553>
- Singh R, Navneet (2021) Green synthesis of silver nanoparticles using methanol extract of Ipomoea carnea Jacq. to combat multidrug resistance bacterial pathogens. *Curr Res Green Sustainable Chem* 4:100152
- Soshnikova V, Kim YJ, Singh P et al (2018) Cardamom fruits as a green resource for facile synthesis of gold and silver nanoparticles and their biological applications. *Artif Cells Nanomed Biotechnol* 46:108–117. <https://doi.org/10.1080/21691401.2017.1296849>
- Sung H, Ferlay J, Siegel RL et al (2021) Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 71:209–249. <https://doi.org/10.3322/caac.21660>
- Tagde P, Najda A, Nagpal K et al (2022) Nanomedicine-based delivery strategies for breast cancer treatment and management. *Int J Mol Sci* 23(5):2856. <https://doi.org/10.3390/ijms23052856>
- Ullah I, Khalil AT, Ali M, Iqbal J, Ali W, Alarifi S, Shinwari ZK (2020) Green-synthesized silver nanoparticles induced apoptotic cell death in MCF-7 breast cancer cells by generating reactive oxygen species and activating caspase 3 and 9 enzyme activities. *Oxid Med Cell Longev* 2020:1215395
- Vanlalveni C, Lallianrawna S, Biswas A et al (2021) Green synthesis of silver nanoparticles using plant extracts and their antimicrobial activities: a review of recent literature. *RSC Adv* 11:2804–2837. <https://doi.org/10.1039/d0ra09941d>
- Vichai V, Kirtikara K (2006) Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nat Protoc* 1:1112–1116. <https://doi.org/10.1038/nprot.2006.179>
- Wu D, Si M, Xue H-Y, Wong H-L (2017) Nanomedicine applications in the treatment of breast cancer: current state of the art. *Int J Nanomedicine* 12:5879–5892. <https://doi.org/10.2147/IJN.S123437>
- Yap YH, Azmi AA, Mohd NK et al (2020) Green synthesis of silver nanoparticle using water extract of onion peel and application in the acetylation reaction. *Arab J Sci Eng* 45:4797–4807. <https://doi.org/10.1007/s13369-020-04595-3>
- Yuan Y-G, Zhang S, Hwang J-Y, Kong I-K (2018) Silver nanoparticles potentiates cytotoxicity and apoptotic potential of camptothecin in human cervical cancer cells. *Oxid Med Cell Longev* 2018:6121328. <https://doi.org/10.1155/2018/6121328>
- Zhang X-F, Liu Z-G, Shen W, Gurunathan S (2016) Silver nanoparticles: synthesis, characterization, properties, applications, and therapeutic approaches. *Int J Mol Sci* 17(9):1534. <https://doi.org/10.3390/ijms17091534>
- Zia Q, Mohammad O, Rauf MA et al (2017) Biomimetically engineered amphotericin B nano-aggregates circumvent toxicity constraints and treat systemic fungal infection in experimental animals. *Sci Rep* 7:11873. <https://doi.org/10.1038/s41598-017-11847-0>
- Zuhrotun A, Oktaviani DJ, Hasanah AN (2023) Biosynthesis of gold and silver nanoparticles using phytochemical compounds. *Molecules* 28(7):3240. <https://doi.org/10.3390/molecules28073240>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.