



Arbuscular mycorrhizal fungi affect early phenological stages of three secondary vegetation species in a temperate forest

Yasmin Vázquez-Santos^{1,2} · Silvia Castillo-Argüero² · Noé Manuel Montaña³ ·
Francisco Javier Espinosa-García⁴ · César Flores-Ortiz⁵ · Yuriana Martínez-Orea²

Received: 12 April 2024 / Accepted: 12 July 2024 / Published online: 22 July 2024
© The Author(s) 2024

Abstract

The relationship between arbuscular mycorrhizal fungi (AMF) and secondary vegetation (SV) species at early phenological stages is critical for the successful establishment of these plants on disturbance sites in temperate forests. The main objective of this research is to evaluate the effect of AMF colonization on the early phenological stages (germination and early growth) of three shrub species present in the SV of a temperate forest in central Mexico. We collected soil from different sites in the *Abies religiosa* forest in central Mexico. We collected seeds of *Acaena elongata*, *Ageratina glabrata*, and *Solanum pubigerum*. We used a controlled experimental design with pasteurized soil (-AMF treatments) and unpasteurized soil (+AMF treatments). We monitored germination percentage, growth (shoot and root weight and total biomass), AMF root colonization, and the mycorrhizal response index (MRI) for each plant species. All three species tested benefited by AMF, showing higher germination rates. Shoot and root weight and total biomass were significantly higher in the +AMF treatment. *Solanum pubigerum* showed greater stem length and *Ageratina glabrata* showed greater root development due to AMF. *Ageratina glabrata* and *Acaena elongata* were the most responsive to AMF as indicated by MRI. This research underscores the critical role of AMF in the early phenological stages of SV and highlights the potential ecological benefits of AMF in supporting plant germination and plant growth. These results suggest that AMF enhance germination and early growth of secondary vegetation species, which can be considered in management plans for forest ecosystems.

Keywords Dependence mycorrhizal · Germination · Growth · Phenology · Secondary forest · Symbiosis

Communicated by Courtney Siebert.

✉ Yasmin Vázquez-Santos
yasminvazquez@ciencias.unam.mx

- ¹ Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México. Unidad de Posgrado, Circuito de Posgrados, Coyoacán, 04510 Mexico City, Mexico
- ² Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Universidad Nacional Autónoma de México. Investigación Científica, Coyoacán, 04510 Mexico City, Mexico
- ³ Departamento de Biología, División de Ciencias Biológicas y de La Salud, Universidad Autónoma Metropolitana Unidad Iztapalapa. Av. Ferrocarril de San Rafael Atlixco 186, Leyes de Reforma 1ª Sección, 09310 Mexico City, Mexico
- ⁴ Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Antigua Carretera a Patzcuaro, 8701 Morelia, Michoacán, Mexico
- ⁵ Facultad de Estudios Superiores Iztacáala, 54090 Tlalnepantla, Mexico

Introduction

Temperate forests are the second most widespread ecosystem in North America, Central America, Northeast Asia, Western Europe and Central Europe, accounting for 16% of the global forest area (665.8 million ha) (Gilliam 2016; FAO 2020). These forests face conservation challenges and have experienced significant degradation due to human activities, extreme climate events, fires, pests and diseases (Rawat et al. 2022; Lin et al. 2024). Consequently, there has been a notable increase in the expansion of secondary temperate forests worldwide (FAO 2020).

Disturbance alters the structure of plant communities, leading to the establishment of secondary vegetation (SV) species, which play a crucial role in increasing the resilience and functionality of temperate forests by mitigating the loss of ecological functions, interactions and processes (van Hall et al. 2017). Structurally, SV species contributes to the forest fragment connectivity (Sayer et al. 2004), and significantly

influences soil formation and structure (Álvarez-Yépiz et al. 2008; van Hall et al. 2017). In addition, SV species actively participate in nutrient cycling and maintain critical biotic interactions through its root system (Post and Kwon 2000; Lesschen et al. 2008; van Hall et al. 2017).

A way to satisfy the low nutrient availability in temperate forest soils, plants are associated with soil microorganisms, such as arbuscular mycorrhizal fungi (AMF). These fungi are obligate symbionts forming arbuscular mycorrhizae with *ca.* 75% of plant families (Brundrett and Tedersoo 2018), and facilitate the efficient transfer of nutrients and water (Smith and Smith 2011). This symbiotic relationship enables host plants to withstand both biotic and abiotic stresses, while gaining advantages in germination (Kumar et al. 2023) and growth in their host plants (Smith and Smith 2012; Cavagnaro et al. 2015). On the other hand, SV species mitigate the loss of taxonomic and functional diversity of AMF by acting as temporary hosts for these fungi until mature forest species colonize the area (Guadarrama 2014).

Plant species of SV exhibit highly competitive ability, rapid germination and growth rates, and a tendency to rapidly colonize disturbed sites (Leck et al. 2008). These traits suggest a significant investment of resources by SV plants, including nutrients, water, and light. Thus, the association of SV species with AMF may serve as a buffer against the stressful environmental conditions they encounter at different life cycle stages in a changing environment due to anthropogenic disturbance.

The AMF are known to favor the germination and growth of their host plants (Sortibrán et al. 2019; Souza et al. 2023; Tian et al. 2023; Yang et al. 2024). The nutrient requirements of plants differ at early phenological stages due to the adaptive dynamics of their physiological and metabolic processes (Milla et al. 2009). This adaptation of nutrient requirements reflects the specific response of plants to changing demands during their germination and early growth phases.

The early phenological stages are critical for plant establishment as they are the periods during which seedlings overcome their vulnerability to abiotic and biotic stresses. This ensures the proper formation of tissues that will support critical processes such as photosynthesis and nutrient uptake (Ismail et al. 2012; Gioria et al. 2018). Recent studies have shown that AMF promote seed germination and seedling growth in the presence of abiotic stresses, such as drought and heavy metal toxicity (Van Der Heijden 2004; Soto-Cruz et al. 2021), and can even modulate plant-parasite interactions to the benefit of the host plant (Maighal et al. 2016).

The role of AMF in enhancing seed germination is of great importance (Kumar et al. 2023) and may have implications for the tolerance and establishment of plant species in disturbance sites. It has been proposed that AMF

may contribute to the establishment of favorable microsites for seed germination through the formation of micro- and macroaggregates in the soil and the production of hyphal exudates that can scarify seeds (Koide 2000). However, the specific physical or chemical mechanisms underlying this enhanced germination remain largely unknown. It is also likely that the interacting plant and fungal species also regulate the effect of AMF on seed germination.

In contrast, during early growth, resource allocation shifts to biomass formation and accumulation, a critical process for tissue development and leaf growth (Lambers et al. 2019). This shift implies different requirements for essential nutrients such as nitrogen (N) and phosphorus (P) for optimal growth, both of which are critical for plant tissue formation. Ågren (2008) showed that nutrient concentrations in the tissues of *Lycopersicum esculentum*, *Betula pendula* and *Selenastrum minutum* vary with time. These variations are related to the plants relative growth rates and affect the stoichiometry of key elements such as carbon (C), N, and P (C:N:P). In particular, N concentrations increase more slowly than P concentrations. In this context, AMF can be an important option to meet the nutrient requirements of plants at each stage (Mosse and Hayman 1971; van der Heijden 2004). Thus, the shift in phenological phase and nutrient demand of the host plant also suggests a functional shift in the mycorrhizal association. During the early growth phase, the primary function of the AMF is nutrient and water transport, which is critical for both aboveground and belowground biomass plant production (Smith and Smith 2011; Chongtham and Pandey 2020; Zhou et al. 2022).

Evolutionarily, some plant species are classified as ‘obligate or facultative mycotrophic’, while others fall into the ‘non-mycotrophic’ category (Li et al. 2016; Wang et al. 2021). AMF colonization shows variability among plant species, with four possible responses: a) no colonization and no growth response, b) no colonization and reduced growth, c) colonization with growth stimulation, and d) colonization with increased P concentration without changes in biomass (Helgason et al. 2002). Responses c and d do not fully capture changes in colonization throughout plant phenology. Investigating these variations may provide a more complete understanding of the dynamics of plant-AMF associations and their effects on different plant phenological phases.

The outcome of this association depends on factors such as the identity of the host plant and fungus, interactions with other beneficial organisms, and the level of abiotic stress (Baslam et al. 2011; Copetta et al. 2011; Smith et al. 2011; Baslam and Goicoechea 2012; Carillo et al. 2020). For example, low P and high N availability in the soil have been observed to promote AM association, whereas in soils with high availability of both nutrients, this association may become parasitic or even inhibited (Johnson 2010). In addition, under limited light conditions, AMF may have an

antagonistic effect on the plant due to their ability to extract carbon from the plant. Therefore, changes in the cost–benefit balance of the mycorrhizal association occur in a continuous mutualism–antagonism (Johnson 2010).

Current knowledge on changes in intraradical colonization during early phenological stages in temperate forest species is still scarce. Researchers often focus on the effect of AMF on later stages of plant growth. In turn, our study aims to contribute to the understanding of this interaction, particularly in temperate forest SV plant species. The hypothesis was that if the nutritional requirements of host plants vary according to their phenological stage, and this will affect arbuscular mycorrhizal colonization, then we could observe different degrees of colonization that vary interspecifically during each phenological stage of the host plant. The main objective of this research is to evaluate the effect of AMF colonization on the early phenological stages (germination and early growth) of three shrub species present in the SV of a temperate forest in central Mexico. The study of arbuscular mycorrhizal association in early phenological stages of SV species will allow a better understanding of the initial establishment and long-term effects of this symbiosis on plant development.

Material and methods

Study area

This research was carried out in the temperate forest dominated by *Abies religiosa* (H.B.K.) Schl. & Cham. (Fir) in the Magdalena river basin (MRB), located in Mexico City (Fig. 1). The climate is temperate subhumid, characterized by summer rains, with an average annual temperature of 18 °C, an average annual rainfall of 1 250 mm, and a minimal thermal oscillation, typically below 5 °C (García 1988). The forest has a strong seasonality, with a rainy season ranging from May to October and a dry season from November to April (Delgadillo-Duran 2011). The predominant soil group is classified as humic Andosol, according to the WRB (2022). This soil is characterized by its acidity, abundant presence of allophane, high organic matter content and remarkable ability to adsorb phosphates on the surface of amorphous mineral clays. This phosphate fixation, documented by Galicia et al. (2016) and INEGI (2011), may limit P availability for plants.




Fig. 1 Geographical location of the *Abies religiosa* forest within the Magdalena River Basin, Mexico City, Mexico showcasing **A)** Mexico, **B)** Mexico City, **C)** the Magdalena River Basin, with the *Abies*

religiosa Forest area highlighted in brown. Prepared by Romero-Romero M.A. (2024)

Study species

In this study, the effect of arbuscular mycorrhizal association on germination and growth of three SV species was addressed. The selected plant species are common shrubs in the SV of the study forest (Santibañez-Andrade et al. 2015): *Acaena elongata* L., *Ageratina glabrata* (Kunth) R.M. King & H. Rob., and *Solanum pubigerum* Dunal. These plants thrive in the understory and contribute significantly to natural regeneration through sources such as seed rain and the seed bank (Argüelles-Mayao 2018). Variations in flower and fruit types, seed size, and reproductive phenology patterns are observed among the three plant species.

Soil collection and processing

From July 2019 to January 2020, 56 kg of soil was collected randomly in the *A. religiosa* forest of the MRB. The soil was collected at 20 cm of depth, previously removing the litter. It was homogenized and divided into two equal parts: the first part was used as inoculum for the “+ AMF” treatments, assuming that sufficient AMF propagules were present, such as spores (70 spores / 50 g soil), fragments of colonized roots and mycelium. The second part was subjected to a pasteurization process; this part of the soil was used for the “-AMF” treatments, where the AMF propagules were eliminated. To pasteurize the soil, we used a pressure cooker heated to 55 kPa at approximately 90 °C (194°F). We maintained these conditions for 45 min before allowing the cooker to cool. This method was repeated on the same soil sample for three consecutive days at a constant temperature.

Inoculum description

Vázquez-Santos et al. (2019) characterized the species richness and spore abundance of arbuscular mycorrhizal fungi (AMF) in 80 soil samples. The study reported 70 spores per 50 g of soil. These samples were collected from established plots within the *A. religiosa* forest in study. Analysis revealed a diverse array of 29 morphospecies representing nine families and ten genera. Notably, the genus *Acaulospora* dominated the AMF community, accounting for 77.7% of the total spores recorded. This was closely followed by *Ambispora* with 9.07%, *Claroideoglomus* with 4.96%, *Funnelformis* with 3.8%, while other genera such as *Archaeospora*, *Diversispora*, *Glomus*, *Rhizophagus*, *Sclerocystis* and *Scutellospora* accounted for less than 2.5% each.

Experiment set up: germination

From 50 individuals we randomly collected ripe fruits directly. For each study species, we had a collection of

eight hundred seeds. The seeds were homogeneously mixed and a subsample of 160 mature seeds was extracted. General characterization (length, width, and weight) was performed. Subsequently, 80 seeds per species were disinfected with 10% of an initial concentration of 3.5% sodium hypochlorite for 10 min and washed with distilled water. The seeds were germinated in pots containing 250 g of soil, with pasteurized soil (– AMF) and unpasteurized soil (+ AMF). We set up a greenhouse (4 × 6 m) in March 2021, where we installed two HOBO DATA Loggers (easy LogUSB-ONSET, Massachusetts, USA) to record the variation in environmental temperature and moisture during the experimental period.

The pots were watered daily at 10:00 a.m. to field capacity while seed germination was monitored by checking for signs of radicle emergence, and once the germination process had begun, radicle or root fragments were made, with a minimum of 5 individuals and a maximum of 8 individuals per extraction. The radicles and/or roots were processed according to Koske and Gemma’s method (1989), in order to obtain the mycorrhizal colonization percentage and the fungal structures characteristics of AM according to the method of McGonigle et al. (1990). Seedling collection was completed when cotyledon emergence was observed. Therefore, the remaining seedlings are destined for the next phase.

Experiment set up: growth

After the previous experiment, another 90 seeds of each species were set to germinate, in pots with a capacity of 250 g, in pasteurized soil (– AMF) and unpasteurized soil (+ AMF). The total number of pots was 360 (12 individuals × 3 species × 2 mycorrhization treatments × 5 crops). The pots were distributed in species × treatment blocks design.

Measurements started in August 2021. We worked with a minimum of 30 individuals/species/treatment, with a total of 180 seedlings (6 individuals × 3 species × 2 treatments × 5 harvests). Five harvests were made each month, with an extraction of six individuals per species and treatment, to determine plant height, root length, stem length, fresh root weight and fresh shoot weight. The seedlings were then dried in a Riossa ® oven at 72 °C for 24 h (until constant weight) and the dry weight of the root and aerial part was determined using an OHAUS ® analytical balance. The fresh weight and the dry weight of the root and stem were combined to give the total dry weight (TDW). These dry tissue samples were used for chemical analysis of the phosphorus and nitrogen content of the plant tissue for each harvest.

Arbuscular mycorrhizal fungi roots colonization

Simultaneously with the seed extraction for growth measurements, we carried out five root extractions from six individuals/species/treatment (6 individuals $\times$ 3 species $\times$ 2 treatments $\times$ 5 harvests), during the same months, to determine the AMF colonization percentage. We extracted the finest roots to process them according to Koske and Gemma (1989) and to obtain the mycorrhizal colonization percentage and per AM structures according to McGonigle et al. (1990).

The effect of mycorrhizal fungi on plants TDW was estimated using the mycorrhization response index (MRI). Treatments were compared where the only difference was the presence of mycorrhizae. This index ranges from $-\infty$ to 100% (Plenchette et al. 1983), and expresses the difference in dry weight between mycorrhizal and non-mycorrhizal plants as a percentage of the dry mass of the mycorrhizal plant.

$$\text{MRI} = \left(\frac{\text{TDWm}(g) - \text{TDWnm}(g)}{\text{TDWm}} \right) \times 100 \quad (1)$$

where: TDWm = total dry weight of the mycorrhizal plant and TDWnm = total dry weight of the non-mycorrhizal plant.

Data analysis

We used a simple random design with a 3×2 factorial arrangement (three species, two mycorrhization treatments), with five replicates. The accumulated germination percentages for each treatment were calculated using the following equation:

$$\text{Germination (\%)} = \left(\frac{\text{Germinated seeds}}{\text{Seeds total}} \right) (100) \quad (2)$$

Two-way analyzes of variance (Two-way ANOVA) was used to determine the effects of species (*Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum*), mycorrhization (+ AMF and – AMF) and the interaction between these factors on germination, growth, phosphorus and nitrogen contents of the plants tissue and arbuscular mycorrhizal colonization.

To meet the assumptions of normality and homogeneity of variances, the cumulative germination percentages were transformed using Eq. 3 (Zar 1974). Similarly, the first day of germination, the maximum day of germination, the average germination time and seed vigor were transformed using the natural logarithm (ln).

$$\left(\arcsin \sqrt{\frac{\text{Germination \%}}{100}} \right) \quad (3)$$

Growth parameters such as root and shoot length, root and shoot dry weight, total dry weight, and root/shoot ratio were transformed with the natural logarithm to meet the assumptions of normality and homogeneity of variances. On the other hand, total colonization percentages by hyphae, arbuscules and spores were transformed with the arcsine function (Eq. 3). For all the parameters evaluated, a multiple comparison of means was performed using Tukey tests, with a significance level of $p < 0.05$. All statistical analyzes were performed with the RStudio software.

Results

Germination

The two-way ANOVA showed significant differences between treatments with mycorrhiza (+ AMF) and without mycorrhiza (– AMF) for the three species studied (Fig. 2). The + AMF treatments showed a higher value of accumulated germination percentages compared to the – AMF treatments.

Mycorrhiza had a positive effect on the accumulated germination values of the species over time. We observed that this value was significantly higher in all species with the (+ AMF) treatment compared to the treatment without mycorrhiza (– AMF) (Fig. 3).

Our results show that the first day of germination, maximum day of germination, and the average germination time differed significantly between the mycorrhization treatments (+ AMF, – AMF) (Table 1). On the other




Fig. 2 Mean value and standard deviation bars ($\pm$ S.D.) of germination percentages for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum* seeds treated with mycorrhiza (+ AMF) and without mycorrhiza (– AMF). Significant differences are indicated by letters ($p < 0.001$)




Fig. 3 Mean values of accumulated germination percentages and standard deviation bars ($\bar{x} \pm \text{S.D.}$) for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum* in two treatments: + AMF (with mycorrhiza) and – AMF (without mycorrhiza), over time in days

Table 1 Statistical summary of the two-way ANOVA for germination percentage=germination, day of first germination=First germination, day of maximum germination=Max germination, average germination time=T50 and seed vigor, for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum*, for treatments with mycorrhiza (+ AMF) and without mycorrhiza (– AMF)

Sources of variation	df	Germination		First germination		Max germination		T50		Vigor	
		F	P	F	P	F	P	F	P	F	P
Species	2	4.972	0.001	4.093	0.029	1.648	0.21	0.207	0.81	39.32	0.001
AMF	1	59.93	0.001	14.24	0.001	9.905	0.004	32.28	0.001	0.063	0.93
Species × AMF	2	0.741	0.487	2.47	0.105	2.13	0.14	4.6	0.0203	0.277	0.75

*Significant differences in bold

hand, differences in seed vigor were observed between species. No differences were observed between treatments (+ AMF, – AMF).

We observed that *Acaena elongata* and *Solanum pubigerum* seeds with mycorrhiza (+ AMF) germinated earlier than seeds without it (– AMF). However, *Ageratina glabrata* seeds showed no significant differences in the first day of germination between the mycorrhiza treatments. The day of maximum germination was significantly earlier for *Ageratina glabrata* in the mycorrhiza treatment. Similarly, the mean germination time occurred earlier in

the + AMF treatment compared to the – AMF treatment, for *Ageratina glabrata* and *Solanum pubigerum* (Table 2).

Growth

According to two-way ANOVA, significant differences were found for shoot length, shoot dry weight, root dry weight, total dry weight and root/shoot ratio between plant species, mycorrhization and the interaction of these factors (Table 3).

Solanum pubigerum showed a higher stem length in the + AMF treatment, but not *Ageratina glabrata* and

Table 2 Means values ($\bar{x} \pm \text{S.D.}$) of germination traits for species and treatments with mycorrhiza (+ AMF) and without mycorrhiza (– AMF)

	<i>Acaena elongata</i>		<i>Ageratina glabrata</i>		<i>Solanum pubigerum</i>	
	+ AMF	– AMF	+ AMF	– AMF	+ AMF	– AMF
Firts germination day	8.4 ± 0.4 ^b	13.7 ± 0.44 ^a	8 ± 0.86 ^b	8.5 ± 0.44 ^a	8.25 ± 0.22 ^b	14 ± 3.07 ^a
Maximum germination day	35.4 ± 4.2 ^b	37.5 ± 2.04 ^a	26.5 ± 4.9 ^b	40 ± 1.9 ^a	31.25 ± 3.6 ^b	40.75 ± 1.1 ^a
T ₅₀	20.3 ± 2.6 ^b	26 ± 4.6 ^a	12.6 ± 1.2 ^b	39.5 ± 5 ^a	14.6 ± 1.2 ^b	32.37 ± 4.4 ^a
Vigor (mm)	14.28 ± 3.9 ^a	15.3 ± 5.0 ^a	5.46 ± 2.7 ^c	5.31 ± 1.6 ^c	11.92 ± 8.4 ^b	12.69 ± 9.2 ^b

*Significant differences are indicated by superscript letters

Table 3 Statistical summary of the two-way ANOVA for stem length, root length, SDW = stem dry weight, RDW = root dry weight, TDW = total dry weight and root/stem ratio for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum*. Var.exp = percentage variance explained

Source of variation	Stem length				Root length			SDW		
	df	F	P	Var. exp	F	P	Var. exp	F	P	Var. exp
Species	2	10.03	0.001	11.29	14.64	0.001	17.25	19.25	0.001	13.02
AMF	1	15.73	0.001	8.85	1.81	0.18	1.06	87.13	0.001	29.46
Species:AMF	2	4.44	0.013	4.99	2.83	0.062	3.33	18.51	0.001	12.52
Source of variation	RDW				TDW			Root/stem		
	df	F	P	Var. exp	F	P	Var. exp	F	P	Var. exp
Species	2	24.813	0.001	19.7	12.66	0.001	7.58	74.36	0.001	46.79
AMF	1	47.56	0.001	18.89	58.28	0.001	17.69	20.04	0.001	6.3
Species × AMF	2	10.8	0.001	8.58	12.36	0.001	7.51	8.03	0.001	5.05

*Significant differences are shown in bolds

Acaena elongata. Root length was greater in *Ageratina glabrata* in the + AMF treatment, but not in the other species. Shoot, root and total biomass were significantly higher in the + AMF treatments for *Acaena elongata* and *Ageratina glabrata*, but not for *Solanum pubigerum*. The root-shoot ratio was significantly higher in the -AMF treatments only for *Ageratina glabrata* and *Solanum pubigerum* (Fig. 4).

Growth-Nutrient content

Concentrations of nitrogen and phosphorus in plant tissues varied significantly among species, between aerial parts and roots, and among mycorrhization treatments. However, no significant differences were observed for the three-way interaction among species, plant part, and mycorrhization treatment, as reported in Table 4 for the species:AMF interaction.

Arbuscular mycorrhizal fungi (AMF) generally increase nutrient levels in plant tissues, although the extent of this increase varies among species. In *Acaena elongata*, the presence of AMF increased nitrogen and phosphorus levels in both shoot and root tissues. In *Ageratina glabrata*, AMF

treatments also increased nitrogen and phosphorus concentrations, with the most notable difference occurring in the roots. In the case of *Solanum pubigerum*, the AMF effect was more muted, but there was a noticeable increase in nitrogen and phosphorus levels in the aerial parts when AMF were present (Fig. 5).

Significant differences were observed in the MRI of the three species ($X^2 = 28.24$, $P < 0.001$) (Fig. 6). *Ageratina glabrata* showed the highest growth response to AMF colonization ($\bar{x} = 77 \pm \text{S.D.} = 13$), followed by *Acaena elongata* ($\bar{x} = 59 \pm 27$), and *Solanum pubigerum* ($\bar{x} = -14 \pm 41$) showed the lowest. However, the species with significant differences in MRI was *Solanum pubigerum*.

Arbuscular mycorrhizal fungi roots colonization

Significant differences between species and between mycorrhization treatments were observed for the total and hyphal colonization percentages (Table 5).

Acaena elongata and *Solanum pubigerum* showed the highest percentages of AMF colonization. *Ageratina glabrata* showed significantly lower percentages of total and




Fig. 4 Mean values of growth parameters and standard deviation bars ($\bar{x} \pm \text{S.D.}$) for the three species from secondary vegetation in a temperate forest under the treatments: with mycorrhiza (+AMF) and

without mycorrhiza (– AMF). Different letters indicated statistical differences between treatments $p < 0.05$

hyphal colonization (Fig. 7). Colonization by arbuscules and spores showed no differences between species.

Discussion


This study provides evidence of the positive effect of AMF on the germination and early growth of three characteristic SV plant species, which are considered as important in natural regeneration sources of this temperate forest: *Acaena*

elongata, *Ageratina glabrata*, and *Solanum puberum*. Germination and growth are key stages in a plant's life cycle that are strongly influenced by the symbiosis with AMF (Smith et al. 2011), as observed in this study. Knowledge of the role of AMF in the early phenological stages of SV plant species is crucial for understanding their ecological dynamics. These fungi are key players in enhancing plant establishment in disturbed ecosystems, contributing to ecosystem resilience and natural regeneration. This has

Table 4 Statistical summary of the two-way ANOVA for nitrogen and phosphorus content in plant tissue for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum* in +AMF and – AMF treatments


Source of variation	df	Nitrogen		Phosphorus	
		<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Species	2	3.89	<0.01	10.359	<0.001
Plant tissue	2	47.37	<0.0001	0.156	0.855
AMF	1	9.63	<0.001	10.483	<0.01
Species:Plant tissue	2	1.99	0.153	6.057	<0.01
Species:AMF	2	1.28	0.291	0.463	0.633
Plant tissue:AMF	1	5.12	<0.05	16.315	<0.001
Species:Plant tissue:AMF	2	3.26	0.052	0.044	0.957

*Significant differences are shown in bolds

**Fig. 5** Mean of Nitrogen and Phosphorus content and standard deviation bars ($\bar{x} \pm S.D.$) in the shoots and roots of *Acaena elongata*, *Ageratina glabrata*, and *Solanum pubigerum*, with (+AMF) and without

been demonstrated in studies of different plant species (Ramos-Zapata et al. 2012, 2013).

The influence of AMF on germination was found to vary between plant species. The greatest benefit was shown by *Ageratina glabrata*, followed by *Solanum pubigerum* and the least for *Acaena elongata* (Fig. 2). This variability can be attributed to differences in biological traits between plant species, such as fruit type, seed dormancy, seed size, and reserve content (Closa and Goicoechea 2011; Orozco-Segovia and Sanchez-Coronado 2013; Zamora-Crescencio et al. 2018). For instance, *Ageratina glabrata* produces the smallest seed size (0.0001 g), while *Solanum pubigerum* has the intermediate seed size (0.002 g). In contrast, *Acaena elongata* produces the largest seed size (0.02 g) (Calderón



(– AMF) arbuscular mycorrhizal fungi treatment. The bar graphs illustrate the percentage of Nitrogen (above) and Phosphorus (below) in both the shoots and roots


Fig. 6 Mycorrhiza response index and standard deviation bars ($\bar{x} \pm S.D.$) of the three secondary vegetation species in a temperate forest

Table 5 Statistical summary of the two-way ANOVA for the colonization by AMF parameters for *Acaena elongata*, *Ageratina glabrata* and *Solanum pubigerum* in treatments with mycorrhiza (+ AMF) and without mycorrhiza (– AMF)

Source of variation	df	AMF Colonization		Hyphae colonization		Arbuscules colonization		Spores colonization	
		<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Species	2	5.701	0.004	5.97	0.003	1.26	0.28	2.31	0.102
AMF	1	330.107	0.001	334.7	0.001	26.13	0.001	6.42	0.01
Species:AMF	2	1.29	0.29	1.28	0.27	1.25	0.28	1.71	0.18

*Significant differences are observed in bolds

and Rzedowski 2005). It is possible to observe a direct correlation between seed size and seed reserve content (Soriano et al. 2011), since smaller seeds (such as those of *Ageratina glabrata*), which have fewer stored reserves, may need to establish arbuscular mycorrhizal associations more quickly to obtain nutritional benefits and be able to allocate them to plant tissue production. In contrast, larger seeds, such as those of *Acaena elongata*, take longer to establish such symbiotic relationships and may benefit less due to their higher reserve content. In this context, Jin et al. (2009) found a strong negative correlation between seed size and AMF colonization. This suggests that species with smaller seeds may be more dependent on AMF for nutrient acquisition, especially during early growth stages, possibly due to their limited seed reserves.

A positive effect of AMF on accumulative germination was observed in the three species studied. Germination rates were significantly higher in treatments with AMF (+ AMF) than without (– AMF). This indicates that these fungi can induce seed germination (Tian et al. 2023). In this context, Varga (2015) reported a negative effect of AMF spores on seed germination, while Rueda-Puente et al. (2010) suggested that AM fungi could induce seed germination, although the underlying mechanisms remain unclear. AMF

may contribute to the establishment of favorable microsites for seed germination through the formation of micro and macro aggregates in the soil and the production of hyphal exudates (Koide 2000). The presence of micro and macro-aggregates can increase nutrient availability and maintain a more stable moisture regime, AMF create a nurturing environment that can enhance germination rates and subsequent growth of plant seeds. Therefore, AMF may play a role in creating moisture microenvironment conditions conducive to seed germination of tree species in temperate forests (Olivera-Morales et al. 2011).

It is also possible that seed exudates may stimulate germination of AMF spores, and as a result, rapid establishment of AM association has been observed. These exudates contain a variety of organic compounds such as sugars, amino acids, and other signaling molecules that may act as chemical cues to induce AMF spores to germinate (Besserer et al. 2008). However, the specific physical or chemical processes of AMF were not examined in this study.

Acaena elongata and *Solanum pubigerum* showed an earlier germination in the + AMF treatment. This early response could give mycorrhizal plants an initial competitive advantage, facilitating early establishment (Van Der Heijden 2004; Jansa et al 2008; Varga 2015). These results highlight the importance of arbuscular mycorrhiza in the early stages of the life cycle of SV plant species in temperate forests. *Ageratina glabrata*, however, did not show a significant difference in the first day of germination in any treatment. It is very important to note that other mechanisms, such as the presence of beneficial fungi and bacteria, may also contribute to seed germination in + AMF treatments. These microorganisms could provide a “priming” effect similar to AMF, increasing nutrient availability and possibly even providing protection against soil pathogens.

Significant differences were found in the time to 50% germination (T_{50}) between treatments, with *Ageratina glabrata* and *Solanum pubigerum* taking longer to germinate 50% of their seeds in the + AMF treatment. This suggests that AMF symbiosis affects not only the germination rate and total percentage, but also the synchrony of germination, as seen in the T_{50} (Table 2). Such synchronization may be advantageous for establishment as it allows seeds to remain viable in the soil for a longer period of time. Seed




Fig. 7 Mean values of AMF hyphae colonization percentage and standard deviation bars ($\bar{x} \pm S.D.$) of the three species from secondary vegetation in a temperate forest. Different letters correspond to statistical differences $p < 0.05$

priming can be facilitated by treating seeds with specific osmotic agents, soaking times, cold or heat to achieve more synchronized germination in a seed population. Marimuthu et al. (2018) propose that AMF may have a priming effect on seeds, and seedlings inoculated with AMF at the seed stage may show increased resistance to water stress. However, the ecophysiological mechanisms underlying these observations remain poorly understood.

For *Acaena elongata*, no significant differences were observed between treatments, suggesting that mycorrhizae do not significantly affect germination synchrony in this species. *Acaena elongata* is typically found in disturbed areas of temperate forests and is the most common species near agricultural fields and open canopy forest sites among the three species studied. High seed viability rates (95% as determined by the tetrazolium test) have been previously documented for this species. However, the presence of high seed heteromorphism within woody Rosaceae species and a non-deep physiological dormancy, as in the case of *Acaena elongata* (Martínez-Camacho et al. 2018) could contribute to a lack of synchrony in germination, even in the presence of AMF. In relation to *A. elongata*'s response to AMF, it has been previously shown that there are negative and significant correlations between colonization and some phenological phases, and this varies seasonally (Vázquez-Santos et al. 2019).

For the growth stage, Closa and Goicoechea (2011) demonstrated interspecific variation in the percentage of colonization by AMF in *Oxalis acetosella* and *Viola riviniana*. In this case, root structure and root hair characteristics are considered to be the fundamental factors regulating the mycorrhizal association (Roumet et al. 2006). *Solanum pubigerum* has a taproot system (Bui et al. 2015), which could correlate with its ability to store nutrients and therefore show a low colonization response during early phenological stages. We suggest that root systems should be considered in future analyses of AMF dynamics.

A recently emerged seedling loses the resistance provided by the seed coat, making this phenological stage particularly vulnerable as it lacks the physical robustness of an adult plant (Kitajima and Fenner 2000). During this critical period, the plant must prioritize rapid growth and the development of a deep and branched root system to ensure adequate water and nutrient supply. At the same time, the allocation of resources for maintaining the AMF association increases (Vázquez-Santos et al. 2021). In our study, colonization percentages were significantly higher in *Acaena elongata* and *Solanum pubigerum* compared to *Ageratina glabrata*. Despite the high colonization rates observed in the first two species, they showed a low growth response.

It has been shown that AMF inoculation significantly enhances seedling growth, biomass, nutrient uptake and improves biomass partitioning (Souza et al. 2023; Tian

et al. 2023; Yang et al. 2024). In this study, the greatest stem length observed in *Solanum pubigerum* treated with AMF, together with the most notable response in total dry weight in *Acaena elongata* and *Ageratina glabrata*, indicates the role of AMF in resource allocation to vegetative structures. This was corroborated by the positive influence of AMF on shoot N y P concentration for *Acaena elongata* and *Ageratina glabrata*, suggesting that association with AMF may be critical for the efficiency of soil nutrient uptake and translocation to shoot. Notably, *Solanum pubigerum* showed the least response to mycorrhizal colonization, with no significant differences in root length, stem dry weight, root dry weight or total dry weight. In addition, this species showed a less pronounced effect of AMF, since only an increase in P accumulation in the aerial part was observed in the presence of AMF, but not of N. This is consistent with its significantly lower rate of mycorrhizal response compared to *Acaena elongata* and *Ageratina glabrata*, suggesting that *Solanum pubigerum* is least dependent on AMF association during early growth.

It is important to clarify that plants can be colonized by different species of arbuscular mycorrhizal fungi (AMF) and that AMF species do not colonize two different plant species with the same intensity (Helgason et al. 2002). This AMF diversity may lead to different responses in germination and growth in different plant species. The specificity of the relationship between plants and their fungal symbionts could reflect adaptations to particular environmental conditions, leading to variation in colonization efficiency and consequently in the benefits derived from mycorrhizal symbiosis, such as improved nutrient uptake and resistance to soil pathogens. Thus, it is important for future research to recognize that intraradical colonization varies between plant and fungal species. This consideration may lead to a more accurate understanding of AMF functions at different phenological stages, and the processes involved in their association.

In our study, there may be an additive effect from the interaction of AMF with other soil microorganisms. For example, bacteria and other organisms associated with the hyphosphere of AMF (Mansfeld-Giese et al. 2002; Jansa and Gryndler 2010; Lecomte et al. 2011) could act as scarifying agents for seeds, thereby enhancing germination in association with these fungi. In addition, the microbiome associated with mycelial networks may have promoted nutrient mineralization processes (Wang et al. 2022, 2024), resulting in increased plant biomass. Therefore, we suggest conducting further experiments to confirm whether other biotic agents, in addition to AMF, influence seed germination and growth of these secondary vegetation species. This may provide a new paradigm for understanding the role of biotic interactions in plant ecophysiology.

We conclude that AMF colonization has a positive and significant influence on the early phenological stages of the SV plant species studied. During early germination, rapid root development and robust vegetative growth are critical and require substantial nutrient supply, in which AMF may play a key role. As the plant grows to later stages, the resource allocation requirements for AMF maintenance may change, reflecting a dynamic association that adapts to the developmental needs of the plant. Also, AMF, directly or through their associated biota, may promote germination and growth of the four SV species studied, although further studies are needed. Moreover, the variability in AMF colonization among host species suggests that specific plant-fungus interactions are finely tuned during the plant life cycle, which is also influenced by environmental conditions and soil properties. This knowledge can be incorporated into forest management programs.

Acknowledgements We are grateful to Brenda Guerrero Bautista, Ariadna Peralta Valencia, and Irene Sanchez Gallén for their help with field work and technical assistance in the laboratory. Romero-Romero M.A. for his help in maps edition. Doctor Sampayo-Maldonado S. for their advice in data analysis. This research is part of the doctoral studies of Y. Vázquez-Santos in the Posgrado en Ciencias Biológicas of the Universidad Nacional Autónoma de México. Y. Vázquez-Santos acknowledges the Consejo Nacional de Humanidades, Ciencia y Tecnología (CONHACyT)-Mexico (No. 818569) for scholarships to pursue her doctoral studies.

Author contributions [Yasmin Vázquez-Santos], [Silvia Castillo-Argüero], [Noé Montaña] and [Francisco J. Espinosa-García]: Conceptualization, reviewing and editing; [Yasmin Vázquez-Santos] [Yuriana Martínez-Orea] and [Silvia Castillo-Argüero]: Methodology and Fieldwork; [Yasmin Vázquez-Santos], and [César Flores-Ortiz]: Data Collection and Data Analysis [César Flores-Ortiz]. [Yasmin Vázquez-Santos]: Writing—Original Draft; All authors Writing—Reviewing and Editing.

Funding This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Competing interests The authors declare that they have no competing financial interests or personal relationships that could have appeared to influence the research reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted

use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ågren GI (2008) Stoichiometry and nutrition of plant growth in natural communities. *Annu Rev Ecol Evol Syst* 39:153–170. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173515>
- Álvarez-Yépiz JC, Martínez-Yrizar A, Búrquez A, Lindquist C (2008) Variation in vegetation structure and soil properties related to land use history of old-growth and secondary tropical dry forests in northwestern Mexico. *For Ecol Manage* 256(3):355–366. <https://doi.org/10.1016/j.foreco.2008.04.049>
- Argüelles-Mayao A (2018) Estudio de las redes ectomicorizas del bosque de Abies religiosa (Kunth) Schlechtendal et Chamisso para determinar su uso potencial en estrategias de restauración. PhD Thesis in Biological Sciences, Instituto de Biología, UNAM, Mexico City
- Baslam M, Goicoechea N (2012) Water deficit improved the capacity of arbuscular mycorrhizal fungi (AMF) for inducing the accumulation of antioxidant compounds in lettuce leaves. *Mycorrhiza* 22:347–359. <https://doi.org/10.1007/s00572-011-0408-9>
- Baslam M, Garmendia I, Goicoechea N (2011) Arbuscular mycorrhizal fungi (AMF) improved growth and nutritional quality of greenhouse-grown lettuce. *J Agric Food Chem* 59(10):5504–5515. <https://doi.org/10.1021/jf200501c>
- Besserer A, Bécard G, Jauneau A, Roux C, Séjalon-Delmas N (2008) GR24, a synthetic analog of strigolactones, stimulates the mitosis and growth of the arbuscular mycorrhizal fungus *Gigaspora rosea* by boosting its energy metabolism. *Plant Physiol* 148(1):402–413. <https://doi.org/10.1104/pp.108.121400>
- Brundrett MC, Tedersoo L (2018) Evolutionary history of mycorrhizal symbioses and global host plant diversity. *New Phytol* 220(4):1108–1115. <https://doi.org/10.1111/nph.14976>
- Bui HH, Serra V, Pagès L (2015) Root system development and architecture in various genotypes of the Solanaceae family. *Botany* 93(8):465–474. <https://doi.org/10.1139/cjb-2015-0008>
- Calderón de Rzedowski G, Rzedowski J (2005) Flora fanerogámica del Valle de México. Pátzcuaro: Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Instituto de Ecología, A.C. México. ISBN-e: 978-607-7607-36-6.
- Carrillo-Saucedo SM, Gavito ME (2020) Resilience of soil aggregation and exocellular enzymatic functions associated with arbuscular mycorrhizal fungal communities along a successional gradient in a tropical dry forest. *Mycorrhiza* 30(1):109–120. <https://doi.org/10.1007/s00572-019-00928-9>
- Cavagnaro TR, Bender SF, Asghari HR, van der Heijden MG (2015) The role of arbuscular mycorrhizas in reducing soil nutrient loss. *Trends Plant Sci* 20(5):283–290. <https://doi.org/10.1016/j.tplants.2015.03.004>
- Chongtham I, Pandey RR (2020) Influence of Arbuscular Mycorrhizal Fungi (AMF) on seedling growth responses of *Cedrela serrata* and *Dipterocarpus tuberculatus*. *J Mycopathol Res* 58(3):181–186
- Closa I, Goicoechea N (2011) Infectivity of arbuscular mycorrhizal fungi in naturally regenerating, unmanaged and clear-cut beech forests. *Pedosphere* 21(1):65–74. [https://doi.org/10.1016/S1002-0160\(10\)60080-X](https://doi.org/10.1016/S1002-0160(10)60080-X)
- Copetta A, Bardi L, Bertolone E, Berta G (2011) Fruit production and quality of tomato plants (*Solanum lycopersicum* L.) are affected by green compost and arbuscular mycorrhizal fungi. *Plant Biosyst* 145(1):106–115

- Delgadillo-Duran E (2011) Productividad primaria neta de los bosques templados de la cuenca del río Magdalena. Master Thesis in Biological Sciences, UNAM, Mexico City. <https://hdl.handle.net/20.500.14330/TES01000669952>
- FAO (2020) Global Forest Resources Assessment 2020: Main report. FAO. Rome [WWWDocument], n.d. www.fao.org. URL <http://www.fao.org/forest-resources-assessment/2020> (accessed 2.2.21)
- Galicia L, Gamboa AM, Cram S, Vergara B, Ramírez VP, Saynes V, Siebe C (2016) Stocks and dynamics of soil organic carbon in temperate forests of Mexico. *Terra Latinoam* 34(1):1–29
- García E (1988) Modificaciones al sistema de clasificación de Koppen para adaptarlo a las condiciones de la República Mexicana (No. 551.59 G37 1988)
- Gilliam FS (2016) Forest ecosystems of temperate climatic regions: from ancient use to climate change. *New Phytol* 212:871–887. <https://doi.org/10.1111/nph.14255>
- Gioria M, Pyšek P, Osborne BA (2018) Timing is everything: does early and late germination favor invasions by herbaceous alien plants? *J Plant Ecol* 11:4–16. <https://doi.org/10.1093/jpe/rtw105>
- Guadarrama P, Castillo S, Ramos-Zapata JA, Hernández-Cuevas LV, Camargo-Ricalde SL (2014) Arbuscular mycorrhizal fungal communities in changing environments: The effects of seasonality and anthropogenic disturbance in a seasonal dry forest. *Pedobiologia* 57(2):87–95. <https://doi.org/10.1016/j.pedobi.2014.01.002>
- Helgason T, Merryweather JW, Denison J, Wilson P, Young JPW, Fitter AH (2002) Selectivity and functional diversity in arbuscular mycorrhizas of co-occurring fungi and plants from a temperate deciduous woodland. *J Ecol* 90(2):371–384. <https://doi.org/10.1046/j.1365-2745.2001.00674.x>
- INEGI (2011) Instituto Nacional de Estadística y Geografía. <https://www.inegi.org.mx/>. Accessed 10 Dec 2019
- Ismail AM, Johnson DE, Ella ES, Vergara GV, Baltazar AM (2012) Adaptation to flooding during emergence and seedling growth in rice and weeds, and implications for crop establishment. *AoB Plants*. <https://doi.org/10.1093/aobpla/pls019>
- Jansa J, Smith FA, Smith SE (2008) Are there benefits of simultaneous root colonization by different arbuscular mycorrhizal fungi? *New Phytol* 177(3):779–789. <https://doi.org/10.1111/j.1469-8137.2007.02294.x>
- Jansa J, Gryndler M (2010) Biotic environment of the arbuscular mycorrhizal fungi in soil. *Arbuscular mycorrhizas: physiology and function*. In: Hinanit K, Yoram K (eds) *Arbuscular mycorrhizas: physiology and function*. Springer, Dordrecht, pp 209–236
- Jin L, Wang S, Wang X, Shen Y (2009) Seed size influences arbuscular mycorrhizal symbiosis across leguminous host-plant species at the seedling stage. *Symbiosis* 49:111–116. <https://doi.org/10.1007/s13199-009-0013-2>
- Johnson NC (2010) Resource stoichiometry elucidates the structure and function of arbuscular mycorrhizas across scales. *New Phytol* 185(3):631–647. <https://doi.org/10.1111/j.1469-8137.2009.03110.x>
- Kitajima K, Fenner M (2000) Ecology of seedling regeneration. *Seeds: the ecology of regeneration in plant communities*. CABI publishing, Wallingford UK, pp 331–359. <https://doi.org/10.1079/9780851994321.03>
- Koide RT (2000) Mycorrhizal symbiosis and plant reproduction. In: Kapulnik Y, Douds DDJ (eds) *Arbuscular Mycorrhizas: physiology and function*. Kluwer academic, Dordrecht, pp 19–46
- Koske RE, Gemma JN (1989) A modified procedure for staining roots to detect VA mycorrhizas. *Mycol Res* 92(4):486–488. [https://doi.org/10.1016/S0953-7562\(89\)80195-9](https://doi.org/10.1016/S0953-7562(89)80195-9)
- Kumar D, Manhas RK, Ohri P (2023) Deciphering the growth promoting and stress curtailing role of nitric oxide in *Meloidogyne incognita* infested *Solanum lycopersicum* seedlings. *Sci Horticul* 319:112147. <https://doi.org/10.1016/j.scienta.2023.112147>
- Lambers H, Oliveira RS, Lambers H, Oliveira RS (2019) Growth and allocation. *Plant Physiol Ecol*. https://doi.org/10.1007/978-3-030-29639-1_10
- Leck MA, Parker VT, Simpson R (eds) (2008) *Seedling ecology and evolution*. Cambridge University Press, Cambridge, Reino Unido, p 514
- Lecomte J, St-Arnaud M, Hijri M (2011) Isolation and identification of soil bacteria growing at the expense of arbuscular mycorrhizal fungi. *FEMS Microbiol Lett* 317(1):43–51. <https://doi.org/10.1111/j.1574-6968.2011.02209.x>
- Lesschen JP, Cammeraat LH, Kooijman AM, van Wesemael B (2008) Development of spatial heterogeneity in vegetation and soil properties after land abandonment in a semi-arid ecosystem. *J Arid Environ* 72(11):2082–2092. <https://doi.org/10.1016/j.jaridenv.2008.06.006>
- Li M, Jordan NR, Koide RT, Yannarell AC, Davis AS (2016) Meta-analysis of crop and weed growth responses to arbuscular mycorrhizal fungi: implications for integrated weed management. *Weed Sci* 64(4):642–652. <https://doi.org/10.1614/WS-D-16-00050.1>
- Lin S, Fan C, Wang J, Zhang C, Zhao X, von Gadow K (2024) Chronic anthropogenic disturbance mediates the biodiversity-productivity relationship across stand ages in a large temperate forest region. *J Appl Ecol* 00:1–11. <https://doi.org/10.1111/1365-2664.14588>
- Maighal M, Salem M, Kohler J, Rillig MC (2016) Arbuscular mycorrhizal fungi negatively affect soil seed bank viability. *Ecol Evol* 6(21):7683–7689. <https://doi.org/10.1002/ece3.2491>
- Mansfeld-Giese K, Larsen J, Bødker L (2002) Bacterial populations associated with mycelium of the arbuscular mycorrhizal fungus *Glomus intraradices*. *FEMS Microbiol Ecol* 41(2):133–140. <https://doi.org/10.1111/j.1574-6941.2002.tb00974.x>
- Marimuthu P, Kumutha K, Sivakumar U (2018) Seed priming effect of arbuscular mycorrhizal fungi against induced drought in rice. *J Pharmacogn Phytochem* 7(2):1742–1746
- Martínez-Camacho YD, Castillo-Argüero S, Martínez-Orea Y, Sánchez-Coronado ME (2018) Ecofisiología de la germinación de *Acaena elongata* (Rosaceae), una especie indicadora de perturbación de un bosque templado, al sur de la Ciudad de México. *Revista Mex De Biodivers* 89(3):806–814
- McGonigle TP, Miller MH, Evans DF, Fairchild GL, Swan JA (1990) A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New Phytol* 115:495–501. <https://doi.org/10.1111/j.1469-8137.1990.tb00476.x>
- Milla R, Giménez-Benavides L, Escudero A, Reich PB (2009) Intra- and interspecific performance in growth and reproduction increase with altitude: a case study with two *Saxifraga* species from northern Spain. *Funct Ecol* 23:111–118. <https://doi.org/10.1111/j.1365-2435.2008.01484.x>
- Mosse B, Hayman DS (1971) Plant growth responses to vesicular-arbuscular mycorrhiza: II. In *Unsterilized Field Soils* *New Phytol* 70(1):29–34. <https://doi.org/10.1111/j.1469-8137.1973.tb02017.x>
- Olivera-Morales D, Castillo-Argüero S, Guadarrama P, Ramos-Zapata J, Álvarez-Sánchez J, Hernández-Cuevas L (2011) Establecimiento de plántulas de *Quercus rugosa* Née inoculadas con hongos micorrizógenos arbusculares en un bosque templado de México. *Boletín De La Sociedad Botánica De México* 89:115–121
- Orozco-Segovia A, Sanchez-Coronado Ma E (2013) Germinación. In: Márquez Guzmán J, Collazo Ortega M, Martínez-Gordillo M, Orozco-Segovia A, Vázquez Santana S (eds) *biología de angiospermas*. Universidad Nacional Autónoma de México. Ciudad de México, México
- Plenchette C, Fortin JA, Furlan V (1983) Growth responses of several plant species to mycorrhizae in a soil of moderate P-fertility. I. mycorrhiza dependency under field conditions. *Plant Soil* 70:199–209

- Post WM, Kwon KC (2000) Soil carbon sequestration and land-use change: processes and potential. *Glob Change Biol* 6(3):317–327. <https://doi.org/10.1046/j.1365-2486.2000.00308.x>
- Ramos-Zapata JA, Marrufo-Zapata D, Guadarrama P, Carrillo-Sánchez L, Hernández-Cuevas L, Caamal-Maldonado A (2012) Impact of weed control on arbuscular mycorrhizal fungi in a tropical agroecosystem: a long-term experiment. *Mycorrhiza* 22:653–661. <https://doi.org/10.1007/s00572-012-0443-1>
- Ramos-Zapata J, Marrufo-Zapata D, Guadarrama-Chávez P, Solís-Rodríguez U, Salinas-Peba L (2013) Ruderal plants: temporary hosts of arbuscular mycorrhizal fungi in traditional agricultural systems? *Trop Subtrop Agroecosyst* 16(3):399–406
- Rawat M, Singh R, Sharma J, Saklani H, Chand T, Bhatt ID, Pandey R (2022) An overview of the functioning of temperate forest ecosystems with particular reference to Himalayan temperate forest. *Trees for People* 8:100230. <https://doi.org/10.1016/j.tfp.2022.100230>
- Roumet C, Urcelay C, Díaz S (2006) Suites of root traits differ between annual and perennial species growing in the field. *New Phytol* 170:357–368. <https://doi.org/10.1111/j.1469-8137.2006.01667.x>
- Rueda-Puente EO, Murillo-Amador B, Castellanos-Cervantes T, García-Hernández JL, Tarazón-Herrera MA, Medina SM, Barrera LEG (2010) Effects of plant growth promoting bacteria and mycorrhizal on *Capsicum annuum* L. var. *aviculare* [Dierbach] D'Arcy and Eshbaugh) germination under stressing abiotic conditions. *Plant Physiol Biochem* 48(8):724–730. <https://doi.org/10.1016/j.plaphy.2010.04.002>
- Santibañez-Andrade G, Castillo-Argüero S, Martínez-Orea Y (2015) Evaluación del estado de conservación de la vegetación de los bosques de una cuenca heterogénea del Valle de México. *Bosque* 36(2):299–313. <https://doi.org/10.4067/S0717-92002015000200015>
- Sayer J, Chokkalingam U, Poulsen J (2004) The restoration of forest biodiversity and ecological values. *For Ecol Manage* 201(1):3–11. <https://doi.org/10.1016/j.foreco.2004.06.008>
- Smith SE, Smith FA (2011) Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annu Rev Plant Biol* 62:227–250. <https://doi.org/10.1146/annurev-arplant-042110-103846>
- Smith SE, Smith FA (2012) Fresh perspectives on the roles of arbuscular mycorrhizal fungi in plant nutrition and growth. *Mycologia* 104(1):1–13. <https://doi.org/10.3852/11-229>
- Smith SE, Jakobsen I, Grønlund M, Smith FA (2011) Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiol* 156(3):1050–1057. <https://doi.org/10.1104/pp.111.174581>
- Soriano D, Orozco-Segovia A, Márquez-Guzmán J, Kitajima K, Gamboa-de Buen A, Huante P (2011) Seed reserve composition in 19 tree species of a tropical deciduous forest in Mexico and its relationship to seed germination and seedling growth. *Ann Bot* 107(6):939–951. <https://doi.org/10.1093/aob/mcr041>
- Sortibrán L, Verdú M, Valiente-Banuet A (2019) A nurse plant benefits from facilitative interactions through mycorrhizae. *Plant Biol* 21(4):670–676. <https://doi.org/10.1111/plb.12948>
- Soto-Cruz FJ, Zorrilla JG, Rial C, Varela RM, Molinillo JM, Igartuburu JM, Macías FA (2021) Allelopathic activity of strigolactones on the germination of parasitic plants and arbuscular mycorrhizal fungi growth. *Agronomy* 11(11):2174. <https://doi.org/10.3390/agronomy11112174>
- Souza GG, Santos SC, Santos CC, Dias AS, Silverio JM, Trovato VW, Flauzino DS (2023) Arbuscular mycorrhizal fungi promote the growth of *Dipteryx alata* Vogel. *Braz J Biol* 83:e275172. <https://doi.org/10.1590/1519-6984.275172>
- Tian F, Wang JC, Bai XX, Yang YB, Huang L, Liao XF (2023) Symbiotic seed germination and seedling growth of mycorrhizal fungi in *Paphiopedilum hirsutissimum* (Lindl. Ex Hook.) Stein from China. *Plant Signal Behav* 18(1):2293405
- Van Der Heijden MG (2004) Arbuscular mycorrhizal fungi as support systems for seedling establishment in grassland. *Ecol Lett* 7(4):293–303. <https://doi.org/10.1111/j.1461-0248.2004.00577.x>
- van Hall RL, Cammeraat LH, Keesstra SD, Zorn M (2017) Impact of secondary vegetation succession on soil quality in a humid Mediterranean landscape. *CATENA* 149:836–843. <https://doi.org/10.1016/j.catena.2016.05.021>
- Varga S (2015) Effects of arbuscular mycorrhizal fungi and maternal plant sex on seed germination and early plant establishment. *Am J Bot* 102(3):358–366. <https://doi.org/10.3732/ajb.1400361>
- Vázquez-Santos Y, Castillo-Argüero S, Martínez-Orea Y, Sánchez-Gallen I, Vega-Frutos R, Camargo-Ricalde SL, Hernández-Cuevas LV (2019) The reproductive phenology of *Acaena elongata* and its relation with arbuscular mycorrhizal fungi. *Symbiosis* 79:129–140. <https://doi.org/10.1007/s13199-019-00629-z>
- Vázquez-Santos Y, Martínez-Orea Y, Álvarez-Sánchez J, Montaña NM, Camargo-Ricalde SL, Castillo-Argüero S (2021) Interaction of *Acaena elongata* L. with arbuscular mycorrhizal fungi under phosphorus limitation conditions in a temperate forest. *Phyton*. <https://doi.org/10.32604/phyton.2021.014208>
- Wang Y, Li Y, Li S, Rosendahl S (2021) Ignored diversity of arbuscular mycorrhizal fungi in co-occurring mycotrophic and non-mycotrophic plants. *Mycorrhiza* 31:93–102. <https://doi.org/10.1007/s00572-020-00997-1>
- Wang F, Zhang L, Zhou J, Rengel Z, George TS, Feng G (2022) Exploring the secrets of hyphosphere of arbuscular mycorrhizal fungi: processes and ecological functions. *Plant Soil* 481(1):1–22. <https://doi.org/10.1007/s11104-022-05621-z>
- Wang L, George TS, Feng G (2024) Concepts and consequences of the hyphosphere core microbiome for arbuscular mycorrhizal fungal fitness and function. *New Phytol* 242(4):1529–1533. <https://doi.org/10.1111/nph.19396>
- WRB, World reference base for soil resources (2022) International soil classification system for naming soils and creating legends for soil maps. FAO. 85–117
- Yang W, Zhuang J, Tian Y, Wans S, Ding S, Zhang M (2024) The effect of arbuscular mycorrhizal fungi on the growth of *cyclocarya paliurus* cutting seedlings. *Turkish J Agric for*. <https://doi.org/10.55730/1300-011X.3166>
- Zamora-Crescencio P, Rico-Gray V, Ramírez-Medina LNG, Barrientos-Medina RC, Plasencia-Vázquez AH, Villegas P, del Domínguez-Carrasco MR, Gutiérrez-Báez C (2018) Composición Y Estructura De La Vegetación Secundaria En Bethania. Composition and structure of secondary vegetation in Bethania, Campeche Mexico. *Polibotánica, Campeche, México*. <https://doi.org/10.18387/polibotanica.45.5>
- Zar JH (1974) Biostatistical analysis. Prentice-Hall Inc, Englewood Cliffs, New Jersey
- Zhou L, Zhou X, He Y, Fu Y, Du Z, Lu M, Thakur MP (2022) Global systematic review with meta-analysis shows that warming effects on terrestrial plant biomass allocation are influenced by precipitation and mycorrhizal association. *Nat Commun* 13(1):4914. <https://doi.org/10.1038/s41467-022-32671-9>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.