

Traditional uses, antimicrobial and acaricidal activities of 20 plants selected among Reunion Island's flora

Emmanuelle Dorla ^{a)}, Isabelle Grondin ^{a)}, Thomas Hue ^{b)}, Patricia Clerc ^{a)}, Séverine Dumas ^{a)}, Anne Gauvin-Bialecki ^{a)}, Philippe Laurent ^{a)*}

^a Laboratoire de Chimie des Substances Naturelles et des Sciences des Aliments (LCSNSA), Université de La Réunion, avenue René Cassin–CS 92003–97744 Saint-Denis Cedex 9, France.

^b Laboratoire de Parasitologie, Institut Agronomique néo-Calédonien, BP 73, 98890 Païta, New Caledonia

* Corresponding author: philippe.laurent@univ-reunion.fr
Tel: +262 96 28 50

1 **Abstract:**

2 The aim of this study was to screen the antimicrobial and acaricidal activity of 20 endemic or
3 indigenous plants from Reunion Island (Indian Ocean). Plants were chosen on the basis of
4 their traditional uses and their biocidal activities found in the literature. A survey was
5 conducted in the local population to assess and supplement knowledge about the selected
6 plants. The collected information confirmed and/or supplemented the data obtained for nine
7 plants. Seven plants were described for the first time for their traditional uses in medicine and
8 ethnoveterinary practices. To evaluate their biocidal activities, leaves or bark were treated
9 with ethyl acetate using an accelerated solvent extraction method. Six bacteria and five fungi,
10 frequently implicated in infectious diseases, were used to assess the antimicrobial activity of
11 these extracts. A preliminary screening using the paper disk diffusion assay showed an
12 effective antibacterial activity of 16 extracts. The minimum inhibitory concentration (MIC) of
13 active plant extracts was then determined using a microdilution method. The leaf extract from
14 *Peperomia borbonensis* displayed the widest spectrum of antibacterial activity and was the
15 only one to act as a fungicide. In parallel, acaricidal bioassays were performed on the larvae
16 of the tick *Rhipicephalus microplus* (Ixodidae), and plant extracts from *Peperomia*
17 *borbonensis* and *Zanthoxylum heterophyllum* were the most effective. The preliminary studies
18 of these plant extracts exhibited biocidal activities that were not described in the literature and
19 that are congruent with traditional uses for some of them. Investigations are currently being
20 conducted to isolate the active compound(s) and evaluate their potential for future
21 developments and applications.

22 **Keywords:** acaricidal, antimicrobial, La Réunion, plant extracts, traditional use, survey
23
24

1. Introduction

The resistance of microorganisms and arthropods to chemicals is becoming a major concern for agriculture and public health issues. According to several reports on antimicrobial resistance, edited by the World Health Organization, new resistance mechanisms have emerged in the last few years and have spread globally, threatening our ability to treat common infectious diseases. The non-availability and the high costs of a new generation of antibiotics have led to an increase in mortality and morbidity. In the European Union, strains of resistant bacteria are responsible for 25,000 deaths and an extra health care cost of €1.5 billion each year (World Health Organization, 2015a, 2015b, 2014). Worldwide, drug-resistant infections cause 700,000 deaths every year, and a recent report has suggested that without policies to stop the spread of antimicrobial resistance, the number of deaths in 2050 would rise to 10 million every year (O’Neil, 2016). Hence, there is an urgent need to react, and one option is to find effective novel molecules to circumvent this resistance phenomenon. In this context, several studies have highlighted the antimicrobial activities of plants from different regions of the world (Agyare et al., 2016; Aumeeruddy-Elalfi et al., 2015; Mgbeahuruike et al., 2017; Mickymaray et al., 2016).

Likewise, for several years, resistance to synthetic acaricides has been frequently reported in the literature (Abbas et al., 2014; Jongejan and Uilenberg, 2004). Among Acaria, the cattle tick *Rhipicephalus (Boophilus) microplus* is considered as the most important parasite of livestock in the world (Estrada-Peña et al., 2006). It was introduced to Reunion Island (also called La Réunion) by the importation of “Moka cattle” from Madagascar (Barré and Uilenberg, 2010). This tick causes enormous losses in cattle production, due to blood loss, stress and irritation, which affect the milk and hide. Moreover, this arthropod can transmit diseases to cattle, such as babesiosis caused by *Babesia bovis* and *Babesia bigemina* and anaplasmosis caused by *Anaplasma marginale* (Estrada-Peña et al., 2006). The main

treatment to control this arthropod is the use of chemical acaricides, such as organochlorines, carbamates and organophosphates, which have negative consequences in the environment and contribute to the development of resistant populations. There is therefore increasing interest by the scientific community to find new potential sources of compounds with biological activity against pests. Several recent studies have suggested the use of plants to fight cattle tick (Abbas et al., 2014; Adenubi et al., 2016; Borges et al., 2011; H  e et al., 2015).

Reunion Island has a great number of assets to find new molecules in the abundant plant biodiversity. Emerging from the Indian Ocean about three million years ago, La R  union (55  3' E and 21  5'S) is a French volcanic island located about 665 km east of Madagascar in the Mascarene archipelago. The island has a rugged topography and has the highest peak in the Indian Ocean (Piton des Neiges; 3070 m), deep valleys and one of the most active volcanoes in the world (Piton de La Fournaise; 2631 m). Its tropical climate is tempered by the prevailing southeast trade winds and is occasionally unsettled by cyclones. The tropical location and the dramatic landform of the island, combined with the moist trade winds, determine strong asymmetry between the eastern windward coast subjected to daily rainfall (up to 4 to 7 m/year) and the much drier western leeward coast (0.5 to 2.5 m/year). A large variety of microclimates is observed, with radical changes in sun light, precipitation and temperature, allowing a large number of vegetal species to develop and evolve. Consequently, Reunion Island is listed among the world's top biodiversity hotspots with an endemic rate approximately of 40%. Isolated from the mainland and other islands, the biodiversity of La R  union results from the colonization of species from Madagascar, Africa, Asia, and Australia by marine currents and birds. The evolution of indigenous species, far from their region of origin, led to the emergence of several endemic species. Almost 840 indigenous species constituted the vascular flora of La R  union, with 236 plants species (28%) being strictly endemic to La R  union and 153 species endemic (18%) to the archipelago of

Mascarene. Since the arrival of the Europeans in the 16th century, Reunion Island flora has experienced major perturbations, such as forest clearance and exotic plant introduction. Nevertheless, 25% of the original forest cover remains, and this allows unique plant communities to develop (Lavergne, 2001; Parc National de La Réunion, 2008; Strasberg et al., 2005; Thébaud et al., 2009). The human inhabitants are composed of people who originated from Europe (mainly France), Africa, Madagascar, India and China. The local traditional medicine was the result of the mixed knowledge of this cosmopolitan population (Pourchez, 2011). For instance *Ayapana triplinervis* (Vahl) R.M. King et H. Rob was introduced from India at the end of 18th century and was used by the local population for digestive problems. Additionally, numerous endemic plants from La Réunion are used in traditional medicine; for instance, Ambaville (*Hubertia ambavilla* Bory) for dermic problems, blood circulation, gastric anti-ulcer and diabetes (“Aplamedom”). The exploration of this empirical knowledge could lead to the discovery of new active molecules.

The screening of 20 plants from La Réunion for biocidal activities constituted the core of this study. Plants were first chosen because of their endemic or indigenous status. Then, we considered their traditional uses in medicine and ethnoveterinary practices, and if no data were available, we selected the plants on the basis of existing biocidal activities in species of the same genus. To complement the fragmentary and limited knowledge of the selected plants, a small-scale survey of the population was conducted. Plant extracts were tested for antimicrobial and acaricidal activities. To our knowledge, this is the first study to report such properties of plants from La Réunion.

2. Material and methods

2.1. Plants and extracts

2.1.1. Selection of plant samples

Samples were chosen from our laboratory's plant collection formed during the extensive research program BIOMOL-TCN, which aimed to find new therapeutic, cosmetic and nutraceutical molecules in the marine, terrestrial and microbial biodiversity from La Réunion.

2.1.2. Preparation of crude extracts

The 20 plants chosen in this study were collected between 2009 and 2013 in different forests of Reunion Island. As described above, the selection criteria of the plants were (1) their biological status, (2) the availability of information concerning traditional practices and/or (3) data for biocidal activity of each species found in the literature or for the genus if no data were available concerning the species. This information is summarized in Table 1. The plant materials were identified by the botanists Jacques Fournel and Professor Dominique Strasberg (Faculty of Science and Technology, University of La Réunion). Voucher specimens are kept and were deposited in the Herbarium of the University.

2.2. Small scale survey

2.2.1. Study area

La Réunion (55°3' E and 21°5'S), with Mauritius and Rodrigues, is one of three islands that composes the Mascarene archipelago located in the Indian Ocean. Discovered in 1507–1512, Reunion Island has been a French overseas department since 1946, and is divided into 24 municipalities (average size: 100 km²) (Strasberg et al., 2005). According to the census of 2012, the island's population was about 833,944 (403,907 males and 430,037 females). The population is mainly composed of people who originated from France, Africa, Madagascar India (Tamil Nadu and Gujarat) and China ("Insee – Département de La Réunion (974)" 2012). Economic activities and around 80% of the population are located in the coastal lowlands due to the rugged topography of the island. Almost 40% of the territory belongs to the National Park, which was established to preserve and conserve the terrestrial biodiversity

(Figure 1). (Strasberg et al., 2005). The core zone of La Réunion National Park coincides with the area of “Pitons, cirques and remparts” in the World Heritage List of the UNESCO.

2.2.2. Questionnaire design and data collection

A small-scale survey of the general population was carried out in Reunion Island from October 2015 to February 2016. The questionnaire was adapted from previous studies (Grønhaug et al., 2008; Samois and Mahomoodally, 2015). Data were recorded via face-to-face interviews in the local language (Créole) or French, depending on the interviewee. The first section concerned general information about gender, age, and place of residence. The second section concerned the knowledge of the selected plants and their traditional uses. Non-specialist people (n = 55) were randomly chosen and interviewed in the street, at their home or in their garden. Nineteen municipalities out of 24 were covered in this study. In addition, four traditional healers, one pharmacist, two ethnobotanists and one farmer were interviewed about the medicinal usages of the 20 studied plants.

2.3. Biological assays

2.3.1. Preparation of crude extracts

Collected plant materials (leaves and/or bark) were air-flow dried (40°C), crushed into fine powder and kept dry at room temperature until use. Crude extracts were obtained with an accelerated solvent extractor (ASE® 300, Accelerated Solvent Extractor, France) using ethyl acetate as solvent ($\geq 99.5\%$, Carlo-Erba, France). Conditions were as follows: temperature, 40°C; pressure, 100 bars; five cycles with static extraction time of 6 min; and flush volume, 100%. The crude extracts were concentrated in a rotary vacuum and were kept at 4°C until further use.

2.3.2. Evaluation of antimicrobial activity

➤ Microorganisms

Antimicrobial activity was evaluated on bacteria and yeasts with sanitary relevance. Three Gram-positive bacteria, *Listeria monocytogenes* (ATCC 1914), methicillin-resistant *Staphylococcus aureus* (NCTC 12493) and *Streptococcus pyogenes* (ATCC 19615), three Gram-negative bacteria, *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 10145) and *Salmonella enterica* (ATCC 13076), two filamentous fungi, *Aspergillus fumigatus* (ATCC 204305) and *Aspergillus niger* (ATCC 1688), and three yeasts, *Candida albicans* (ATCC 10231), *Candida tropicalis* (ATCC 1369) and *Cryptococcus neoformans* (ATCC 76484) (Humeau Laboratory, France) were used. *E. coli*, *S. aureus*, *P. aeruginosa* and *S. enterica* were grown on Mueller-Hinton medium (Sigma Aldrich, France), whereas *L. monocytogenes* and *S. pyogenes* were grown on Brain Heart medium (bioMérieux, France). Fungi were grown on Sabouraud medium (Sigma Aldrich, France). The media were prepared according to the manufacturer's instructions.

➤ Disc diffusion test

The assay was conducted by a modified disk diffusion method of the Clinical and Laboratory Standard Institute (Cavalieri et al., 2009). Suspensions of microorganisms were made in sterile medium and adjusted spectrophotometrically between 1×10^5 and 1×10^6 CFU/mL for bacteria and between 1×10^3 and 1×10^4 spores/mL for fungi. A 150 mL volume of sterile medium was added to Petri dishes (245 × 245 × 25 mm) to a thickness of 4 mm. Once the surface of media was inoculated with 1 mL of the microbial suspension, the disks (6 mm diameter) were placed on the surface of the medium. Each extract was dissolved in dimethylsulphoxide (DMSO ≥ 99.5 %, Sigma-Aldrich, France) to have a final concentration of 10 mg/mL and was tested at 20 µL/disk. Positive controls were chloramphenicol (BDH

Chemicals, England) for bacteria and amphotericine B (Sigma-Aldrich, France) for fungi. The plates were left for 30 min at room temperature to allow diffusion of the extract into the agar. They were then incubated at 37°C for 24 h and 48 h for bacteria and fungi, respectively. Antimicrobial activity was determined by measuring the inhibition zone with a caliper in millimeters. The experiment was repeated three times.

➤ **Determination of minimum inhibitory concentration by the microdilution method**

The minimal inhibition concentration (MIC) determination method was applied to extracts that had demonstrated their efficiency against microorganisms by the disk diffusion method (Kuate et al., 2009). The extracts and antibiotics were dissolved in DMSO (max 6.2% of DMSO per well). The initial concentration of the extracts was 1000.00 µg/mL and they were serially diluted two-fold in order to obtain a concentration range from 1000.00 to 0.98 µg/mL in sterile nutrient broth. A negative control was run in parallel to study the impact of the solvent on the microorganism growth. Each well was inoculated with 50 µL of suspension containing between 1×10^5 and 1×10^6 CFU/mL for the bacteria and between 1×10^3 and 1×10^4 spores/mL for the fungi. After incubation (37°C, 24 h and 48 h for bacteria and fungi, respectively), 20 µL (0.2 mg/mL) of *p*-iodonitrotetrazolium violet (INT, Sigma-Aldrich, France) was added to each well and the plates were incubated for an additional hour. MIC was determined as the lowest concentration of plant extract inhibiting microbial growth, indicated by a decrease in the intensity of the red colour of the formazan product. Positive controls were chloramphenicol (BDH Chemicals, England) for bacteria and amphotericine B (Sigma-Aldrich, France) for fungi. Three replicates were made for each extract.

2.3.3. Evaluation of acaricidal activity

➤ **Preparation of ticks**

R. microplus engorged females were collected after detachment from cattle. The females were incubated in the laboratory of the Agronomic Institute of New Caledonia (IAC) at a temperature of 27°C and a relative humidity (RH) of 85% for one week. Eggs were then collected and placed in the same conditions until larvae were 2–3 weeks old.

➤ Larval packet test

The acaricidal activity of plant extracts was evaluated using the modified larval packet test (LPT) (Stone and Haydock, 1962) on 14–21-day-old larvae. Extracts were diluted in ethanol to a 5% solution. A 7.5 × 8.5-cm nylon paper (Anowo Ltd, Switzerland) was impregnated with the different extracts and placed in a fume hood for 1 h to allow ethanol evaporation, before being folded into packets using bulldog clips. Approximately 100 *R. microplus* larvae were placed into each treated nylon paper packet, which were then sealed with additional bulldog clips and placed in an incubator (27°C, 85% Relative Humidity) for 24 h. Two replicates for each plant extracts, a negative control (nylon paper with ethanol) and a positive control (amitraz 1 g/L) were used. After 24 h, the numbers of live and dead larvae were counted to calculate the percentage of larval mortality.

➤ Statistical analysis

Mean values of mortality and standard deviation of the mean were calculated for each plant extracts. The LC₅₀ (50% lethal concentration) was calculated using the Probit method (Finney, 1971), generated by the Probit POLOPC program (LeOra Software, 1987, Berkeley, CA, USA).

3. Results and Discussion

3.1. Traditional use of the selected plants

Despite traditional plants uses being an important cultural component of Reunion Island, surprisingly, no survey had been made in the population to collect information about these plants and their uses until our work.

Of the people interviewed, 58.2% were male and 41.8% were female; 32.7% were aged 18–35 years old, 49.1% were 36–60 years old and 18.2% were >60 years old. Finally, 65.5% of them lived in a rural area. A substantial proportion of the interviewed non-specialists (78.2%) cultivated and used plants mainly from their gardens as medicinal remedies. Among them, the proportion of women (81.8%) who used traditional remedies was slightly higher than the proportion of men (75.7%). This result is aligned with other studies around the world (Samoisy and Mahomoodally, 2015). In La Réunion, as in Mauritius and Rodrigues, people are particularly attached to traditional practices, and women are often the principal source of knowledge transmission (Pourchez, 2011). More than 85% of herbal medicine traditional uses were indeed transmitted through generations within the families. The knowledge of traditional uses of plants is indeed held by older people, and it is likely that the transfer of this information is now greatly affected by lifestyle modernization, as in many regions of the world. The preliminary literature analysis showed that only a few traditional uses were reported concerning the 20 selected plants (Table 1). During the survey, no traditional use in relation to antimicrobial or acaricidal properties was cited by the non-specialist population even though one or more of the 20 studied plants were recognized by the interviewees. For example, *Terminalia bentzoe* (L.) Pers. (benjoin) was recognized by 89% of interviewed persons, *Zanthoxylum heterophyllum* (Lam.) Sm. (bois de poivre) 50% and *Calophyllum tacamahaca* Willd. (takamaka) was recognized by 40%. This knowledge was observed only for some representative plants from La Réunion because the general population did not know the selected plants of this study well. For instance, the *Psiadia* species were the least known probably because of their low distribution and scarcity in the island. *Psiadia amygdalina*

Cordem. and *Psiadia boivinii* B.L.Rob. were recognized by 6 and 4% of interviewed people, respectively.

We also questioned eight specialized people (four traditional healers, one pharmacist, two ethnobotanists and one farmer) about the traditional uses of the 20 selected plants. These people acted daily for the valorization, conservation and preservation of the biodiversity of Reunion Island and the associated traditional knowledge, which is endangered by the lack of verbal transfer to new generations. As shown in Table 1, this small scale-survey allowed the collection of helpful information about the traditional use of little-known plants. This is the first time that seven plants were described for their traditional uses in medicine and ethnoveterinary practices: *Psiadia dentata* DC., *Psiadia retusa* DC., *Vernonia fimbrillifera* Less., *Antidesma madagascariense* Lam., *Stillingia lineata* (Lam.) Müll.Arg., *Indigofera amoxylum* (DC.) Polhill and *Peperomia borbonensis* Miq. Furthermore, the collected information confirmed and/or supplemented the data obtained from the literature for nine plants (*Poupartia borbonica* J.F.Gmel, *Carissa spinarum* L., *Secamone volubilis* (Lam.) Marais., *Calophyllum tacamahaca* Willd, *Terminalia bentzoe* (L.) Pers., *Croton mauritanus* Lam., *Sophora denudata* Bory, *Zanthoxylum heterophyllum* (Lam.) Sm., *Nuxia verticillata* Lam.). Finally, no traditional use was identified for *Psiadia amygdalina* Cordem, *Psiadia boivinii* B.L.Rob., *Psiadia laurifolia* Cordem and *Monimia rotundifolia* Thouars. These results reinforce bioprospecting of plants and herbal traditional uses to discover new substances. They should help us to direct our bioactivity tests and phytochemical studies.

3.2. Antimicrobial activities of the selected plants

Crude extracts were prepared using ethyl acetate as solvent, and were tested against the different microorganisms. Table 2 shows the results obtained for plants displaying antibacterial activity for at least one bacteria/fungi. The antimicrobial activity varied greatly

266 according to the tested species. The leaves of *Carissa spinarum*, *Indigofera amnoxylum*,
 267 *Psiadia amygdalina*, *Sophora denudata*, *Vernonia fimbrillifera*, *Zanthoxylum heterophyllum*
 268 displayed no antimicrobial activity. The same result was found for the bark of *Psiadia dentata*
 269 and *P. retusa* and for the leaves and bark of *P. laurifolia*. Sixteen extracts inhibited at least
 270 one microbial species. Among these positive extracts, 12 had a zone of growth inhibition
 271 greater than 10 mm: *Antidesma madagascariense* (bark), *Callophylum tacamahaca* (leaves),
 272 *Croton mauritianus* (leaves), *Monimia rotundifolia* (leaves), *Psiadia amygdalina* (bark),
 273 *Poupartia borbonica* (leaves), *Psiadia boivinii* (leaves), *Psiadia retusa* (leaves), *Secamone*
 274 *volubilis* (leaves), *Sophora denudata* (bark), *Stillingia lineata* (leaves) and *Terminalia bentzoe*
 275 (leaves). The minimal inhibition concentration (MIC) was determined for the positive extracts
 276 following the disk diffusion test (Table 3). It is generally considered that antimicrobial
 277 activity is good for extracts with a MIC less than 100 µg/mL; from 100 to 500 µg/mL, the
 278 antimicrobial activity is moderate; from 500 to 1000 µg/mL, the antimicrobial activity is
 279 weak; and over 1000 µg/mL, the extract is considered inactive (Holetz et al., 2002). Only
 280 seven MIC values were >1000 µg/mL. The active extracts displayed a MIC ranged between
 281 1000 µg/mL and 15.62 µg/mL. Gram-positive bacteria were more sensitive than Gram-
 282 negative bacteria, as was frequently found in the literature (Ríos and Recio, 2005). This is
 283 probably due to cell-wall differences, as Gram-negative bacteria have an outer membrane
 284 known to act as a barrier to many molecules. The most susceptible bacterium was the
 285 methicillin-resistant *Staphylococcus aureus* strain, since 12 extracts were active against it.
 286 The best activities against this bacterium were observed with *Callophylum tacamahaca*
 287 (leaves) and *Psiadia dentata* (leaves) with a MIC of 62.50 µg/mL. The activity of
 288 chloramphenicol was 37.50 µg/mL on this strain. No traditional uses as an antimicrobial were
 289 reported for *Callophylum tacamahaca*. According to our survey's results, *Psiadia dentata* is
 290 traditionally used to treat dermic problems, such as mycoses in La Réunion, but the ethyl

acetate extract of this plant was not active against the tested fungi. Other extraction methods and a wider range of fungi should be used to evaluate the fungicide activity of this plant in the future. In Mauritius, five endemic *Psiadia* species used traditionally to treat pulmonary infections, wounds and burns were evaluated for their antimicrobial activity. They did not show any or showed moderate activity against the studied strains (Govinden-Soulange et al., 2004). The *Terminalia bentzoe* leaf extract showed a weak activity against *Staphylococcus aureus* and *Streptococcus pyogenes* with a MIC of 1000.00 µg/mL. Additionally, the current findings showed good antimicrobial activity of the extract of *Sophora denudata* bark against *Listeria monocytogenes* and *Streptococcus pyogenes* with a MIC value of 125.00 µg/mL and 15.62 µg/mL respectively. Indeed, the antimicrobial activity of *Sophora* species has often been reported being due to flavonoids (Sohn et al., 2004; Tsuchiya et al., 1996). *Psiadia dentata* (leaves) and *Peperomia borbonensis* (leaves) extracts both displayed the broadest spectrum of antibacterial activities. Although MICs were not very low, *P. borbonensis* was active against four bacteria among the six tested, and was especially effective against all the Gram-negative species. Moreover, *P. borbonensis* was the only species active against fungi (one out of five, *Aspergillus fumigatus*). The traditional use of *P. borbonensis* in La Réunion as an antimicrobial is not listed. Likewise, the *Peperomia* species around the world have not been known for this traditional use, except *P. tetraphylla* (G.Forst.) Hook. & Arn., which was used to fight microbial infections in India (Nishanthi et al., 2012). Nevertheless, several studies on the antimicrobial activity of *Peperomia* species were reported in the literature (Ferreira et al., 2014; Langfield et al., 2004; Mbah et al., 2002; Saga Kitamura et al., 2006). The butanolic fraction of *Peperomia pellucida* (L.) Kunth, composed of tannins, flavonoids and saponins, showed good inhibition diameters against *E. coli* and *P. aeruginosa* (Khan and Omoloso, 2002). Patuloside A isolated from *P. pellucida* was tested on four Gram-positive bacteria and six Gram-negative bacteria and a low minimal inhibition concentration (MIC = 8

μg/mL) was obtained against *Staphylococcus aureus* and *Streptococcus β-haemolyticus* (Khan et al., 2010). Finally, Malquichagua Salazar et al. (2005) showed the activity of two compounds from *Peperomia villipetiola* C.D.C. against the fungus *Cladosporium sphaerospermum*.

Concerning the three indigenous species of the study, only *Antidesma madagascariense* and *Stillingia lineata* extracts showed antibacterial activity with the disk diffusion method and only against *S. aureus*. The MIC of these extracts were 125.00 μg/mL and >1000.00 μg/mL, respectively. Unlike *S. lineata*, the antimicrobial activity of *A. madagascariense* was already evaluated and the methanol leaf extracts showed activity against *Enterococcus faecalis* (MIC = 60.00 μg/mL), *S. aureus* (MIC = 500.00 μg/mL), methicillin resistant *S. aureus* (MIC = 250.00 μg/mL) and *Candida albicans* (MIC = 500.00 μg/mL) (Mahomoodally et al., 2015). In a study conducted by Rangasamy et al. (2007), the *A. madagascariense* crude methanol extract was active against some of the tested microorganisms of our study. Among them, *S. aureus* (MIC = 500.00 μg/mL) and *S. enteridis* (MIC = 125.00 μg/mL) were the most susceptible strains (Rangasamy et al., 2007). Finally, the species *Carissa spinarum* (leaves) was not active against the tested microorganisms; however, the methanolic extract of the roots of *C. spinarum* from India was active against *E. coli* (MIC = 125 ±10 μg/mL), *S. aureus* (MIC = 110 ± 28 μg/mL) and *A. niger* (MIC = 256 ±30 μg/mL) in a study by Sanwal and Chaudhary (2011).

That being said, plant extracts with a MIC of 1000 μg/mL should not be neglected, as they could contain interesting antimicrobial molecules. From this point of view, the leaf extracts from *Peperomia borbonensis* should be further explored, since it displayed the widest spectrum of antibacterial activity and also acted as a fungicide. Indeed, bioactive compound concentrations in plant extracts vary depending on the polarity of solvents. Ethyl acetate was used in our work to maximize the collection of a wide range of molecules in order to select

interesting plants for further research. Other solvents should now be gradually used to optimize active compound extraction and their isolation by bioguided-fractionation.

3.3. Acaricidal activities of the selected plants

Five extracts showed acaricidal activity on the tick *Rhipicephalus microplus* larvae (Figure 2).

At a concentration of 5 %, the most active samples, with a mortality rate of 100%, were

extracted from the leaves of *Peperomia borbonensis* (Piperaceae) and the bark of

Zanthoxylum heterophyllum (Rutaceae). The extracts obtained from *Zanthoxylum*

heterophyllum leaves showed weaker activity (63.8%) than those obtained from the bark

(100%). Medium activity was also observed for *Monimia rotundifolia* (65.7% of mortality).

Finally, *Psiadia amygdalina* leaf extract had weak acaricidal activity with 31.8% of mortality.

No previous study has reported acaricidal activity of these four plants. However, several

studies have reported this property in plants belonging to the Piperaceae and Rutaceae

families. Solvent extracts and essential oils from *Piper* (Piperaceae) species have been widely

studied for their acaricidal properties (de Souza Chagas et al., 2012; Ferraz et al., 2010; Lima

et al., 2014; Silva et al., 2009). Likewise, several species of the genus *Zanthoxylum* were

studied for their biocidal activity against arthropods (Moussavi et al., 2015; Prieto et al.,

2011). The essential oil of *Zanthoxylum caribaeum* Lam. was assessed for its acaricidal

activity against cattle tick. This volatile extract acted on engorged females and inhibited

oviposition and egg eclosion (Nogueira et al., 2014). Before this work, no reports about

biocidal activity (*in vitro* or traditional) against arthropods was recovered for *Peperomia* spp.,

Psiadia spp. and *Monimia* spp. As previously said, other solvents should now be gradually

used to optimize active compound extraction and their isolation by bioguided-fractionation.

Furthermore, these efforts should be complemented by the use of other extraction methods

and the implementation of tests on a wide spectrum of biological targets. This is best

illustrated by another work conducted in parallel on *Peperomia borbonensis* in our laboratory, which led to demonstrating insecticidal activity of the leaf essential oil of *Peperomia borbonensis* and of its isolated major components against the melon fly *Bactrocera cucurbitae* (Dorla et al., 2017).

4. Conclusion

To conclude, this *in vitro* study corroborated the acaricidal activity of *Peperomia borbonensis* traditionally used by a few farmers on the island to protect their cattle from ticks. The crude extract could possess one or several bioactive molecules acting in combination. Further research is currently being conducted to isolate the active component(s) by bioguided-fractionation. Furthermore, this study also demonstrated, for the first time, the acaricidal activity of *Zanthoxylum heteropyllum* and the antimicrobial activity of *Psiadia dentata* (MIC of 62.50 µg/mL against *S. aureus*) and *Sophora denudata* (MIC of 15.62 µg/mL against *S. pyogenes*). These results will further be used in bioguided phytochemical studies. Lastly, a large-scale survey should be carried out to collect more information throughout the territory from older people. It could be especially interesting to dedicate a part of this future survey to the three mountain cirques Mafate, Cilaos and Salazie (Figure 1), and more particularly Mafate, which is accessible only by a pedestrian path network. In this landlocked area, the traditional lifestyle is more preserved than in coastal areas. Many potential bioactive plants remain unexplored among the Reunion Island biodiversity. Our multi-criteria approach would also allow the discovery of many other plants with interesting properties. Moreover, several plants selected in this study are endangered and protected in La Réunion. Raising the awareness and knowledge about such plants and improving their valorization by researching biological properties could allow their preservation.

389

390 **Conflict of interest**

391 The authors declare no conflict of interest.

392 **Acknowledgements**

393 The authors are very grateful to D. Strasberg and J. Fournel (UMR C_53 PVBMT), H.
394 Thomas (Parc National de La Réunion) and G. Begue for their contributions to species
395 collection and plant material identification. Thanks to the following students for their
396 technical contributions: L. Rivière, A. Fabre, M. Marchand and E. Grondin. The authors
397 would like also to thank Anowo Ltd. for providing nylon paper.

398 **Funding**

399 This work was supported by the European Commission and the Regional Council of Reunion
400 Island through the BIOMOL-TCN programme and ERDF (European Regional Development
401 Fund, Grant No. 2012–326).

References

- Abbas, R.Z., Zaman, M.A., Colwell, D.D., Gilleard, J., Iqbal, Z., 2014. Acaricide resistance in cattle ticks and approaches to its management: The state of play. *Veterinary Parasitology* 203, 6–20.
- Ademola, I.O., Ibrinke, A.A., Aghanu, N.V., Chris, C.O., 2014. Evaluation of acaricidal activity of different seed oils on *Psoroptes cuniculi* (Delafond, 1859) (Cheyletidae) in rabbits. *Journal of Pharmacy and Biological Sciences* 9, 94–97.
- Adenubi, O.T., Fasina, F.O., McGaw, L.J., Eloff, J.N., Naidoo, V., 2016. Plant extracts to control ticks of veterinary and medical importance: A review. *South African Journal of Botany* 105, 178–193.
- Adesina, J.M., Ofuya, T.I., Afolabi, L.A., 2012. Insecticidal activity of *Secamone afzelii* (Schult) K. *schum* powder in the control of *Stiphilous zeamais* (Mots) (Coleoptera: Curculionidae). *International Research Journal of Pharmacy* 8, 117–124.
- Agrawal, I.L., Mall, S.B., 1988. Studies on the insecticidal and antifeedant activity of some plant extracts on Bihar hairy caterpillar, *Diacrisia obliqua* Walker (Lep., Arctiidae). *Journal of Applied Entomology* 105, 529–532.
- Agyare, C., Boakye, Y.D., Bekoe, E.O., Hensel, A., Dapaah, S.O., Appiah, T., 2016. Review: African medicinal plants with wound healing properties. *Journal of Ethnopharmacology* 177, 85–100.
- Akinnibosun, H.A., Akinnibosun, F.I., German, B.E., 2008. Antibacterial activity of aqueous and ethanolic leaf extracts of *Peperomia pellucida* (L.) HB & K. (Piperaceae) on three Gram-negative bacteria isolates. *Science World Journal* 3, 33–36.
- Alexander, I.C., Pascoe, K.O., Manchard, P., Williams, L.A.D., 1991. An insecticidal diterpene from *Croton linearis*. *Phytochemistry* 30, 1801–1803.

425 Alkhamaiseh, S.I., Taher, M., Ahmad, F., Qaralleh, H., Althunibat, O.Y., Susanti, D., Ichwan, S.-
426 J.A., 2012. The phytochemical content and antimicrobial activities of Malaysian *Calophyllum*
427 *canum* (stem bark). Pakistan Journal of Pharmaceutical Sciences 25, 555–563.

428 Aplamedom. Association des Plantes Aromatiques et Médicinales de La Réunion.
429 <http://www.aplamedom.org/4.html> (accessed 4.14.16).

430 Aumeeruddy-Elalfi, Z., Gurib-Fakim, A., Mahomoodally, F., 2015. Antimicrobial, antibiotic
431 potentiating activity and phytochemical profile of essential oils from exotic and endemic
432 medicinal plants of Mauritius. Industrial Crops and Products 71, 197–204.

433 Barré, N., Uilenberg, G., 2010. Spread of parasites transported with their hosts: Case study of two
434 species of cattle tick. Revue scientifique et technique (International Office of Epizootics) 29,
435 135–147.

436 Belmi, R.M., Giron, J., Tansengo, M.L., 2014. *Antidesma buniis* (bignay) fruit extract as an organic
437 pesticide against *Epilachna* spp. Journal of Asian Scientific Research 4, 320–327.

438 Borges, L.M.F., Sousa, L.A.D. de, Barbosa, C. da S., 2011. Perspectives for the use of plant extracts
439 to control the cattle tick *Rhipicephalus (Boophilus) microplus*. Revista Brasileira de
440 Parasitologia Veterinária 20, 89–96.

441 Cavalieri, S.J., Harbeck, R.J., McCarter, Y.S., Ortez, J.H., Rankin, I.D., Sharp, S.E., Spiegel, C.A.,
442 2009. Manual of antimicrobial susceptibility testing. American Society for Microbiology,
443 Washington, DC.

444 Cota, B.B., Johann, S., Oliveira, D.M., Siqueira, E.P., Souza-Fagundes, E.M., Cisalpino, P.S., Alves,
445 T.M.A., Zani, C.L., 2011. Biological potential of *Stillingia oppositifolia*. Revista Brasileira de
446 Farmacognosia 21, 70–77.

447 Cuesta-Rubio, O., Oubada, A., Bello, A., Maes, L., Cos, P., Monzote, L., 2015. Antimicrobial
448 assessment of resins from *Calophyllum antillanum* and *Calophyllum inophyllum*. Phytotherapy
449 Research 29, 1991–1994.

450 Dahot, M.U., 1999. Antibacterial and antifungal activity of small protein of *Indigofera oblongifolia*
 451 leaves. *Journal of Ethnopharmacology* 64, 277–282.

452 de Souza Chagas, A.C., de Barros, L.D., Cotinguiba, F., Furlan, M., Giglioti, R., de Sena Oliveira,
 453 M.C., Bizzo, H.R., 2012. In vitro efficacy of plant extracts and synthesized substances on
 454 *Rhipicephalus (Boophilus) Microplus* (Acari: Ixodidae). *Parasitology Research* 110, 295–303.

455 Dechayont, B., Hansakul, P., Itharat, A., 2012. Comparison of antimicrobial, antioxidant activities
 456 and total phenolic content of *Antidesma thwaitesianum* fruit extracts by different methods.
 457 *Journal of the Medical Association of Thailand* 95 Suppl 1, S147–S153.

458 Dharmaratne, H.R.W., Wijesinghe, W.M.N.M., Thevanasem, V., 1999. Antimicrobial activity of
 459 xanthones from *Calophyllum* species, against methicillin-resistant *Staphylococcus aureus*
 460 (MRSA). *Journal of Ethnopharmacology* 66, 339–342.

461 Dorla, E., Gauvin-Bialecki, A., Deucher, Z., Allibert, A., Grondin, I., Deguine, J.-P., Laurent, P.,
 462 2017. Insecticidal activity of the leaf essential oil of *Peperomia borbonensis* Miq. (Piperaceae)
 463 and its major components against the melon fly *Bactrocera cucurbitae* (Diptera: Tephritidae).
 464 *Chemistry and Biodiversity* 14, 1–9.

465 Erasto, P., Grierson, D.S., Afolayan, A.J., 2006. Bioactive sesquiterpene lactones from the leaves of
 466 *Vernonia amygdalina*. *Journal of Ethnopharmacology* 106, 117–120.

467 Estrada-Peña, A., García, Z., Sánchez, H.F., 2006. The distribution and ecological preferences of
 468 *Boophilus microplus* (Acari: Ixodidae) in Mexico. *Experimental & Applied Acarology* 38, 307–
 469 316.

470 de B.F. Ferraz, A., Balbino, J.M., Zini, C.A., Ribeiro, V.L.S., Bordignon, S.A.L., von Poser, G.,
 471 2010. Acaricidal activity and chemical composition of the essential oil from three *Piper* species.
 472 *Parasitology Research* 107, 243–248.

473 Ferreira, E.A., Juliana B., R., Correia, M.V., Young, M.C.M., Guimarães, E.F., Franchi, G.C.,
 474 Nowill, A.E., Lago, J.H.G., Yamaguchi, L.F., Kato, M.J., 2014. Antifungal and cytotoxic 2-

475 Acylcyclohexane-1,3-diones from *Peperomia alata* and *P. trineura*. Journal of Natural Products
 476 27, 1377–1382.

477 Finney D. J., 1971. Probit Analysis, 3rd ed. Cambridge University Press, Cambridge. Journal of
 478 Pharmaceutical Sciences 60, 1432–1432.

479 Gemed, N., Mokonnen, W., Lemma, H., Tadele, A., Urga, K., Addis, G., Debell, A., Getachew,
 480 M., Tek, F., Yirsaw, K., Mudie, K., Gebre, S., 2014. Insecticidal activity of some traditionally
 481 used Ethiopian medicinal plants against sheep ked *Melophagus ovinus*. Journal of Parasitology
 482 Research 2014, 1–7.

483 Govinden-Soulange, J., Magan, N., Gurib-Fakim, A., Gauvin, A., Smadja, J., Kodja, H., 2004.
 484 Chemical composition and in vitro antimicrobial activities of the essential oils from endemic
 485 *Psiadia* species growing in Mauritius. Biological & Pharmaceutical Bulletin 27, 1814–1818.

486 Grønhaug, T.E., Glæserud, S., Skogsrud, M., Ballo, N., Bah, S., Diallo, D., Paulsen, B.S., 2008.
 487 Ethnopharmacological survey of six medicinal plants from Mali, West-Africa. Journal of
 488 Ethnobiology and Ethnomedicine 4, 1–11.

489 Holetz, F.B., Pessini, G.L., Sanches, N.R., Cortez, D.A.G., Nakamura, C.V., Filho, D., Prado, B.,
 490 2002. Screening of some plants used in the Brazilian folk medicine for the treatment of
 491 infectious diseases. Memórias do Instituto Oswaldo Cruz 97, 1027–1031.

492 Hüb, T., Cauquil, L., Fokou, J.B.H., Dongmo, P.M.J., Bakarnga-Via, I., Menut, C., 2015. Acaricidal
 493 activity of five essential oils of *Ocimum* species on *Rhipicephalus (Boophilus) microplus* larvae.
 494 Parasitology Research 114, 91–99.

495 Insee – Département de La Réunion (974). 2012.
 496 http://www.insee.fr/fr/themes/dossier_complet.asp?codegeo=DEP-974 (accessed 4.11.16).

497 Islam, A., Sayeed, A., Bhuiyan, M.S.A., Mosaddik, M.A., Islam, M.A.U., Astaq Mondal Khan,
 498 G.R.M., 2001. Antimicrobial activity and cytotoxicity of *Zanthoxylum budrunga*. Fitoterapia 72,
 499 428–430.

500 Jongejan, F., Uilenberg, G., 2004. The global importance of ticks. *Parasitology* 129, S3–S14.

501 Jonville, M.C., Kodja, H., Strasberg, D., Pichette, A., Ollivier, E., Frédérick, M., Angenot, L.,
502 Legault, J., 2011. Antiplasmodial, anti-inflammatory and cytotoxic activities of various plant
503 extracts from the Mascarene Archipelago. *Journal of Ethnopharmacology* 136, 525–531.

504 Kadir, R., Awang, K., Khamaruddin, Z., Soit, Z., 2015. Chemical compositions and termiticidal
505 activities of the heartwood from *Calophyllum inophyllum* L. *Annals of the Brazilian Academy of*
506 *Sciences* 87, 743–751.

507 Kamal, R., Mangla, M., 1993. In vivo and in vitro investigations on rotenoids from *Indigofera*
508 *tinctoria* and their bioefficacy against the larvae of *Anopheles stephensi* and adults of
509 *Callosobruchus chinensis*. *Journal of biosciences* 18, 93–101.

510 Khan, A., Rahman, M., Islam, M.S., 2010. Isolation and bioactivity of a xanthone glycoside from
511 *Peperomia pellucida*. *Life Sciences and Medicine Research* 2010, 1–10.

512 Khan, M.R., Omoloso, A.D., 2002. Antibacterial activity of *Hygrophila stricta* and *Peperomia*
513 *pellucida*. *Fitoterapia* 73, 251–254.

514 Kitonde, C.K., Fidahusein, D.S., Lukhoba, C.W., Jumba, M.M., 2012. Antimicrobial activity and
515 phytochemical study of *Vernonia glabra* (Steetz) Oliv. & Hiern. in Kenya. *African Journal of*
516 *Traditional, Complementary, and Alternative Medicines* 10, 149–157.

517 Krishna, P.M., KNV, R., S, S., Banji, D., 2012. A review on phytochemical, ethnomedical and
518 pharmacological studies on genus *Sophora*, Fabaceae. *Revista Brasileira de Farmacognosia* 22,
519 1145–1154.

520 Küçükboyacı, N., Özkan, S., Adigüzel, N., Tosun, F., 2011. Characterisation and antimicrobial
521 activity of *Sophora alopecuroides* L. var. *alopecuroides* alkaloid extracts. *Turkish Journal of*
522 *Biology* 35, 379–385.

523 Kuete, V., Nana, F., Ngameni, B., Mbaveng, A.T., Keumedjio, F., Ngadjui, B.T., 2009.
 524 Antimicrobial activity of the crude extract, fractions and compounds from stem bark of *Ficus*
 525 *ovata* (Moraceae). *Journal of Ethnopharmacology* 124, 556–561.
 526 Langfield, R.D., Scarano, F.J., Heitzman, M.E., Kondo, M., Hammond, G.B., Neto, C.C., 2004. Use
 527 of a modified microplate bioassay method to investigate antibacterial activity in the Peruvian
 528 medicinal plant *Peperomia galioides*. *Journal of Ethnopharmacology* 94, 279–281.
 529 Lavergne, R., 2001. *Le grand livre des tisaneurs et plantes médicinales indigènes de La Réunion*,
 530 Orphie. ed. France.
 531 Leite, S.P., Vieira, J.R.C., de Medeiros, P.L., Leite, R.M.P., de Menezes Lima, V.L., Xavier, H.S., de
 532 Oliveira Lima, E., 2006. Antimicrobial activity of *Indigofera suffruticosa*. *Evidence-based*
 533 *Complementary and Alternative Medicine* 3, 261–265.
 534 Lima, A. da S., Filho, J.G. do N.S., Pereira, S.G., Guillon, G.M.S.P., Santos, L. da S., Júnior, L.M.C.,
 535 2014. Acaricide activity of different extracts from *Piper tuberculatum* fruits against
 536 *Rhipicephalus microplus*. *Parasitology Research* 113, 107–112.
 537 Lima, G.P.G. de, Souza, T.M. de, Freire, G. de P., Farias, D.F., Cunha, A.P., Ricardo, N.M.P.S.,
 538 Morais, S.M. de, Carvalho, A.F.U., 2013. Further insecticidal activities of essential oils from
 539 *Lippia sidoides* and *Croton* species against *Aedes aegypti* L. *Parasitology Research* 112, 1953–
 540 1958.
 541 Lindsay, E.A., Berry, Y., Jamie, J.F., Bremner, J.B., 2000. Antibacterial compounds from *Carissa*
 542 *lanceolata* R.Br. *Phytochemistry* 55, 403–406.
 543 Mahomoodally, M.F., Korumtollee, H.N., Chady, Z.Z.B.K., 2015. Ethnopharmacological uses of
 544 *Antidesma madagascariense* Lam. (Euphorbiaceae). *Journal of Intercultural Ethnopharmacology*
 545 4, 86–89.
 546 Malquichagua Salazar, K.J., Delgado Paredes, G.E., Lluncor, L.R., Young, M.C.M., Kato, M.J.,
 547 2005. Chromenes of polyketide origin from *Peperomia villipetiola*. *Phytochemistry* 66, 573–579.

548 Mao, L., Henderson, G., 2007. Antifeedant activity and acute and residual toxicity of alkaloids from
 549 *Sophora flavescens* (leguminosae) against formosan subterranean termites (Isoptera:
 550 Rhinotermitidae). Journal of Economic Entomology 100, 866–870.

551 Masoko, P., Picard, J., Eloff, J.N., 2005. Antifungal activities of six South African *Terminalia*
 552 species (Combretaceae). Journal of Ethnopharmacology 99, 301–308.

553 Mbah, J.A., Ngemenya, M.N., Abawah, A.L., Babiaka, S.B., Nubed, L.N., Nyongbela, K.D., Lemuh,
 554 N.D., Efange, S.M., 2012. Bioassay-guided discovery of antibacterial agents: in vitro screening
 555 of *Peperomia vulcanica*, *Peperomia fernandopoioana* and *Scleria striatinux*. Annals of Clinical
 556 Microbiology and Antimicrobials 11, 1–10.

557 Mbah, J.A., Tchuendem, M.H., Tane, P., Sterner, O., 2002. Two chromones from *Peperomia*
 558 *vulcanica*. Phytochemistry 60, 799–801.

559 Mgbeahuruike, E.E., Yrjönen, T., Vuorela, H., Holm, Y., 2017. Bioactive compounds from medicinal
 560 plants: Focus on *Piper* species. South African Journal of Botany 112, 54–69.

561 Mickymaray, S., Al Aboody, M.S., Rath, P.K., Annamalai, P., Nooruddin, T., 2016. Screening and
 562 antibacterial efficacy of selected Indian medicinal plants. Asian Pacific Journal of Tropical
 563 Biomedicine 6, 185–191.

564 Misra, L.N., Wouatsa, N.A.V., Kumar, S., Venkatesh Kumar, R., Tchoumboungang, F., 2013.
 565 Antibacterial, cytotoxic activities and chemical composition of fruits of two Cameroonian
 566 *Zanthoxylum* species. Journal of Ethnopharmacology 148, 74–80.

567 Moussavi, N., Malterud, K.E., Mikolo, B., Dawes, D., Chandre, F., Corbel, V., Massamba, D.,
 568 Overgaard, H.J., Wangenstein, H., 2015. Identification of chemical constituents of *Zanthoxylum*
 569 *heitzii* stem bark and their insecticidal activity against the malaria mosquito *Anopheles gambiae*.
 570 Parasites and Vectors 8, 1–8.

571 Mwanauta, R.W., Mtei, K.A., Ndakidemi, P.A., 2014. Prospective bioactive compounds from
 572 *Vernonia amygdalina*, *Lippia javanica*, *Dysphania ambrosioides* and *Tithonia diversifolia* in
 573 controlling legume insect pests. Agricultural Sciences 5, 1129–1139.

574 Mwangomo, D.T., Moshi, M.J., Magadula, J.J., 2012. Antimicrobial activity and phytochemical
 575 screening of *Antidesma venosum* root and stem bark ethanolic extracts. International Journal of
 576 Research in Phytochemistry and Pharmacology 2, 90–95.

577 Ngemenya, M.N., Mbah, J.A., Tane, P., Titanji, V.P., 2006. Antibacterial effects of some
 578 Cameroonian medicinal plants against common pathogenic bacteria. African Journal of
 579 Traditional, Complementary and Alternative medicines 3, 84–93.

580 Nishanthi, M., Mohanambal, E., Narendiran, S., Shankar, M., Aanandhi, M.V., Vijayakumar, B.,
 581 2012. Extraction and preliminary phytochemical investigation on whole plant of *Peperomia*
 582 *tetraphylla* (G. Forst. Hook & Arn). International Journal of Novel Trends in Pharmaceutical
 583 Sciences 2, 35–38.

584 Nogueira, J., Vinturelle, R., Mattos, C., Tietbohl, L.A.C., Santos, M.G., Junior, I.S.V., Mourão, S.C.,
 585 Rocha, L., Folly, E., 2014. Acaricidal properties of the essential oil from *Zanthoxylum*
 586 *caribaeum* against *Rhipicephalus microplus*. Journal of Medical Entomology 51, 971–975.

587 Nyahangare, E.T., Mvumi, B.M., Mutibvu, T., 2015. Ethnoveterinary plants and practices used for
 588 ecto-parasite control in semi-arid smallholder farming areas of Zimbabwe. Journal of
 589 Ethnobiology and Ethnomedicine 11, 1–16.

590 Obey, J.K., von Wright, A., Orjala, J., Kauhanen, J., Tikkanen-Kaukanen, C., Obey, J.K., von
 591 Wright, A., Orjala, J., Kauhanen, J., Tikkanen-Kaukanen, C., 2016. Antimicrobial activity of
 592 *Croton macrostachyus* stem bark extracts against several human pathogenic bacteria,
 593 antimicrobial activity of *Croton macrostachyus* stem bark extracts against several human
 594 pathogenic bacteria. Journal of Pathogens 2016, 1–5.

595 Olila, D., Olwa-Odyek, Opuda-Asibo, J., 2001. Antibacterial and antifungal activities of extracts of
 596 *Zanthoxylum chalybeum* and *Warburgia ugandensis*, Ugandan medicinal plants. African Health
 597 Sciences 1, 66–72.

598 Oloyede, G.K., Onocha, P.A., Olaniran, B.B., 2011. Phytochemical, toxicity, antimicrobial and
 599 antioxidant screening of leaf extracts of *Peperomia pellucida* from Nigeria. *Advances in*
 600 *Environmental Biology* 5, 3700–3709.

601 O’Neil, J., 2016. Infection prevention, control and surveillance: Limiting the development and spread
 602 of drug resistance.

603 Parc National de La Réunion, 2008. Dossier de candidature au Patrimoine Mondial de l'UNESCO
 604 Pitons, Cirques et Remparts de l’île de La Réunion, France.

605 Peres, M.T.L., Monache, F.D., Cruz, A.B., Pizzolatti, M.G., Yunes, R.A., 1997. Chemical
 606 composition and antimicrobial activity of *Croton urucurana* Baillon (Euphorbiaceae). *Journal of*
 607 *Ethnopharmacology* 56, 223–226.

608 Poullain, C., Girard-Valenciennes, E., Smadja, J., 2004. Plants from Reunion Island: Evaluation of
 609 their free radical scavenging and antioxidant activities. *Journal of Ethnopharmacology* 95, 19–
 610 26.

611 Pourchez, L., 2011. *Savoirs des femmes : médecine traditionnelle et nature: Maurice, Réunion,*
 612 *Rodrigues. UNESCO, Paris.*

613 Prieto, J.A., Patiño, O.J., Delgado, W.A., Moreno, J.P., Cuca, L.E., 2011. Chemical composition,
 614 insecticidal, and antifungal activities of fruit essential oils of three Colombian *Zanthoxylum*
 615 species. *Chilean Journal of Agricultural Research* 71, 73–82.

616 Rabe, T., Mullholland, D., van Staden, J., 2002. Isolation and identification of antibacterial
 617 compounds from *Vernonia colorata* leaves. *Journal of Ethnopharmacology* 80, 91–94.

618 Rangasamy, O., Raelison, G., Rakotoniriana, F.E., Cheuk, K., Urverg-Ratsimamanga, S., Quetin-
 619 Leclercq, J., Gurib-Fakim, A., Subratty, A.H., 2007. Screening for anti-infective properties of
 620 several medicinal plants of the Mauritian flora. *Journal of Ethnopharmacology* 109, 331–337.

621 Rani, P.U., Venkateshwaramma, T., Devanand, P., 2011. Bioactivities of *Cocos nucifera* L.
 622 (Arecales: Arecaceae) and *Terminalia catappa* L. (Myrtales: Combretaceae) leaf extracts as

623 post-harvest grain protectants against four major stored product pests. *Journal of Pest Science* 84,
624 235–247.

625 Ríos, J.L., Recio, M.C., 2005. Medicinal plants and antimicrobial activity. *Journal of*
626 *Ethnopharmacology* 100th Volume Special Section: Perspectives of Ethnopharmacology 100,
627 80–84.

628 Saga Kitamura, R.O., Romoff, P., Young, M.C.M., Kato, M.J., Lago, J.H.G., 2006. Chromenes from
629 *Peperomia serpens* (Sw.) Loudon (Piperaceae). *Phytochemistry* 67, 2398–2402.

630 Samois, A.K., Mahomoodally, M.F., 2015. Ethnopharmacological analysis of medicinal plants used
631 against non-communicable diseases in Rodrigues Island, Indian Ocean. *Journal of*
632 *Ethnopharmacology* 173, 20–38.

633 Sanwal, R., Chaudhary, A.K., 2011. Wound healing and antimicrobial potential of *Carissa spinarum*
634 Linn. in albino mice. *Journal of Ethnopharmacology* 135, 792–796.

635 Selowa, S.C., Shai, L.J., Masoko, P., Mokgotho, M.P., Magano, S.R., 2010. Antibacterial activity of
636 extracts of three *Croton* species collected in Mpumalanga region in South Africa. *African*
637 *Journal of Traditional, Complementary, and Alternative Medicines* 7, 98–103.

638 Silva, W.C., de Souza Martins, J.R., de Souza, H.E.M., Heinzen, H., Cesio, M.V., Mato, M.,
639 Albrecht, F., de Azevedo, J.L., de Barros, N.M., 2009. Toxicity of *Piper aduncum* L. (Piperales:
640 Piperaceae) from the Amazon forest for the cattle tick *Rhipicephalus (Boophilus) microplus*
641 (Acari: Ixodidae). *Veterinary Parasitology* 164, 267–274.

642 Smadja, J., Vera, 1991. Essais préliminaires sur quelques plantes de La Réunion. *Plantes médicinales*
643 *et phytothérapie* 25, 212–241.

644 Sohn, H.-Y., Son, K.H., Kwon, C.-S., Kwon, G.-S., Kang, S.S., 2004. Antimicrobial and cytotoxic
645 activity of 18 prenylated flavonoids isolated from medicinal plants: *Morus alba* L., *Morus*
646 *mongolica* Schneider, *Broussonetia papyrifera* (L.) Vent, *Sophora flavescens* Ait and
647 *Echinosophora koreensis* Nakai. *Phytomedicine* 11, 666–672.

648 Stone, B.F., Haydock, K.P., 1962. A method for measuring the acaricide-susceptibility of the cattle
649 tick *Boophilus microplus* (Can.). Bulletin of Entomological Research 53, 563–578.

650 Strasberg, D., Rouget, M., Richardson, D.M., Baret, S., Dupont, J., Cowling, R.M., 2005. An
651 assessment of habitat diversity and transformation on La Reunion Island (Mascarene Islands,
652 Indian Ocean) as a basis for identifying broad-scale conservation priorities. Biodiversity and
653 Conservation 14, 3015–3032.

654 Thébaud, C., Warren, B., Strasberg, D., Cheke, A., 2009. Mascarene islands, biology, in: Gillespie,
655 R.G. and Clague, D.A., (eds.), The Encyclopedia of Islands. University of California Press, 612-
656 619.

657 Toyang, N.J., Ateh, E.N., Keiser, J., Vargas, M., Bach, H., Tane, P., Sondengam, L.B., Davis, H.,
658 Bryant, J., Verpoorte, R., 2012. Toxicity, antimicrobial and anthelmintic activities of *Vernonia*
659 *guineensis* Benth. (Asteraceae) crude extracts. Journal of Ethnopharmacology 144, 700–704.

660 Tsuchiya, H., Sato, M., Miyazaki, T., Fujiwara, S., Tanigaki, S., Ohyama, M., Tanaka, T., Iinuma,
661 M., 1996. Comparative study on the antibacterial activity of phytochemical flavanones against
662 methicillin-resistant *Staphylococcus aureus*. Journal of Ethnopharmacology 50, 27–34.

663 Valente, P.P., Amorim, J.M., Castilho, R.O., Leite, R.C., Ribeiro, M.F.B., 2013. In vitro acaricidal
664 efficacy of plant extracts from Brazilian flora and isolated substances against *Rhipicephalus*
665 *microplus* (Acari: Ixodidae). Parasitology Research 113, 417–423.

666 Vera, R., Smadja, J., Conan, J., 1990. Preliminary studies on some alkaloid plants in the Reunion
667 Island. French. Plantes Medicinales et Phytotherapie 24, 50-65.

668 Wang, C.-F., Yang, K., You, C.-X., Zhang, W.-J., Guo, S.-S., Geng, Z.-F., Du, S.-S., Wang, Y.-Y.,
669 2015. Chemical composition and insecticidal activity of essential oils from *Zanthoxylum*
670 *dissitum* leaves and roots against three species of storage pests. Molecules 20, 7990–7999.

671 Wang, Y., Hamburger, M., Gueho, J., Hostettmann, K., 1989. Antimicrobial flavonoids from *Psiadia*
672 *trinervia* and their methylated and acetylated derivatives. Phytochemistry 28, 2323–2327.

673 Wei, L.S., Wee, W., Siong, J.Y.F., Syamsumir, D.F., 2011. Characterization of anticancer,
 674 antimicrobial, antioxidant properties and chemical compositions of *Peperomia pellucida* leaf
 675 extract. *Acta Medica Iranica* 49, 670–674.

676 World Health Organization, 2015a. Global action plan on antimicrobial resistance.

677 World Health Organization, 2015b. Worldwide country situation analysis: Response to antimicrobial
 678 resistance.

679 World Health Organization (Ed.), 2014. Antimicrobial resistance: global report on surveillance.
 680 World Health Organization, Geneva, Switzerland.

681 Yimdjo, M.C., Azebaze, A.G., Nkengfack, A.E., Meyer, A.M., Bodo, B., Fomum, Z.T., 2004.
 682 Antimicrobial and cytotoxic agents from *Calophyllum inophyllum*. *Phytochemistry* 65, 2789–
 683 2795.

684 Zhang, Y., Luo, Z., Wang, D., He, F., Li, D., 2014. Phytochemical profiles and antioxidant and
 685 antimicrobial activities of the leaves of *Zanthoxylum bungeanum*. *Scientific World Journal* 2014,
 686 1–13.

688 **Table 1.** The studied plants: type, habitat, traditional uses and biocidal activity for the genus.

Family		Traditional uses in medicine and ethnoveterinary practices		Biocidal activities known in the genus	
Botanical name	Status	Found in literature	Collected in this study	Against arthropods	Against microbes
Vernacular Créole name					
Plant type/habitat					
Anacardiaceae					
<i>Poupartia borbonica</i> J.F.Gmel.	End. R.	contraceptive, furuncle, nephritis	menopause, blood	nd	nd
Bois de poupart, Zévi marron	M. P.	(Lavergne, 2001; Smadja and Vera, 1991)	circulation disorders, fungi, insects		
Shrub/semi-dry forests					
Apocynaceae					
<i>Carissa spinarum</i> L.	Ind. P.	skin disorders, wounds, gonorrhoea, stomach disorders, tonic, nephritis.	typhoid fever, fever	<i>C. edulis</i> (Nyahangare et al., 2015)	<i>C. lanceolata</i> (Lindsay et al., 2000)
Bois amer		(Poullain et al., 2004; Vera et al., 1990)			<i>C. spinarum</i> (Sanwal and Chaudhary, 2011).
Shrub/semi-dry forests					
<i>Secamone volubilis</i> (Lam.) Marais	End. R.	hernia, diarrhoea, fever, diabetes, hypertension.	cancer	<i>S. afzelli</i> (Adesina et al., 2012)	nd
Liane bois d'olive	M.	(Poullain et al., 2004; Smadja and Vera, 1991)			
Liana/semi-dry forests					
Asteraceae					
<i>Psiadia amygdalina</i> Cordem.	End. R	nd	nd	nd	<i>P. trinervia</i> (Wang et al., 1989) <i>P. arguta</i> , <i>P. lithospermifolia</i> , <i>P. penninervia</i> , <i>P. terebinthina</i> , <i>P. viscosa</i> (Govinden-Soulangue et al., 2004) nd
Bois collant, Ti mangue					
Shrub/cloud forests					
<i>Psiadia boivinii</i> B.L.Rob.	End. R	nd	nd	nd	
Bouillon blanc					
Shrub/dense cloud forests					
<i>Psiadia dentata</i> DC.	End. R	nd	skin disorders (mycoses), insects	nd	<i>P. trinervia</i> (Wang et al., 1989) <i>P. arguta</i> , <i>P. lithospermifolia</i> , <i>P. penninervia</i> , <i>P. terebinthina</i> , <i>P. viscosa</i> (Govinden-Soulangue et al., 2004) nd
Bois collant, Ti mangue					
Shrub/dense cloud forests					
<i>Psiadia laurifolia</i> Cordem.	End. R.	nd	nd	nd	<i>P. trinervia</i> (Wang et al., 1989) <i>P. arguta</i> , <i>P. lithospermifolia</i> , <i>P. penninervia</i> , <i>P. terebinthina</i> , <i>P. viscosa</i> (Govinden-Soulangue et al., 2004) nd
Bois de tabac, Bois de chenille					
Tree/dense cloud forests					
<i>Psiadia retusa</i> DC.	End. R.	nd	source of mineral salts	nd	<i>P. trinervia</i> (Wang et al., 1989) <i>P. arguta</i> , <i>P. lithospermifolia</i> , <i>P. penninervia</i> , <i>P. terebinthina</i> , <i>P. viscosa</i> (Govinden-Soulangue et al., 2004) nd
Saliette	P.				
Shrub/rocky coastal areas					

End: endemic, Ind.: Indigenous, R: Reunion Island, M: Mauritius Island, Mas: Mascarenes, P: Protected species, nd: no data

<i>Vernonia fimbrillifera</i> Less. Bois de source Shrub/rainforests	End. R.	nd	blood circulation, cancer, wound healing	<i>V. phosphorea</i> (Valente et al., 2013) <i>V. amygdalina</i> (Mwanauta et al., 2014) <i>V. auriculifera</i> (Gemedi et al., 2014)	<i>V. colorata</i> (Rabe et al., 2002) <i>V. amygdalina</i> (Erasto et al., 2006) <i>V. glabra</i> (Kitonde et al., 2012) <i>V. guineensis</i> (Toyang et al., 2012)
Clusiaceae					
<i>Calophyllum tacamahaca</i> Willd. Takamaka Tree/lowland rainforests	End. R. M.	eye diseases, rheumatism, headache, gout, arthritis, dermic problems (Lavergne, 2001)	skin disorders, memory troubles, rheumatism, blood circulation	<i>C. inophyllum</i> (Ademola et al., 2014; Agrawal and Mall, 1988; Kadir et al., 2015)	<i>C. moonii</i> , <i>C. thwaitesii</i> (Dharmaratne et al., 1999) <i>C. inophyllum</i> (Yimdjo et al., 2004) <i>C. canum</i> (Alkhamaiseh et al., 2012) <i>C. antillanum</i> (Cuesta-Rubio et al., 2015)
Combretaceae					
<i>Terminalia bentzoe</i> (L.) Pers. Faux benjoin Tree/dry lowland forests	End. Mas.	fever, cold, cough, influenza, asthma, dysmenorrhoea, pleuritis paludism. (Lavergne, 2001; Poullain et al., 2004; Smadja and Vera, 1991)	reproductive disorders (spermatozooids), flu, bronchitis, cold	<i>T. catappa</i> (Rani et al., 2011)	<i>T. brachystemma</i> , <i>T. gazensis</i> , <i>T. mollis</i> , <i>T. prunioides</i> , <i>T. sambesiaca</i> , <i>T. sericea</i> , (Masoko et al., 2005)
Euphorbiaceae					
<i>Antidesma madagascariense</i> Lam. Bois de cabri Shrub/medium altitude forests	Ind.	nd	skin disorders, urine secretion	<i>A. bunius</i> (Belmi et al., 2014)	<i>A. thwaitesianum</i> (Dechayont et al., 2012) <i>A. venosum</i> (Mwangomo et al., 2012)
<i>Croton mauritanicus</i> Lam. Ti bois de senteur Shrub/coastal areas	End. R. P.	fever (Poullain et al., 2004; Vera et al., 1990)	fever, cold, muscle pains	<i>C. linearis</i> (Alexander et al., 1991) <i>C. argyrophyllodes</i> <i>C. nepetaefolius</i> , <i>C. sonderianus</i> <i>zehntneri</i> (Lima et al., 2013)	<i>C. urucurana</i> (Peres et al., 1997) <i>C. megalobotrys</i> (Selowa et al., 2010). <i>C. macrostachyus</i> (Obey et al., 2016)
<i>Stillingia lineata</i> (Lam.) Müll.Arg. Bois de lait Tanguin de pays Tree/lowland dry forests	Ind. P.	nd	chikungunya virus, furuncles	nd	nd
Fabaceae					
<i>Indigofera amoxylum</i> (DC.) Polhill Bois de sable, Bois de rose Tree/steep gorges	End. R. P.	nd	hypercholesterolemia, diabetes	<i>I. tinctoria</i> (Kamal and Mangla, 1993)	<i>I. oblongifolia</i> (Dahot, 1999) <i>I. suffruticosa</i> (Leite et al., 2006)

End: endemic, Ind.: Indigenous, R: Reunion Island, M: Mauritius Island, Mas: Mascarenes, P: Protected species, nd: no data.

<i>Sophora denudata</i> Bory Petit tamarin des hauts Tree/high-altitude forests	End. R. M.	skin cancer (Poullain et al., 2004; Vera et al., 1990)	skin disorders (psoriasis, eczema)	<i>S. flavescens</i> (Mao and Henderson, 2007)	<i>S. alopecuroides</i> (Küçükboyacı et al., 2011) <i>S. oppositifolia</i> (Cota et al., 2011) <i>S. exigua</i> , <i>S. flavescens</i> (Krishna et al., 2012)
Monimiaceae					
<i>Monimia rotundifolia</i> Thouars Mapou Tree/lowland rainforests	End. R.	nd	nd	nd	nd
Piperaceae					
<i>Peperomia borbonensis</i> Miq. Pourpier Epiphyte succulent/highland rainforests	End. R.	nd	ticks	nd	<i>P. villipetiola</i> (Malquichagua Salazar et al., 2005) <i>P. fernandopoina</i> (Mbah et al., 2012; Ngemenya et al., 2006) <i>P. Pellucida</i> (Akinnibosun et al., 2008; Khan and Omoloso, 2002; Oloyede et al., 2011; Wei et al., 2011)
Rutaceae					
<i>Zanthoxylum heterophyllum</i> (Lam.) Sm. Poivrier des hauts Tree/rain/semi-dry forests	End. Mas. P.	back pain, toxins, fever (Lavergne, 2001; Poullain et al., 2004)	tooth lesion, local anaesthetic (mooth)	<i>Z. caribaeum</i> (Nogueira et al., 2014) <i>Z. dissitum</i> (Wang et al., 2015) <i>Z. heitzii</i> (Moussavi et al., 2015)	<i>Z. budrungea</i> (Islam et al., 2001) <i>Z. chalybeum</i> (Olila et al., 2001) <i>Z. zanthoxyloides</i> , <i>Z. leprieurii</i> (Misra et al., 2013) <i>Z. bungeanum</i> (Zhang et al., 2014)
Stillbaceae					
<i>Nuxia verticillata</i> Lam. Bois maigre Tree/medium altitude forests	End. R. M.	toxins, albuminuria, venereal diseases, intestinal transit disorders. (Jonville et al., 2011; Lavergne, 2001; Poullain et al., 2004; Smadja and Vera, 1991)	hypercholesterolemia, stomach problems, malaria, urine secretion, albuminuria	nd	nd

End: endemic, Ind.: Indigenous, R: Reunion Island, M: Mauritius Island, Mas: Mascarenes, P: Protected species, nd: no data.

689

690

691

692 **Table 2.** Antimicrobial activity of ethyl acetate extracts of 16 plants from Reunion Island.
693

Plants	Parts used ^a	Gram-negative bacteria			Gram-positive bacteria			Fungi
		<i>Salmonella enterica</i>	<i>Pseudomonas aeruginosa</i>	<i>Escherichia coli</i>	<i>Streptococcus pyogenes</i>	<i>Listeria monocytogenes</i>	<i>Staphylococcus aureus</i>	<i>Aspergillus fumigatus</i>
<i>Antidesma madagascariense</i>	B	-	-	-	-	-	13.15 ± 0.21	-
<i>Callophylum tacamahaca</i>	L	-	-	-	-	-	10.20 ± 0.28	-
<i>Croton mauritianus</i>	L	-	-	-	-	-	10.25 ± 1.18	-
<i>Monimia rotundifolia</i>	L	-	-	-	-	-	10.70 ± 0.23	-
<i>Nuxia verticillata</i>	L	-	-	-	-	9.67 ± 0.58	-	-
<i>Peperomia borbonensis</i>	L	8.77 ± 0.49	8.77 ± 0.40	9.87 ± 0.81	-	9.20 ± 1.11	-	7.97 ± 0.67
<i>Poupartia borbonica</i>	L	-	-	-	-	-	15.00 ± 0.60	-
<i>Psiadia amygdalina</i>	B	-	-	10.33 ± 0.58	-	-	-	-
<i>Psiadia boivinii</i>	L	-	-	-	-	-	10.67 ± 0.83	-
<i>Psiadia dentata</i>	L	-	9.87 ± 0.23	9.60 ± 0.40	-	9.70 ± 0.52	9.15 ± 1.20	-
<i>Psiadia retusa</i>	L	-	-	-	-	9.77 ± 0.38	10.50 ± 0.71	-
<i>Secamone volubilis</i>	L	-	-	-	-	-	11.70 ± 2.01	-
<i>Sophora denudata</i>	B	-	-	-	8.00 ± 0.00	10.27 ± 0.46	-	-
<i>Stillingia lineata</i>	L	-	-	-	-	-	10.15 ± 0.21	-
<i>Terminalia bentzoe</i>	L	-	-	-	11.70 ± 0.42	9.23 ± 0.06	8.80 ± 0.28	-
<i>Zanthoxylum heterophyllum</i>	B	-	-	-	-	-	9.70 ± 0.42	-
Chloramphenicol		31.30 ± 2.70	8.40 ± 0.53	21.60 ± 2.27	29.20 ± 0.81	29.57 ± 3.19	25.50 ± 0.81	-
Amphotericine B		-	-	-				10.67 ± 0.61

694
695 The growth inhibition was determined by paper disk diffusion. Inhibition diameters were given in mm ± standard deviation obtained in three
696 replicates, ‘-’ means not active. Chloramphenicol was used as standard for bacteria, amphotericine B was the standard used for fungi.

697 ^a L = leaves, B = bark.

698

699

700

701

702 **Table 3.** Minimal inhibitory concentration (MIC, µg/mL) of 16 plants extracts from Reunion Island.
703

Plants	Parts used ^a	Gram-negative bacteria			Gram-positive bacteria			Fungi
		<i>Salmonella enterica</i>	<i>Pseudomonas aeruginosa</i>	<i>Escherichia coli</i>	<i>Streptococcus pyogenes</i>	<i>Listeria monocytogenes</i>	<i>Staphylococcus aureus</i>	<i>Aspergillus fumigatus</i>
<i>Antidesma madagascariense</i>	B	-	-	-	-	-	125.00	-
<i>Callophylum tacamahaca</i>	L	-	-	-	-	-	62.50	-
<i>Croton mauritianus</i>	L	-	-	-	-	-	>1000.00	-
<i>Monimia rotundifolia</i>	L	-	-	-	-	-	1000.00	-
<i>Nuxia verticillata</i>	L	-	-	-	-	1000.00	-	-
<i>Peperomia borbonensis</i>	L	1000.00	1000.00	1000.00	-	1000.00	-	500.00
<i>Poupartia borbonica</i>	L	-	-	-	-	-	125	-
<i>Psiadia amygdalina</i>	B	-	-	>1000.00	-	-	-	-
<i>Psiadia boivinii</i>	L	-	-	-	-	-	>1000.00	-
<i>Psiadia dentata</i>	L	-	1000.00	1000.00	-	500.00	62.50	-
<i>Psiadia retusa</i>	L	-	-	-	-	500.00	125.00	-
<i>Secamone volubilis</i>	L	-	-	-	-	-	>1000.00	-
<i>Sophora denudata</i>	B	-	-	-	15.62	125.00	-	-
<i>Stillingia lineata</i>	L	-	-	-	-	-	>1000.00	-
<i>Terminalia bentzoe</i>	L	-	-	-	1000.00	>1000	1000.00	-
<i>Zanthoxylum heterophyllum</i>	B	-	-	-	-	-	>1000.00	-
Chloramphenicol		18.75	150.00	37.50	4.68	nt ^b	37.50	-
Amphotericine B		-	-	-	-	-	-	0.50

704

705 The minimum inhibitory concentration (MIC) expressed in µg/mL was determined by broth dilution. Chloramphenicol was used as standard for
706 bacteria, amphotericine B was the standard used for fungi. Three replicates were made for all extracts tested. ‘-’ means not active.

707 ^a L = leaves, B = bark.

708 ^b nt = not tested

Figure captions

Figure 1. a) Map of the south-west Indian Ocean. b) The National Park and the repartition of the Reunion Island population.

The 24 municipalities are represented as circles. The three mountains cirques appear in italic font.

Figure 2. Percentage mortality for *Rhipicephalus microplus* larvae exposed to plant extracts (dilution 5%). The Standard Deviation of the mean values are represented by error bars.

Table captions

Table 1. The studied plants: type, habitat, traditional uses and biocidal activity for the genus.

Table 2. Antimicrobial activity of ethyl acetate extracts of 16 plants from Reunion Island.

The growth inhibition was determined by paper disk diffusion. Inhibition diameters were given in mm $\pm$ standard deviation obtained in three replicates, ‘-’ means not active. Chloramphenicol was used as standard for bacteria, amphotericine B was the standard used for fungi.

^a L = leaves, B = bark.

Table 3. Minimal inhibitory concentration (MIC, μ g/mL) of 16 plant extracts from Reunion Island.

The minimum inhibitory concentration (MIC) expressed in μ g/mL was determined by broth dilution. Chloramphenicol was used as standard for bacteria, amphotericine B was the standard used for fungi. Three replicates were made for all extracts tested. ‘-’ means not active.

^a L = leaves, B = bark.

^b nt = not tested



