

Comprehensive quantitative evaluation and mechanism analysis of influencing factors on yield and quality of cultivated *Gastrodia elata*

Xiaojun Jiang

Sichuan College of Traditional Chinese Medicine

Jing Dai

Sichuan College of Traditional Chinese Medicine

Yi Huang

Southwest University of Science and Technology

Dan Zhao

Mianyang Academy of Agricultural Sciences

Yao Yin

Sichuan College of Traditional Chinese Medicine

Qiao Lin

Sichuan College of Traditional Chinese Medicine

Mengmeng Hou

China Academy of Chinese Medical Sciences

Hong Jin

jh1120@139.com

Sichuan College of Traditional Chinese Medicine

Article

Keywords: *Gastrodia elata*, yield, quality, influencing factors, mechanism

Posted Date: November 14th, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-5289918/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on May 27th, 2025. See the published version at <https://doi.org/10.1038/s41598-025-99706-1>.

Abstract

Gastrodia elata (*G. elata*) is a dual-purpose herb for medicine and food. Wild resources are depleted, and there is a significant decrease in yield or quality when they are cultivated artificially. However, what factors led to the decline is still unclear. In this study, based on comprehensive data under multiple production regions, hierarchical partitioning and partial least squares path modeling were used for the first time to quantitatively evaluate the dominant influencing factors and mechanism for the yield and quality of cultivated *G. elata*. The results showed that *Armillaria* relative abundance, soil available potassium and temperature seasonality were the key factors. Their independent effects were 74.14%, 24.78%, and 20.36% on yield, and 36.83%, 25.63%, and 21.30% on quality, respectively. Plant subspecies directly determined the yield and quality ($P < 0.01$). Soil physical properties affected chemical properties, which in turn affected biological properties and ultimately yields ($P < 0.05$). Meanwhile, soil physical properties affected quality by influencing soil chemical properties ($P < 0.01$). Our study contributes novel insight to optimize cultivation strategies of *G. elata*, providing a more comprehensive knowledge on improving the yield and quality of *G. elata* and achieving large-scale greenhouse cultivation.

Introduction

G. elata is a perennial herb in Orchidaceae, with a wide geographical range in China, Japan, Korea, Bhutan, and India, which is both an important medicinal plant and a popular traditional food^[1]. Its dried tubers, *Gastrodiae Rhizoma* (also known as Tianma in Chinese), are often used as valuable traditional Chinese medicines^[2]. Modern pharmacological studies have proved that isolated components or crude extracts of *G. elata* possess neuroprotection, learning and memory improvement effects, cardioprotection, vasomodulatory effects, anti-depression, anti-cancer, and other activities^[3–6]. Due to immense medicinal value, high market demand, and over-excavation^[7], *G. elata* has been included in the list of rare and endangered plants, and cultivated *G. elata* has become the main source of obtaining them^[8]. However, there is a significant decrease in yield or quality when they are cultivated artificially^[9].

The yield and quality of *G. elata* are greatly influenced by genetic and environmental factors. As one of the heterotrophic plants without chlorophyll, the growth and development of *G. elata* require symbiosis with *Armillaria* and *Mycena*^[10–13]. Among them, the fungi belonging to the genus *Armillaria* imposed a significant impact on the yield of *G. elata*^[14]. At the same time, there are great differences in the content of active components in different plant subspecies^[15]. For instance, *G. elata* Bl. f. *viridis* was found to have a higher total sugar content than *G. elata* f. *glauca* and *G. elata* f. *elata*^[16]. Compared with the population from Yunnan Province, the population from Guizhou Province has a higher gastrodin (GAS) content^[17]. In addition, soil properties also affect the yield and quality of *G. elata*. Soil physical properties are the main aspects affecting soil chemical properties^[18], and they together emerge as key drivers of microbiome structure and microbial activities^[19]. Furthermore, altitude can affect the water content of fresh tubers of *G. elata*, which in turn affects its quality^[20]. Low temperature treatment can promote a stable production of immature rhizomes by breaking dormancy^[21]. When the temperature is 20–25°C, it

is beneficial to the development of protocorm in vitro in the achlorophyllous orchid *Gastrodia pubilabiata* Sawa^[22].

Although these findings emphasize their important roles in the formation of yield and quality of *G. elata* from single aspects such as plant materials, soil properties, and climatic factors, under actual cultivation conditions, the yield and quality of *G. elata* can be affected by both plant material, soil properties, and climatic factors^[23]. Moreover, their influences on the yield and quality of *G. elata* are extremely complex^[24]. However, few studies have linked plant materials, soil properties, and climatic factors to explore the independent effects as well as their interaction on the yield and quality of *G. elata*. Therefore, it is of great significance to determine the major factors affecting the yield and quality of *G. elata* and their mechanisms in order to optimize cultivation strategies of *G. elata*.

In this study, seven representative environments in China with prominent *G. elata* production were selected to collect *G. elata* tubers and soil. Based on the hierarchical partitioning (HP) and partial least squares path modeling (PLS-PM), plant subspecies, soil physical properties, soil chemical properties, soil biological properties, temperature, precipitation, and altitude were evaluated and analyzed simultaneously. The present paper not only comprehensively identified the significant factors affecting the yield and quality of *G. elata* for the first time, but also further clarified the mechanism of these influencing factors. The research findings could offer a theoretical foundation for the breeding of *G. elata*, and provide valuable insights for improving the yield and quality of *G. elata* and achieving large-scale greenhouse cultivation.

Materials and methods

Sample collection

Seven representative environments (named GSKX, GZDF, HBYL, SCPW, SXLY, YNYL, and YNZX) with prominent *G. elata* production (geo-authentic product region) were selected to collect healthy tubers at harvest (Table S1). Three sampling points were randomly selected in each environment. Removing the topsoil and peripheral soil using a shovel, the tubersphere soil was pooled into 50 mL sterile tubes from 1 cm away from the tuber. The soil was sieved through a 2 mm sieve after removing debris and stored at -80°C. The sample collection was discontinuously conducted between October and November 2023 (Table S1). Before the sample collection, there had been no rain in the environments for at least one week. A total of 21 *G. elata* samples and their soil were collected. Meanwhile, locality and GPS data of each sampling point were obtained (Table S1).

Evaluation of yield and quality

After completing the collection of soil and *G. elata* samples, all *G. elata* tubers were collected within 1 square meter of each sampling point and were weighed for assessment of the yield of *G. elata*. Fresh single *G. elata* were cleaned with water and steamed based on single tuber fresh weight (Table S2). After

steaming at atmospheric pressure, *G. elata* tubers were cut lengthwise into 2 mm thick slices and were dried at 50°C for 4 hours until the water content was less than 15%. Each *G. elata* sample was powdered and passed through a 0.355 mm sieve. Water content before high performance liquid chromatography (HPLC) analysis was measured after drying at 105°C for 10 min. The contents of GAS and p-hydroxybenzyl alcohol (HBA) were detected and calculated based on the methods in the Chinese Pharmacopoeia 2020^[25]. The standard samples were purchased from Sichuan Weikeyi Biological Technology Co., Ltd. (Chengdu, China).

G. elata subspecies identification and phylogenetic analysis

DNA was extracted from the arrow shoot of *G. elata*. Six DNA fragments were selected as DNA barcoding^[26–31], including ITS, ITS2, *rbcl*, *matK*, *psbA-trnH*, and *trnL-F* (Table S3). A final volume of 25 µL was amplified, containing 2 µL of genomic DNA, 12.5 µL of 2×Taq RCR Master Mix, 1.0 µL of each primer, and 8.5 µL of distilled deionized water. The amplicons of six DNA barcodings were sequenced bidirectionally on SANGER ABI3730XL (Tsingke Biotech Co., Ltd., Chengdu, China). The peak map was spliced by Contig Express version 3.0 (Codon Code Co., USA), and then the primer region and low region were removed. The hidden Markov model was used to remove the 5.8S and 28S segments to obtain the ITS2 sequences^[32]. The final ITS2 sequence has been transmitted to the GenBank database (<http://www.ncbi.nlm.nih.gov/genbank>), and all sequence names and the accession numbers were listed in Table S4. At the same time, other ITS2 sequences of *G. elata* were downloaded from the GenBank database. All ITS2 sequences were imported into MEGA 7.0 software. Based on the Kimura2-parameter model, the phylogenetic tree of *G. elata* subspecies was constructed by the neighbor-joining method, and the support rate of each branch was tested by bootstrapping 1000 iterations^[33].

Measurement of soil physical, chemical and biological properties

Soil physical properties are mainly soil particle size, which was measured using a laser particle size analyzer based on the previous report^[34]. The percentages of soil clay (< 0.002 mm), silt (0.002–0.05 mm), and sand (0.05–2 mm) were calculated based on the soil particle size following the United States Department of Agriculture (USDA) particle size classification system^[35]. In this study, the percentage of clay content was employed as an indicator of soil texture.

To measure soil chemical properties, soil samples were naturally dried, ground, and sieved through a 0.25 mm sieve. The soil organic matter was determined by the oven heating method^[36]. Soil pH was measured based on a soil-to-water ratio of 1:2.5 using a pH meter (F20, Shanghai Mettler-Toledo International Trading Co., Ltd., Shanghai, China). Total N, total P, total K, available N, available P, and available K in soil were determined by Sanshu Biotechnology Co., LTD. (Nantong, China) according to the previous literature^[37–38]. Soil moisture content was determined by the drying method, namely, soil samples were dried in an oven at 105°C until they reached a constant weight, and then SMC was calculated according to the weight change.

For soil biological properties, soil DNA was extracted with PowerSoil® DNA Isolation Kits (MOBIO Laboratories, Carlsbad, California, USA). Amplification and sequencing were performed at Beijing Tsingke Biotech Co., Ltd. (Beijing, China). The bacterial 16S rRNA gene V3–V4 region was amplified using universal primers 338F (5'–ACTCCTACGGGAGGCAGCA–3') and 806R (5'–GGACTACHVGGGTWTCTAAT–3'), and the fungal ITS1 region was amplified using universal primers ITS1F (5'–CTTGGTCATTTAGAGGAAGTAA–3') and ITS2R (5'–GCTGCGTTCTTCATCGATGC–3'). Sequencing of the V3–V4 and ITS1 regions was performed via Illumina NovaSeq 6000 PE250. Raw reads were filtered using Trimmomatic v0.33 software, and clean reads were obtained by identifying and removing primer sequences using Cutadapt v1.9.1 software. The amplicon sequence variants (ASVs) were obtained by de-noising using the DADA2 method in QIIME2 software^[39–40]. To conduct taxonomic annotation, the ribosome sequences V3–V4 were aligned to the sequences in the SILVA 138 database (<http://www.arbsilva.de>), and ITS1 were aligned to the sequences in the UNITE 8.0 database (<https://unite.ut.ee/>). Colony abundance indexes (Chao1 and Ace) and diversity indexes (Shannon and Simpson) were calculated based on ASVs.

Ecologically suitable analysis

By searching the related literature and major botanical specimen databases (Chinese Virtual Herbarium, <http://www.cvh.org.cn/>; Global Biodiversity Information Facility, <http://www.gbif.org/>; Royal Botanic Gardens, Kew, <http://www.kew.org/>; Discover Life, <http://www.discoverlife.org/>), the information was collected on the distribution points of cultivated *G. elata* and wild *G. elata* in China. Combining with the latitude and longitude obtained from the field surveys of the present study, all distribution points of *G. elata* were obtained. Using the Global Medicinal Plant Geographic Information System (GMPGIS)^[41], the ecological suitability of *G. elata* in China was studied to obtain the range of suitability values of its major ecological factors, major soil types, and the regional distribution map of its maximum ecological similarity in China.

Statistical analysis

The data of climatic factors were obtained from the WorldClim database (https://www.worldclim.org/data/bioclim.html#google_vignette), including 11 temperature parameters (bio1–bio11) and eight precipitation parameters (bio12–bio19)^[42]. All statistical analyses were performed using R version 4.0.2^[43]. The Shapiro-Wilk test was used to assess normality before testing for significance, and the homogeneity of variance was analyzed with an F-test. One-way analysis of variance (ANOVA) were conducted to assess the differences in active component content and soil properties in different environments. Pearson correlation was employed to measure the associations among active component content, soil properties, and climatic factors with the “corrplot” package in R^[44]. The HP was performed using the “glmm.hp” package in R^[45–46]. The PLS-PM framework was employed to examine the hypothesized causal connections among active component content, plant subspecies, soil properties, and climatic factors with the “plspm” package in R^[47]. A bootstrap method

(1000 iterations) was used to validate the estimates of the path coefficients and the coefficients of determination (R^2). Finally, eliminating loading less than 0.7, the goodness-of-fit index was used to evaluate the overall predictive performance of the model^[48]. Figures were drawn with the “ggplot2” package in R^[49].

Results

The yield and quality of *G. elata*

The morphological features of cultivated *G. elata* from different environments were depicted in Fig. S1. For the yield of *G. elata*, it was 13.22 kg/m², 5.12 kg/m², 11.80 kg/m², 5.33 kg/m², 14.00 kg/m², 6.64 kg/m², and 6.50 kg/m², respectively (Fig. 1). The yields of *G. elata* from GSKX, HBYL, and SXLY were significantly higher than GZDF, SCPW, YNYL, and YNZX ($P < 0.05$). The water content of all samples ranged from 6.97–14.73% before HPLC analysis (Table S5), which meets the requirements of the Chinese Pharmacopoeia 2020 of less than 15%. For the quality of *G. elata*, GAS content was 0.28%, 0.39%, 0.31%, 0.46%, 0.34%, 0.56%, and 0.49%, respectively (Fig. 1), which differed significantly between the seven environments ($P < 0.05$). The content of HBA is 0.10%, 0.08%, 0.01%, 0.09%, 0.03%, 0.08%, and 0.05%, respectively (Fig. 1). The content of HBA from HBYL, SXLY, and YNZX were significantly lower than GSKX, GZDF, SCPW, and YNYL ($P < 0.05$). The sum of GAS and HBA content was a minimum of 0.32% (HBYL) and a maximum of 0.66% (YNYL, Fig. 1). Similarly, their content met the requirements of the Chinese Pharmacopoeia 2020, that is, the sum of the contents is not less than 0.25%.

G. elata subspecies and phylogenetic tree

Six DNA markers (ITS, ITS2, rbcL, matK, psbA–trnH and trnL–F) were selected as DNA barcoding in this study. The PCR success rate ranged from 10–100% (Fig. S2). Among the six DNA barcodings, only the ITS2 was amplified with a success rate of 100%. Based on ITS2, all *G. elata* were categorized into two cultivated subspecies (Fig. 2), including *G. elata* Bl. f. *elata* (known as hongtianma in Chinese) and *G. elata* Bl. f. *glauca* (known as wutianma in Chinese). The phylogenetic tree of *G. elata* was also depicted in Fig. 2. Three *G. elata* Bl. f. *elata* from HBYL and eight sequences from the GenBank database were clustered into one group, and its bootstrap support was 100%. Six *G. elata* Bl. f. *elata* from SXLY and GSKX were clustered into one group, and its bootstrap support was 80%. Twelve *G. elata* Bl. f. *glauca* from SCPW, GZDF, YNZX, and YNYL were clustered into one group, and its bootstrap support was 96%.

Soil physical, chemical and biological properties

The soil texture classification based on USDA was depicted in Fig. 3. The highest soil clay content was 33.27% (GSKX), and the lowest is 1.86% (SXLY). The highest soil silt content was 47.17% (YNYL), and the lowest is 17.84% (SXLY). The highest soil sand content was 80.30% (SXLY), and the lowest is 20.77% (YNYL). The soil from SXLY and HBYL belonged to loamy sand. The soil from GZDF, SCPW, and YNZX belonged to loam. The soil from GSKX and YNYL belonged to clay loam.

Nine soil chemical components were detected in this study (Fig. 4), including soil moisture content (SMC), soil pH, soil organic matter (SOM), soil total nitrogen (STN), soil total phosphorus (STP), soil total potassium (STK), soil available nitrogen (SAN), soil available phosphorus (SAP), and soil available potassium (SAK). The results showed that SMC ranged from 10.5% (SXLY) to 39.6% (YNYL), soil pH ranged from 4.5 (SCPW) to 5.9 (SXLY), SOM ranged from 6.6 g/kg (HBYL) to 74.7 g/kg (YNZX), STN ranged from 0.4 g/kg (HBYL) to 3.4 g/kg (YNZX), STP ranged from 0.2 g/kg (SCPW) to 1.1 g/kg (SXLY), STK ranged from 7.0 g/kg (GZDF) to 23.6 g/kg (YNZX), SAN ranged from 29.6 mg/kg (HBYL) to 295.7 mg/kg (YNYL), SAP ranged from 0.2 mg/kg (GSKX) to 12.5 mg/kg (SXLY), and SAK ranged from 85.4 mg/kg (SCPW) to 282.6 mg/kg (GSKX), respectively. There were significant differences in each soil chemical composition among different environments ($P < 0.05$).

After merging of paired-end raw reads, their quality control, and filtration of chimeric sequences, a total of 969,195 active amplicon sequences were obtained belonging to the 16S rRNA region. Similarly, a total of 1,259,591 active sequences were obtained for the ITS1 region from 21 soil samples. At a threshold of 0.97, the sequences were clustered to 32,555 ASVs of bacteria and 11,772 ASVs of fungi. A total of 876,732 clean reads of bacteria belonging to 40 phylum and a total of 1,174,888 clean reads of fungi belonging to 19 phylum. The sparse curve of the ASVs gradually fattened out (Fig. S3), which indicated that the sequencing depth covered all the species in the sample.

The taxonomic annotation of ASVs was performed from phylum to species level. In the case of bacteria, Proteobacteria (33.6%) was the most dominant phylum, followed by Acidobacteriota (22.5%) and Bacteroidota (6.4%). Approximately 6.2% of OTUs remained unclassified (Fig. 5(a)). Unclassified bacteria (6.2%) was the dominant genus, followed by unclassified Xanthobacteraceae (3.5%) and unclassified Acidobacteriales (2.8%). Approximately 72.3% of OTUs belong to others (Fig. 5(b)). Similarly, in the case of fungi, Ascomycota (47.0%) was the most frequent phylum, followed by Basidiomycota (22.0%) and Mortierellomycota (12.6%). The unclassified ASVs accounted for up to 6.4% (Fig. 5(c)). Mortierella (4.8%) was the dominant genus, followed by unclassified Ascomycota (6.9%) and Fusarium (2.8%), with 14.4% remaining unidentified (Fig. 5(d)).

In order to clarify the effect of *Armillaria* and *Mycena* on the yield and active component content, their relative abundances were detected (Fig. S4). For *Armillaria*, the relative abundance was 0.78%, 0.08%, 0.66%, 0.16%, 0.94%, 0.30, and 0.24%, respectively ($P < 0.05$). For *Mycena*, only were detected in GZDF, HBYL, YNYL, and YNZX. Their relative abundances were 0.02%, 0.07%, 0.12%, and 0.01%, respectively ($P < 0.05$).

Colony abundance indexes (ACE index and Chao1 index) and diversity indexes (Shannon index and Simpson index) of bacteria and fungi were depicted in Fig. S5. From the result, in bacteria, the trends of ACE index and Chao1 index were completely consistent. Namely, GZDF, SXLY, and YNYL were significantly higher than SCPW and YNZX ($P < 0.05$). Meanwhile, GZDF and YNYL had significantly higher Shannon index and Simpson index than the other five environments ($P < 0.05$). In fungi, the ACE index

and Chao1 index were essentially the same as those of the bacteria. SCPW and YNZX had significantly lower Shannon index and Simpson index than the other environments ($P < 0.05$).

Ecologically suitable distributions

Using the GMPGIS, 337 distribution points of *G. elata* in China were located as far west as Tibet, as far east and north as Jilin, and as far south as Yunnan and Taiwan (Fig. S6). Based on the sampling points of *G. elata*, the key threshold values of the ecological factors of the species were extracted from the GMPGIS (Table S6). The biological range of annual mean temperature was -0.3°C – 21.4°C , the mean diurnal range was 6.5°C – 14.3°C , temperature seasonality was 2.8–13.4%, annual precipitation was 543.0 mm–2832.0 mm, and precipitation seasonality was 36.0–107.0%. The distribution map of maximum ecological similarity areas of *G. elata* in China mainly included Tibet, Sichuan Province, Xinjiang, Yunnan Province, Heilongjiang Province, Inner Mongolia, Qinghai Province, Guizhou Province, Shaanxi Province, etc. (Fig. 6). Especially in Sichuan Province, Yunnan Province, and Guizhou Province, almost the whole province was suitable for cultivation of *G. elata*.

Relationship among yield, quality, plant subspecies, soil properties and climatic factors

The yield of *G. elata* refers to the weight of all *G. elata* tubers per unit area, while the quality of *G. elata* refers to the weight proportion of active components in dried products. In *G. elata*, GAS and HBA are the main active components, especially GAS. Therefore, GAS content can represent the quality of *G. elata*. The HP and PLS-PM were performed using the yield and GAS content. The results of the HP were shown in Fig. 7. For the yield of *G. elata*, *Armillaria* relative abundance (ARA), SAK, and BIO4 were the most important influencing factors of soil biological properties, soil chemical properties, and climate factors, with the explained variation of 74.14%, 24.78%, and 20.36%, respectively (Fig. 7(a)). The R^2 for soil biological properties, soil chemical properties, and climate factors were 0.98, 0.74, and 0.97, respectively. When the first three important variables of soil biological properties, soil chemical properties and climate factors were selected, and plant subspecies (PS) and soil texture were added, ARA, PS, and BIO4 were the most important influencing factors, explaining yield variation of 21.22%, 19.41%, and 18.10%, respectively (Fig. 7(a)). The R^2 for the 11 most important variables was 0.97.

For GAS content, ARA, SAK, and BIO4 were also the most important influencing factors of soil biological properties, soil chemical properties, and climate factors, with the explained variation of 36.83%, 25.63%, and 21.30%, respectively (Fig. 7(b)). The R^2 for soil biological properties, soil chemical properties, and climate factors were 0.85, 0.90, and 0.99, respectively. When the first three important variables of soil biological properties, soil chemical properties and climate factors were selected, and PS and soil texture were added, PS, BIO4, and STN were the most important influencing factors, explaining the variation of 15.37%, 12.77%, and 11.74%, respectively (Fig. 7(b)). The R^2 for the 11 most important variables was 0.98.

Pearson correlation analysis revealed the relationship between yield, active components, soil physical properties, soil chemical properties, soil biological properties, and climatic factors using the 11 most important variables (Fig. 8). The results confirmed that yield was significantly and positively correlated with ARA, BIO4, BIO7 ($P < 0.001$), and STK ($P < 0.05$), but negatively correlated with ALT ($P < 0.001$) and ST ($P < 0.05$). GAS content was significantly and positively correlated with SAN, STN ($P < 0.001$), ALT ($P < 0.01$), and MRA ($P < 0.05$), but negatively correlated ($P < 0.05$) with ARA, BIO4, BIO7 ($P < 0.001$), and SAK ($P < 0.01$). In addition, ARA was significantly and positively correlated with BIO4 and BIO7 ($P < 0.001$), but negatively correlated with ALT ($P < 0.001$). For MRA, it was significantly and positively correlated with FSI ($P < 0.05$).

The constructed PLS-PM models can explain 98% and 92% of the variability in yield and GAS content, respectively (Fig. 9(a) and (b)). Regarding the PLS-PM model of the yield of *G. elata* (Fig. 9(a)), plant subspecies and soil biological properties had a direct positive effect, with the path coefficients of 0.62 ($P < 0.01$) and 0.64 ($P < 0.01$), respectively. Soil physical properties had a direct positive effect on soil chemical properties, with a path coefficient of 0.68 ($P < 0.001$). Meanwhile, soil chemical properties had a direct positive impact on soil biological properties, with a path coefficient of 0.47 ($P < 0.05$). The goodness-of-fit of the yield of *G. elata* was 0.70. Regarding the PLS-PM model of the GAS content of *G. elata* (Fig. 9(b)), it revealed plant subspecies and soil chemical properties had a direct positive effect, with the path coefficients of 0.95 ($P < 0.001$) and 0.56 ($P < 0.01$), respectively. In addition, soil physical properties had a direct positive effect on soil chemical properties, with a path coefficient of 0.71 ($P < 0.001$). The goodness-of-fit of the GAS content of *G. elata* was 0.72.

Indirect effects and total effects were shown in Fig. 9(c) and (d). Soil physical properties had an indirect effect (0.32) on soil biological properties, leading to a total effect of 0.48. Soil physical properties had also an indirect effect (0.33) on the yield of *G. elata*, leading to a total effect of 0.39. Moreover, soil chemical properties had an indirect effect (0.30) on the yield of *G. elata*, leading to a total effect of 0.33. For the PLS-PM model of the GAS content of *G. elata*, soil physical properties had an indirect effect (-0.11) on soil biological properties, leading to a total effect of -0.48. Soil physical properties had also an indirect effect (0.26) on the GAS content of *G. elata*, leading to a total effect of 0.15. Soil chemical properties had an indirect effect (-0.04) on the GAS content of *G. elata*, leading to a total effect of 0.52.

Discussion

The establishment conditions and quantitative process for the models for the relationship among yield, quality and influencing factors

Previous studies have mostly explored the influence of soil properties or climate factors on the yield and quality of *G. elata*, ignoring the effects of different plants^[50–51]. However, the effect of plant subspecies is usually greater than other factors^[52]. Therefore, incorporating plant subspecies into the influencing factors and analyzing them at the same time will be closer to the actual production, which will be more scientific and practical. In order to identify *G. elata* subspecies, the DNA barcoding technique was used,

which created advantageous conditions for analyzing the relationship among plant subspecies, yield, and quality. In the past, the authentication of *G. elata* was mainly based on its phenotypic characteristics or personal experience^[53], while DNA barcoding has become an extremely widely used technology in molecular marker-based species authentication technologies, given its standardization, minimization, and scalability^[54]. In particular, ITS2 could be identified at the subspecies taxonomic unit and had the fastest growth in the number of major plant DNA barcoding sequences in the Genbank database^[55]. However, until now, the identification and taxonomy of *G. elata* subspecies in only two provinces (Yunnan and Hubei) have been performed by SSR markers, SNP markers, and SLAF sequencing^[56–57]. Our results showed that *G. elata* from seven provinces in China's core producing regions was categorized into *G. elata* Bl. f. *elata* (known as hongtianma in Chinese) and *G. elata* Bl. f. *glauca* (known as wutianma in Chinese) cultivated subspecies based on ITS2 (Fig. 2). There were significant differences in yield and quality of *G. elata* among different plant subspecies. The classification result from Yunnan Province and Hubei Province is in line with previous reports^[56–57]. Moreover, *G. elata* subspecies also showed its geographical distribution characteristics. Specifically, *G. elata* from Yunnan, Guizhou and Sichuan were one type, while those from Hubei, Shaanxi and Gansu were another type. These results are consistent with the history of artificial cultivation of *G. elata*^[14].

Quantifying the influences of plant subspecies, soil properties, and climate factors on the yield and quality of *G. elata* is a premise for determining key influencing factors and optimizing cultivation strategies. The application of HP and PLS-PM achieved a quantitative evaluation process for the influences of these influencing factors on the yield and quality of *G. elata*. Past studies have reported that the yield and quality of *G. elata* were significantly correlated with soil properties and climatic factors^[21, 51], but few studies quantitatively evaluated the influence because the multicollinearity caused by strong correlations among soil properties or climatic factors usually resulted in the errors of evaluation (Fig. 8). Fortunately, the emergence of HP successfully solved the problem, which has become an important statistical method to analyze the independent effects of the predictor variables on the response variable^[45–46]. In this study, HP was used to preliminarily quantify the independent effects of plant subspecies, soil properties, and climatic factors on the yield and quality of *G. elata* separately (Fig. 7).

Although HP could identify the most important soil properties and climatic factors affecting the yield and quality of *G. elata*, it ignored the direct and indirect effects between different types of environmental factors. Comparing with HP, the structural equation model (SEM) could directly quantify the causalities between multiple variables, but a large number of variables usually resulted in a low degree of freedom and accuracy of SEM^[58]. At present, SEM mainly includes structural equation model based on maximum likelihood estimation and structural equation model based on partial least squares estimation. The latter representative model is PLS-PM, which has lower requirements for samples and is more suitable for comprehensive analysis of a newly constructed structure^[47]. Therefore, based on the preliminary results of HP, the PLS-PM was successfully conducted to analyze the interaction patterns between plant subspecies, key soil properties and climatic factors, and quantify their comprehensive influences on the

yield and quality of *G. elata* in this study (Fig. 9). These results could identify the most important influencing factors for the yield and quality of *G. elata* and further clarify their mechanisms, which were also a premise for improving the yield and quality of *G. elata*.

The differences in the effects of *Armillaria* and *Mycena* on the yield and quality of *G. elata*

Unlike other plants, *G. elata* is one of the heterotrophic plants without chlorophyll. Its growth and development require symbiosis with *Armillaria* and *Mycena*^[10–13]. *Armillaria* play important roles in the nutrient supply and growth modulation of *G. elata*, and they will undergo severe competition with native soil organisms before colonization and become symbiotic with *G. elata*^[59]. For *Mycena*, because seeds of *G. elata* have no endosperm, their sexual reproduction depends on the nutrients provided by the embryo cells infected by *Mycena* to complete seed germination^[60–61]. Our results showed that *Armillaria* could be detected in all soil environments at harvest, while *Mycena* could only be detected in four soil environments (Fig. S4). This result was consistent with their role in the growth and development of *G. elata*^[59–61]. Namely, the main role of *Mycena* was to promote the germination of *G. elata* seeds in the early stage, while the main role of *Armillaria* was to promote the growth of *G. elata* tubers in the later stage.

The ARA was the most important factor in soil biological properties based on HP, which explained variation of 74.14% for the yield of *G. elata* (Fig. 7(a)), which was approximately three times the sum of all other soil biological factors. Meanwhile, ARA was also extremely correlated with the yield of *G. elata*, whose correlation coefficient was 0.98 ($P < 0.001$, Fig. 8(a)), and ARA had a direct positive effect on yield with the path coefficients of 0.64 ($P < 0.01$). Compared with the effect of *Armillaria* on yield, its independent effect and correlation on GAS content were lower. Even ARA and GAS showed a negative correlation ($P < 0.001$, Fig. 8(b)), because the higher the yield of *G. elata*, the lower the GAS content. Based on this, it can be inferred that the main role of *Armillaria* was to increase the yield of *G. elata* rather than quality, which is the same as previous studies^[14]. Interestingly, our study also found that the soil environment of high-quality *G. elata* contained more *Mycena*. HP analysis also showed that MRA was the second important variable (explained variation of 34.45%, Fig. 7(b)), and correlation analysis showed that MRA and GAS content had significant correlation ($P < 0.05$, Fig. 8(b)). Although *Mycena* cannot directly improve the GAS content of *G. elata*, they may promote GAS synthesis by enhancing other beneficial microorganisms or soil nutrients^[51, 59, 62]. These results showed the importance and difference of soil microorganisms on the yield and quality formation of *G. elata*, especially excellent *Armillaria* and *Mycena* should be selected in the cultivation of *G. elata*.

The main mechanism affecting the yield and quality of *G. elata* and the direction of optimizing cultivation strategies

Combining the above analysis, the mechanism of influencing factors to the yield of *G. elata* could be summarized as two aspects: direct effect and indirect effect (Fig. 9(a)). On the one hand, plant subspecies (mainly *G. elata* Bl. f. *elata*) as the main factor directly determined the yield of *G. elata* from

the genetic. On the other hand, soil physical properties affected chemical properties, which in turn affected biological properties and ultimately yields. Among them, the ARA was the most important soil biological factor, while STP and STK were the most important soil chemical factors for the yield of *G. elata*. Similarly, the mechanism of influencing factors to the quality of *G. elata* could also be summarized as two aspects (Fig. 9(b)). On the one hand, plant subspecies (mainly *G. elata* Bl. f. *glauca*) as the main factor directly determined the quality of *G. elata* from the genetic. On the other hand, soil physical properties affected quality by influencing soil chemical properties. Among them, SAN and STN were the most important soil chemical factors for the quality of *G. elata*. To sum up, the specific regulation direction for improving the yield and quality of *G. elata* should include as follows: (1) introduce superior varieties for cultivation, which is the dominant factor that determines the yield and quality; (2) select the soil texture with higher clay content, such as loam or clay loam; (3) pay attention to the improvement of soil organic matter, because it is closely related to soil N, P, and K, and the cultivation of *G. elata* cannot use any fertilizer; (4) cultivate excellent *Armillaria* and *Mycena* strains, which play an important role in the seeds germination and growth of *G. elata*; (5) cultivate *G. elata* in suitable ecological areas based on GMPGIS, which can ensure that the climate is fit for their growth.

In general, the factors affecting the yield and quality of *G. elata* include plant subspecies, soil properties, environmental factors, processing methods, and tree species inoculated with *Armillaria*^[63–65]. However, it is difficult to quantify all factors because of the differences in processing methods and tree species inoculated with *Armillaria*. In this study, the models were only conducted using plant subspecies, soil properties, environmental factors, yield and quality, and selected the most important influencing factors. This study established a model for the first time between plant subspecies, soil properties, environmental factors, and the yield and quality of *G. elata*. In the future, more comprehensive mechanisms and models of plant subspecies, soil properties, environmental factors, processing methods, and tree species inoculated with *Armillaria* affecting the yield and quality of *G. elata* should be further studied continuously.

Declarations

Competing interests

The authors declare no competing interests.

Author Contribution

X.J.J.: data curation, writing-original draft preparation. H.J., D.Z.: writing-reviewing and editing. J.D., Y.H.: conceptualization and methodology. Y.Y., Q.L.: software, validation. H.J.: supervision. M.M.H.: investigation. All authors reviewed the manuscript.

Acknowledgement

This research was supported by the Science and Technology Achievement Transfer and Transformation Guidance Project (2022ZHXC0101) and the Key Discipline Construction Project of Traditional Chinese Medicine (2020ZD07).

Data Availability

Data is provided within the manuscript or supplementary information files.

References

1. Gong, M. Q. et al. Traditional uses, phytochemistry, pharmacology, applications, and quality control of *Gastrodia elata* Blume: A comprehensive review. *J. Ethnopharmacol.* **319**, 22. <https://doi.org/10.1016/j.jep.2023.117128> (2024).
2. Zhan, H. D. et al. The rhizome of *Gastrodia elata* Blume - An ethnopharmacological review. *J. Ethnopharmacol.* **189**, 361–385. <https://doi.org/10.1016/j.jep.2016.06.057> (2016).
3. Ye, P. et al. A small molecule 20C from *Gastrodia elata* inhibits α -synuclein aggregation and prevents progression of Parkinson's disease. *Cell. Death Dis.* **14**. <https://doi.org/10.1038/s41419-023-06116-0> (2023).
4. Chao-Qi, Z. et al. Sleep promoting and omics exploration on probiotics fermented *Gastrodia elata* Blume. *npj Sci. Food.* **8** <https://doi.org/10.1038/s41538-024-00277-8> (2024).
5. Huang, Y. J. et al. *Gastrodia elata* Blume water extract modulates neurotransmitters and alters the gut microbiota in a mild social defeat stress-induced depression mouse model. *Phytother Res.* **35**, 5133–5142. <https://doi.org/10.1002/ptr.7091> (2021).
6. Kim, N. H. et al. Antitumor and immunomodulatory effect of *Gastrodia elata* on colon cancer *In Vitro* and *In Vivo*. *Am. J. Chin. Med.* **45**, 319–335. <https://doi.org/10.1142/s0192415x17500203> (2017).
7. Liu, Y. & Huang, G. The chemical composition, pharmacological effects, clinical applications and market analysis of *Gastrodia elata*. *Pharm. Chem. J. +.* **51**, 211–215. <https://doi.org/10.1007/s11094-017-1584-5> (2017).
8. Yang, S. & Li, J. Effect of *Gastrodia elata* Bl cultivation under forest stands on runoff, erosion, and nutrient loss. *Forests.* **15** (7), 1127. <https://doi.org/10.3390/f15071127> (2024).
9. Wang, X. et al. Decrease in beneficial bacteria and increase in harmful bacteria in *Gastrodia* seedlings and their surrounding soil are mainly responsible for degradation of *Gastrodia* asexual propagation. *Front. Plant. Sci.* **15**. <https://doi.org/10.3389/fpls.2024.1334958> (2024).
10. Yuan, Y. et al. The *Gastrodia elata* genome provides insights into plant adaptation to heterotrophy. *Nat. Commun.* **9**. <https://doi.org/10.1038/s41467-018-03423-5> (2018).

11. Xu, Y. et al. A chromosome-scale *Gastrodia elata* genome and large-scale comparative genomic analysis indicate convergent evolution by gene loss in mycoheterotrophic and parasitic plants. *Plant. J.* **108**, 1609–1623. <https://doi.org/10.1111/tpj.15528> (2021).
12. Camprubi, A. et al. Plant performance and metabolomic profile of loquat in response to mycorrhizal inoculation, *Armillaria mellea* and their interaction. *Agronomy-Basel* **10**. (2020). <https://doi.org/10.3390/agronomy10060899>
13. Yu, E. et al. An exploration of mechanism of high quality and yield of *Gastrodia elata* Bl. f. *glauca* by the isolation, identification, and evaluation of *Mycena*. *Front. Microbiol.* **14**. <https://doi.org/10.3389/fmicb.2023.1220670> (2023).
14. Yu, E. et al. An exploration of mechanism of high quality and yield of *Gastrodia elata* Bl. f. *glauca* by the isolation, identification and evaluation of *Armillaria*. *BMC Plant. Biol.* **22**. <https://doi.org/10.1186/s12870-022-04007-8> (2022).
15. Chen, W. C. et al. Method development and validation for the high-performance liquid chromatography assay of gastrodin in water extracts from different sources of *Gastrodia elata* Blume. *J. Food Drug Anal.* **23**, 803–810. <https://doi.org/10.1016/j.jfda.2015.02.006> (2015).
16. Ji, N., Liu, P., Zhang, N., Yang, S. & Zhang, M. Comparison on bioactivities and characteristics of polysaccharides from four varieties of *Gastrodia elata* Blume. *Front. Chem.* **10**. <https://doi.org/10.3389/fchem.2022.956724> (2022).
17. Tao, J. et al. Relationships among genetic makeup, active ingredient content, and place of origin of the medicinal plant *Gastrodia* tuber. *Biochem. Genet.* **47**, 8–18. <https://doi.org/10.1007/s10528-008-9201-7> (2009).
18. Yang, Y. et al. Biochar stability and impact on soil organic carbon mineralization depend on biochar processing, aging and soil clay content. *Soil. Biol. Biochem.* **169** <https://doi.org/10.1016/j.soilbio.2022.108657> (2022).
19. Philippot, L., Chenu, C., Kappler, A., Rillig, M. C. & Fierer, N. The interplay between microbial communities and soil properties. *Nat. Rev. Microbiol.* **22**, 226–239. <https://doi.org/10.1038/s41579-023-00980-5> (2024).
20. Wang, C. H. et al. Effects of *Armillaria* species on growth and quantity of active medicinal components of *G. elata* Bl. f. *elata* tubers along an altitude gradient: Evidence from empirical experiments. *Phytochem Lett.* **54**, 101–106. <https://doi.org/10.1016/j.phytol.2023.01.016> (2023).
21. Kim, C., Kim, D., Kim, H., Song, Y. & Lee, W. Production of immature rhizomes by breaking dormancy under low temperature of *Gastrodia elata* Blume. *Planta Med.* **82**. <https://doi.org/10.1055/s-0036-1596606> (2016).
22. Godo, T., Hashimoto, T., Nakata, M. & Miyoshi, K. The effects of illumination, temperature and 6-benzylaminoprine on asymbiotic seed germination and protocorm development in vitro in the achlorophyllous orchid *Gastrodia pubilabiata* Sawa. *Vitro Cell. Dev-Pl.* **56**, 230–235. <https://doi.org/10.1007/s11627-020-10061-4> (2020).

23. Li, Y., Kong, D., Fu, Y., Sussman, M. R. & Wu, H. The effect of developmental and environmental factors on secondary metabolites in medicinal plants. *Plant. Physiol. Biochem.* **148**, 80–89. <https://doi.org/10.1016/j.plaphy.2020.01.006> (2020).
24. Chen, L. et al. Dynamics of fungal communities during *Gastrodia elata* growth. *BMC Microbiol.* **19**. (2019). <https://doi.org/10.1186/s12866-019-1501-z>
25. Chinese Pharmacopoeia Committee. *Pharmacopoeia of the People's Republic of China* 59–60 (China Medical Science and Technology, 2020).
26. White, T. J. *Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR Protocols: A guide to methods and applications* 315–322 (Academic, 1990).
27. Gao, T. et al. Evaluating the feasibility of using candidate DNA barcodes in discriminating species of the large Asteraceae family. *BMC Evol. Biol.* **10**. (2010). <https://doi.org/10.1186/1471-2148-10-324>
28. Kress, W. J., Wurdack, K. J., Zimmer, E. A., Weigt, L. A. & Janzen, D. H. Use of DNA barcodes to identify flowering plants. *P Natl Acad. Sci. USA.* **102**, 8369–8374. <https://doi.org/10.1073/pnas.0503123102> (2005).
29. Hollingsworth, P. M. et al. A DNA barcode for land plants. *P Natl Acad. Sci. USA.* **106**, 12794–12797. <https://doi.org/10.1073/pnas.0905845106> (2009).
30. Sang, T., Crawford, D., Stuessy, T. & Chloroplast, D. N. A. phylogeny, reticulate evolution, and biogeography of *Paeonia* (Paeoniaceae). *Am. J. Bot.* **84**, 1120–1120. <https://doi.org/10.2307/2446155> (1997).
31. 38 Taberlet, P., Gielly, L., Pautou, G. & Bouvet, J. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant. Mol. Biol.* **17**, 1105–1109. <https://doi.org/10.1007/bf00037152> (1991).
32. Keller, A. et al. 5.8S-28S rRNA interaction and HMM-based ITS2 annotation. *Gene.* **430**, 50–57. <https://doi.org/10.1016/j.gene.2008.10.012> (2009).
33. Kumar, S., Stecher, G. & Tamura, K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* **33**, 1870–1874. <https://doi.org/10.1093/molbev/msw054> (2016).
34. Yang, X. et al. Determination of soil texture by laser diffraction method. *Soil. Sci. Soc. Am. J.* **79**, 1556–1566. <https://doi.org/10.2136/sssaj2015.04.0164> (2015).
35. Minasny, B. & McBratney, A. B. The Australian soil texture boomerang: a comparison of the Australian and USDA/FAO soil particle-size classification systems. *Soil. Res.* **39** (6), 1443–1451. <https://doi.org/10.1071/SR00065> (2001).
36. Su, J. et al. Soil conditions and the plant microbiome boost the accumulation of monoterpenes in the fruit of *Citrus reticulata* 'Chachi'. *Microbiome* **11**. (2023). <https://doi.org/10.1186/s40168-023-01504-2>
37. Muscarella, S. M. et al. Water reuse of treated domestic wastewater in agriculture: Effects on tomato plants, soil nutrient availability and microbial community structure. *Sci. total environ.* **928**, 172259–172259. <https://doi.org/10.1016/j.scitotenv.2024.172259> (2024).

38. Zhang, L. et al. NRT1.1B improves selenium concentrations in rice grains by facilitating selenomethionine translocation. *Plant. Biotechnol. J.* **17**, 1058–1068. <https://doi.org/10.1111/pbi.13037> (2019).
39. Bolyen, E. et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857. <https://doi.org/10.1038/s41587-019-0209-9> (2019).
40. Callahan, B. J. et al. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods.* **13**, 581–. <https://doi.org/10.1038/nmeth.3869> (2016).
41. Wu, J. et al. A new GIS model for ecologically suitable distributions of medicinal plants. *Chin. Med-UK.* **14**. <https://doi.org/10.1186/s13020-019-0226-0> (2019).
42. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* **37**, 4302–4315. <https://doi.org/10.1002/joc.5086> (2017).
43. R Core Team. R: A language and environment for statistical computing. Foundation for Statistical Computing, Vienna, Austria. (2013).
44. Wei, T. et al. Package ‘corrplot’. *Statistician.* **56** (316), e24 (2017).
45. Lai, J., Zou, Y., Zhang, S., Zhang, X. & Mao, L. glmm.hp: an R package for computing individual effect of predictors in generalized linear mixed models. *J. Plant. Ecol.* **15**, 1302–1307. <https://doi.org/10.1093/jpe/rtac096> (2022).
46. Lai, J., Zhu, W., Cui, D., Mao, L. & Liao, J. Extension of the glmm.hp package to zero-inflated generalized linear mixed models and multiple regression. *J. Plant. Ecol.* **16**. <https://doi.org/10.1093/jpe/rtad038> (2023).
47. Henseler, J., Ringle, C. M. & Sinkovics, R. R. *The use of partial least squares path modeling in international marketing* 277–319 (Emerald Group Publishing Limited, 2009).
48. Tang, S. et al. Microbial community succession in soil is mainly driven by carbon and nitrogen contents rather than phosphorus and sulphur contents. *Soil. Biol. Biochem.* **180** <https://doi.org/10.1016/j.soilbio.2023.109019> (2023).
49. Wickham, H. ggplot2. *Wires Comput. Stat.* **3** (2), 180–185. (2011). <https://doi.org/10.1002/wics.147>
50. Xie, X. et al. Changes in soil chemistry and microbial communities in rhizospheres of planted *Gastrodia elata* on a barren slope and under a forest. *Forests.* **15** <https://doi.org/10.3390/f15020331> (2024).
51. Xinyu, Y. et al. Discovery of seed germinating fungi (*Mycetinis scorodoni*) from *Gastrodia elata* Bl. f. *glauca* S. chow in Changbai Mountain and examination of their germination ability. *Sci. Rep-UK.* **14** <https://doi.org/10.1038/s41598-024-63189-3> (2024).
52. Luo, H. et al. Research progress on quality assurance of genuine Chinese medicinal in Sichuan. *Chin. Med-UK.* **16**. <https://doi.org/10.1186/s13020-021-00428-z> (2021).
53. Yang, P. et al. Rapid authentication of variants of *Gastrodia elata* Blume using near-infrared spectroscopy combined with chemometric methods. *J. Pharm. Biomed. Anal.* **235**. <https://doi.org/10.1016/j.jpba.2023.115592> (2023).

54. Ren, F. M. et al. DNA barcoding of *Corydalis*, the most taxonomically complicated genus of Papaveraceae. *Ecol. Evol.* **9**, 1934–1945. <https://doi.org/10.1002/ece3.4886> (2019).
55. Zhu, S., Liu, Q., Qiu, S., Dai, J. & Gao, X. DNA barcoding: an efficient technology to authenticate plant species of traditional Chinese medicine and recent advances. *Chin. Med-UK*. **17**. <https://doi.org/10.1186/s13020-022-00655-y> (2022).
56. Wang, Y., Shahid, M. Q., Ghouri, F., Ercisli, S. & Baloch, F. S. Development of EST-based SSR and SNP markers in *Gastrodia elata* (herbal medicine) by sequencing, de novo assembly and annotation of the transcriptome. *3 Biotech* **9**. (2019). <https://doi.org/10.1007/s13205-019-1823-4>
57. Xu, Z. et al. Genome-wide assessment of genetic variation and differentiation for *Gastrodia elata* germplasm based on SLAF sequencing. *Genet. Resour. Crop Evol.* **70**, 1971–1984. <https://doi.org/10.1007/s10722-023-01548-1> (2023).
58. Fan, Y. et al. Applications of structural equation modeling (SEM) in ecological studies: an updated review. *Ecol. Process.* **5**, 1–12. <https://doi.org/10.1186/s13717-016-0063-3> (2016).
59. Wang, Y. et al. Effect of symbiotic fungi-*Armillaria gallica* on the yield of *Gastrodia elata* Bl. and insight into the response of soil microbial community. *Front. Microbiol.* **14** <https://doi.org/10.3389/fmicb.2023.1233555> (2023).
60. Zeng, X. et al. Transcriptomic analyses reveal clathrin-mediated endocytosis involved in symbiotic seed germination of *Gastrodia elata*. *Bot. Stud.* **58**. <https://doi.org/10.1186/s40529-017-0185-7> (2017).
61. Li, Y. Y., Guo, S. X. & Lee, Y. I. Ultrastructural changes during the symbiotic seed germination of *Gastrodia elata* with fungi, with emphasis on the fungal colonization region. *Bot. Stud.* **61**. <https://doi.org/10.1186/s40529-019-0280-z> (2020).
62. Barazetti, A. R. et al. Formulations of arbuscular mycorrhizal fungi inoculum applied to soybean and corn plants under controlled and field conditions. *Appl. Soil. Ecol.* **142**, 25–33. <https://doi.org/10.1016/j.apsoil.2019.05.015> (2019).
63. Subramaniam, S., Wen, X. Y. & Jing, P. One-step microwave curing-dehydration of *Gastrodia elata* Blume: Relationship between phytochemicals, water states and morphometric characteristics. *Ind. Crop Prod.* **153**. <https://doi.org/10.1016/j.indcrop.2020.112579> (2020).
64. Park, E. J. & Lee, W. Y. Quantitative effects of various tree species on tuber growth and pharmacological compositions of *Gastrodia elata*. *Hortic. Environ. Biote.* **54**, 357–363. <https://doi.org/10.1007/s13580-013-0030-1> (2013).
65. Xie, Y. K. et al. Effects of steam and water blanching on drying characteristics, water distribution, microstructure, and bioactive components of *Gastrodia elata*. *Plants-Basel* **12**. (2023). <https://doi.org/10.3390/plants12061372>

Figures




Figure 1

The yield and content of active components in different environments.




Figure 2

Phylogenetic tree of cultivated *G. elata* using ITS2 based on neighbor joining.




Figure 3

The soil texture classification based on the United States Department of Agriculture.




Figure 4

The soil chemical properties in different environments.




Figure 5

The annotation of top ten relative abundance of bacteria and fungi.




Figure 6

The distribution map of maximum ecological similarity areas of *G. elata* in China based on the Global Medicinal Plant Geographic Information System.




Figure 7

Quantitatively evaluate the effects of plant subspecies, soil properties and climatic fac-tors on the yield and gastrodin content of cultivated *G. elata* based on hierarchical partitioning.




Figure 8

Pearson correlation analysis among yield, gastrodin content, soil properties and climatic factors.




Figure 9

The interaction relationship among yield, gastrodin content, plant subspecies, soil properties and climatic factors based on the PLS-PM model.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterial.docx](#)