

Genome-wide comparative analysis of the BAHD superfamily in seven Rosaceae species and expression analysis in pear (*Pyrus bretschneideri*)

Chunxin Liu

Nanjing Agricultural University

Xin Qiao

Nanjing Agricultural University

Qionghou Li

Nanjing Agricultural University

Weiwei Zeng

Nanjing Agricultural University

Shuwei Wei

Nanjing Agricultural University

Xin Wang

Nanjing Agricultural University

Yangyang Chen

Nanjing Agricultural University

Xiao Wu

Nanjing Agricultural University

Jun Wu

Nanjing Agricultural University

Hao Yin (✉ yinhao@njau.edu.cn)

Nanjing Agricultural University

Shaoling Zhang

Nanjing Agricultural University

Research article

Keywords: BAHD, pear, evolution, Rosaceae, transcriptome, volatile esters

Posted Date: November 5th, 2019

DOI: <https://doi.org/10.21203/rs.2.13258/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 8th, 2020. See the published version at <https://doi.org/10.1186/s12870-019-2230-z>.

Abstract

Background: The BAHD acyltransferase superfamily exhibits various biological roles in plants, including regulating fruit quality, catalytic synthesizing of terpene, phenolics and esters, and improving stress resistance. However, the copy numbers, expression characteristics and associations with fruit aroma formation of the BAHD genes remain unclear. **Results:** In total, 717 BAHD genes were obtained from the genomes of seven Rosaceae, (*Pyrus bretschneideri* , *Malus domestica* , *Prunus avium* , *Prunus persica* , *Fragaria vesca* , *Pyrus communis* and *Rubus occidentalis*). Based on the classifications in model plants, we divided the BAHD family genes into seven groups, I-a, I-b, II-a, II-b, III-a, IV and V. An intra-species synteny analysis detected 78 syntenic gene pairs among the seven Rosaceae species. Dispersed gene duplication occurred frequently in all the investigated species. Different modes of duplicated gene pairs identified in each investigated species revealed that the Ka/Ks ratios were less than one, indicating that they evolved through purifying selection. Based on the correlation analysis between the ester content and expression levels of BAHD genes at different developmental stages, we selected five genes for verification as assessed by qRT-PCR. Pbr020016.1 , Pbr019034.1 , Pbr014028.1 and Pbr029551.1 are important candidate genes involved in aroma formation during pear fruit development. **Conclusion:** We have thoroughly annotated the BAHD superfamily genes and performed a comprehensive comparative analysis of their colinearity, phylogenetic relationships and gene duplication patterns in seven Rosaceae species, and we also obtained four candidate genes involved in aroma synthesis in pear fruit. These results provide a theoretical basis for future studies of the specific biological functions of BAHD superfamily members and the improvement of pear fruit quality. **Keywords :** BAHD, pear, evolution, Rosaceae, transcriptome, volatile esters

Background

During growth, plants can produce not only common carbohydrates, fats and proteins, but also a multitude of secondary metabolites, such as phenolics, terpenoids, alkaloids, and miscellaneous compounds [1]. The diversity of these compounds results from hydroxylation, decarboxylation, glycosylation, oxidation/reduction, acylation and methylation, which expand the basic skeleton and modifying enzymes [2]. These secondary metabolites have many significant effects. Phenolic compounds are important dietary antioxidants that can prevent various oxidative stress-associated diseases [3]. The volatile chemicals disperse at ambient temperatures, playing different roles in organisms, and these secondary metabolites have long been used as flavor compounds in spices (fragrance) and preservatives in foods [4, 5]. Moreover, sugar content, fruit size, fruit aroma (fruit volatile organic compounds) are important indicators that affect fruit quality, and ester compounds are the key substances that determine fruit aromas [6]. Therefore, it is important to identify genes related to ester synthesis in pear fruit development.

The *BAHD* (benzylalcohol O-acetyl transferase, anthocyanin O-hydroxycinnamoyl transferase, N-hydroxycinnamoyl anthranilate benzoyl transferase and deacetylindoline 4-O-acetyltransferase [7]) superfamily is composed of enzymes having two common domains (HXXXD and DFGWG) and similar

amino acid sequences [8]. The HXXXD motif region is located in the reaction channel center and participates in catalysis. As an indispensable motif for the reaction, the DFGWG motif is located far from the active site [9]. Benzylalcohol O-acetyl transferase from the Californian wildflower *Clarkia breweri* can produce the floral volatile benzylacetate [10, 11], deacetylindoline 4-O-acetyltransferase from *Catharanthus roseus* is related to the synthesis of the alkaloid vindoline [10, 12], N-hydroxycinnamoyl anthranilate benzoyl transferase from *Dianthus caryophyllus* is responsible for producing a class of phytoalexins known as anthramides [13], and anthocyanin O-hydroxycinnamoyl transferases from *Gentiana triflora* can catalyze anthocyanin synthesis [14, 15]. *BAHD* superfamily genes play various roles during plant growth, such as the regulation of fruit quality improvement of stress resistance, as well as in the synthesis of plant phenolic compounds, anthocyanin, lignin and volatile esters [16–18]. In addition, several genes of this superfamily play vital roles in the synthesis of morphine, taxol and vindoline, which are precursors to anti-cancer drugs (vincalacoblastine and vincristine) [19]. Alcohol acyltransferase (*AAT*) from the *BAHD* superfamily plays roles in the synthesis of various plant secondary metabolites, such as volatile esters, anthocyanins and terpenes [20]. Furthermore, *AAT* is associated with the production of plant aroma compounds in apple, kiwi and strawberry [21–24]. Given the potential roles of *AAT* genes in fruit aroma biosynthesis, we aimed to identify the *BAHD* gene family on a genome-wide scale [20, 25].

Pear is an important temperate Rosaceae fruit tree worldwide. According to historical records, pears originated between 55 and 65 million years ago, and their cultivation history can be traced back for more than 30,000 years [25]. In 2012, the genome sequence of pear ('Dangshan sul', *Pyrus × bretschneideri*) was released, providing a resource for studying omics and molecular biology in pear. At present, genome sequences of six other Rosaceae species are also available, allowing comparative genomics studies among economically important Rosaceae species. In this study, we aimed to identify the *BAHD* superfamily members in the genomes of *P. bretschneideri* and six other Rosaceae species. To comprehensively identify and elaborate upon the *BAHD* superfamily, and unravel their functional roles in plant development, especially in fruit aroma biosynthesis, we investigated their evolutionary and expression patterns using genome and transcriptome resources. Gene structure, conserved motifs, phylogeny, selection pressure and spatiotemporal expression profiles of *BAHD* genes were analyzed in this study. Furthermore, we verified gene expression patterns by qRT-PCR, and several candidate *BAHD* genes that are closely associated with the pear volatile ester content and belong to Clades I and IV. These results provide insights into the evolution and expansion of the *BAHD* genes and will aid in further studies of their molecular functions in these fruit species.

Results

Identification of *BAHD* genes in seven Rosaceae species

The *BAHD* superfamily's characteristic domain (Pfam: PF02458) and the *BAHD* Hidden Markov Model (HMM) configuration file (PF02458) were used to identify the *BAHD* members. The online site SMART (<http://smart.embl-heidelberg.de/>) was used to analyze protein sequences of candidate genes and to

determine the presence of the *BAHD* domain. Ultimately, we obtained 773 putative *BAHD* family candidate genes in the seven species with E-values < 1e⁻¹⁰. Furthermore, a multiple sequence alignment was conducted to verify the presence of two characteristic conserved domains (HXXXD and DFGWG) in the *BAHD* family genes [8]. Because they lacked the two domains, three, six, six, sixseven, eight and 20 genes were removed from *Prunus persica* (peach), *P. bretschneideri* (Chinese white pear), *Pyrus communis* (European pear), *Rubus occidentalis* (black raspberry), *Fragaria vesca* (strawberry), *Malus × domestica* (apple) and *Prunus avium* (sweet cherry), respectively. Finally, 56 sequences were removed that lacked both domains. Ultimately, 717 *BAHD* genes were identified and analyzed (Table 1). Detailed information on the features of *BAHD* genes is available in Additional file 1: Table S1.

Table 1 Genomic information and identified *BAHD* gene numbers in Rosaceae species

Common name	Scientific name	Chromosome number	Release version	Genome gene number	Identified <i>BAHD</i> genes
Chinese white pear	<i>Pyrus bretschneideri</i>	34	NJAU, v1.0	42,341	114(120)
Apple	<i>Malus domestica</i>	34	JGI, v1.1	63,541	141(149)
Strawberry	<i>Fragaria vesca</i>	14	GDR, v4.0	32,831	89(96)
European pear	<i>Pyrus communis</i>	34	GDR, v2.0	37445	97(103)
Sweet cherry	<i>Prunus avium</i>	16	GDR, v1.0	43,679	125(145)
Peach	<i>Prunus persica</i>	16	JGI, v1.1	27,864	82(85)
Black raspberry	<i>Rubus occidentalis</i>	14	GDR, v3.0	33,286	69(75)

The database addresses were listed below: NJAU (<http://peargenome.njau.edu.cn/>); GDR (<http://www.rosaceae.org/>); JGI (<http://www.jgi.doe.gov/>); The numbers in parentheses show the count of genes before filtering for unanchored and missing conserved domain genes.

Phylogenetic and conserved motif analyses of *BAHD* genes in Chinese white pear

The amino acid sequences encoded by *BAHD* genes in Chinese white pear, European pear, *Arabidopsis* and *Populus* were used to construct a phylogenetic tree. The Chinese white pear *BAHD* family protein genes were classified into five clades (I, II, III-a, IV and V; Fig. 1). Clade I consisted of two subclades (clades I-a and I-b). The *Arabidopsis* genes belonging to clade I-a are involved in modifying aromatic and aliphatic alcohols in *Arabidopsis* and *Populus* [26, 27]; therefore, we speculated that the *BAHD* genes clustered into clade I-a in Chinese white pear might encode proteins with similar functions. The members of clade I-b had functions related to the biosynthesis of lignin monomeric intermediates [28, 29], including

tobacco and *Arabidopsis* shikimate hydroxycinnamoyltransferases [30]. Clade II consisted of two subclades (clades II-a and II-b). The functions of the *Arabidopsis* genes belonging to clade II-a are unknown. Clade II-b contained two *Arabidopsis* genes, *AT3G29590.1* (*At5MAT*) and *AT1G03940.1* (*At3AT1*), which are associated with anthocyanin biosynthesis [31, 32]. Clade III-a contained few members, including four genes from Chinese white pear, three genes from *Arabidopsis*, and one gene from poplar. Clade V contained nine members, including one well-studied *Arabidopsis* gene *AT4G24510.1* (*CER2*) involved in regulating the cuticular wax biosynthesis [3]. Clade IV contained many members involved in catalyzing the acetylation of aromatic alcohols and acetylating small- or medium-chain alcohols [27]. Additionally, only three and four members clustered in clades III-a and V. However, ~28.1% (32 of 114) *BAHD* genes were in clade I and 36.0% (41 of 114) were in clade IV, respectively. *BAHD* genes closely related to pear volatile ester contents, such as *AAT*, belonged to clades I and IV. The maximum-likelihood tree of *BAHDs* for European pear with *Arabidopsis* and *Populus* as outgroups was also constructed. (Additional file 2: Figure S1). Paralleling the phylogenetic tree built for Chinese white pear, the European pear's *BAHD* genes were divided into five clades. Clades V and III-a only contained four members, respectively. However, ~36.1% (35 of 97) genes clustered in clade IV, and 26.8% (26 of 97) clustered in clade I. Additionally, five function-known acyltransferases of *Arabidopsis* [*AT3G0480.1* (*CHAT*), *AT2G23510.1* (*SOT*), *AT5G48930.1* (*AtHCT*), *AT3G29590.1* (*At5MAT*) and *AT1G03490.1* (*At3AT1*)] were classified into two subgroups (I-a and II-b). These results provided putative candidates for the study of gene functions. We detected 20 conserved motifs in the *BAHD* proteins of Chinese white pear using the online software MEME (Fig. 2). All the *BAHD* family members contained motif1 or motif3, and ~65.8% (75 of 114) of members contained them both. Based on a gene structural analysis, we determined that motif1 and motif3, with sequences CGGFAIGLSMSHKVADGSSSLSTFINSWAE and FYEADFGWGKP, respectively, correspond to domains HXXXD and DFGWG, respectively. Members of subclades I-a, I-b, II-a, II-b, III-a and V do not contain motif17, except for *Pbr005916.1*, and members of the subclades I-a, I-b, II-a, II-b and III-a, as well as clade V, do not contain motif19, except for *Pbr010925.1*. Except for *Pbr005746.1*, *Pbr014025.1*, *Pbr035166.1* and *Pbr036245.1*, members of clade IV do not contain motif10. Motif14 and motif16 were only detected in the Clade II-b. The type and distribution of the conservative motifs of the same subclades were similar, further supporting the evolutionary tree's classification (Fig. 2). Information on conservative motifs is shown in Additional file 3: Table S2.

Gene duplication events identified in the pear *BAHD* superfamily and a *BAHD* collinearity analysis of seven Rosaceae species

Different patterns of gene replication have jointly promoted the evolution of the *BAHD* family, including whole-genome duplication (WGD) or segmental duplication, tandem duplication (TD), proximal duplication (PD), transposed duplication (TRD) and dispersed duplication (DSD) [34, 35]. We used DupGen_finder software [36] to detect duplicated *BAHD* family gene pairs in seven Rosaceae genomes. All the *BAHD* gene family members were assigned to WGD, PD, TD, TRD or DSD. The number of WGD duplications in Chinese white pear and apple were 29 and 59, respectively, but there were only three in strawberry and peach, four in black raspberry and sweet cherry, and 23 in European pear. The number of DSDs in Chinese white pear, European pear and sweet cherry were 113, 95 and 142, respectively.

Additionally, there were 91 in strawberry and 81 in apple, which are more than in peach (76) and black raspberry (70). Genomic rearrangements and gene loss may lead to the large proportion of DSDs in these species. Moreover, the RNA- and DNA-based TRD event can also produce this result [34]. WGDs and DSDs impacted the evolution of the *BAHD* superfamily in Chinese white pear, apple and European pear (Fig. 3). In peach and strawberry TDs and DSDs were the main forces, while PDs and DSDs played major roles in the evolution of black raspberry and sweet cherry. In pear, ~57.1% (113 of 198) *BAHD* genes were involved in DSD events, while there were 66.9% (91 of 136) in strawberry, 44.0% (81 of 184) in apple, 60.3% (76 of 126) in peach, 67.3% (70 of 104) in black raspberry, 66.4% (142 of 214) in sweet cherry and 55.2% (95 of 172) in European pear (Additional file 4: Table S3). The results indicated that DSDs were ubiquitous in all the investigated species.

In addition, we identified intra-genomic synteny blocks for each species [34]. As shown in Fig. 4a, the *BAHD* genes of Chinese white pear are randomly distributed on 17 chromosomes and there is only one gene on chromosome 13. Similarly, the *BAHD* genes were detected as randomly distributed in the other species. We found 78 syntenic gene pairs among the seven Rosaceae species. Of these, 17, 21, and 29 syntenic pairs were identified in European pear (Fig. 4g), Chinese white pear (Fig. 4a) and apple (Fig. 4b), compared with only three in strawberry (Fig. 4f), peach (Fig. 4d) and black raspberry (Fig. 4e) and two in sweet cherry (Fig. 4c) (Additional file 5: Table S4).

Nonsynonymous (Ka) and synonymous (Ks) substitutions per site, and a Ka/Ks analysis for *BAHD* family genes

The stage of evolution for the WGD is usually estimated using Ks [37–39]. In addition to the original WGD [Ks = 1.5–1.8, ~140 million years ago (Mya)] (denoted as a γ -paleohexaploidization event) that was shared by core eudicots [40], a more recent WGD was detected in pear and dated to 30–45 Mya (Ks = 0.15 to 0.3) [25]. As shown in Additional file 6: Table S5, Ks values of WGD-derived gene pairs in Chinese white pear ranged from 0.006 to 3.909, and the ranges of Ks values for gene pairs derived from TD, PD, TRD and DSD were 0.001–4.247, 0.07–3.670, 0.029–4.381 and 0.005–5.066, respectively. Similar results were found in apple. In Chinese white pear, there are nine WGD-derived genes pairs with Ks values that ranged from 0.15 to 0.30, demonstrating that they may be derived from the current WGD (30–45 Mya) [25]. Some other duplicated gene pairs possessed higher Ks values (1.992–3.909), implying that they probably originated from more ancient duplication events. The Ks values of the WGD-derived gene pairs in black raspberry, European pear, peach and sweet cherry were 1.356–2.965, 0.145–4.288, 1.416–4.357 and 1.469–4.210, respectively. The higher Ks values of WGD-derived gene pairs in peach, black raspberry and sweet cherry suggested that they were duplicated and retained from more ancient WGD events, supporting the absence of more recent WGD events in these species.

Deleterious mutations can be removed by negative selection (purifying selection). Conversely, new favorable mutations can be accumulated by positive selection (Darwinian selection) and spread through the population [6]. To detect the selection pressure acting on *BAHD* genes, we analyzed the Ka and Ka/Ks values in the seven Rosaceae species (Additional file 6: Table S5). The direction and magnitude of the

selection pressure were inferred based on Ka/Ks ratio (Ka/Ks > 1: positive selection; Ka/Ks = 1: neutral evolution; and Ka/Ks < 1: purifying selection) [42]. The Ka/Ks values of all the *BAHD* gene pairs in strawberry (Fig. 5d), peach (Fig. 5c) and European pear (Fig. 5b) were less than one, indicating that these genes evolved through purifying selection (Fig. 5). Similar results were found in the other four Rosaceae species [the Chinese pear (Fig. 5f), sweet cherry (Fig. 5a), black raspberry (Fig. 5g) and apple (Fig. 5e)], except for a few gene pairs with Ka/Ks values greater than one. The box plots also indicated that the data distributions were concentrated, especially in Chinese white pear, sweet cherry and apple.

Expression pattern of *BAHD* genes in Chinese white pear

Based on transcriptome data (Additional file 7: Table S6) from different pear tissues, we determined that most genes in Chinese white pear showed higher expression in roots (Fig. 6), and we discovered that 37 of the *BAHD* genes were expressed in all four stages of fruit development. *Pbr014238.1* was only expressed in the four stages of pollen-tube development, while *Pbr020016.1*, *Pbr027303.1*, *Pbr029551.1*, *Pbr014028.1* and *Pbr006821.1* were highly expressed in the late stage of fruit development (Fruit_S4). Most members of the *BAHD* superfamily showed no expression during the four stages of pollen-tube development.

Gene expression analyses with qRT-PCR

Based on the transcriptome expression profiles and the ester content analysis, we selected five potential Chinese white pear genes (*Pbr020016.1*, *Pbr019034.1*, *Pbr014028.1*, *Pbr006821.1* and *Pbr029551.1*) that showed strong correlations with total ester content changes during fruit development (Fig. 7f). We used qRT-PCR to examine these candidate genes. The expression patterns of several individual genes were highly correlated with the ester content changes during pear fruit development (Fig. 7). Our results indicated that the expression level of *Pbr014028.1* (Fig. 7c) decreased from S1 (45DAF) to S2 (75DAF), increased sharply from S3 (105DAF) to S4 (145DAF), and then reached a peak value. Surprisingly, three indices of *Pbr014028.1* (Fig. 7c), the relative expression level, the RNA-seq data and the changes in total ester content at all stages exhibited correlated trends. In addition, *Pbr019034.1* (Fig. 7a), *Pbr029551.1* (Fig. 7d) and *Pbr020016.1* (Fig. 7e) showed similar expression patterns. They each had a sharp increase from S3 (105DAF) to S4 (145DAF) and reached their peak value in the last period. Moreover, the relative expression levels of these genes and the changes in total ester contents at all the stages showed consistent trends. The expression pattern of *Pbr006821.1* (Fig. 7b) presented a different trend. It decreased first, then increased, reached a peak value in the S3 (105DAF) stage, and then decreased sharply from the S3 (105DAF) to S4 (145DAF) stage. However, the overall expression level in S4 (145DAF) was still greater than that in S1 (45DAF). Thus, except for *Pbr006821.1*, the expression levels of these genes, the RNA-seq data and the changes in total ester contents at all the stages exhibited correlated trends. Therefore, the four genes (*Pbr020016.1*, *Pbr019034.1*, *Pbr029551.1* and *Pbr014028.1*) appear to be important candidate genes for ester synthesis.

To further investigate this result, five *BAHD* genes (*Pbr020016.1*, *Pbr019034.1*, *Pbr014028.1*, *Pbr006821.1* and *Pbr029551.1*) from Chinese white pear and 11 biochemically characterized *AAT* genes from other

species were used to construct a maximum-likelihood tree. As seen in Fig. 8, we found that these pear *BAHD* genes shared a high homology with the reported *AAT* genes. Thus, we speculated that these genes might have a strong correlation with ester synthesis.

Discussion

The *BAHD* superfamily has members in various ester synthetic pathways. The functions of several *BAHD* genes have been reported [43–45]. Here, by detecting their HXXXD and DFGWG motifs using a HMMER search, we identified 717 *BAHD* genes from seven Rosaceae species. Similar to the results of the classification of *BAHD* genes in *Arabidopsis* [27], *BAHD* family protein-encoding genes of pear were classified into five clades (I, II, III-a, IV and V). In addition, a conserved motif analysis of these *BAHD* amino acid sequences showed that the type and distribution of the conservative motifs among the same subfamily were similar, indicating that the classification results are authentic. All the *BAHD* family members contained motif1 (HXXXD) or motif3 (DFGWG), and ~65.8% (75 of 114) of members contained them both. Thus, we hypothesized that the motifs (1 and 3) may be related to catalytic activity. The HXXXD catalytic motif is located in the solvent channel, and it can also deprotonize oxygen or nitrogen atoms on specific receptor substrates [8, 10]. The DFGWG motif is essential to the reaction but located far from the active site [8, 10, 19, 46, 47]. Combining the phylogenetic tree classification with the results of previous studies, we speculated that the *BAHD* genes that are closely related to the pear volatile ester content, such as *AAT*, belonged to clades I and IV, and the *AAT* are involved in the synthesis of esters, which are involved in determining the aromas of flowers and fruits [48]. At present, the functions of many *AAT* genes in fruit trees have been identified, such as *PpAAT1* in peach [49], *FaAAT2* and *SAAT* in cultivated strawberry [23, 24], *MpAAT1* in apple [9] and *CmAAT1,3,4* in melon [50, 51].

Gene duplication patterns can be generally divided into five types, WGD, PD, TD, DSD and TRD [34, 35]. Each pattern of gene duplication contributes differentially to gene family expansion [52]. It is estimated that 90% of the gene increase in the *Arabidopsis thaliana* pedigree is due to the result of WGDs [53]. WGD, TD and DSD are the main features of eukaryotic genome evolution and mainly drive the development of new functions in the genome and genetic evolutionary systems [54, 55]. Gene families, such as *SWEET* and Hydroxycinnamoyl transferases, expanded primarily through WGD and DSD [56, 57]. TD was the major force in the expansion of the *AP2/ERF* and *WRKY* gene families [58, 59]. Two gene families, Aluminum-activated malate transporters and Heat-shock transcription factors were amplified by WGD and DSD [34, 39]. The numbers of *BAHD* genes identified in seven plants from the rose family are different because of recent WGD events and ancient polyploid events. In this study, DSD and WGD were determined to be the main forces that expanded the *BAHD* genes in apple, European pear and Chinese white pear. The numbers of *BAHD* genes that had undergone WGD and DSD were the largest, while the numbers of genes that underwent other replication modes were relatively small. In addition, because peach, sweet cherry, strawberry and sweet cherry have not undergone recent WGD events, the number of DSDs account for a large proportion of the gene duplications. Accordingly, WGD patterns account for large proportions in apples and pears, perhaps because they have experienced two rounds of WGD [41]. In addition, based on the K_a , K_s and K_a/K_s analysis, we discovered that in peach, black raspberry and sweet

cherry the K_s values of WGD-derived gene pairs were greater than 1.3, suggesting that they were duplicated and retained from more ancient WGD events. This further supported the absence of more recent WGD events in these species. We also found that all the *BAHD* gene pairs in strawberry, peach and European pear had $K_a/K_s < 1$, indicating that they have experienced strong purifying selection.

The diversity of the biochemical functions of the *BAHD* genes has been determined in many species. For example, the *BAHD* acyltransferase gene can enhance saccharification [44], catalyzes the last step in cocaine biosynthesis [60], and the protein plays an important role in plant innate immunity [61]. Furthermore, *AAT* gene functions have been determined in several species. For instance, the *AAT* gene in apple, *MdAAT2*, may provide resistance to stress and can change the volatile compound profile [62]. In Mountain papaya (*Vasconcellea pubescens*), *VpAAT1* is involved in ester biosynthesis [63]. Despite the diverse functions of the *BAHD* gene family members, we focused on their roles in ester synthesis. Volatile esters produced by these *AAT* genes usually promote the recognition of plants as food because they contribute to the “fruity” aroma of edible fruits, and some esters are also the key flavor or smell component of a particular plant [21]. In this study, our RNA-seq data revealed that 37 *BAHD* genes were detected at various expression levels in the four stages of pear fruit development. Surprisingly, though lacking motif3 (DFGWG), the *Pbr019034.1* and *Pbr020016.1* genes were both still expressed at all the stages. Based on transcriptome expression profiles combined with the ester content analysis, five potential genes (*Pbr020016.1*, *Pbr019034.1*, *Pbr014028.1*, *Pbr006821.1* and *Pbr029551.1*) belonging to clades I and IV and that showed strong correlations with ester content changes during fruit ripening, were selected for further qRT-PCR analysis. The assessed expression patterns of four of the genes, *Pbr020016.1*, *Pbr029551.1*, *Pbr019034.1*, and *Pbr014028.1*, were basically consistent with the RNA-seq data. The peak of *Pbr006821.1* appeared in the third stage, and increased sharply from S2 (75DAF) to S3 (105DAF); therefore, we speculate that *Pbr006821.1* may be related to intermediate ester synthesis in the fruit.

Conclusions

Here, we identified 717 *BAHD* genes from seven species, and among them, 114 belonged to Chinese white pear. We divided the *BAHD* superfamily into five large clades according to the classification results from model plants. DSDs occurred frequently in all the investigated species. All the *BAHD* members of strawberry, peach and European pear experienced purifying selection. Similar results were found in the other four Rosaceae species (Chinese pear, sweet cherry, black raspberry and apple), except for a few gene pairs with K_a/K_s values greater than one. Finally, after a qRT-PCR analysis, we speculated that *Pbr020016.1*, *Pbr029551.1*, *Pbr019034.1* and *Pbr014028.1* are candidate genes related to ester synthesis. These results provide a foundation for further studies of the molecular mechanisms underlying aroma biosynthesis and release.

Methods

Source of Plants

The ‘Dangshan suli’ [25] fruit were picked at four different developmental stages from the pear germplasm orchard of the Center of Pear Engineering Technology Research located at Jiangpu in Nanjing. The fruit samples were collected on 26th May, 27th June, 28th July and 6th Sep in 2017. Previously published RNA-seq data was used to analyze the expression patterns of ‘Dangshan suli’ *BAHD* [64, 65]. All the samples were ground in liquid nitrogen and stored at -80°C . The genome annotation files and genome sequences of pear were collected from the Nanjing Agricultural University pear genome project website (<http://peargenome.njau.edu.cn>), and the other four Rosaceae species sequences (*Fragaria vesca*, *Pyrus communis*, *Prunus avium* and *Rubus occidentalis*) were downloaded from the Genome Database for Rosaceae (<http://www.rosaceae.org>). The sequences of apple (*M. domestica*) and peach (*P. persica*) were downloaded from Joint Genome Institute (<http://www.jgi.doe.gov/>).

Sequence identification and collection

To identify putative *BAHD* genes from pear and the six other species, peach, apple, strawberry, European pear, sweet cherry and black raspberry, several approaches were employed. Using a *BAHD* family characteristic domain (Pfam: PF02458) as a query sequence in accordance with the HMM configuration file (PF02458) of *BAHD*, we searched for candidate genes using the HMM with E-values $< 1e^{-10}$. We downloaded the HMM profile (PF02458) from the Pfam website (<https://pfam.xfam.org/>). The online site SMART (<http://smart.embl-heidelberg.de/>) was used to determine the domain PF02458 of *BAHD*, and then their HXXXD and DFGWG motifs were inspected. The online website MAFFT (<https://mafft.cbrc.jp/alignment/server/>) was used for multiple sequence alignments, and the comparison results were download to GeneDoc.

Phylogenic analysis

We use the online website MAFFT for multiple sequence alignments, and then upload the comparison results to IQ-TREE (<http://www.cibiv.at/software/iqtree>) [67]. The maximum-likelihood trees were constructed with bootstraps of 1,000 using IQ-TREE.

Conservative motif analysis of *BAHD* family members

The conservative motifs were analyzed using the online software MEME (http://meme.sdsc.edu/meme4_3_0/intro.html). The maximum value of the motif was set to 20, and the motif length was set between six and 200.

Identification of gene duplication modes and a collinearity analysis

A gene duplication analysis among seven Rosaceae genomes was performed, and the method developed by PGDD (<http://chibba.agtec.uga.edu/duplication/>) was carried out locally [68]. First, the chromosomal locational information of the *BAHD* gene family members and related gene pair information were obtained from the genome annotation files of each species, and then the localization and synteny of the

BAHD genes were determined using TB-tools software [69]. MCSscanX was further used to identify different duplication patterns in the *BAHD* superfamily [70].

Calculating Ka and Ks

Ka, Ks and the Ka/Ks ratio were calculated using the calculate_Ka_Ks_pipeline (https://github.com/qiaoxin/Scripts_for_GB/tree/master/calculate_Ka_Ks_pipeline) [36]. In brief, the coding sequences and gene pairs were prepared first. Then, we ran the computing_Ka_Ks_pipe.pl script to automatically perform multiple alignments using MAFFT software and to convert the AXT format, which was used as input for KaKs_Calculator [71], to calculate Ka and Ks using the GMYN model. Finally, the readable results, including Ka, Ks, Ka/Ks and P-value, were generated.

Expression analysis of *BAHD* in 15 samples of *P. bretschneideri* and a qRT-PCR analysis

The RNA-seq data of 15 ‘Dangshan suli’ samples were used to analyze the expression patterns of ‘Dangshan suli’ *BAHD*, including Fruit_S1, Fruit_S2, Fruit_S3, Fruit_S4 (obtained from the NCBI bioproject PRJNA563942), Root (data not shown), Stem, Leaf, Bud, Petal, Sepal, Ovary, Mature pollen, Hydrated pollen, Pollen tube and Stop growth pollen [64, 65]. Then, the gene expression heatmap was drawn using TB-Tools [69].

Based on the correlation analysis between transcriptome and ester content data of pear fruit during four developmental stages, we selected five possible AAT genes for qRT-PCR. The primers of all the possible genes were designed using Primer Premier 6.0. Total RNA was extracted using a Plant Total RNA Isolation Kit Plus (Fuji, China), and cDNA was synthesized using TransScript One-Step gDNA Removal and cDNA Synthesis SuperMix (TransGen, China). The PCR mixture was as follows: 1 µl of sense and anti-sense primer (10 µM), 5 µl fluorescent dye, 1 µl template and 3 µl sterilized water. qRT-PCR was performed using a LightCycler 480 SYBRGREEN I Master (Roche, USA). We ran the PCR reaction under the following conditions: 10 min at 95°C, followed by 45 cycles of 95°C for 3 s, 60°C for 10 s and 72°C for 30 s. Then, the $2^{-\Delta\Delta Ct}$ method was used to determine the relative expression with the reference genes being tubulin and WD-repeat protein (Additional file 8: Table S7).

Analysis of volatile aroma compounds

The SPME-GC method was used to determine volatile aromas from Chinese white pear. The concentrations of components (µg/g) = [peak area of component/peak area of internal standard × the concentration of internal standard (g/ml) × 5µl]/mass of sample (g). The concentration of the internal standard 3-nonanone was 0.82×10^{-3} g/ml [6].

List Of Abbreviations

WGD: Whole-genome duplication; TD: Tandem duplication; PD: Proximal duplication; TRD: Transposed duplication; DSD: Dispersed duplication; AAT: Alcohol acyltransferase; Mya: Million years ago; HMM:

Declarations

Acknowledgments

The authors would like to thank everyone who contributed to this article.

Funding

This study was supported by the National Key Research and Development Program [2018YFD1000200], the National Natural Science Foundation of China (31701890, 31601708), Fundamental Research Funds for the Central Universities (KJQN201818). These fundings provided the financial support to the research projects, but did not involve in project design, data collection, analysis, or preparation of the manuscript.

Authors' contributions

Conceived and designed the experiments: HY and SZ. Performed the experiments: CL, QL, XW, YC and JW. Analyzed the data: CL, QL, WZ, SW, HY and XQ. Wrote the paper: CL, XQ, HY and SZ. And all authors have read and approved the manuscript.

Ethics approval and consent to participate

All the pear materials used and analyzed for this study were collected from the Center of Pear Engineering Technology Research, Nanjing Agricultural University, which were public and available for non-commercial purpose. This article did not contain any studies with human participants or animals performed by any of the authors.

Availability of data and materials

The pear genome datasets used during the current study are available in our pear center website (<http://peargenome.njau.edu.cn/>), the other four Rosaceae species sequences (*Fragaria vesca*, *Pyrus communis*, *Prunus avium* and *Rubus occidentalis*) were downloaded from the GDR (<http://www.rosaceae.org>). The sequence of apple (*Malus domestica*) and peach (*Prunus persica*) were collected from JGI (<http://www.jgi.doe.gov/>). The RNA-seq data were obtained from the NCBI database (<https://www.ncbi.nlm.nih.gov/>)

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Song CK, Hartl K, McGraphery K, Hoffmann T, Schwab W. Attractive but Toxic: Emerging Roles of Glycosidically Bound Volatiles and Glycosyltransferases Involved in Their Formation. *Mol Plant*. 2018, 11(10):1225-1236.
2. Pichersky E, Noel JP, Dudareva N. Biosynthesis of plant volatiles: Nature's diversity and ingenuity. *Science*. 2006, 311(5762):808-811.
3. Legrand G, Delporte M, Khelifi C, Harant A, Vuylsteker C, Morchen M, Hance P, Hilbert JL, Gagneul D. Identification and Characterization of Five BAHD Acyltransferases Involved in Hydroxycinnamoyl Ester Metabolism in Chicory. *Front Plant Sci*. 2016, 7.
4. Dudareva N, Pichersky E, Gershenzon J. Biochemistry of plant volatiles. *Plant Physiol*. 2004, 135(4):1893-1902.
5. Schwab W, Davidovich-Rikanati R, Lewinsohn E: Biosynthesis of plant-derived flavor compounds. *Plant J*. 2008, 54(4):712-732.
6. Chen YY, Yin H, Wu X, Shi XJ, Qi KJ, Zhang SL. Comparative analysis of the volatile organic compounds in mature fruits of 12 Occidental pear (*Pyrus communis* L.) cultivars. *Sci Hortic-Amsterdam*. 2018, 240:239-248.
7. St-Pierre B, Luca VD. Chapter Nine Evolution of acyltransferase genes: Origin and diversification of the BAHD superfamily of acyltransferases involved in secondary metabolism. *Recent Advances in Phytochemistry*. 2000, 34(00):285-315.
8. Isabel M, Dylan K. Role of HXXXD-motif/BAHD acyltransferases in the biosynthesis of extracellular lipids. *Plant Cell Rep*. 2015, 34(4):587-601.
9. El-Sharkawy I, Manriquez D, Flores FB, Regad F, Bouzayen M, Latche A, Pech JC. Functional characterization of a melon alcohol acyl-transferase gene family involved in the biosynthesis of ester volatiles. Identification of the crucial role of a threonine residue for enzyme activity. *Plant Mol Biol*. 2005, 59(2):345-362.
10. D'Auria JC. Acyltransferases in plants: a good time to be BAHD. *Curr Opin Plant Biol*. 2006, 9(3):331-340.
11. Dudareva N, D'Auria JC, Nam KH, Raguso RA, Pichersky E. Acetyl-CoA : benzylalcohol acetyltransferase - an enzyme involved in floral scent production in *Clarkia breweri*. *Plant J*. 1998, 14(3):297-304.
12. St-Pierre B, Laflamme P, Alarco AM, De Luca V. The terminal O-acetyltransferase involved in vindoline biosynthesis defines a new class of proteins responsible for coenzyme A-dependent acyl transfer. *Plant J*. 1998, 14(6):703-713.
13. Yang Q, Reinhard K, Schiltz E, Matern U. Characterization and heterologous expression of hydroxycinnamoyl/benzoyl-CoA : anthranilate N-hydroxycinnamoyl/benzoyltransferase from elicited cell cultures of carnation, *Dianthus caryophyllus* L. *Plant Mol Biol*. 1997, 35(6):777-789.

14. Fujiwara H, Tanaka Y, Yonekura-Sakakibara K, Fukuchi-Mizutani M, et al. cDNA cloning, gene expression and subcellular localization of anthocyanin 5-aromatic acyltransferase from *Gentiana triflora*. *Plant J*. 1998, 16(4):421-431.
15. Fujiwara H, Tanaka Y, Fukui Y, Ashikari T, et al. Purification and characterization of anthocyanin 3-aromatic acyltransferase from *Perilla frutescens*. *Plant Sci*. 1998, 137(1):87-94.
16. Nevarez DM, Mengistu YA, Nawarathne IN, et al. An N-Aroyltransferase of the BAHD Superfamily Has Broad Aroyl CoA Specificity in Vitro with Analogues of N-Deaoylpaclitaxel. *J Am Chem Soc*. 2009, 131(16):5994-6002.
17. Bontpart T, Cheynier V, Ageorges A, et al. BAHD or SCPL acyltransferase? What a dilemma for acylation in the world of plant phenolic compounds. *New Phytol*. 2015, 208(3):695-707.
18. Gonzalez M, Gaete-Eastman C, Valdenegro M, et al. Aroma Development during Ripening of *Fragaria chiloensis* Fruit and Participation of an Alcohol Acyltransferase (FcAAT1) Gene. *J Agr Food Chem*. 2009, 57(19):9123-9132.
19. Ma XY, Koepke J, Panjekar S, et al. Crystal structure of vinorine synthase, the first representative of the BAHD superfamily. *J Biol Chem*. 2005, 280(14):13576-13583.
20. Srivastava S, Sangwan RS. Analysis of *Artemisia annua* transcriptome for BAHD alcohol acyltransferase genes: Identification and diversity of expression in leaf, stem and root. *J Plant Biochem Biot*. 2012, 21(1):S108-S118.
21. Gunther CS, Chervin C, Marsh KB, et al. Characterisation of two alcohol acyltransferases from kiwifruit (*Actinidia* spp.) reveals distinct substrate preferences. *Phytochemistry*. 2011, 72(8):700-710.
22. Souleyre EJF, Greenwood DR, Friel EN, et al. An alcohol acyl transferase from apple (cv. Royal Gala), MpAAT1, produces esters involved in apple fruit flavor. *Febs J*. 2005, 272(12):3132-3144.
23. Cumplido-Laso G, Medina-Puche L, Moyano E, et al. The fruit ripening-related gene FaAAT2 encodes an acyl transferase involved in strawberry aroma biogenesis. *J Exp Bot*. 2012, 63(11):4275-4290.
24. Aharoni A, Keizer LCP, Bouwmeester HJ, et al. Identification of the SAAT gene involved in strawberry flavor biogenesis by use of DNA microarrays. *Plant Cell*. 2000, 12(5):647-661.
25. Wu J, Wang ZW, Shi ZB, Zhang S, et al. The genome of the pear (*Pyrus bretschneideri* Rehd.). *Genome Res*. 2013, 23(2):396-408.
26. D'Auria JC, Pichersky E, Schaub A, Hansel A, et al. Characterization of a BAHD acyltransferase responsible for producing the green leaf volatile (Z)-3-hexen-1-yl acetate in *Arabidopsis thaliana*. *Plant J*. 2007, 49(2):194-207.
27. Yu XH, Gou JY, Liu CJ. BAHD superfamily of acyl-CoA dependent acyltransferases in *Populus* and *Arabidopsis*: bioinformatics and gene expression. *Plant Mol Biol*. 2009, 70(4):421-442.
28. Hoffmann L, Maury S, Martz F, Geoffroy P, et al. Purification, cloning, and properties of an acyltransferase controlling shikimate and quinate ester intermediates in phenylpropanoid metabolism. *J Biol Chem*. 2003, 278(1):95-103.

29. Niggeweg R, Michael AJ, Martin C. Engineering plants with increased levels of the antioxidant chlorogenic acid. *Nat Biotechnol.* 2004, 22(6):746-754.
30. Grienemberger E, Besseau S, Geoffroy P, Debayle D, et al. A BAHD acyltransferase is expressed in the tapetum of *Arabidopsis* anthers and is involved in the synthesis of hydroxycinnamoyl spermidines. *Plant J.* 2009, 58(2):246-259.
31. Luo J, Nishiyama Y, Fuell C, Taguchi G, et al. Convergent evolution in the BAHD family of acyl transferases: identification and characterization of anthocyanin acyl transferases from *Arabidopsis thaliana*. *Plant J.* 2007, 50(4):678-695.
32. D'Auria JC, Reichelt M, Luck K, et al. Identification and characterization of the BAHD acyltransferase malonyl CoA: Anthocyanidin 5-O-glucoside-6 "-O-malonyltransferase (At5MAT) in *Arabidopsis thaliana*. *Febs Lett.* 2007, 581(5):872-878.
33. Xia YJ, Nikolau BJ, Schnable PS. Developmental and hormonal regulation of the *Arabidopsis* CER2 gene that codes for a nuclear-localized protein required for the normal accumulation of cuticular waxes. *Plant Physiol.* 1997, 115(3):925-937.
34. Qiao X, Li M, Li L, Yin H, et al. Genome-wide identification and comparative analysis of the heat shock transcription factor family in Chinese white pear (*Pyrus bretschneideri*) and five other Rosaceae species. *BMC Plant Biol.* 2015, 15(1):12.
35. Maher C, Stein L, Ware D. Evolution of *Arabidopsis* microRNA families through duplication events. *Genome Res.* 2006, 16(4):510-519.
36. Qiao X, Li Q, Yin H, Qi KJ, et al. Paterson AH: Gene duplication and evolution in recurring polyploidization-diploidization cycles in plants. *Genome Biol.* 2019, 20.
37. Lemoine F, Lespinet O, Labedan B. Assessing the evolutionary rate of positional orthologous genes in prokaryotes using synteny data. *BMC Evol Biol.* 2007, 7.
38. Jun J, Mandoiu II, Nelson CE. Identification of mammalian orthologs using local synteny. *BMC Genomics.* 2009, 10.
39. Xu LL, Qiao X, Zhang MY, Zhang SL. Genome-Wide analysis of aluminum-activated malate transporter family genes in six rosaceae species, and expression analysis and functional characterization on malate accumulation in Chinese white pear. *Plant Sci.* 2018, 274:451-465.
40. Fawcett JA, Maere S, Van de Peer Y. Plants with double genomes might have had a better chance to survive the Cretaceous-Tertiary extinction event. *P Natl Acad Sci.* 2009, 106(14):5737-5742.
41. Starr TK, Jameson SC, Hogquist KA. Positive and negative selection of T cells. *Annu. Rev. Immunol.* 2003, 21(1):139-176.
42. Yang ZH. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol.* 2007, 24(8):1586-1591.
43. Li GT, Jones KC, Eudes A, et al. Overexpression of a rice BAHD acyltransferase gene in switchgrass (*Panicum virgatum* L.) enhances saccharification. *BMC Biotechnol.* 2018, 18.

44. Zhang ZQ, Xu LP. Arabidopsis BRASSINOSTEROID INACTIVATOR2 is a typical BAHD acyltransferase involved in brassinosteroid homeostasis (vol 69, pg 1925, 2018). *J Exp Bot.* 2018, 69(15):3787-3787.
45. de Souza WR, Pacheco TF, Duarte KE, Sampaio BL, et al. Silencing of a BAHD acyltransferase in sugarcane increases biomass digestibility (vol 12, pg 111, 2019). *Biotechnol Biofuels.* 2019, 12.
46. Unno H, Ichimaida F, Suzuki H, Takahashi S, et al. Structural and mutational studies of anthocyanin malonyltransferases establish the features of BAHD enzyme catalysis. *J Biol Chem.* 2007, 282(21):15812-15822.
47. Morales-Quintana L, Moya-Leon MA, Herrera R. Computational study enlightens the structural role of the alcohol acyltransferase DFGWG motif. *J Mol Model.* 2015, 21(8).
48. Defilippi BG, Kader AA, Dandekar AM. Apple aroma: alcohol acyltransferase, a rate limiting step for ester biosynthesis, is regulated by ethylene. *Plant Sci.* 2005, 168(5):1199-1210.
49. Cao XM, Xie KL, Duan WY, Zhu YQ, et al. Peach Carboxylesterase PpCXE1 Is Associated with Catabolism of Volatile Esters. *J Agr Food Chem.* 2019, 67(18):5189-5196.
50. Galaz, S, Morales-Quintana, L, Moya-León, MA, et al. Structural analysis of the alcohol acyltransferase protein family from Cucumis melo shows that enzyme activity depends on an essential solvent channel. *FEBS J.* 2013, 280(5), 1344–
51. Lucchetta L, Manriquez D, El-Sharkawy I, Flores FB, et al. Biochemical and catalytic properties of three recombinant alcohol acyltransferases of melon. Sulfur-containing ester formation, regulatory role of CoA-SH in activity, and sequence elements conferring substrate preference. *J Agr Food Chem.* 2007, 55(13):5213-5220.
52. Freeling M. Bias in Plant Gene Content Following Different Sorts of Duplication: Tandem, Whole-Genome, Segmental, or by Transposition. *Annu Rev Plant Biol.* 2009, 60:433-453.
53. Maere S, De Bodt S, Raes J, Casneuf T, et al. Modeling gene and genome duplications in eukaryotes. *P Natl Acad Sci.* 2005, 102(15):5454-5459.
54. Friedman R, Hughes AL. Pattern and timing of gene duplication in animal genomes. *Genome Res.* 2001, 11(11):1842-1847.
55. Moore RC, Purugganan MD. The early stages of duplicate gene evolution. *Proc. Natl. Acad. Sci.* 2003, 100(26): 15682-15687.
56. Li JM, Qin MF, Qiao X, Cheng YS, et al. A New Insight into the Evolution and Functional Divergence of SWEET Transporters in Chinese White Pear (*Pyrus bretschneideri*). *Plant Cell Physiol.* 2017, 58(4):839-850.
57. Ma C, Zhang HP, Li JM, Tao ST, Qiao X, et al. Genome-wide analysis and characterization of molecular evolution of the HCT gene family in pear (*Pyrus bretschneideri*). *Plant Syst Evol.* 2017, 303(1):71-90.
58. Du DL, Hao RJ, Cheng TR, Pan HT, et al. Genome-Wide Analysis of the AP2/ERF Gene Family in *Prunus mume*. *Plant Mol Biol Rep.* 2013, 31(3):741-750.

59. Guo CL, Guo RR, Xu XZ, Gao M, Li XQ, et al. Evolution and expression analysis of the grape (*Vitis vinifera* L.) WRKY gene family. *J Exp Bot.* 2014, 65(6):1513-1528.
60. Schmidt GW, Jirschitzka J, Porta T, Reichelt M, et al. The Last Step in Cocaine Biosynthesis Is Catalyzed by a BAHD Acyltransferase. *Plant Physiol.* 2015, 167(1):89-101.
61. Zheng ZY, Qualley A, Fan BF, Dudareva N, et al. An important role of a BAHD acyl transferase-like protein in plant innate immunity. *Plant J.* 2009, 57(6):1040-1053.
62. Li DP, Shen J, Wu T, Xu YF, et al. Overexpression of the apple alcohol acyltransferase gene alters the profile of volatile blends in transgenic tobacco leaves. *Physiol Plant.* 2008, 134(3):394-402.
63. Balbontin C, Gaete-Eastman C, Fuentes L, et al. VpAAT1, a Gene Encoding an Alcohol Acyltransferase, Is Involved in Ester Biosynthesis during Ripening of Mountain Papaya Fruit. *J Agr Food Chem.* 2010, 58(8):5114-5121.
64. Li QH, Qiao X, Yin H, Zhou YH, et al. Unbiased subgenome evolution following a recent whole-genome duplication in pear (*Pyrus bretschneideri* Rehd.). *Hortic Res.* 2019, 6.
65. Zhou HS, Qi KJ, Liu X, Yin H, et al. Genome-wide identification and comparative analysis of the cation proton antiporters family in pear and four other Rosaceae species. *Mol Genet Genomics.* 2016, 291(4):1727-1742.
66. Nicholas KB. Genedoc: a tool for editing and annotating multiple sequence alignments. <http://www.pscedu/biomed/genedoc>, 1997.
67. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol.* 2015, 32(1):268-274.
68. Lee TH, Tang HB, Wang XY, Paterson AH. PGDD: a database of gene and genome duplication in plants. *Nucleic Acids Res.* 2013, 41(D1):D1152-D1158.
69. Chen C, Xia R, Chen H, et al. TBtools, a Toolkit for Biologists integrating various biological data handling tools with a user-friendly interface. *BioRxiv.* 2018: 289660.
70. Wang Y, Tang H, DeBarry J D, et al. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res.* 2012, 40(7): e49-e49.
71. Wang DP, Zhang YB, Zhang Z, et al. KaKs_Calculator 2.0: A Toolkit Incorporating Gamma-Series Methods and Sliding Window Strategies. *Genomics, Proteomics Bioinf.* 2010, 8(1):77-80.

Additional Files

Additional file 1: Table S1. Gene lists and basic feature of all identified *BAHD* genes.

Additional file 2: Figure S1. Phylogenetic analysis of *BAHD* from *Arabidopsis*, *Pyrus communis* and *Populus*. Using the maximum-likelihood method to construct the phylogenetic tree. The amino-acid sequences of *Arabidopsis* and *Populus* were obtained from phytozome (<https://phytozome.jgi.doe.gov/pz/portal.html#>).

Additional file 3: Table S2. Motif sequences identified by MEME tools in pear *BAHDs*.

Additional file 4: Table S3. Information of gene duplication events identified in all identified *BAHD* genes.

Additional file 5: Table S4. Collibearity relationship among *BAHD* genes in the same species.

Additional file 6: Table S5. Ka, Ks and Ka/Ks value of duplication gene pairs among *BAHD*.

Additional file 7: Table S6. The TPM value of *BAHD* genes in pear.

Additional file 8: Table S7. Primers for qRT-PCR of candidate genes in Chinese white pear *BAHD* gene family.

Figures




Figure 1

Phylogenetic analysis of BAHDs from Arabidopsis, Chinese white pear and Poplar. The maximum-likelihood method was used to construct the phylogenetic tree.




Figure 2

Phylogeny and conserved motifs of BAHD genes in pear. MEME tools were used to identify motifs. Different colors represent different motifs.




Figure 3

Different modes of gene duplication in BAHD families. The x-axis represents the species. The y-axis represents the number of duplicated gene pairs.




Figure 4

Gene location and a collinearity analysis of the BAHD gene family. (a): Chinese white pear; (b): apple; (c): sweet cherry; (d): peach; (e): black raspberry; (f): strawberry and (g): European pear. The genes were located on different chromosomes. Red lines represent the syntenic gene pairs.




Figure 5

Ka/Ks ratio of seven Rosaceae species. We analyzed the Ka/Ks values using coding sequences. The x-axis represents five different duplication categories. The y-axis indicates the Ka/Ks ratio. (a): sweet cherry; (b): European pear; (c): peach; (d): strawberry; (e): apple; (f): Chinese white pear and (g): black raspberry.




Figure 6

Heat map analysis of BAHD genes in pear. RNA-seq data was used to determine the expression patterns of BAHD gene. The classes IV–II-b on the left indicate the different clades. Fruit_S1–Fruit_S4 indicate the four distinct stages: on 16th May 2015 (Fruit_S1); 1st July 2015 (Fruit_S2); 31st July 2015 (Fruit_S3); and 29th August 2015 (Fruit_S4). Dark red indicates high expression, and light pink indicates low expression.




Figure 7

The relative expression levels of several BAHD genes among different periods. Tubulin and WD-repeat protein were used as the reference genes to measure expression levels in each period. The x-axis indicates the four distinct periods (45DAF, 75DAF, 105DAF and 145DAF). The y-axis indicates the relative expression and TPM (Transcripts per million reads) value. Data represent the means $\pm$ SDs (n = 3). (a): Pbr019034.1; (b): Pbr006821.1; (c): Pbr014028.1; (d): Pbr029551.1; (e): Pbr020016.1 and (f): Content of ester.




Figure 8

Phylogenetic analysis of candidate genes and 11 function-known AAT genes from other species. The maximum-likelihood method was used to construct the phylogenetic tree. The GenBank accession numbers are as follows: HCBT (Z84383), BanAAT (AX025506), CM-AAT1 (CAA94432), PCAAT1 (AAS48090), MdAAT2 (AAS79797), MpAAT1 (AY707098), RhAAT (AAW31498), FcAAT1 (FJ548610), SAAT (AAG13130), FvAAT (AAN07090), VAAT (AX025504).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.pdf](#)
- [AdditionalfilesTableS1S7.xlsx](#)