



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Paper

Investigation of the Anti-inflammatory Potential of Novel Phytotherapeutic Formulations for the Management of Arthritis, Gout, and Inflammation

Tisha Gupta*, Sourabh Yadav, Satyam Pandey

CSMU School of Pharmacy, Chhatrapati Shivaji Maharaj University, Panvel, Navi Mumbai-410221, India

ARTICLE INFO

Published: 18 Feb. 2025

Keywords:

Arthritis, Gout,
Inflammation, Phytotherapy,
Anti-inflammatory

DOI:

10.5281/zenodo.14888824

ABSTRACT

Arthritis and gout are examples of chronic inflammatory disorders that represent significant global health challenges that require effective long-term management. Based on this review, the therapeutic prospect of advanced phytotherapeutic formulations derived from ancient ethnomedicinal plants is appraised focusing on the anti-inflammatory properties of *Costus lappa* (Costus root), *Eucalyptus* spp. (*Eucalyptus* oil), and *Withania coagulans*. Phytotherapeutic interventions require the standardization of herbal formulations and strict quality control. Such natural remedies are extensively documented and with precise research could offer replacement for inflammation treatment along with of hope for billions battling them. As shown in our analysis of preclinical and limited clinical studies, they are effective in modulating cytokine production and major pathways of inflammation. These results underscore the importance of performing rigorous research and well-designed clinical trials to substantiate these potential therapeutic advantages and ensure safety.

INTRODUCTION

Millions of people around the world suffer from arthritis, gout, and other inflammatory diseases, representing a major global health burden. Traditional therapies provide partial relief at best, come with a host of unwanted side effects, and have limits to their effectiveness, especially when it comes to chronic use. This includes investigating

potential novel therapeutic avenues with natural origins. Herbal medicines have been used in traditional medicine for inflammatory disease treatment. Investigating the Anti-Inflammatory Potential of a New Phytotherapeutic Formulation Containing Three Medicinal Plants Used in Traditional Medicine.

***Corresponding Author:** Tisha Gupta

Address: CSMU School of Pharmacy, Chhatrapati Shivaji Maharaj University, Panvel, Navi Mumbai-410221, India

Email ✉: tishagupta1303@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



- Paneer doda: A medicinal herb reputed for its anti-inflammatory properties owing to the presence of alkaloids and flavonoids.
- Costus root: A traditional remedy showing reported anti-arthritic and anti-inflammatory properties which is attributed mainly to the compounds costunolide and dehydrocostus lactone
- Eucalyptus leaf: Eucalyptus leaf used for analgesic and reduce fever, such as are there actively present eucalyptol and cineole. [16]

The study postulates that the combined synergistic effect of all three of these plant extracts will demonstrate an improved anti-inflammatory activity to that of their individual components.

MATERIALS AND METHODS

Plant Materials: -

1) Costus root (Costus lappa)

Costus root (aka Saussurea lappa) is similarly used for different inflammatory diseases in traditional Ayurvedic and Chinese medicine. Early studies indicate that Costus root may have anti-inflammatory [1] and analgesic [2] effects. One study looked into the anti-inflammatory properties of Costus root extract in adjuvant arthritis induced rats. The findings suggested that

the extract markedly alleviated inflammation, enhanced joint function, and lowered inflammatory marker levels [4]. In addition, costus root is thought to reduce uric acid levels by some traditional practitioners, making it potentially helpful for gout management. Researchers are exploring traditional medicine in their search for new, long-lasting and side effect-minimizing anti-inflammatory drugs. Such exploration has the potential to revolutionize treatment approaches and improve patient outcomes. [5, 6]. Saussurea lappa root is a common medicinal plant that can be used as a remedy for various diseases. Studies indicate that they have anti-inflammatory, anti-ulcer, anticancer, antiviral, and hepatoprotective properties, indicating their traditional medicinal applications. [10,11]

● Anti-Inflammatory Action:

- Potent anti-inflammatory action in a wide range of animal models, including carrageenan-induced paw edema and adjuvant-induced arthritis.
- Mechanisms include 1) inhibition of pro-inflammatory cytokines; 2) suppression of leukocyte infiltration; and 3) stabilization of lysosomal membranes.

Scientific Classification	
Kingdom	plantae
Clade	Tracheophytes,Angiosperm,Monocots,Commelinids
Order	Zingiberales
Family	Costaceae
Genus	Costus
Synonyms	1.Cadalvena fenzi 2.Gissanthe salisb 3.Jacuanga T.Lestib



Costus Root

2)Withania coagulans (Paneer Doda):

A plant used in various ailments, including inflammation, in traditional medicine includes *Withania coagulans*, otherwise, known as paneer doda[3]. While there is little data specifically about it for gout and arthritis anti-inflammatory actions, some studies note that it can modify inflammatory pathways, which may or may not include anti-inflammatory effects. More research is needed to determine its efficacy and safety in the treatment of gout and arthritis, as well as the exact bioactive compounds responsible for these benefits.

The hydroalcoholic extract of *Withania coagulans* fruits demonstrated significant anti-inflammatory activity in the carrageenan-induced edema model suggesting that it may serve as a novel therapeutic agent in some inflammatory disorders. (Rajurkar et al 2001)[9]

The alcoholic extract of *Withania coagulans* showed an intense anti-inflammatory action in acute inflammation induced by egg albumin, indicating that it could be a very potent therapeutic agent. (Budhiraja et al 1977)[7]

Constituents of *Withania coagulans* are reported to contain potent anti-inflammatory withanolide, which alter the production of acute inflammatory markers. (Budhiraja et al 1984) [8]

- **Anti-Inflammatory Activity:**

- Anti-inflammatory effects in animal models have been observed, reportedly due to inhibition of pro-inflammatory mediators (e.g. cytokines (TNF- α , IL-1 β) and enzymes (cyclooxygenase, lipoxygenase).
- Withanolides modulate inflammatory pathways, including nuclear factor-kappa B (NF- κ B) and mitogen-activated protein kinases (MAPKs).

Scientific Classification	
Kingdom	plantae
Clade	Angiosperm, Tracheophytes, Eudicots, Asterids
Order	Solanales
Family	Solanaceae
Genus	<i>Withania</i>
Synonyms	<i>Puneeria coagulans</i> Stocks
Binomial name	<i>Withania Coagulans</i>



Paneer Doda (*Withania Coagulans*)

3) Eucalyptus spp. (Eucalyptus Oil):

Eucalyptus oil is extracted from the leaves of eucalyptus trees, and is known for its {musculoskeletal [19], rheumatoid arthritis [19], neuropathic[17]8, respiratory[18]} amelioration. According to Brazilian folk medicine, eucalyptus

essential oil is an anti-inflammatory, analgesic, and antipyretic agent[20]. 1,8-cineole, its main ingredient often referred to as eucalyptol, has repeatedly shown strong anti-inflammatory properties in various studies. As per the studies, eucalyptol interferes with significant, pain

mediating pathways and efficiently down-regulates the up-regulation of pro-inflammatory cytokines such as TNF- α and IL-1 β [15]. Sadlon and McKay cite applications of eucalyptus oil for musculoskeletal pain and respiratory infections^a. Zymosan-induced arthritis in mice was improved by an oral administration of 1,8-cineole (400 mg/kg), which decreased the pro-inflammatory cytokines and joint pain. [12]

The hTRPA1 inhibitory and hTRPM8/hTRPV3 stimulant actions of 1,8-cineole suggest that its analgesic and anti-inflammatory effects correlate with these mechanisms. [13]

In a monosodium urate (MSU)-induced inflammatory and pain model of gout arthritis in

mice, the anti-inflammatory and anti-nociceptive effects of eucalyptol (1,8-cineole) were evaluated. Furthermore, it diminished the infiltration of inflammatory cells, pro-inflammatory cytokines, and oxidative stress in ankle tissues, and suppressed the activation of NLRP3 inflammasome and TRPV1 expression. [14]

• **Anti-inflammatory Activity:**

- Exhibits anti-inflammatory actions through the inhibition of pro-inflammatory mediators including prostaglandins and leukotrienes.
- May also have an antioxidant and anti-nociceptive action that contribute to its general anti-inflammatory action.

Scientific Classification	
Kingdom	plantae
Clade	Angiosperm, Tracheophytes, Eudicots, Rosids
Order	Myrtales
Family	Myrtaceae
Subfamily	Myrtoideae
Tribe	Eucalypteae
Genus	Eucalypus
Synonyms	1. Aromadendrum W.Anderson ex R.Br.,1810 2. Eudesmia R.Br.,1814 3. Symphyomyrtus



Eucalyptus Leaf

Methodology: -

Extraction: The extraction will be done using 80% Methanol as solvent by Soxhlet extraction. The extracts will be evaporated under low pressure through a rotary evaporator.

Formulation: Based on preliminary studies, extracts of Paneer doda, Costus root and Eucalyptus leaf will be combined in a pre-determined ratio to prepare a standardized formulation.

HPLC Analysis: HPLC analysis shall be separated on suitable reverse-phase C18 column and on this reactant, we will use gradient mobile phase. Characterization will be performed using standards of known bioactive compounds (e.g. flavonoids, terpenoids).

In vitro Assay: We will stimulate RAW 264.7 macrophages using LPS for inducing inflammation. In this regard, at different concentrations this formulation will be tested for inhibition of NO production which will be measured using Griess reagent.

In vivo Assay: The rats will be subjected to carrageenan-induced paw edema. Dosage: The formulation will be given orally in ascending doses and the anti-edema activity will be performed by estimating changes of paw volume using a plethysmometer.

Data Analysis: Statistical analysis: All data will be analyzed by appropriate statistical methods (such as SPSS, ANOVA and t-tests) to identify significant differences between groups.

CONCLUSION

After analyzing the various research and review papers we found that the phytotherapeutic compounds hold a profound promise for establishing anti-inflammatory drugs in the treatment of arthritis, gout, and other inflammatory ailments. The potential suicide herbaceous of *Eucalyptus* spp., *Costus lappa* and *Withania coagulans* need to be verified by scientific studies as well for the clinically proven efficacy, mechanism of action and safety. The concerns related to standardization and quality control must be addressed to formulate effective interventions. Paneer doda, *Costus* root and *Eucalyptus* leaf are traditionally used for inflammatory disorders. The study concerns with, determining the anti-inflammatory potential of the newly formulated containing these plant extracts ultimately, could be useful for establishment of

safe and effective phytotherapeutics for arthritis and similar ailments.

REFERENCES

1. Rajkumar, K., et al. "Anti-inflammatory and anti-arthritic activity of *Costus speciosus* rhizome extract." *Asian Journal of Pharmaceutical and Clinical Research*, 2017.
2. inny, K., et al. "Anti-inflammatory and antipyretic properties of the rhizome of *Costus speciosus*." *Asian Journal of Pharmaceutical and Clinical Research*, 2010.
3. Mirjalili, M.H., et al. "Withanolides from *Withania* species: A review on their pharmacological properties and biosynthesis." *Phytochemistry Reviews*, 2009. [DOI: 10.1007/s11101-009-9135]
4. Srivastava S, Singh P, Jha KK, Mishra G, Srivastava S, Khosa RL. Evaluation of anti-arthritic potential of the methanolic extract of the aerial parts of *Costus speciosus*. *J Ayurveda Integr Med* 2012; 3:204–8.
5. Ekambaram S, Perumal SS, Subramanian V. Evaluation of antiarthritic activity of *Strychnos potatorum* Linn seeds in Freund's adjuvant induced arthritic rat model. *BioMol Concept Complement Altern Med* 2010; 10:1–9.
6. Patil RB, Nanjwade BK, Manvi FV. Effect of *Sesbania grandiflora* and *Sesbania sesban* Bark on carrageenan induced acute inflammation and adjuvant-induced arthritis in rats. *Pharmacol Sci Monit Int J Pharm Sci* 2010; 1:75–89. Rajurkar, S.M., Thakre, P.N., Waddukar, S.G., *Phytochemical and Pharmacological screening of W. coagulans berries as anti-inflammatory*. 53 IPC 2001. New Delhi, Dec. Sci. Abst. CP 38. P. no. 215.
7. Budhiraja RD, Sudhir S, Garga KN (1977) Pharmacological investigations on fruits of *Withania coagulans* Dunal. *Planta Medica* 32: 154-57



8. Budhiraja RD, Sudhir S, Garg KN, Arora BC (1984) Anti-inflammatory activity of 3 β -Hydroxy-2, 3-dihydro-withanolide F. *Planta Medica* 50 (2): 134-136
9. . Rajurkar, S.M., Thakre, P.N., Waddukar, S.G., Phytochemical and Pharmacological screening of W. coagulans berries as anti-inflammatory. 53 IPC 2001. New Delhi, Dec. Sci. Abst. CP 38. P. no. 215.
10. Pandey MM, Govindarajan R, Rawat AK, Pushpangadan P. Free radical scavenging potential of *Saussurea costus*. *Acta Pharmacol* 2005; 55:297–304.
11. Kumar A, Kumar B, Kumari S. Diabetes mellitus and its herbal treatment. *Int J Biol Med Res* 2014; 5:4180–5.
12. Costa DVDS, Nunes RDM: Oral administration of eucalyptol reduces cell migration and pain-like behavior in zymosan-induced arthritis mice. *Braz J Pharm Sci* 2022;58: e21189.
13. Takaishi M, Fujita F, Uchida K, Yamamoto S, Sawada SM, Hatai UC, et al: 1,8-cineole, a TRPM8 agonist, is a novel natural antagonist of human TRPA1. *Mol Pain* 2012; 8:86-99.
14. Yin C, Liu B, Wang P, Li X, Li Y, Zheng X, et al: Eucalyptol alleviates inflammation and pain responses in a mouse model of gout arthritis. *Br J Pharmacol* 2020; 177:2042-57.
15. Zhao C, Cao Y, Zhang Z, Nie D, Li Y: Cinnamon and Eucalyptus oils suppress the inflammation induced by lipopolysaccharide in vivo. *Molecules* 2021; 26:7410-25.
16. International Organization for Standardization: Crude or rectified oils of *Eucalyptus globulus* (*Eucalyptus globulus* Labill.). ISO 770:2002. Geneva, ISO. 2002.
17. Ridouh I, Hackshaw KV: Essential oils and neuropathic pain. *Plants* 2022; 11:1797-808.
18. Horváth G, Ács K: Essential oils in the treatment of respiratory tract diseases highlighting their role in bacterial infections and their anti-inflammatory action: a review. *Flavour Fragr J* 2015; 30:331-41.
19. Varkaneh ZK, Karampourian A, Oshvandi K, Basiri Z, Mohammadi Y: The effect of eucalyptus inhalation on pain and the quality of life in rheumatoid arthritis. *Contemp Clin Trials Commun* 2022; 29:100976-81.
20. Lee G, Park J, Kim MS, Seol GH, Min SS: Analgesic effects of eucalyptus essential oil in mice. *Korean J Pain* 2019; 32:79-86

HOW TO CITE: Tisha Gupta*, Sourabh Yadav, Satyam Pandey, Investigation of the Anti-inflammatory Potential of Novel Phytotherapeutic Formulations for the Management of Arthritis, Gout, and Inflammation, *Int. J. of Pharm. Sci.*, 2025, Vol 3, Issue 2, 1548-1553. <https://doi.org/10.5281/zenodo.14888824>

