

Qualitative analysis of secondary metabolites in *Cordia myxa* L. fruits: a multipurpose tree species

Arjun Patra¹, Emdad Hossain^{2*}, Bidhan Chandra Roy³, Sohan Lal⁴, Nibedita Jena⁵ and Sanjeet Kumar⁵

¹Department of Botany, P.K. College, Contai, West Bengal, India

²Department of Pharmaceutical Technology, Jadavpur University, Kolkata, West Bengal, India

³Department of Botany, Dinabandhu Mahavidyalaya, Bongaon, West Bengal, India

⁴School of Studies in Botany, Shaheed Mahendra Karma Vishwavidhyalaya, Bastar, Jagdalpur, Chhattisgarh, India

⁵Ambika Prasad Research Foundation, Odisha, India

*Email-id: emdad@rediffmail.com; ORCID: <https://orcid.org/0000-0001-9110-6449>



Licensed under a Creative Commons Attribution 4.0 International License

Abstract

Cordia myxa L., a widely used multipurpose tree species, has been employed in traditional medicine for its diverse therapeutic properties. The present study aimed to investigate the qualitative composition of secondary metabolites in *C. myxa* fruits. Phytochemical analysis of aqueous, methanol, and ethanol extracts revealed the presence of tannins, saponins, flavonoids, phenolic compounds, steroids, and alkaloids. The identified bioactive compounds suggest potential antioxidant, anti-inflammatory, antimicrobial, and antidiabetic properties, supporting the use of this plant in traditional medicine. This study provides a foundation for further research into the pharmacological and therapeutic applications of *C. myxa* fruits.

Keywords: Anti-inflammatory, antioxidant, multipurpose tree, pharmacological potential

Introduction

Cordia myxa L., a species of flowering plant in the Boraginaceae family, is a widely used multipurpose tree species found in various parts of the world (Aleem *et al.*, 2017). The plant has been used in traditional medicine for centuries, with its different parts being used to treat a range of ailments (Kumar *et al.*, 2021). Recent studies have shown that the studied plant possesses various biological activities, including antioxidant, anti-inflammatory, and antimicrobial properties (Singh *et al.*, 2022). The plant extract was helpful for treating cognitive dysfunction in rodents (Al-Musawi *et al.*, 2022; Kendir *et al.*, 2022). Secondary metabolites, such as alkaloids,

flavonoids, and phenolic acids, are known to play a crucial role in the medicinal properties of plants (Al-Khafaji *et al.*, 2021). Therefore, it is essential to investigate the preliminary qualitative composition of secondary metabolites in *C. myxa* fruits to understand their potential health benefits. The present study aims to investigate the preliminary qualitative composition of secondary metabolites in fruits of the studied plant using various solvent extracts. The study will provide valuable information on the phytochemical composition of *C. myxa* fruits, which can be used to develop new drugs and therapies. Additionally, the study will contribute to the conservation and sustainable use of *C. myxa*, a valuable multipurpose tree species.




Figure 1: Fruits of of *C. myxa*

Materials and methods

The fruits of *C. myxa* were collected from the bank of the Mahanad River areas in Cuttack in October 2024 (Figure 1; Kumar *et al.*, 2021). Authors have identified the plant specimen. Fruit extracts of aqueous, ethanol, and methanol were prepared, and nine different bioactive compounds were detected using standard methods (Devi *et al.*, 2023; Marndi *et al.*, 2024; Devi *et al.*, 2024). Detection of nine secondary metabolites was conducted using standard methods and as followed by previous work of our laboratories (Kokate *et al.*, 2006; Jena *et al.*, 2024; Pradhan *et al.*, 2024).

Test for tannin: About 1 ml of the fruit extract was taken. 3-5 drops of 10% lead acetate solution were added to it. The gelatinous precipitate formation confirmed the presence of tannin.

Test for saponin: About 1 ml of the fruit extract was taken and 1 ml of distilled water was added and shaken well. The formation of persistent froth was observed confirming the presence of saponin.

Test for flavonoids: About 1 ml of the fruit extract was taken. 2 ml of 2% NaOH solution and 3 to 4 drops of dilute hydrochloric acid were added to it. The colour initially turned to an intense yellow colour with NaOH solution and later became colourless. This colour change appearance confirmed the presence of flavonoids.

Test for terpenoids: About 1 ml of the filtrate was added with 6 drops of chloroform and placed in the water bath for a few minutes. Then 6 drops of concentrated H₂SO₄ were added. The reddish- brown interface confirmed the presence of terpenoids.

Test for phenolic groups: About 1 ml of the filtrate extract was taken. A few drops of 5% Ferric chloride solution were added. The dark bluish-black appearance confirmed the presence of phenolic compounds.

Test for reducing sugars: About 1 ml of the fruit extract was taken and 2 drops of Fehling's solution A followed by Fehling's solution B were added and kept in the water bath for some time. The presence of red-orange precipitate confirmed the presence of reducing sugar.

Test for steroids: About 1 ml of the fruit extract was taken. 1 ml of chloroform and 1 ml of concentrated sulphuric acid was added gently into it

touching test tube mouth. The appearance of upper red and lower yellow with green fluorescence provides the presence of steroids.

Test for alkaloids: About 1 ml of the fruit extract was taken and added 3 to 4 drops of Dragendroff's reagent. The formation of a reddish-brown precipitate confirmed the presence of alkaloids.

Test for carbonyl compounds: About 1 ml of the fruit extract was taken and added 3 to 4 drops of 2,4- Dinitrophenylhydrazine (DNPH) reagent. The formation of yellow crystal confirmed the presence of carbonyl compounds.

Results and discussion

The phytochemical analysis of *C. myxa* fruits revealed the presence of various bioactive compounds. The aqueous extract contained tannins, saponins, flavonoids, phenolic compounds, steroids, and alkaloids (Table 1; Figure 2). The methanol extract contained saponins, flavonoids, phenolic compounds, and steroids, but no tannins or alkaloids. The ethanol extract contained saponins, flavonoids, phenolic compounds, steroids, and alkaloids, but no tannins (Figure 3). The presence of these bioactive compounds suggests that *C. myxa* fruits may have various pharmacological properties (Tak *et al.*, 2024). The saponins, flavonoids, and phenolic compounds present in all three extracts may have a major pharmacological role in treating various health problems. Table 1 provides a comprehensive overview of the phytochemical analysis of *C. myxa* fruits. The table shows that the different extracts (aqueous, methanol, and ethanol) contain varying levels of bioactive compounds.

Table 1: Qualitative phytochemical analysis of *C. myxa* using different extracts

Bioactive compounds	Extracts		
	Aqueous	Methanol	Ethanol
Tannin	+	-	-
Saponin	+	+	+
Flavonoids	+	+	+
Terpenoids	-	-	-
Phenolic compounds	+	+	+
Reducing sugars	-	-	-
Steroids	+	+	+

Alkaloids	+	-	+
Carbonyl compounds	-	-	-

(+: Positive; -: Negative)




Figure 2: Detection of secondary metabolites of *C. myxa* using aqueous extract




Figure 3: Detection of secondary metabolites of *C. myxa* using ethanol extract




Figure 4: Pharmacological potentials of *C. myxa* fruits

The result suggests that the choice of extract may be important for specific applications. Table 1 and Figure 4 show that the fruits may have a range of biological activities, including neuroprotective, fungicidal, anticarcinogenic, immunomodulatory, and antidiabetic effects. This suggests that fruits may be a valuable source of bioactive compounds with potential therapeutic applications. The presence of tannins, saponins, flavonoids, phenolic compounds, steroids, and alkaloids in the fruit extracts suggests that *C. myxa* may possess antioxidant, anti-inflammatory, antimicrobial, and antidiabetic properties, among others (Figure 4). The fruits are edible. With high nutrition and medicinal value, the fruits can be utilized as functional fruit (Tak *et al.*, 2024).

Conclusion

The present study has successfully identified and characterized the secondary metabolites present in *C. myxa* fruits, revealing a diverse range of bioactive compounds with potential therapeutic applications. These findings provide a scientific basis for the traditional use of *C. myxa* in medicine and highlight its potential as a valuable source of natural products for the development of new drugs and therapies.

References

Aleem ERA, Seddik FEF, Samy MN and Desoukey SY. (2017). Botanical studies of the leaf of *Cordia myxa* L. Journal of Pharmacognosy and Phytochemistry. 6(6): 2086-2091.

- Al-Khafaji SA, Alsaadawi MA, Yasari AMA and Al-Saadawe MA. (2021). Article review: *Cordia myxa* L.: The gift of the nature, a review. Basrah Journal of Agricultural Sciences. 34(2): 267-277.
- Al-Musawi MH, Ibrahim KM, Albukhaty S. (2022). *In vitro* study of antioxidant, antibacterial, and cytotoxicity properties of *Cordia myxa* fruit extract. Iranian Journal of Microbiology. 14 (1): 97-103.
- Devi RS, Satapathy KB and Kumar S. (2023). Validation of tribal claims for formulation of future drugs through evaluation of ethno-pharmacological values of *Ludwigia adscendens*. Medicinal Plants. 15(4): 691-697.
- Devi RS, Satapathy KB, Manjula BL, Sharma BP and Kumar S. (2024). Evaluation of ethnopharmacological potential of *Oenanthe javanica* DC. Annals of Biology. 40(1): 135-139.
- Jena N, Rout S, Devi RS and Kumar S. (2024). Phytochemical and cytotoxicity analysis of fruits of *Zanthoxylum asiaticum* (L.) Appelhans, Groppo & J. Wen: a minor fruit plant of Odisha, India. E-planet. 22(1): 62-70.
- Kendir G, Bae HJ, Kim J, Jeong Y, Bae HJ, Park K, Yang X, Cho YJ, Kim JY, Jung SY, Köroğlu A, Jang DS, Ryu JH. (2022). The effects of the ethanol extract of *Cordia myxa* leaves on the cognitive function in mice. BMC Complementary Medicine and Therapies. 22:215. DOI: 10.1186/s12906-022-03693-z.
- Kokate CK, Purohit AP, and Gokhale SB. (2006). Pharmacognosy. 35th Ed. Nirali Prakashan, Pune. ISBN 81-85790-00-1.
- Kumar SN, Mishra S and Kumar S. (2021). Documentation of indigenous traditional knowledge (ITK) on commonly available plants in Koira range, Bonai Forest Division, Sundargarh, Odisha, India. Asian Plant Research Journal. 8(4): 83-95.
- Marndi S, Barik M and Kumar S. (2024). Phytochemical and cytotoxicity analysis of *Eulophia explanata* Lindl. (Tuber extracts). African Journal of Biological Sciences. 6(4): 2751-2756.
- Pradhan I, Rani JJ, Pattanayak S, Jacob M, Rout S, Kumar S, and Hossain E (2024). Phytochemical profiling and cytotoxic evaluation of *Schrebera swietenoides* fruits. In: Sharma BP, Suple SD, Mishra S and Rath SK. Plants & Secondary Metabolites, Volume 3. APRF Publishers, India. Pp 80-90. ISBN: 978-81-973870-3-6.
- Singh V, Sood A, Pruthi S, Singh M, Saini B, Singh TG and Kumar A. (2022). Chemical constituents and biological activities of *Cordia myxa* L.: a review. The Natural Products Journal. 12(2): 30-41.
- Tak Y, Samota MK, Meena NK, Kaur G, Jain MC, Kumar R, Kaur M and Amarowicz R. (2024). Underutilized fruit lasoda (*Cordia myxa* L.): review on bioactive compounds, antioxidant potentiality and applications in health bioactivities and food. Fitoterapia. 175: 105898. DOI: 10.1016/j.fitote.2024.105898.