

Targeted mutagenesis of *Medicago truncatula* Nodule-specific Cysteine-rich (NCR) genes using the *Agrobacterium rhizogenes*-mediated CRISPR/Cas9 system

Berivan Güngör^{1,2*}, János Barnabás Biró^{1*}, Ágota Domonkos², Beatrix Horváth², Péter Kaló^{1,2}

1 Institute of Plant Biology, Biological Research Centre, Eötvös Lóránd Research Network, Szeged, Hungary

2 Institute of Genetics and Biotechnology, Hungarian University of Agriculture and Life Sciences, Gödöllő, Hungary

* B.G. and J.B.B. contributed equally to this work.

Author for correspondence: Péter Kaló, Email: kalo.peter@brc.hu

ORCID IDs:

Peter Kalo: 0000-0002-0404-8904

Berivan Güngör: 0000-0002-5612-1130

János Barnabás Biró: 0000-0001-8851-0387

Ágota Domonkos: 0000-0003-4017-0605

Beatrix Horvath: 0000-0001-8499-568X

44 **Abstract**

45

46 The host-produced nodule specific cysteine-rich (NCR) peptides control the terminal
 47 differentiation of endosymbiotic rhizobia in the nodules of IRLC legumes. Although the
 48 *Medicago truncatula* genome encodes about 700 NCR peptides, only few of them have been
 49 proved to be crucial for nitrogen-fixing symbiosis. In this study, we applied the CRISPR/Cas9
 50 gene editing technology to generate knock-out mutants of *NCR* genes for which no genetic or
 51 functional data were previously available. We have developed a workflow to analyse the
 52 mutation and the symbiotic phenotype of individual nodules formed on *Agrobacterium*
 53 *rhizogenes*-mediated transgenic hairy roots. The selected *NCR* genes were successfully edited
 54 by the CRISPR/Cas9 system and nodules formed on knockout hairy roots showed wild type
 55 phenotype indicating that peptides NCR068, NCR089, NCR128 and NCR161 are not
 56 essential for symbiosis between *M. truncatula* Jemalong and *Sinorhizobium medicae*
 57 WSM419. We regenerated stable mutants edited for the *NCR068* from hairy roots obtained
 58 from *A. rhizogenes*-mediated transformation. The analysis of the symbiotic phenotype of
 59 stable *ncr068* mutants showed that peptide NCR068 is not required for symbiosis with *S.*
 60 *meliloti* strains 2011 and FSM-MA either. Our study reports that gene editing can help to
 61 elicit the role of particular NCRs in symbiotic nitrogen fixation.

62

63 Introduction

64

65 Most legumes establish nitrogen-fixing symbiotic interaction with soil bacteria termed
 66 rhizobia. Compatible interaction between the host legumes and their rhizobia partner induces
 67 the formation of a specialised organ on the root, referred to as nodules. Rhizobia colonize the
 68 symbiotic cells of the nodules and differentiate into their symbiotic form, the bacteroids
 69 which convert the atmospheric nitrogen into ammonia¹. The indeterminate nodules formed on
 70 the roots of temperate legumes, such as the model legume *Medicago truncatula*, possess a
 71 persistent apical meristem². The continuous activity of the meristem (zone I) produces cells
 72 creating a developmental gradient in the nodules, whereupon these nodules are cylindrical in
 73 shape. The new cells mature, many of them become infected in zone II (infection zone)
 74 wherein bacteria are engulfed by a plant-derived membrane forming new organelles, the
 75 symbiosomes. Infected host cells and rhizobia differentiate in interzone (IZ) and the reduction
 76 of nitrogen takes place in zone III (nitrogen fixation zone). The completely developed and
 77 effectively nitrogen-fixing nodules produce leghemoglobin which gives wild type nodules
 78 their pink colour.

79 Rhizobia in the nodules of legumes belonging to the inverted repeat-lacking clade (IRLC),
 80 including *M. truncatula*, and some Dalbergoid species undergo irreversible differentiation
 81 mediated by nodule-specific cysteine-rich (NCR) peptides³⁻⁵. These host produced peptides
 82 contain four or six cysteine residues in conserved positions and possess signal peptides which
 83 target them to the symbiosomes through a nodule-specific protein secretory pathway^{6,7} where
 84 they govern the transition of rhizobia into bacteroids. The *M. truncatula* genome encodes a
 85 large family of NCR peptides with more than 700 members^{8,9}. Although these high number of
 86 peptides show remarkable amino acid sequence variation, their function was initially
 87 presumed redundant. Nevertheless, recent studies have identified some peptides, NCR169,
 88 NCR211, NCR247, NCR343 and NCR-new35, crucial for the bacteroid differentiation and
 89 persistence in *M. truncatula* nodules but also found peptides (NCR178, NCR341, NCR344
 90 and NCR345) which are not essential for nitrogen-fixing symbiosis¹⁰⁻¹⁵. The identification of
 91 most of these essential peptides was random during the forward genetic analyses of
 92 mutagenized *M. truncatula* populations. Therefore, the reverse genetic approach provides a
 93 comprehensive and targeted way to analyse the function of the NCR gene family members.
 94 The recently developed CRISPR/Cas9 (Clustered Regularly Interspaced Short Palindromic
 95 Repeats/CRISPR-associated protein 9) system is a simple and precise tool for targeted gene
 96 editing and analyse gene function in various plants¹⁶. Several reports described the application

of CRISPR/Cas9 gene editing system in legume species (soybean¹⁷; peanut¹⁸; pea¹⁹; *Lotus japonicus*²⁰ including *M. truncatula*^{21–24}. Unfortunately, most legume species are largely recalcitrant to somatic embryogenesis and *in vitro* plant regeneration, therefore generating stable transformant legume plants using the *Agrobacterium tumefaciens*-mediated transformation system is a cumbersome and time-consuming process. In contrast, the *A. rhizogenes*-mediated hairy root transformation does not need complete plant regeneration, therefore it provides an efficient tool for a fast and large-scale functional analysis of root-related traits, such as symbiotic nodule development. Despite its robustness, the *A. rhizogenes*-mediated hairy root transformation has some disadvantages. For instance, the transformation efficiency varies, and gene transfer is not achieved in all root cells²⁵, or the emerged hairy roots develop from multiple cells with different gene editing events which can result in mosaicism of transformed and untransformed cells or even multiple genotypes. This could cause trouble during gene editing of a wild-type genome to generate knockouts because a small amount of residual wild-type cells could affect the phenotype to be analysed. Therefore, the removal of untransformed or partially transformed roots and analysing the genotype of the whole hairy root is particularly important.

The CRISPR/Cas9 gene editing system, have been applied successfully to knockout and analyse the requirement of peptides NCR247, NFS1 and NFS2^{12,26,27} using *A. rhizogenes*-mediated hairy root transformation. In these reports, the targeted *NCR* genes were selected based on preliminary genetic or functional studies. In this study, we optimized the CRISPR/Cas9 system using hairy root transformation to generate mutation in selected *NCR* genes and developed a workflow to conduct their functional analysis for symbiosis.

Results

Testing the efficiency of a modified vector for CRISPR/Cas9 gene editing system to target *NCR169*

To analyse the function of *NCR* genes, the CRISPR/Cas9 gene editing technology was used to generate composite *Medicago truncatula* plants mutated in hairy roots. In order to monitor the presence of CRISPR constructs, delivered into the genome of *M. truncatula* root cells using the *Agrobacterium rhizogenes*-mediated hairy-root system²⁸, the gene coding for the DsRed fluorescent transformation marker was introduced into the CRISPR/Cas9 plant transformation vector pKSE401 resulting in the vector pKSE401-RR. The constitutively active red fluorescent marker allowed us to remove non-transgenic roots at the early stages to promote

the growth of transgenic hairy roots. To test the efficiency of targeted mutagenesis of pKSE401-RR, we designed three different single guide RNA (sgRNA) constructs targeting the *NCR169* gene, previously proved to be essential for effective nitrogen-fixing symbiosis¹⁰. Each sgRNA (NCR169a, b and c) was designed directing to the first half of the coding sequence of *NCR169* but the initial 5' nucleotides for the protospacer sequences of NCR169b and NCR169c were C instead of the optimal G(/A) which is preferred by the AtU6 promoter (Fig. S1a;²⁹). Nevertheless, we used these two suboptimal sgRNA constructs as well because the short, 189 bp long coding sequence of *NCR169* constrained the available optimal target sequences. The three constructs carrying each sgRNAs were introduced into roots of wild-type *M. truncatula* 2HA genotype using *A. rhizogenes*-mediated hairy root transformation. At least ten transformed roots for each construct, detected by the DsRed fluorescent marker, were sampled and used for genomic DNA isolation. The region surrounding the targeted sequence in *NCR169* was amplified by PCR. The sequence analysis of the amplicons detected only wild-type genotypes on all the NCR169b-transformed roots and roots transformed with the construct NCR169c were heterogeneous of wild-type and various mutant genotypes. However, the construct NCR169a generated mutations at higher efficiency and beside chimeric roots having both wild-type and distinct mutations, we could identify roots with homozygous or biallelic mutations for *NCR169* (Fig. S1d). The results indicated that the presence of G at the 5' end of the protospacer sequence (G(/A)N19NGG) is crucial and using C in this position greatly reduced the efficiency of genome editing which restricts the number of possible target sites in the case of small genes such as *NCRs*. In addition, the sequence analysis of amplicons containing the targeted region revealed that the sequences obtained by Sanger method composed of multiplex sequence traces around the break site (Fig. S1b). An online tool, TIDE, enabling the decomposition of mixed sequence signals was applied³⁰ that predicted the types of the mutation and the size of the deletion or insertion. For instance, the decomposition of one of the amplicons of NCR169c disclosed the mixture of wild-type and CRISPR/Cas9 edited sequences containing a 7 bp deletion and/or a one bp insertion (Fig. S1c).

In agreement with a previous report²², we also found that different genome editing events occurred in distinct hairy roots of a single plant which necessitates the analysis of individual nodules separately. These findings led us to the conclusions that the effectiveness of genome editing and the reliability of the identification of the nature of genome editing events needed to be improved in our system. Therefore (i) we replaced the Arabidopsis U6-26 promoter with the *M. truncatula* U6 (MtU6.6) promoter driving sgRNA synthesis, (ii) we developed a

preparation procedure allowing phenotypic and genotypic analysis of individual nodules, and (iii) applied an NGS amplicon sequencing approach to assess allele distribution in individual nodules.

Inducing mutations in *NCR169* and *NCR-new35* using the optimized CRISPR/Cas9-mediated genome editing system

Previous studies found that the *M. truncatula* U6 promoter enhanced the expression efficiency of guide or hairpin RNAs in legume species^{21,31}. To increase the efficacy of targeted mutagenesis by CRISPR/Cas9, the Arabidopsis U6-26 promoter was replaced with MtU6.6 in the pKSE401-RR resulting in the vector pKSE466-RR. We inserted the sgRNAs NCR169a and another one, targeting the *NCR-new35*, of which requirement for effective symbiosis has been recently demonstrated¹³, into pKSE466-RR, respectively. To validate the efficiency of the pKSE466-RR vector construct, the empty vector and pKSE466-RR-NCR169a were introduced into wild-type *M. truncatula* 2HA plants using *A. rhizogenes*-mediated hairy root transformation and plants were inoculated with *S. medicae* WSM419. Nodules developed on DsRed fluorescent transformed roots were harvested 3-4 weeks post inoculation (wpi) with rhizobia, individually fixed and sections were stained with nucleic acid-binding dye SYTO13. Following image capturing, genomic DNA was purified from nodule sections and the regions surrounding the targeted sequences were amplified by PCR.

The elongated nodules developed on empty vector-transformed roots were pink and showed the characteristic zonation of indeterminate nodules (Fig. 1a and d). The nodules on the roots of *M. truncatula* 2HA transformed with the pKSE466-RR-NCR169a were either pink and elongated, suggesting that these nodules were functional wild-type (Fig. 2g, i and k), or white undeveloped, implying that these were ineffective symbiotic nodules (Fig. 1b and 2b, d and f). To determine the nature of the mutations at the targeted region, the sequence analysis of the amplicons, generated from randomly selected elongated and undeveloped nodules, was carried out. The targeted amplicon sequences were determined using a next-generation sequencing (NGS) approach and the quantitative allele composition and the verification of genome editing mutations were analysed using the online tool CRISPResso2³². The analysis revealed that all the white undeveloped nodules carried few base pair biallelic insertions or deletions at a high ratio and the very few reads of wild-type alleles identified in these nodules were probably derived from sporadic wild-type cells or the trace amount of wild-type contamination (Fig. 1b, Fig. 2a-f). The morphology and bacterial occupancy of these SYTO13-stained undeveloped nodules were very similar to the nodulation phenotype of the

deletion mutant *dnf7-2* showing no fluorescence in the nodule region proximal to the root (Fig. 1e). In addition, these plants, compared with empty-vector-transformed plants, showed the symptoms of nitrogen starvation (yellowish leaves and stunted growth habit) indicating the defect of symbiotic nitrogen fixation of these nodules (Fig. 1b). Contrary to these, the sequence analysis of the randomly selected elongated pink nodules showed that they either carried almost exclusively wild-type alleles (Fig. 2g and h) or they were heterozygous for example for a 4 bp deletion and the wild-type alleles (Fig. 2i and j). We also found a wild-type nodule that predominantly carried a homozygous deletion of 3 base pairs resulting in a deletion of the first single Glu residue of the mature peptide of NCR169 which mutation did not abolish the symbiotic function of NCR169 (Fig. 2k and l).

We also applied the developed CRISPR/Cas9 toolkit to target *NCR-new35*, a recently identified *NCR* gene which is crucial for the persistence of bacteroids in *M. truncatula* cv. Jemalong nodules¹³. The empty vector and the pKSE466-RR-NCRnew35 constructs were introduced into the roots of *M. truncatula* 2HA and *M. truncatula* ecotype R108 plants using *A. rhizogenes*-mediated hairy root transformation. Nodules on empty vector-transformed roots displayed the characteristic zonation of indeterminate nodules showing cells occupied by rhizobia (Figure S2e and m). In contrast, undeveloped nodules were observed on roots of *M. truncatula* 2HA and R108 plants transformed with the editing construct of *NCR-new35* (Figures S2 and Fig. S3g-r and Fig. 4e-h). The sequence analysis showed that undeveloped nodules, randomly collected from transformed roots of *M. truncatula* 2HA and R108 plants, carried biallelic or monoallelic homozygous mutations in the amplicons containing the target sequence in *MtNCR-new35* (Fig. 4q and r). SYTO13-staining revealed that these nodules did not show the typical zonation and only their apical part fluoresced indicating that the region corresponding to the prospective nitrogen fixation zone was devoid of rhizobia (Figure S2f-h and n-p). The morphology and bacterial occupancy of genome edited nodules were in agreement with the nodulation phenotype of *Mtsym19* and *Mtsym20* nodules defective in *NCR-new35*¹³. These results also revealed that *NCR-new35* is crucial for the symbiotic interactions not only between *M. truncatula* 2HA and *S. medicae* WSM419 but between *M. truncatula* R108 and *S. meliloti* FSM-MA as well.

Selecting of *NCR* genes to be targeted by CRISPR/Cas9 editing

The *M. truncatula* genome contains a large gene family coding for more than 700 *NCR* peptides but only few of them, NCR169, NCR211, NCR247, NCR343 and *NCR-new35*, have been proved to be crucial for symbiosis¹⁰⁻¹⁵. The sequences of the essential *NCR* genes are

highly variable, and their common feature is that they contain four cysteine residues in conserved positions. The peptide NCR247 is unique among the crucial and most of the other NCR peptides because it contains five residues between the third and fourth conserved cysteines compared with the canonical four internal residues¹³. In addition, no common feature could be identified among few NCR peptides found to be non-essential for symbiosis^{10,11,13}. Therefore, it was not straightforward to ascertain which additional NCR peptides are essential and should be targeted. Despite all that, we have set up a selection pipeline to identify *NCR* genes to be targeted. We have analysed several features of more than 700 predicted *NCR* genes identified in the *M. truncatula* genome⁹. The signal peptides of NCRs are crucial for delivering the peptides to the bacteroids^{4,6,7}, therefore, (i) first we selected 655 *NCR* genes coding for peptides possessing a predicted signal peptide (Fig. 3a). The pool of *NCR* genes to be targeted (ii) were further enriched by selecting the encoded peptides with four cysteine residues in conserved positions. (iii) We have analysed the expression profile of the diminished list of 248 *NCR* genes in the database obtained by laser-capture microdissection of *M. truncatula* nodules³³. We identified 11 *NCRs* with no reads, evidently seem to be pseudogenes, and 65 *NCRs* which are expressed at low level (< 10 000 reads) in nodules compared with other *NCRs*. The recently identified *NCR-new35* shows the lowest expression level among the crucial *NCRs*¹³ and thus, we have selected genes from the 172 *NCRs* showing similar or higher expression level as *NCR-new35* (Fig. 3c and d). (iv) These *NCR* genes were analysed to predict whether they had contained sequences which were amenable to CRISPR-based editing. The cysteine residues are essential for the function of *NCR* peptides *in planta*^{10,13}, therefore we have prioritised those *NCRs* which could be targeted prior to the fourth conserved cysteines to generate presumably non-functional peptides. We have further searched the *M. truncatula* genome using an online prediction tool CRISPOR³⁴ for off-target sites that could be recognized by sgRNAs designed for *NCR* genes. The isoelectric points (pI) of crucial *NCR* peptides vary between 4.78 and 10.15 implying that anionic, neutral and cationic *NCR* peptides could be essential for nitrogen-fixing symbiosis in *M. truncatula*. For this reason, the gene of *NCR* peptides with wide range of charges were selected to be edited (Fig. 3b). (v) Finally, we have selected four *NCR* genes, *NCR068*, *NCR089*, *NCR128* and *NCR161*, to be targeted by CRISPR/Cas9 editing. The pI of the encoded mature peptides varied between 5.75 and 8.34 (Fig. 3b). *NCR128* and *NCR161* are induced in the late infection zone and showed expression in the interzone but *NCR068* and *NCR089* are preferentially expressed in the interzone and at lower level in the nitrogen fixation zone based on the LCM transcription data (Fig. 3c,³³). The transcription activity of

the four selected *NCRs* was lower than *NCR169* and *NCR211* but similar or higher than *NCR-new35* (Fig. 3d).

Inducing mutations in selected *NCR* genes using the CRISPR/Cas9 system

To induce mutations in genes *NCR068*, *NCR089*, *NCR128* and *NCR161* by CRISPR/Cas9-mediated cleavage, sgRNA constructs targeting a single site of each gene were designed. The designed sgRNAs targeted either the upstream region of the gene close to the sequence coding for the first residues of the mature peptide or at least at a site prior the sequence coding for the last cysteine residue (Fig. 3e). These sgRNA constructs along with the control empty vector and the constructs targeting *NCRnew-35* and *NCR169* were introduced into *M. truncatula* 2HA roots using *A. rhizogenes*-mediated hairy root transformation. Transformed plants were inoculated with *S. medicae* WSM419 at 2-3 weeks after removal of non-transformed roots and the symbiotic phenotype of transformed plants were scored at 4-7 wpi. The aerial part of plants transformed with empty vector or the constructs targeting *NCR068* did not show the symptoms of nitrogen deficiency (Fig. 1a and c) and plants transformed with constructs targeting genes *NCR089*, *NCR128* and *NCR161* displayed the similar growth habit (data not shown) indicating the effective symbiotic nitrogen fixation capacity of these plants. Nodules formed on roots transformed with sgRNA constructs of the four selected *NCRs* were elongated and pink suggesting that they were functional nodules (Fig. 4j, l, n and p). The staining of nodule sections with SYTO13 revealed that the bacterial colonization of nodules targeted for mutagenesis of genes *NCR068*, *NCR089*, *NCR128* and *NCR161* was identical to the wild-type indeterminate nodules developed on empty-vector transformed roots (Fig. 4i-p). Contrary, 2HA plants carrying mutations in *NCR169* or *NCR-new35* in each hairy root displayed retarded growth, showed the symptoms of nitrogen starvation characteristically (Fig. 1b) and slightly elongated white nodules were found on their roots (Fig. 4f, Figures S2 and S3). In addition, the occupancy of nodules mutagenized for *NCR169* and *NCRnew-35* showed the phenotype characteristic for the previously described deletion mutants^{10,13} (Fig. 1b, 4e-h, Figures S2 and S3).

To verify the result of CRISPR/Cas9 genome editing, the targeted regions were amplified from nodule sections following microscopic analysis and sequenced using a MiSeq platform. The sequence analysis of the amplicons was carried out with the online tool CRISPResso2. The Cas9-induced mutations were predominantly indels, with frequencies between 47 and 100% of total reads (Table 1.). The genotyping also revealed that individual nodules were often chimeric containing more than two alleles or the ratio of the two alleles deviated from

the expected 1:1 ratio indicating that the tissues of these nodules on hairy roots originated from multiple founder cells with different genetic background. The identified indels generated frameshift mutations in the targeted *NCR* genes presumably producing malfunctioning peptides. Nevertheless, all the nodules carrying biallelic mutations in the targeted *NCRs* were elongated and pink, indicating the presence of leghemoglobin, and were fully colonized by rhizobia indicating that peptides NCR068, NCR089, NCR128 and NCR161 are not required for the effective symbiotic nitrogen fixation between *M. truncatula* 2HA and *S. medicae* WSM419 (Fig. 4i-p. and S4).

The nodulation phenotype of stable mutant of *NCR068* regenerated from gene edited hairy roots is not strain dependent

To analyse the symbiotic phenotype of genome editing mutants in more detail, we regenerated transgenic plants from hairy roots mutagenized for *NCR* genes. As a proof of concept, we cut root segments that had been transformed with the editing construct of *NCR068* and stable transformants were regenerated using a slightly modified formerly published protocol³⁵. DNA was purified from the leaves of hairy root-derived regenerated plants and the sequence analysis revealed that all the plants were homozygous containing a single adenine insertion in *NCR068* (Fig. 5g). The one bp insertion generates a frame shift in the coding sequence of *NCR068* and the mutant sequence encodes a chimera peptide containing 27 NCR068-specific and 23 non-NCR068-specific residues before termination. The fourth cysteine residue in the conserved position is lacking in the chimeric NCR068 peptide and based on the necessity of cysteines for the function of NCRs *in planta*^{10,13}, we reckoned that the mutant NCR068 peptide does not function correctly.

To confirm the symbiotic phenotype of regenerated *ncr068* mutant, T1 seeds were synchronously germinated and planted into substrate with no nitrogen content. Three wpi with *S. medicae* WSM419, *ncr068* stable transformants did not show symptoms of nitrogen starvation and developed like wild type *M. truncatula* 2HA plants (Fig. 5a). Nodules, formed on the roots of stable transformant *ncr068* mutant plants were elongated similarly what we found on the hairy roots of composite plants induced by the transformation construct targeting *NCR068* (Fig. 5c and e). The staining of *ncr068* nodule sections with SYTO13 revealed infected cells in interzone and nitrogen fixation zone fully occupied with elongated bacteria, similarly as it was found in wild-type nodules and *NCR068*-targeted hairy roots (Fig. 1d and f, 5b and d). The nodulation phenotype of the regenerated mutant *ncr068* confirmed that the

peptide NCR068 is not essential for the nitrogen-fixing symbiosis between *M. truncatula* and *S. medicae* WSM419.

Different symbiotic efficiency has been reported for *Sinorhizobium* sp.^{36,37}, therefore we assayed the strain-dependent nodulation phenotype of regenerated *ncr068* mutant in response to *S. meliloti* strains 2011 and FSM-MA compared with *S. medicae* WSM419. The growth habit of wild-type *M. truncatula* 2HA and *ncr068* mutant plants was similar following inoculation with any of the tested strains (Fig. 6). We observed a similar shoot dry weight of wild-type and *ncr068* plants inoculated with *S. meliloti* FSM-MA or *S. medicae* WSM419 (Figure S5). In agreement with published results³⁸, the performance of wild-type and, in our study *ncr068* plants as well, inoculated with *S. medicae* WSM419 superseded the plants inoculated with *S. meliloti* 2011 (Figure S5). The *dnf7-2* mutant showed the previously described ineffective symbiotic phenotype with all three tested rhizobia strains (Fig. 6). In line with the growth habit, the nodules developed on wt and *ncr068* roots inoculated with the highly compatible strains *S. medicae* WSM419 and *S. meliloti* FSM-MA were cylindrical, pink and fully colonized by rhizobia but nodules elicited by the less effective strain *S. meliloti* 2011 were only slightly elongated (Fig. 6). In brief, no difference in plant development and nodule structure was noticed between wt and *ncr068* plants with the tested rhizobia which suggests that the symbiotic phenotype of plants lacking NCR068 is not strain-dependent.

Discussion

NCR peptides, present solely in IRLC clade and Dalbergoid legumes, modulate the differentiation of nitrogen-fixing endosymbiont bacteria^{4,5,8}. NCRs compose a large gene family with more than 700 members⁹ in *M. truncatula* and because this large number of NCRs, their function was thought to be redundant. The identification of few individual NCRs crucial for nitrogen-fixing symbiosis disproved this concept¹⁰⁻¹⁵. These essential NCRs except for NCR247, which was analysed by reverse genetic approach, were identified fortuitously using deletion mutant collections of *M. truncatula*. In addition, the large deletions in mutants *dnf4-1* and NF-FN9363 unveiled NCR peptides, NCR178, NCR341, NCR344 and NCR345, which are not crucial for the effective nitrogen-fixing symbiosis with the studied rhizobia strains^{11,13}. Moreover, no specific chemical properties, charge or amino acid composition have been observed for the NCRs which would discriminate between crucial and non-crucial NCR peptides. Therefore, the targeted gene editing is the most promising tool to

ascertain the need of a particular NCR peptide for nitrogen-fixing symbiosis. In this study, we used the CRISPR/Cas9 technology in the model legume *M. truncatula* to knock-out four selected *NCR* genes encoding for peptides with four conserved cysteine residues representing the anionic, neutral and cationic members of the gene family.

The *A. rhizogenes*-mediated hairy root transformation produces chimeric root system composed of different ratio of transformed and non-transformed roots depending on the efficiency of transformation. To identify transformed roots in a convenient and non-destructive way and improve gene editing efficiency, a modified vector system carrying the DsRed transformation visible marker and single-site gRNAs under the control of the *M. truncatula* U6 promoter were applied to generate mutations in the selected *NCR* genes in *M. truncatula* 2HA and R108 plants. Despite of its robustness, the *A. rhizogenes*-mediated hairy root transformation system has the disadvantage of occasionally emerged chimeric roots of a mixture of transformed and -non-transformed cells. In addition, the CRISPR/Cas9 system is not necessarily competent to edit the target sites in every cell or independent gene editing events occur in several transformed cells. Our results provided evidence of both cases; we found chimeric roots containing the mixture of wild-type and mutant alleles with different ratio, these roots mostly retained wild-type nodulation phenotype, and we detected roots having multi-allelic mutations at target sites. To eliminate these interfering effects and reveal the association between CRISPR/Cas9-induced mutations and the symbiotic phenotype without a doubt, we have sequenced the amplified fragment population of the targeted sites of SYTO13-stained and examined sections of individual nodules. This enabled the precise analysis of a large set of nodules and gene editing events could be coupled with the phenotype of each nodule.

We used two essential NCRs, NCR169 and NCR-new35, to prove the efficiency of the CRISPR/Cas9 gene editing tool in our *A. rhizogenes*-mediated hairy root transformation system and selected four NCRs (NCR068, NCR089, NCR128 and NCR161) to assay their requirement for symbiotic nitrogen fixation. We generated mutant versions of the four previously non-investigated *NCR*s and the analysis of the colonization of the corresponding nodules indicated that these four NCRs are not essential for the symbiotic interaction between *M. truncatula* 2HA and *S. medicae* WSM419. The removal of variable number of non-transgenic roots from each plant and the potential presence of root cells with non-edited genome did not enable the reliable analysis of the nitrogen starvation phenotype (stunted growth, yellow leaves) of the shoots of transformed plants. The nodules of all the previously identified mutants defective in essential *NCR*s were white and slightly elongated and showed

defects in colonization of the region corresponding to the nitrogen fixation zone^{10,11,13–15}. Therefore, we assessed the symbiotic phenotype of genome edited nodules mostly based on their size, colour, and the colonization of the nitrogen fixation zone. The coordinated sequence and phenotype analyses which we carried out on every single nodule, clearly demonstrated that CRISPR/Cas9-induced mutations generating corrupted open reading frames in the coding sequence of *NCR068*, *NCR089*, *NCR128* and *NCR161* did not affect the nodulation phenotype of gene edited roots, thus these NCR peptides are not essential for the nitrogen-fixing symbiosis between *M. truncatula* 2HA and *S. medicae* WSM419. Although the four target NCR genes were selected based on the common features of the encoded essential peptides (four cysteine residues in conserved positions, predicted signal peptide, nodule-specific expression, unique sequence, various pI;^{10,11,13}), these criteria were not adequate to identify essentials NCR peptides. This conclusion was supported by the genetic studies that identified non-essential peptides, NCR341, NCR344 and NCR345, possessing these features. The four NCR peptides targeted in this study extend the group of NCRs proved to be non-required for nitrogen fixation under the symbiotic conditions used in this study. Our findings show that peptides NCR068, NCR089, NCR128 and NCR161 are either not crucial or have minor or redundant roles in symbiosis. Alternatively, these peptides may be required under specific symbiotic or environmental conditions or essential for strain-specific symbiotic interaction.

We have demonstrated that the *A. rhizogenes*-mediated hairy root transformation was successfully adapted to screen the requirement of NCR peptides for nitrogen fixation. We have regenerated plants of gene edited knockout of *NCR068* gene directly from transformed hairy root segments to carry out a more detailed analysis of CRISPR/Cas9-induced mutations in stable transformants. A previous proteomic study identified peptide NCR068 in bacteroids³⁹ and we anticipated that NCR068 may have a crucial role in symbiosis. The mutant plants carrying a frame-shift mutation, induced by the insertion of a single base pair, in *NCR068* in their germline showed similar nodulation phenotype to nodules found on *A. rhizogenes*-mediated mutant hairy roots proving the efficiency and usability of CRISPR/Cas9-mediated mutagenesis using hairy root transformation. The stable *ncr068* mutant enabled us to assay the symbiotic phenotype with *S. medicae* strain WSM419 and two *S. meliloti* strains, 2011, a laboratory reference strain and FSM-MA, which is highly effective symbiotic partner of *M. truncatula* A17 and R108³⁷. All the three tested rhizobia strains induced the formation of wild-type nodules on the root of *ncr068*, indicating that its symbiotic phenotype is not strain-specific.

Our study demonstrates that an optimized gene editing CRISPR/Cas9 vector construct and the developed workflow (Fig. S6) to analyse nodules developed on hairy roots mediated by *A. rhizogenes*- transformation is an effective system to analyse the symbiotic function of *NCR* genes. Although the selected *NCR* peptides were found to be non-essential for nitrogen-fixing symbiosis, the application of the technology can identify additional indispensable and crucial *NCR* peptides for nitrogen-fixing symbiosis.

Materials and Methods

Plant material, growth conditions, bacterial strains and nodulation assay. *Medicago truncatula* cv. Jemalong 2HA line⁴⁰ (2HA) and the *M. truncatula* R108 accession (R108) were used for transformation experiments. For sterilization and scarification, seeds were soaked in concentrated sulfuric acid for 8 minutes, then rinsed with distilled water. Sterilized seeds were vernalized for 3 days on inverted 0.8 % (w/v) water agar plates at 4 °C and germinated overnight at room temperature before transformation. Seedlings were grown in zeolite substrate (Geoproduct Kft., Mád, Hungary) for five days before inoculation with rhizobia. Plants were grown in SANYO HM350 environmental test chambers at 23 °C at 16/8 hours light/dark periods.

M. truncatula 2HA plants were inoculated with *Sinorhizobium (Ensiferi) medicae* strain WSM419 (SmWSM419) carrying pXLGD4 plasmid⁴¹, constitutively expressing the *lacZ* reporter gene under the control of the *hemA* promoter. To test strain specific symbiotic phenotype of *ncr068* mutants, T1 plants were inoculated with *S. meliloti* 2011, *S. meliloti* FSM-MA and SmWSM419 (pXLGD4) strains. *M. truncatula* R108 plants were inoculated with *S. meliloti* FSM-MA. The rhizobium strains were grown overnight in liquid TA medium. The bacteria were pelleted, resuspended and diluted to OD₆₀₀ 0.1 in nitrogen free liquid Fahraeus medium⁴² and used for inoculation.

Construction of the CRISPR/Cas9 binary vector used for gene editing

For CRISPR/Cas9-based gene editing experiments, a modified version of the pKSE401 vector⁴³ was generated. DsRed fluorescent protein under the control of the 830 bp promoter of *UBQ10* gene from *Arabidopsis thaliana* was inserted into the *EcoRI* digested vector to allow the identification of transformed roots based on their fluorescence. This generated construct (pKSE401_RR) was further modified by replacing the AtU6 promoter with MtU6.6 promoter^{31,44} resulting the construct pKSE466_RR. The MtU6.6 fragment was amplified from the genomic DNA of *M. truncatula* A17 using the primers MtU6.6_prom_InF_F1 and

MtU6.6_prom_InF_R along with the guide cassette fragment amplified from pKSE401_RR using the primers pKSE401_empty_InF_F2 and pKSE401_InF_R2 (Table S1) and the amplified fragments were assembled into the *Hind*III digested pKSE401_RR vector backbone using In-Fusion cloning (Takara Bio, Kusatsu, Japan).

Selection of targeted genes and generating CRISPR/Cas9 constructs

To select the *M. truncatula* NCR genes to be targeted, we have analysed the previously collected and described NCR peptide sequences⁹. During the selection, first the signal peptide prediction was repeated with the SignalP 5.0 tool⁴⁵. If the prediction did not detect signal peptide, the whole peptide sequence was considered as mature peptide. Further on, NCR peptides possessing four cysteine residues in conserved position (X_n -C- X_5 -C- X_n -C- X_4 -C- X_n)⁸ were selected. The transcriptional activity of this set of NCR genes was analysed in the RNAseq data of *M. truncatula* nodules³³ and genes with more than 10 000 reads were retained. The final set of 157 NCR genes was classified based on their isoelectric point⁹ and few genes showing low similarity to other NCRs were selected from each category and were selected to design sgRNA constructs. All sgRNAs were designed using the online tool available at <http://crispor.tefor.net/>³⁴ using the *M. truncatula* A17 (MtrunA17r5.0) genome as reference⁴⁶. In each case, the targeted region was in the first half of the coding sequence of the NCR gene and upstream from the last conserved cysteine residue. To minimize the risk of off-target mutations all selected sgRNA had at least two mismatches comparing with potential off-target sites (Table S2). Targets with GN₁₉ NGG motif were preferred for optimal expression controlled by the MtU6.6 promoter. Plasmid constructs were assembled following the protocol described by Xing at al.⁴³

***Agrobacterium rhizogenes*-mediated hairy root transformation**

CRISPR constructs were introduced into the ARqua1 strain of *Agrobacterium rhizogenes* and used for hairy root transformation of *M. truncatula* 2HA and R108 seedlings as described by Boisson-Dernier et al.²⁸. Each CRISPR construct was transformed into the roots of at least 35 plants in transformation and experiments were repeated at least twice. Following transformation, the seedlings were grown for one week in square petri dishes containing antibiotic-free Fahraeus medium and then seedlings were transferred to selective Fahraeus medium supplemented with 25 mg/l kanamycin. To prevent root growth into the agar, the roots were carefully positioned between two layers of Whatman filter paper. Following the plant growth for additional two weeks, non-transgenic roots, identified based on the absence

of red fluorescence, were excised. After then, plants were transferred into zeolite supplemented with Fahraeus medium containing 2 mM NH_4NO_3 . Plants were further grown in plastic boxes to preserve rhizobium-free conditions and inoculated with rhizobia after 2-3 weeks. Prior inoculation with rhizobia, newly emerged non-transgenic roots were excised again.

Embryonic induction and plant regeneration from hairy-root cultures

Plantlets were regenerated from transformed hairy roots grown on plate under sterile conditions using a slightly modified protocol described previously³⁵. Three-week-old transgenic hairy roots were identified by DsRed fluorescence using a Leica MZ10-F fluorescence stereo microscope (Leica Microsystems GmbH, Wetzlar, Germany) and cut into approximately 0.5-1 cm segments and placed on modified MTR-2 medium (Murashige and Skoog medium including vitamins supplemented with 5.0 mg/l 2,4-Dichlorophenoxyacetic acid, 0.5 mg/l 6-Benzylaminopurine (BAP), 3 % (w/v) sucrose, 50 mg/l kanamycin, 250 mg/l cefotaxime solidified with 0.8 % (w/v) agar) to induce primary calli. After 2-3 weeks of incubation in dark at room temperature, calli were transferred to the modified MTR-3 medium supplemented with BAP (Murashige and Skoog medium including vitamins supplemented with 0.5 mg/l BAP, 2 % (w/v) sucrose, 50 mg/l kanamycin, 250 mg/l cefotaxime solidified with 0.8 % (w/v) agar) to induce embryonic calli and plates were kept in plant growing chambers with a photoperiod of 16 h of light and 8 h darkness at 24 °C. The calli were transferred to fresh medium in 1-2-week intervals. Shoots developed on MTR-3 + BAP medium were put on MTR-3 medium (without selective agent) in glass tubes and transferred to fresh medium on every 2-5 weeks to promote development and induce roots. Well-grown shoots were propagated by cutting the apical 2-3 cm of the shoots and they were transferred into separate tubes. After rooting, the T0 regenerated plants were transferred into jars containing zeolite, watered with Fahraeus medium supplemented with 2 mM NH_4NO_3 . The jars were kept in growth chambers under the same condition as described above and humidity in the jars was gradually decreased. Once new healthy leaves emerged, the plantlets were inoculated and tested for nodulation phenotype or alternatively, they were transferred to nutrient-rich soil in the greenhouse for seed collection.

Phenotypic characterization of transformed plants and microscopy analysis

To observe intact roots and nodules and detect fluorescence of transformed roots, a Leica MZ10F fluorescent dissecting microscope equipped with ET DsRED, and GFP LP filters was

used (Leica Microsystems). Transgenic nodules were harvested, promptly immersed into ice-cold 1xPBS (pH 7.4) and fixed with 4% paraformaldehyde. Fixed nodules were embedded in 5% (w/v) agarose gel prepared in 1xPBS and 100 μ m thick longitudinal sections were prepared with Leica VT 1200S vibratome. Nodule sections were stained in 1xPBS solution containing 5 μ M SYTO13 dye (Thermo Fisher Scientific, Waltham, MA, USA) for 30 minutes at room temperature. Images of stained nodule sections were captured using a Leica SP5 AOBS confocal laser scanning microscope (Leica Microsystems).

DNA isolation from sections of transgenic nodules

To purify DNA from fixed nodule sections, 2-8 sections per nodule sample were collected after SYTO13 staining and microscopic analysis and transferred into nodule lysis buffer [100 mM Tris-HCl (pH 8), 0.5M EDTA (pH 8), 0.2 % SDS, 5 mM NaCl, Proteinase K 0.02 μ g/ μ l]. The samples were homogenized using glass pestles and incubated at 56 °C overnight. Proteinase K was inactivated by incubating the samples at 95 °C for 15 minutes. Samples were spin down by microcentrifuge and cooled them to room temperature. Subsequently 2 μ L RNase A (from 10 mg/ml stock) was added and incubated at room temperature for 2 min. After that, 200 μ L binding buffer [2.55M Guanidium-HCl, 10 % Triton X-100, 37 % ethanol] was added and briefly vortexed, and 200 μ L of 96 % ethanol was added to the mix and vortexed immediately. The mixture was centrifuge at 8000 rpm for 30 second to pellet the tissue debris. The supernatants were then transferred to QIAprep Spin Miniprep columns (Qiagen, Hilden, Germany). Following centrifugation at 8000 rpm for 1 min, the flow-through was discarded and columns were washed two times with 500 μ L washing solution (10 mM Tris-HCl pH 7.5, 80% ethanol). After the second washing step flow-through was discarded and columns were centrifuged at 13,000 rpm for 3 min. DNA elution was performed by incubating the columns with 35-50 μ L of de-ionized water at 50 °C for 3 minutes and eluted DNA was collected by centrifugation at 13,000 rpm for 1 min.

Sequence analysis of targeted genes in gene-edited nodules

The analysis of the result of target site mutagenesis of transgenic roots targeted by the sgRNAs NCR169a, NCR169b and NCR169c was performed by Sanger sequencing of PCR fragments amplified from the target region using ncr169_F1 and ncr169_R17 primer pair. To analyse the edits of targeted regions of genes NCR169, NCR-new-35, NCR068, NCR128, NCR089 and NCR161 in nodule samples, the PCR amplified fragments were analysed by next-generation sequencing (NGS). The genomic regions of approx. 200 bp, including the target

sites, were amplified using GoTaq™ G2 polymerase (Promega corp. Madison WI, USA) according to the manufacturer's instructions and gene specific primer pairs containing 5' adaptor sequences (Table S1) The samples were sequenced on Illumina platform, generating 10 000 reads per sample. NGS sequence analysis was carried out by using CRISPresso2 program³² using default parameters.

Acknowledgements

This work was supported by the Hungarian National Research Fund/National Research, Development and Innovation Office grants OTKA 119652, 129547 and 132646 as well as by the Collaborative Research Programme ICGEB Research Grant HUN17-03. We thank H. Cs. Tolnainé for her skillful technical assistance and Z. Tóth for support in confocal microscopy imaging.

Author contributions

J.B.B. and P.K conceived the project. B.G., J.B.B., D.Á. and B.H. performed the experiments and analysed the data J.B.B. and P.K. wrote the manuscript that all authors reviewed.

Data availability

All data supporting the findings of this study are available within the paper and the corresponding supplementary materials published online.

Competing interests

The authors declare no competing interest.

References

1. Jones, K. M., Kobayashi, H., Davies, B. W., Taga, M. E. & Walker, G. C. How rhizobial symbionts invade plants: The Sinorhizobium - Medicago model. *Nat. Rev. Microbiol.* **5**, 619–633 (2007).
2. Vasse, J., De Billy, F., Camut, S. & Truchet, G. Correlation between ultrastructural differentiation of bacterioids and nitrogen fixation in alfalfa nodules. *J. Bacteriol.* **172**, 4295–4306 (1990).
3. Mergaert, P. *et al.* Eukaryotic control on bacterial cell cycle and differentiation in the

- Rhizobium –legume symbiosis. *Proc. Natl. Acad. Sci.* **103**, 5230–5235 (2006).
4. Van de Velde, W. *et al.* Plant Peptides Govern Terminal Differentiation of Bacteria in Symbiosis. *Science (80-.)*. **327**, 1122–1126 (2010).
5. Czernic, P. *et al.* Convergent Evolution of Endosymbiont Differentiation in Dalbergioid and Inverted Repeat-Lacking Clade Legumes Mediated by Nodule-Specific Cysteine-Rich Peptides. *Plant Physiol.* **169**, 1254–1265 (2015).
6. Wang, D. *et al.* A Nodule-Specific Protein Secretory Pathway Required for Nitrogen-Fixing Symbiosis. *Science (80-.)*. **327**, 1126–1129 (2010).
7. Yang, J. *et al.* A signal peptide peptidase is required for ER-symbiosome proximal association and protein secretion. *Nat. Commun.* **14**, 4355 (2023).
8. Mergaert, P. *et al.* A novel family in *Medicago truncatula* consisting of more than 300 nodule-specific genes coding for small, secreted polypeptides with conserved cysteine motifs. *Plant Physiol.* **132**, 161–173 (2003).
9. Montiel, J. *et al.* Morphotype of bacteroids in different legumes correlates with the number and type of symbiotic NCR peptides. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5041–5046 (2017).
10. Horvath, B. *et al.* Loss of the nodule-specific cysteine rich peptide, NCR169, abolishes symbiotic nitrogen fixation in the *Medicago truncatula* *dnf7* mutant. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 15232–15237 (2015).
11. Kim, M. *et al.* An antimicrobial peptide essential for bacterial survival in the nitrogen-fixing symbiosis. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 15238–15243 (2015).
12. Sankari, S. *et al.* A haem-sequestering plant peptide promotes iron uptake in symbiotic bacteria. *Nat. Microbiol.* **7**, 1453–1465 (2022).
13. Horváth, B. *et al.* The *Medicago truncatula* nodule-specific cysteine-rich peptides, NCR343 and NCRnew35 are required for the maintenance of rhizobia in nitrogen-fixing nodules. *New Phytol.* **239**, 1974–1988 (2023).
14. Zhang, R. *et al.* Nodule-Specific Cysteine-Rich Peptide 343 is required for symbiotic nitrogen fixation in *Medicago truncatula*. *Plant Physiol.* (2023) doi:10.1093/plphys/kiad454.
15. Gao, F. *et al.* NCR343 is required to maintain the viability of differentiated bacteroids in nodule cells in *Medicago truncatula*. *New Phytol.* (2023) doi:10.1111/nph.19180.
16. Zhang, Y., Malzahn, A. A., Sretenovic, S. & Qi, Y. The emerging and uncultivated potential of CRISPR technology in plant science. *Nat. Plants* **5**, 778–794 (2019).
17. Le, H. *et al.* CRISPR/Cas9-Mediated Knockout of Galactinol Synthase-Encoding

- Genes Reduces Raffinose Family Oligosaccharide Levels in Soybean Seeds. *Front. Plant Sci.* **11**, (2020).
18. Shu, H., Luo, Z., Peng, Z. & Wang, J. The application of CRISPR/Cas9 in hairy roots to explore the functions of *AhNFR1* and *AhNFR5* genes during peanut nodulation. *BMC Plant Biol.* **20**, 417 (2020).
19. Li, G. *et al.* Development of an *Agrobacterium*-mediated CRISPR/Cas9 system in pea (*Pisum sativum* L.). *Crop J.* **11**, 132–139 (2023).
20. Wang, L. *et al.* CRISPR/Cas9 knockout of leghemoglobin genes in *Lotus japonicus* uncovers their synergistic roles in symbiotic nitrogen fixation. *New Phytol.* **224**, 818–832 (2019).
21. Meng, Y. *et al.* Targeted mutagenesis by CRISPR/Cas9 system in the model legume *Medicago truncatula*. *Plant Cell Rep.* **36**, 371–374 (2017).
22. Zhang, H. *et al.* Efficient Generation of CRISPR/Cas9-Mediated Homozygous/Biallelic *Medicago truncatula* Mutants Using a Hairy Root System. *Front. Plant Sci.* **11**, 1–11 (2020).
23. Confalonieri, M. *et al.* CRISPR/Cas9-Mediated Targeted Mutagenesis of *CYP93E2* Modulates the Triterpene Saponin Biosynthesis in *Medicago truncatula*. *Front. Plant Sci.* **12**, 1–16 (2021).
24. Jain, D., Jones, L. & Roy, S. Gene editing to improve legume-rhizobia symbiosis in a changing climate. *Curr. Opin. Plant Biol.* **71**, 102324 (2023).
25. Limpens, E. *et al.* RNA interference in *Agrobacterium rhizogenes*-transformed roots of *Arabidopsis* and *Medicago truncatula*. *J. Exp. Bot.* **55**, 983–992 (2004).
26. Wang, Q. *et al.* Host-secreted antimicrobial peptide enforces symbiotic selectivity in *Medicago truncatula*. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6854–6859 (2017).
27. Yang, S. *et al.* Microsymbiont discrimination mediated by a host-secreted peptide in *Medicago truncatula*. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6848–6853 (2017).
28. Boisson-Dernier, A. *et al.* *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol. Plant-Microbe Interact.* **14**, 695–700 (2001).
29. Wang, D. *et al.* Optimized CRISPR guide RNA design for two high-fidelity Cas9 variants by deep learning. *Nat. Commun.* **10**, 1–14 (2019).
30. Brinkman, E. K., Chen, T., Amendola, M. & Van Steensel, B. Easy quantitative assessment of genome editing by sequence trace decomposition. *Nucleic Acids Res.* **42**, 1–8 (2014).

31. Jacobs, T. B., LaFayette, P. R., Schmitz, R. J. & Parrott, W. A. Targeted genome modifications in soybean with CRISPR/Cas9. *BMC Biotechnol.* **15**, 1–10 (2015).
32. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence analysis. *Nat. Biotechnol.* **37**, 224–226 (2019).
33. Roux, B. *et al.* An integrated analysis of plant and bacterial gene expression in symbiotic root nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J.* **77**, 817–837 (2014).
34. Concordet, J. P. & Haeussler, M. CRISPOR: Intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res.* **46**, W242–W245 (2018).
35. Crane, C., Wright, E., Dixon, R. A. & Wang, Z. Y. Transgenic *Medicago truncatula* plants obtained from *Agrobacterium tumefaciens*-transformed roots and *Agrobacterium rhizogenes*-transformed hairy roots. *Planta* **223**, 1344–1354 (2006).
36. Terpolilli, J. J., O’Hara, G. W., Tiwari, R. P., Dilworth, M. J. & Howieson, J. G. The model legume *Medicago truncatula* A17 is poorly matched for N₂ fixation with the sequenced microsymbiont *Sinorhizobium meliloti* 1021. *New Phytol.* **179**, 62–66 (2008).
37. Kazmierczak, T. *et al.* Specific Host-Responsive Associations Between *Medicago truncatula* Accessions and *Sinorhizobium* Strains. *Mol. Plant-Microbe Interact.* **30**, 399–409 (2017).
38. Larrainzar, E., Gil-Quintana, E., Seminario, A., Arrese-Igor, C. & González, E. M. Nodule carbohydrate catabolism is enhanced in the *Medicago truncatula* A17-*Sinorhizobium medicae* WSM419 symbiosis. *Front. Microbiol.* **5**, (2014).
39. Durgo, H. *et al.* Identification of nodule-specific cysteine-rich plant peptides in endosymbiotic bacteria. *Proteomics* **15**, 2291–2295 (2015).
40. Rose, R. J., Nolan, K. E. & Bicego, L. The development of the highly regenerable seed line Jemalong 2HA for transformation of *Medicago truncatula* - Implications for regenerability via somatic embryogenesis. *J. Plant Physiol.* **155**, 788–791 (1999).
41. Leong, S. A., Williams, P. H. & Ditta, G. S. Analysis of the 5’ regulatory region of the gene for δ -aminolevulinic acid synthetase of *Rhizobium meliloti*. *Nucleic Acids Res.* **13**, 5965–5976 (1985).
42. Fahraeus, G. The Infection of Clover Root Hairs by Nodule Bacteria Studied by a Simple Glass Slide Technique. *J. Gen. Microbiol.* **16**, 374–381 (1957).
43. Xing, H. L. *et al.* A CRISPR/Cas9 toolkit for multiplex genome editing in plants. *BMC*

Plant Biol. **14**, (2014).

44. Kim, G. B. & Nam, Y. W. Isolation and Characterization of *Medicago truncatula* U6 Promoters for the Construction of Small Hairpin RNA-Mediated Gene Silencing Vectors. *Plant Mol. Biol. Report.* **31**, 581–593 (2013).
45. Almagro Armenteros, J. J. *et al.* SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat. Biotechnol.* **37**, 420–423 (2019).
46. Pecrix, Y. *et al.* Whole-genome landscape of *Medicago truncatula* symbiotic genes. *Nat. Plants* **4**, 1017–1025 (2018).

Figure legends

Figure 1. Plant gross and nodulation phenotype of *Agrobacterium rhizogenes*-mediated transgenic *M. truncatula* Jemalong 2HA hairy-roots mutagenized with CRISPR/Cas9 gene editing. No sign of nitrogen deficiency was observed on plants transformed with empty vector (a) or the construct targeting *NCR068* (c) 4 weeks post inoculation with *S. medicae* WSM419. Transgenic roots were identified by fluorescence of DsRed protein. Elongated pink nodules, showing the characteristic features of indeterminate nodules, developed on transgenic roots. These nodules were fully colonized by rhizobia visualized by the fluorescent dye SYT013 (d and f). (b) Targeted mutagenesis of *NCR169* resulted in transgenic plants showing the symptoms of nitrogen starvation. Small, spherical and white ineffective nodules were formed on *NCR169*-targeted roots. (e) *ncr169* mutant nodules did not contain infected cells in the region corresponding to the nitrogen fixation zone of wild-type plants. Inset images in (d-f) show the higher magnification of cells in nodule regions indicated by white rectangles. (g and h) The targeted regions were amplified and sequenced from the transgenic nodules. The sequence analysis revealed biallelic mutations causing frameshifts in edited *NCR169* (g) and *NCR068* (h) genes. ZII: infection zone; IZ: interzone; ZIII: nitrogen fixation zone; EV, empty vector

Figure 2. The nodulation phenotype of hairy roots transformed with the construct pKSE466-RR-MtNCR169a was associated with the proportion of wild-type *NCR169* allele or with the presence of mutant *NCR169* lacking a single residue. White (Fix-) and pink (Fix+) nodules were randomly harvested 3-5 wpi with *S. medicae* WSM419 and the result of gene editing was identified by the sequencing of the targeted region using an Illumina platform. The ratios of reads assigned to mutant (brown ochre) and wild-type (blue) alleles are presented in pie

chart. Fix- nodules (**b**, **d**, and **f**) carried almost exclusively insertions or deletions resulting in frameshift in the coding sequence of *NCR169* (**a**, **c** and **e**). Contrary, pink nodules (**g**, **i** and **k**) carried either unmodified alleles on a large scale (>40%) (**h** and **j**) or a deletion of 3 base pairs that did not result in the loss of function of *NCR169* (**l**).

Figure 3. Selection of *NCR* genes to be targeted by CRISPR/Cas9 mutagenesis. (**a**) *M. truncatula* *NCR* genes were grouped based on the presence of a signal peptide in their encoded peptide sequence and containing four cysteine residues in conserved positions. (**b**) the 248 *NCR* peptides complying with all the three features (**a**) were classified based on the charge of the encoded mature peptides. *NCR* peptides crucial for symbiosis are highlighted in red, peptides targeted in this study by gene editing are in black and peptides found to be non-essential in the former study¹³ are in purple. (**c**) Relative spatial expression of four *NCR* genes essential for symbiosis and four *NCRs* mutagenized using CRISPR/Cas9 method in this study. Expression was determined by RNAseq of nodule zones obtained with laser-capture microdissection described in the study³³. (**d**) The total read number of the *NCR* transcripts identified by RNAseq and analysed in panel (**c**). All the four *NCRs* selected for gene editing show higher transcriptional activity than *NCR-new35* showing the lowest expression activity among the crucial *NCR* genes. (**e**) The cleavage sites of Cas9 on the sequences of *NCR* genes targeted by CRISPR/Cas6 gene editing in this study. The target sites of each guideRNA were designed upstream of the nucleotides coding for the fourth conserved cysteines of each *NCR* peptides. ZIII: nitrogen fixation zone; IZ: interzone; IID: distal part of infection zone; IIP: proximal part of infection zone; pI: isoelectric point

Figure 4. (a-p) Nodulation phenotype of *M. truncatula* 2HA (**e**, **f**, **i-p**) and R108 (**g** and **h**) plants mutagenized using the CRISPR/Cas9 method with *A. rhizogenes*-mediated hairy root transformation. Transformation with gene editing construct designed to target *NCR-new35* (**e-h**) resulted in ineffective white and slightly elongated nodules while the homozygous or biallelic mutant alleles of *NCR068* (**i** and **j**), *NC089* (**k** and **l**), *NCR128* (**m** and **n**) and *NCR161* (**o** and **p**) were similar to the elongated and pink nodules formed on empty vector-transformed roots of *M. truncatula* 2HA (**a** and **b**) and R108 (**c** and **d**) plants. Images of nodule sections following SYTO13 staining and whole nodules were taken 4-5 weeks post inoculation with *S. medicae* WSM419 used for *M. truncatula* 2HA and *S. meliloti* FSM-MA for R108 plants. (**q-v**) Allele-specific results of CRISPR/Cas9 editing of the targeted *NCR*

genes identified by next generation sequencing on Illumine platform. Black and brown rectangles show deleted and inserted nucleotides, respectively.

Figure 5. (a) Stable *ncr068* mutants regenerated from segments of transgenic hairy roots with induced mutations in *NCR068* using the CRISPR method retained wild type symbiotic phenotype similar wild wild-type (WT) *M. truncatula* 2HA plants 3 wpi with *S. medicae* WSM419. (b and c) The gene edited nodules developed on *A. rhizogenes*-transformed hairy-roots were elongated and pink with colonized cells in the nitrogen fixation zone. (d and e) Nodules developed on the roots of regenerated stable *ncr068* mutants were similar to the nodules found on hairy-roots mediated by *A. rhizogenes*. (f) The selected hairy root (NCR068-5_S138) was chimeric and composed of cells with homozygous insertions of A or G in the targeted region of *NCR068*. (g) The regenerated stable *ncr068* mutants (Ac2211) carried a homozygous A insertion causing a frameshift in the coding sequence of *NCR068*. (h) The alignment of WT and mutant NCR068 peptide sequences encoded by the mutated gene in the stable *ncr068* mutant. Identical residues are highlighted by black background, conserved Cys residues are in red. SP, signal peptide; MP, mature peptide

Figure 6. Stable *ncr068* T1 mutant plants inoculated with three rhizobia strains show wild-type symbiotic phenotype. Wild-type *M. truncatula* 2HA (WT), the deletion mutant deficient in *NCR169* (*dnf7-2*) and *ncr068* mutant plants were inoculated with *S. medicae* WSM419, *S. meliloti* FSM-MA or *S. meliloti* 2011 and scored for symbiotic phenotype at 5 wpi. WT and *ncr068* plants showed similar growth habit but presented less effective symbiotic interaction with *S. meliloti* 2011 compared with the two other strains. Mutant *dnf7-2* showed nitrogen deficiency with all three tested rhizobia strains. SYTO13-staining revealed a similar colonization of WT and *ncr068* nodules while only bacteria in infection threads and sporadic saprophytic bacteria fluoresced in the basal part of *dnf7-2* nodules.

Table 1. Editing efficiency of different targeted *NCR* genes in nodules harvested from hairy roots mediated by *A. rhizogenes* transformation. Randomly selected nodules were harvested 3-5 wpi with *S. medicae* WSM419 and total DNA was isolated from fixed nodule sections. The PCR amplified target regions were sequenced on Illumina platform and the reads were analysed by using the CRISPresso2 tool. The gene *NCR169* was targeted for gene editing as a control. Three pinkish elongated nodules were selected randomly from hairy roots and also

three small white nodules were picked randomly from plants which did not form any pink nodules on their roots.

*One of the pinkish nodules carried mutant alleles on large proportion (94%). This mutation was a three bp deletion that resulted in an in frame deletion of the first Glu residue of the mature peptide NCR169. The lack a of this single residue probably did not abolish the function of peptide NCR169.

Additional information

Supplementary information

Fig. S1. Analysis of three sgRNAs targeting *NCR169*.

Fig. S2. The symbiotic phenotype of nodules formed on *M. truncatula* Jemalong and R108 hairy roots edited for *NCR-new35*.

Fig. S3. The symbiotic phenotype of nodules formed on *M. truncatula* Jemalong and R108 hairy roots edited for *NCR169* and *NCR-new35*.


Fig. S4. The nodulation phenotype of hairy roots edited for *NCR089*, *NCR0128* and *NCR161*.


Fig. S5. The shoot biomass of stable *ncr068* mutant plants inoculated with three rhizobia strains.


Fig. S6. The workflow applied for simultaneous genetic and phenotypic analysis of gene edited individual nodules.




Table S1. List of primers used in this study.

Table S2. Off-target prediction using the CRISPOR tool.









S. medicae WSM419

WT *dnf7-2* *ncr068* T1




S. meliloti FSM-MA

WT *dnf7-2* *ncr068* T1



S. meliloti 2011

WT *dnf7-2* *ncr068* T1



	NCR169		NCR068	NCR089	NCR128	NCR161
<i>Number of sequenced nodules</i>	3 white	3 pink	15	13	10	20
<i>Number nodules carrying max. 50% WT allele</i>	3	2	15	13	9	19
<i>Number of fully mutant nodules (mismatch/indel)</i>	3	0*	14	13	9	12
<i>Number of fully mutant nodules (only indel)</i>	3	0*	7	13	9	12
<i>Nodule samples carrying only indel alleles</i>	100%	0%	47%	100%	90%	60%