

Original Article

A Review on Ficus racemosa Linn: Medicinal Herb

Corresponding author: Thorat Sheela S *

Kumbhar Ujwala M, Mali Ramling D, Patil Poonam J, Tarlekar Sneha T

S.D. Patil Institute of Pharmacy, Urun-Islampur, Annasaheb Dange College of B. Pharmacy, Ashta
Department of Pharmaceutical Chemistry

*ssheelathorat@gmail.com

ujwalakumbhar897@gmail.com, ramlingmali@gmail.com

Received on: 17-02-2020; Revised and Accepted on: 28-02-2020

ABSTRACT

From ancient time, Herbal medicines are used in all over India. Ficus racemosa L. is widely used plant which belongs to family Moraceae. This is popular in all over India and these are especially popular in indigenous system of medicines. the F.racemosa L. widely distributed in Asia, Africa, America. The various parts of plant are used such as leaves, bark, seed, root, stems and fruits. The plant consist of different chemical constituent such flavanoid glycoside, alkaloids, phenolic acids , steroids , saponins , coumarins , tannins , triterpenoids , flavonoids, vitamin C. The chemical constituent presents in plants are responsible for pharmacological activity. The various parts of plant shows different pharmacological activity such as ,anticancer, anthelmintic antibacterial, antipyretic, wound healing anticarcinogenic ,antidiuretic, antitussive, anti-inflammatory, antidiarrheal, antifilarial, hepatoprotective ,antiulcer, antiasthmatic activity.

Keywords : Ficus racemosa L., Chemical constituent ,Pharmacological activity

1. Introduction

From ancient times, Herbal medicines were used in India. World Health Organization has estimated that 80% of population of india use the herbal medicines.^{1, 2} Many plant species have been identified to shows potent pharmacological activity against disease.

The Ficus racemosa L. plant is widely used plant belonging to family Moraceae. These plant commonly known as Cluster fig., gular fig., also known as Umber. These are popular in indigenous system of medicine like Ayurveda, Siddha, Unani and Homoeopathy. This plant consist of various chemical constituent such as alkaloids, flavanoid glycoside, phenolic acids, saponin, coumarines, tannins, vitamin C triterpenoids, the stem bark contains tannin, wax, and steroid such as β -sitosterol, lupeol, ceryl behenate, lupeol acetate. the unripe fruits are used as a source of dietary fiber, The different parts of plants are responsible for different pharmacological activity such as hypolipidemic, hypoglycemic, antidiuretic, antitussive, anti-inflammatory, antiulcer, antifungal ,antidiarrhoeal, Hepatoprotective and radioprotective. the bark of plant material is shows anti-inflammatory activity and anticancer activity, used in treatment of swelling of gums, swelling of neck, gonorrhea.

The Comprehensive account of morphology, geographical region and microscopical characters, and macroscopical characters, determination of impurity in different parts of plant, chemical constituents, uses, Pharmacological action are included in view of finding importance of plant.

Synonym: Ficus glomerata Roxb.

Kingdom : Plantae

Division: Magnoliophyta

Class: Magnolipsida

Order: Urtales

Family: Moraceae

Genus: Ficus

Species: racemosa

***Corresponding author:**

Thorat Sheela S, Department of Pharmaceutical Chemistry, S.D. Patil Institute of Pharmacy, Urun-Islampur Annasaheb Dange College of B. Pharmacy, Ashta
Email: ssheelathorat@gmail.com

Ayurvedic Properties of *F. racemosa* L.⁷

Rasa : Guru

Virya : Sheet

Vipak : Katu

Doshagnata : Kapha, Pitta



Fig no.1 Images of fruits of *Ficus Racemosa* Linn.

Geographical Source:

This plant of *Ficus racemosa* L. is cultivated in Australia, Malaysia, South East Asia, Pakistan, Shrilanka and different regions of India.

Macroscopical Character:

The leaves are dark green, ovate and elliptic in shape, the fruits are found in large bunches, the fruits are green in colour which turn to orange, which gives reddish or dark crimson colour on ripening. the roots are long and brown in colour. It has characteristic odour and slightly bitter in taste.

Microscopical Character:

The cork is made up of polygonal cells. The phellogen is made up of one to two layers of thin layers of cells. The phelloderm contains small groups of sclereids so This sclereids are lignified in nature with simple pits. the parenchymatous cells contains calcium oxalate crystal(prism) and brownish matter. The phloem rays may be uniseriate, multiseriate, homocellular or heterocellular. The Phloem fibres are non-lignified in nature. Starch grains are ovoid and spherical in nature¹⁰.

Chemical constituent:

Leaf	Sterols, tannins and flavonoids, triterpenoids (Lanosterol) and alkaloids. A new tetracyclic triterpene glauanol acetate, racemosic acid were isolated from the leaves.
-------------	---

Stem-Bark	Tannin, wax, saponin glauanol acetate, β -sitosterol, lupeol, ceryl behenate, lupeol acetate, α -amyrin acetate, leucoanthocyanidin, and leucoanthocyanin from trunk bark, lauanol acetate, lupeol, β -sitosterol and stigmasterol were isolated from stem bark
Trunk-Bark	Upenol, β -sistosterol and stigmasterol.
Fruit	Glauanol, β sitosterol, glauanolacetate, glucose, tiglic acid, esters of taraxasterol, lupeolacetate, friedelin, higher hydrocarbons and other phytosterol
Root	Cycloartenol, euphorbol and its hexacosanoate, taraxerone, tinyatoxin; bark euphorbol and its hexacosanate, ingenol and its triacetate, taraxerone.
Latex	a-amyrin, β -sitosterol, cycloartenol, cycloeuphordenol, 4-deoxyphorbol and its esters, euphol, euphorbinol, isoeuphorbol, palmitic acid, taraxerol, tinyatoxin, tirucallol, trimethyl ellagic acid.

The concentration of minerals in Bark of *F. racemosa* Linn.:

The various minerals found in bark of *F. racemosa* Linn. It contains abundant concentration of potassium, calcium, and chloride of minerals. The bark acts as good source of metal element which includes zinc, copper, phosphorous, nickel, iron, magnesium.⁶

Preliminary Phytochemical Analysis of *F. racemosa* L. plant:

The different extract has been prepared by using various solvent system and such extract have been employed for phytochemical test in identification of chemical constituent.

Sr.No.	Chemical constituent	Chemical Test ¹⁷
1.	Alkaloid	Mayer's test, Hager's test, Picric acid test
2.	Flavanoid	Shinoda test
3.	Triterpenoids	Libermann-Burchard Test
4	Glycosides	Saponin,
5	Tannin	Goldbeater skin test, Ferric chloride test

Uses:

Roots: Dysentery, pectoral complaints, hydrophobia, diabetes, treatment of mumps

Barks: Gynecological disorder, cooling, acrid, treatment of asthma, piles.

Fruits: Astringent, stomachic, treatment of Leucorrhoea, blood disorder, burning sensation, fatigue, urinary discharge, leprosy, menorrhagia, removing of intestinal worms, dry cough, loss of voice, kidney disease, spleen disease, miscarriage.

Latex: Healing of wounds, aphrodisiac, ant diabetic, antidiarrheal. Leaf buds: Improving of skin complexion.

Leaves: ulcer, wounds, glandular swelling, mouthwashes preparation, abscess¹⁶⁻¹⁸

Identity, Purity and Strength of *Ficus racemosa* Linn

The following parameters were measured to identify presence of foreign matters in *F. racemosa* Linn. The different extract of *F. racemosa* Linn. has been used for following test.^{6,15}

Foreign matter	Not more than 2%
Total Ash	Not more than 14%
Acid insoluble ash	Not more than 1%
Alcohol soluble extractive	Not more than 7%
Water soluble extractive	Not more than 9%

Pharmacological activity:

Antidiuretic: The decoction of the bark of *F. racemosa* L. claimed for antidiuretic and the three doses was administered in rats.(250, 500, 1000mg/kg). It has rapid onset of action although, it has greater peak plasma concentration so it also reduces excretion of the urinary Na⁺ level.^{11, 40}

Antitussive: The methanolic extract of stem bark was employed in antitussive activity against cough induced model so, sulphur dioxide gas usually used in mice. the maximum inhibition of 56.9% at dose of 200 mg/kg for 90 min after administration of drug.²⁴

Anthelmintic: The crude extracts of bark were evaluated for anthelmintic activity so, in this case The bark were extracted with methanol and ethanol and water as solvent. Aqueous extract of *F. racemosa* L. was evaluated by using adult earthworms such as *Pheretima posthuma*. therefore they exhibit a dose - dependent inhibition of spontaneous paralysis and these response was compared with 3% piperazine citrate. From the all the observation it was concluded that , The aq extract of drug was showed vermifugal activity.¹¹

Antibacterial: The hydro alcoholic extract of leaves was found effective against *Actinomyces viscosus*. From the observation it was concluded that, the hydro alcoholic extract was showed significant antibacterial activity, the minimum inhibitory concentration was found to be 0.08mg/ml.

Antipyretic: Methanolic extract of stem bark was evaluated on normal body temperature and yeast - induced pyrexia in albino rats, at doses of 100, 200 and 300 mg/kg body wt. p.o. It was a showed significant dose - dependent reduction in normal body temperature and yeast - provoked elevated temperature which extended up to 5 h after drug administration. The anti - pyretic effect of methanolic extract was compared with standard drug paracetamol.

Wound healing:

Ethanol extract of stem bark showed wound healing in excised and incised wound model in rats.

Ant filarial: The alcoholic and aqueous extract of plant has been evaluated for antifilarial activity so, Alcoholic as well as aqueous extracts caused inhibition of spontaneous motility of whole worm and nerve muscle preparation of *Setaria cervi* characterized by increase in amplitude and tone of contractions. Both extracts caused death of microfilariae in vitro LC50 and LC90 were 21 and 35 ng/ml, respectively for alcoholic, which were 27 and 42 ng/ml for aqueous extracts.

Antidiarrheal: Ethanolic extract of stem bark was studied for anti - diarrhoeal activity against different experimental models of diarrhoea in rats. It showed significant inhibitory activity against castor oil induced diarrhoea and PGE2 induced enteropooling in rats. These extracts also showed a significant reduction in gastrointestinal motility in charcoal meal tests in rats. The results obtained established its efficacy as anti - diarrheal agent.

Antifungal: The *F. racemosa* L. was employed for antifungal activity. The methanolic extract of *F. racemosa* shows inhibitory activity against different fungi such as *Trichophyton mentagrophytes*, *Trichophyton rubrum*, and *Candida albicans*.^{26, 41,42}

Larvicidal : The different solvent extract of bark and leaf were employed for larvicidal activity The extract were prepared by using different solvent such as methanol, pet ether, acetone, hexane. The larvicidal activity of methanolic, hexane, pet ether, acetone extract of bark and leaf were used for toxicity studies against Larvae of *Culex quinquefasciatus*. from all observation it was showed that, all the extract of plant showed significant larvicidal activity however, acetone extract of bark showed highest larvicidal activity. The potent chemical constituent was separated by fractionating the column, the gluanol acetate was tetra peptide terpene and it was a new mosquito repellent drug and it was also effective against *A. aegypti*.

Anti - inflammatory: The anti -inflammatory activity of *F. racemosa* L. leaf extract was studied on carrageenan, serotonin, histamine and dextran - induced rat hind paw edema models. The extract (400 mg/kg) exhibited maximum anti-inflammatory effect of 30.4, 32.2, 33.9 and 32.0% with carrageenan, serotonin, histamine, and dextran -induced rat paw oedema, respectively. In a chronic test, the extract (400 /kg) showed 41.5% reduction in granuloma weight, The anti-inflammatory activity was compared with standard drug such as phenylbutazone. The ethanolic extract of leaves of *F. racemosa* L. showed significant inhibitory activity against COX-1 and LOX-2 with invitro IC50 values 90 and 18 micro meter.²⁰

Antiulcer: The 50 % ethanol extract of fruits was studied in different gastric ulcer models, viz pylorus ligation, ethanol and cold restraint stress induced ulcers in rats at a dose of 50, 100 and 200 mg/kg body weight p.o. for 5 days twice daily. The extract showed dose dependent inhibition of ulcer index in all three models of ulcer.

Analgesic activity: The ethanol extract of bark and leaves was evaluated for analgesic activity by analgesiometer at 100, 300 and

500 mg/kg and was found to possess dose dependent analgesic activity.²¹

Hepatoprotective: An ethanolic extract of the leaves was evaluated for hepatoprotective activity in rats by inducing chronic liver damage by subcutaneous injection of 50% v/v carbon tetrachloride in liquid paraffin at a dose of 3 mL/kg on alternate days for a period of 4 weeks. The biochemical parameters SGOT, SGPT, serum and alkaline phosphates were estimated to assess the liver function. In other study, the methanol extract of stem bark at the doses of 250 and 500 mg/kg was evaluated for its hepatoprotective activity in rats against carbon tetrachloride induced liver damage with silymarin as standard.²²

Hypercholesteromic activity:

The isolation of alpha amylin acetate was done from aerial roots of *F. bengalensis* and this isolated alpha amylin was fed to mice for 10 consecutive days after 10 days the decreased in the triglyceride and cholesterol, and LDL by 21.5 % 24.1% 21.2%, increased HDL to TC by 21.0% and 59.1% respectively hence from all above observation it was concluded that the alpha amylin acetate improve the plasma lipid profile and lowering of total plasma cholesterol.

Anticancer Activity: The methanolic extract of *F. racemosa* L. were used for anticancer activity when drug were administered by oral route of administration at dose of 200 to 300 mg /kg, it considerable decreases the xanthine oxidase synthesis and decrease in concentration of alpha glutaryl, lipid peroxidation, generation of hydrogen peroxide, DNA synthesis. The enhancement of renal glutathione so the *F. racemosa* was act as good chemo preventive agents, it was also analysed that this drug inhibit the KBr induced carcinomas.

Anti-asthmatic Activity: An ethanolic bark extract of *F. racemosa* L. were employed for Anti-asthmatic activity. This study were performed by using swiss albino mice, total 6 mice were used and divided into three groups. This activity studied by using two methods clonidine induced catalepsy in mice and Mast cell degranulation; hence finally it was observed that, An ethanolic extract of bark of *F. racemosa* showed significant anti-asthmatic activity.

Antioxidant: The petroleum ether, ethyl acetate, butanol, ethanolic extract of stem bark of *Ficus racemosa* Linn. Were subjected for antioxidant activity. This activity was investigated by using total phenolic content, Inhibitory effect on 1, 1 diphenyl 2 picrylhydrazyl radical assay. From the observation it was showed that, ethyl acetate, ethanolic stem bark extract of *Ficus racemosa* Linn. was showed significant antioxidant activity.

Antioxidant: The water and ethanolic extract were used for antibacterial activity by time resolved and steady state methods. The ethanolic extract of *F. racemosa* L. exhibited a dose dependent activity by steady state method. The free radical scavenging activity were done by measuring DPPH, hydroxyl radical scavenging activity and when standard drug compared with ethanolic extract of *F. racemosa* L. it showed significant lipid peroxidation activity.

Memory enhancing activity: The aqueous bark extract of *Ficus racemosa* Linn, used for investigation of memory enhancing activity. This activity improves the level of acetylcholine in brain however it was concluded that, it is used in Alzheimer disease.³⁹

Conclusion

From ancient times, The study of herbal medicine is important. The herbal medicine has fewer side effects as compared to synthetic medicines. The *Ficus racemosa* Linn. is a herbal plant which is versatile in nature and has many medicinal properties and before some decades the many research has been done on different part of *Ficus racemosa* Linn. The purity of *F. racemosa* of plant material has been analyzed by using different parameter such as ash value, acid soluble extractive value, alkaline insoluble extractive value. The detail description of geographical and microscopical and macroscopical characteristics has been discussed. The different parts of *F. racemosa* L. plant have many pharmacological properties which is beneficial to human being. the leaves, fruits, stem, bark has different pharmacological activity such as anti-inflammatory, antiulcer, antidiabetic, antidiarrheal, ant filarial, anthelmintic, antibacterial, antitussive, hepatoprotective, analgesic, antidiuretic, anticancer, Anti-asthmatic activity.

Acknowledgment:

The authors expressed their grateful to Principal of Annasaheb Dange College of Pharmacy, Ashta Dr. R. B. Jadhav.

Conflict of Interest:

We declare that we have no conflict of interest.

Reference:

1. Raut S.P., Choudhary K.A., Kar D.M. Plants in Traditional Medicinal system- future source of New drugs International Journal of pharmacy and Pharmaceutical Sciences 2009; 1(1)
2. Sumi Salma et.al, Investigation of the key Pharmacological activities of *Ficus racemosa* Linn. and analysis of its major bioactive polyphenols by HPLC-DAD Evidence based Complementary and alternative medicines 2016.
3. Khare C.P. Encyclopedia of Indian Medicinal Plants, Springer Publication, 2004: 216-217.
4. Nayar SL, Bisth BS The Pharmacognosy of the bark of *Ficus glomerata* Roxb. J.Sci. 1959; 18(1): 15-17.
5. Shah Sunil kumar et.al, *Ficus racemosa* Linn.: Its Potential food security and Rural Medicinal Management J. Pharm. Sci. and Res 2016; 8(5): 317-322.
6. Borhade Praveen et.al, Pharmacological Review on *Ficus racemosa* Linn. IJPRBS 2012; 1(6): 51-66.
7. Novotny V. et.al, Host specialization of leaf chewing insects' in new guinea rain forest Journal of Animal Ecology; 71: 400-412.
8. Khare C.P. Encyclopdia of Indian Medicinal Plants Springer Publications, 2004: 216-217.
9. Bhalerao Satish et.al, *Ficus racemosa* Linn.: A Comprehensive Review Journal of Applicable Chemistry 2014; 3(4) :1423-1431.
10. Dhasade et.al, A Review on Phytochemical and Pharmacological Potential of *Ficus racemosa* Linn., EJPMR, 2019; 6(3): 443-446.
11. Parakh Padma, *Ficus racemosa* Linn-An Overview, Natural Product Radiance 2009; 8(1):84-90.
12. C. Suresh et.al, Chemical Examination of the fruits of *Ficus glomerata*, J. Indian. Chem. Soc 1979; 56(12): 1269-1270.

13. Chalam Aruna Et.al, A Pharmacognostical properties of *Ficus racemosa* Linn Pharmacie globale 2010; 5(4) : 1-4.
14. Chopra R.N., Chopra I.C., Handa K.I., Kapur I.D. Indigenous drugs of India, U.N. dhur, Kolkata 1958; 674-675.
15. Vedavathy S., Rao D.N. Herbal folk medicines of Tirumala and Tirupati regions of chitor, andhrapradesh, Fitoterapia 1995; 66: 167-171.
16. Kirtikar KR, Basu BD, Indian Medicinal Plants International Book Distribution publisher, Rajpur 1991; 2(3): 551-552.
17. Trivedi, P., Hinde S and Sharma R.C. Preliminary Phytochemical and pharmacological studies on *Ficus racemosa*. Linn Journal Medicinal Research: 56: 1070-1074.
18. Mitra R., Bibliography on Pharmacognosy of Medicinal plants National Botanical Research Institute, Luknow 1985: 249-250.
19. The Ayurvedic Pharmacopoeia of India, Ministry of Health and Family Welfare, Government of India, New Delhi, 1989, 1(1): 117-118.
20. Mandal SC, Maity TK, Das J, Saba BP, Pal M. Anti-inflammatory Evaluation of *Ficus racemosa* Linn. leaf extract. Journal of Ethnopharmacology 2000; 72:87-92.
21. Mahajan MS, Gulecha VS, Khandare RA, Upaganlawar AB, Gangurde HH, Upasani CD. Anti-edematogenic and analgesic activities of *Ficus benghalensis*. International Journal of Nutrition, Pharmacology, Neurological Diseases 2012; 2:100-104.
22. Mandal SC, Maity TK, Das J, Pal M, Saha BP, 1999. Hepatoprotective activity of *Ficus racemosa* leaf extract on liver damage caused by carbon tetrachloride in rats. Phytotherapy Research, 13: 430-432.
23. Bhalerao Sachin, Deepa verma, Nikhil veli, *Ficus racemosa* Linn.: A Comprehensive Review, Journal of Applicable Chemistry 2014, 3(4)-1423-1431.
24. Rao BR, Murugesan T, Pal M, Saha BP, Mandal SC (2003): Antitussive potential of methanol extract of stem bark of *Ficus racemosa* Linn. Phytotherapy Research 17: 1117-1118.
25. Prakash deep, Amrit Kr.Singh, MdTahir Ansari, Prashant Raghav Pharmacological Potential of *Ficus racemosa*-A Review, International Journal of pharmaceutical Sciences Review and research 2013, 22(1) 29-34.
26. Vonshak A. Barazani et.al, Screening of South Indian Medicinal Plants for Antifungal activity against cutaneous pathogenic, Phytother Research, 2003, 17(9):1123-1125.
27. Joseph S and Justin R, 2010. Phytopharmacological Properties of *Ficus racemosa* Linn- An overview. International Journal of Pharmaceutical Science Review and Research, 3(2): 134-138.
28. Singh AB, Yadav DK, Maurya R, Srivastava AK. Antihyperglycaemic activity of α -amyrin acetate in rats and db/db mice. Natural Product Research, 2009; 23:876-882.
29. Edwin E, Sheeja E, Chaturvedi M, Sharma S, Gupta V. A Comparative study on anti hyperglycemic activity of *Ficus bengalensis*, Linn aerial roots and barks. Phcog Mag, 2008; 4:95-97.
30. Honda M, Hara Y. Inhibition of rat small intestinal sucrase and α -glucosidase activities by tea polyphenols. Bioscience, Biotechnology, and Biochemistry, 1993; 57:123-124.
31. Ahmed F, Chavan S, Satish A, Punith KR. Inhibitory activities of *Ficus benghalensis* bark against carbohydrate hydrolyzing enzymes-An in vitro study. Pharmacognosy Journal 2011; 3:33-37.
32. Gayathri M, Kannabiran K. Antidiabetic and ameliorative potential of *Ficus bengalensis* bark extract in streptozotocin induced diabetic rats. Indian Journal of Clinical Biochemistry 2008; 23:394-400.
33. Gayathri M, Kannabiran K. The Effects of oral administration of an aqueous extract of *Ficus bengalensis* stem bark on some hematological and biochemical parameters in rats with streptozotocin-induced diabetes. Turkish Journal of Biology 2009; 33:9-13.
34. Singh RK, Mehta S, Jaiswal D, Rai PK, Watal G. Antidiabetic effect of *Ficus bengalensis* aerial roots in experimental animals. Journal of Ethnopharmacology 2009; 123:110-114.
35. Shukla R, Anand K, Prabhu K, Murthy PS. Hypoglycaemic effect of the water extract of *Ficus bengalensis* in alloxan recovered, mildly diabetic and severely diabetic rabbits. International Journal of Diabetes in Developing Countries 1994; 14:78-81.
36. Sharma S, Chaturvedi M, Edwin E, Shukla S, Sagrawat H. Evaluation of the phytochemicals and antidiabetic activity of *Ficus bengalensis*. Int J Diab Dev Ctries 2007; 27:56-59.
37. Shinde et.al, Isolation of Phytochemical and Evaluation of Anti-asthmatic potency of *F.racemosa*; 2019 9(6) : 107-109.
38. Byahatti V.V., Rajvaidhya Saurabh, Invitro antioxidant activity of various extract of bark of *Ficus racemosa* Linn. Int J Pharm Sci and Res 2019; 10(3) : 1534-1539.
39. Ahmed F, Chandra J.N., Manjunaths, Acetylcholine and memory enhancing activity of bark extract of *Ficus racemosa* L. Pharmacognosy Research ; 2011; 3(4): 246-249.
40. Ratnasooriya WD, Jayakody JR, Nadarajah T, Antidiuretic activity of aqueous bark extract of

Srilankan *Ficus racemosa* L. in rat, Acta Bio Hung; 2003,54; 3(4) : 357-363.

41. Husain A, Virmani OP, et.al Dictionary of Indian Medicinal plants, CIMAP, Lucknow, 1992, India.
42. Sen AB, Choudhary BK, Chemical investigation of *Ficus glomerata* Roxb, J Indian Chem Soc 1971; 48: 1165-1168.

Article Citation:

Authors Name. Thorat Sheela S. A Review on *Ficus racemosa* Linn.: Medicinal Herb. JPR 2020;9(2): 7-11
DOI: