Phytochemical profiling of two wild unexplored flowers of India

Madhuri Vajha[1], Anuradha Das[2], Subhalakshmi Rout[3] and Sanjeet Kumar[3*]

[1]Department of Biosciences and Biotechnology, Krishna University, Machilipatnam, Andhra Pradesh, India

[2]Department of Botany, Sindri College, Sindri, BBMK University, Dhanbad, Jharkhand, India

[3]Ambika Prasad Research Foundation, Odisha, India

*Email-Id: sanjeetaprf@gmail.com

Licensed under a Creative Commons Attribution 4.0 International License

Abstract

Aristolochia indica and *Dioscorea bulbifera* are two plant species traditionally used in folk medicine for various ailments. Despite their widespread use, the phytochemical composition of flowers of these plants remains poorly understood. Present study aimed to investigate the bioactive compounds present in the flowers of *A. indica* and *D. bulbifera* using hydroethanolic extraction and phytochemical screening. The results revealed the presence of tannins, saponins, and phenolic compounds in both plant extracts, with *A. indica* flowers also containing reducing sugars. These findings provide valuable insights into the phytochemical composition of these plants and suggest their potential for use in the development of natural remedies.

Keywords: Wild flowers, phytochemistry, pharmacology

INTRODUCTION

Plants have been an integral part of human life since ancient times, serving as a source of food, medicine, shelter, and spiritual inspiration (Kumar, 2022; Kumar *et al.,* 2022; Manjula *et al.,* 2022; Mety *et al.,* 2023; Nayak and Kumar, 2023; Sinha *et al.,* 2023; Kumar *et al.,* 2023; Mety *et al.,* 2023; Dimri *et al.,* 2023; Singh and Kumar, 2024; Vimala *et al.,* 2024; Sharma *et al.,* 2024; Dimri *et al.,* 2024). The use of plants for medicinal purposes dates back to ancient civilizations, with evidence of plant-based remedies found in traditional systems of medicine such as Ayurveda, Unani, and Chinese medicine (Sinha *et al.,* 2024; Sethi *et al.,* 2024). Despite the advent of modern medicine, plants continue to play a vital role in healthcare, with many modern medicines derived from plant sources (Basole *et al.,* 2024; Jaiswal *et al.,* 2024). *Aristolochia indica* (Fig. 2) and *Dioscorea bulbifera* (Fig. 1) are two wild

plant species that have been used in traditional medicine for various ailments (Mathew *et al.,* 2011; Kumar *et al.,* 2012; Kumar *et al.,* 2021; Agarwal *et al.,* 2023). *A. indica*, commonly known as Indian birthwort, is a perennial climber used in Ayurvedic medicine to treat fever, rheumatism, and snake bites (Mathew *et al.,* 2011; Bhattacharjee and Bhattacharyya, 2013). *D. bulbifera*, also known as air potato, is a perennial vine used in traditional medicine to treat various conditions, including skin infections and birth control agents (Kumar *et al.,* 2012; Kumar *et al.,* 2013). Despite their widespread use in traditional medicine, the phytochemical composition of *A. indica* and *D. bulbifera* remains poorly understood. Phytochemical screening is a crucial step in the discovery of new medicinal plants, as it helps to identify the bioactive compounds responsible for their therapeutic effects (Kumar and Jena, 2017; Kumar *et al.,* 2017; Mohanty *et al.,* 2021; Swain *et al.,* 2022). Understanding the phytochemical composition of these plants can provide valuable insights into their potential for use in the development of new medicines. The present study aims to investigate the phytochemical composition of *A. indica* and *D. bulbifera* flowers using hydroethanolic extraction and phytochemical screening. The study will provide valuable information on the bioactive compounds present in these plants, which can be used to develop new medicines and healthcare products.

MATERIALS AND METHODS

The flowers of *A. indica* and *D. bulbifera* were collected from Sundargarh district in September 2024. Hydroethanolic extract (50 % ethanol & 50 % distilled water) of flowers were obtained and nine different bioactive compounds were detected using standard methods (Devi *et al.,* 2023; Marndi *et al.,* 2024).

Qualitative phytochemical analysis for secondary metabolites

Test for tannin: 2-5 drops of 0.1 % lead acetate solution were added to 1 ml of the filtrate of the flower extracts. The white gelatinous precipitate provided a positive result for the presence of tannins.

Test for saponin:1 ml of distilled water was added to 1 ml of the filtrate of the flower extracts and shaken vigorously. The persistent froth formation after shaking indicated the presence of saponin.

Test for flavonoids: 2 ml of 2 % NaOH solution and 3 drops of dilute HCl were added to 1 ml of the filtrate of the flower extracts. The colour of the filtrate initially turned to an intense yellow colour with NaOH solution and later turned colourless. This colour change confirmed the presence of flavonoids.

Test for terpenoids:1 ml of the filtrate of flower extracts was added with 6 drops of chloroform and placed in the water bath for a few minutes. Then 6 drops of concentrated H_2SO_4 were added. The reddish-brown interface confirmed the positive result for the presence of terpenoids.

Test for phenolic compounds:1 ml of the filtrate of flower extracts was taken and a few drops of 5 % Ferric chloride solution were added. The bluish-black colour provided a positive result of the phenolic compounds.

Test for reducing sugars: 1 ml of the filtrate of flower extracts was taken and 2 drops of Fehling's solution A followed by Fehling's solution B were added and kept in the water bath for a few minutes. The presence of red-orange precipitate confirmed the presence of reducing sugars.

Test for steroids: 1 ml of the filtrate of the flower extracts was added with 1 ml of chloroform and 1 ml of concentrated H_2SO_4 was added to it. The appearance of upper red and lower yellow with green fluorescence provided the positive result of the test for steroids.

Test for alkaloids: 1 ml of the filtrate of the flower extracts was taken and 3-4 drops of Dragendroff's reagent was added to it. The formation of a reddish-brown precipitate confirmed the presence of alkaloids.

Test for carbonyl compounds: 1 ml of the filtrate of the flower extracts was taken and 3-4 drops of 2,4-Di-nitrophenylhydrazine reagent was added to it. The formation of yellow crystals confirmed the presence of carbonyl compounds.

RESULTS AND DISCUSSION

The phytochemical screening of hydroethanolic extracts of *A. indica* and *D. bulbifera* flowers revealed the presence of several bioactive compounds. Both plant extracts were found to contain tannins, saponins, and phenolic compounds, indicating their potential antioxidant and antimicrobial properties (Table 1). *A. indica* flowers were also found to contain reducing sugars, which was not observed in *D. bulbifera* flowers. This difference in composition may be indicative of distinct biological activities and potential uses of the two plants. Notably, neither extract contained flavonoids, terpenoids, steroids, or alkaloids. The absence of these compounds suggests that the bioactive properties of *A. indica* and *D. bulbifera* flowers may be attributed to the presence of tannins, saponins, and phenolic compounds. Further studies are needed to fully elucidate the biological activities and potential applications of these plants.

Table 1: Phytochemical screening of hydroethanolic extract of *A. indica* and *D. bulbifera* flowers

Bioactive compounds	*A. indica* flowers	*D. bulbifera* flowers
Tannin	+ve	+ve
Saponin	+ve	+ve
Flavonoids	-ve	-ve
Terpenoids	-ve	-ve
Phenolic compounds	+ve	+ve
Reducing sugar	+ve	-ve
Steroids	-ve	-ve
Alkaloids	-ve	-ve

(+ve: positive; -ve: negative)



Fig. 1: Flowers of *D. bulbifera*



Fig. 2: Flowers of *A. indica*

The phytochemical screening of *A. indica* and *D. bulbifera* flowers revealed the presence of tannins, saponins, and phenolic compounds, which are known for their antioxidant, anti-inflammatory, and antimicrobial properties. The presence of these

bioactive compounds suggests that the flowers of these plants may have potential therapeutic applications, such as treating oxidative stress-related disorders, inflammation, and infections. The differences in phytochemical composition between the two plants also highlight the importance of further studies to fully elucidate their bioactive properties and potential uses. Future studies will focus on isolating and characterizing the bioactive compounds present in *A. indica* and *D. bulbifera* flowers, as well as evaluating their pharmacological activities using in vitro and in vivo models. Additionally, further research will be conducted to explore the potential uses of these plants in traditional medicine, including their efficacy and safety in treating various ailments. The findings of this study provide a foundation for further research and development of new medicines and healthcare products from these plants, which could contribute to the improvement of human health and well-being.

CONCLUSION

In conclusion, the phytochemical screening of *A. indica* and *D. bulbifera* flowers revealed the presence of bioactive compounds with potential therapeutic applications. The study highlights the importance of exploring traditional medicinal plants for their phytochemical composition and pharmacological activities. The findings of this study provide a foundation for further research and development of new medicines and healthcare products from these plants, which could contribute to the improvement of human health and well-being, and underscore the need for conservation and sustainable use of these valuable plant resources.

REFERENCES

Agarwal B, Rathore S, Krishnan R, Jaiswal A, Panda A, Hegde L and Kumar S. (2023). A check list on wild nutraceutical tuberous plants of India. Indian Forester. 149(5): 587-590.

Basole S, Jena N and Kumar S. (2024). Indian Medicinal Plants, Volume I. Ambika Prasad Research Foundation, Odisha, India. DOI: https://doi.org/10.5281/zenodo.14523569

Bhattacharjee P and Bhattacharyya D. (2013). Characterization of the aqueous extract of the root of *Aristolochia indica*: evaluation of its traditional use as an antidote for snake bites. Journal of Ethnopharmacology. 145(1): 220–226.

Devi RS, Satapathy KB and Kumar S. (2023). Validation of tribal claims for formulation of future drugs through evaluation of ethno-pharmacological values of *Ludwigia adscendens*. Medicinal Plants. 15(4): 691-697.

Dimri R, Sharma BP, Kumar A, Behera B and Kumar S. (2023). Medico Bio-wealth of India, Volume 8. APRF, Odisha, India. https://doi.org/10.5281/zenodo.8022779

Dimri R, Sinha MK, Moharana A, Ofoeze MA and Kumar S. (2024). Wild Nutraceutical Plants. APRF Publishers, India. pp 1-69. DOI: https://doi.org/10.5281/zenodo.10574104.

Guchhait KC, Lal S, Kumar S and Mishra S. (2024). Medicinal Asteraceae of India, Volume 1. Ambika Prasad Research Foundation, Odisha, India. DOI: https://doi.org/10.5281/zenodo.14005857.

Jaiswal A, Hossain E, Kumar N and Kumar S. (2024). Plants & Secondary Metabolites, Volume II. Ambika Prasad Research Foundation, Odisha, India. DOI: https://doi.org/10.5281/zenodo.13995335

Kumar A, Manas P, Pradhan I and Kumar S. (2023). Fruiting Plants & Wildlife. APRF Publishers, Odisha.

Kumar S and Jena PK. (2017). Tools from Biodiversity: Wild Nutraceutical Plants. Ed: James N Furze et al.: Identifying Frontier Research Integrating Mathematic Approaches to Diverse Systems / Sustainability. Springer, Switzerland. DOI: 10.1007/978-3-319-43901-3-9.

Kumar S, Mahanti P, Rath SK and Patra JK. (2017). Qualitative phytochemical analysis and antibacterial activity of *Dioscorea alata* L.: a nutraceutical tuber crops of rural Odisha. Journal Alt Med Res. 3(1): 122-122.

Kumar S, Mishra AK, Singh RK, Kumar SN and Marndi S. (2022). Orchids of Bonai Forest Division. APRF Publishers, India.

Kumar S, Parida AK and Jena PK. (2013). Ethno-Medico-Biology of Ban Aalu (*Dioscorea* species): a neglected tuber crops of Odisha, India. International Journal of Pharmacy and Life Sciences. 4(12): 3143-3150.

Kumar S, Tripathy PK and Jena PK. (2012). Study of wild edible plants among tribal groups of Simlipal Biosphere Reserve Forest, Odisha, India; with special reference to *Dioscorea* species. International Journal of Biological Technology. 3(1): 11-19.

Kumar S. (2022). Medicinal Plants. IntechOpen Publisher, London, United Kingdom.

Kumar SN, Mishra S and Kumar S. (2021). Documentation of Indigenous Traditional Knowledge (ITK) on Commonly Available Plants in Koira Range, Bonai Forest Division, Sundargarh, Odisha, India. Asian Plant Research Journal. 8(4): 83-95.

Manjula BL, Kumar A, Sahu JK, Dash SS and Kumar S. (2022). Yam. APRF Publisher, India. DOI: https://doi.org/10.5281/zenodo.7147528.

Marndi S, Barik M and Kumar S. (2024). Phytochemical and cytotoxicity analysis of *Eulophia explanata* Lindl. (Tuber extracts). African Journal of Biological Sciences. 6(4): 2751-2756. DOI: 10.48047/AFJBS.6.Si4.2024.2751-2756

Mathew JE, Kaitheri SK, Dinakaranvachala S and Jose M. (2011). Anti-inflammatory, Antipruritic and Mast Cell Stabilizing Activity of *Aristolochia Indica*. Iran Journal of Basic Med Sciences. 14(5):422-427.

Mety SS, Dash SS, Saradar B, Pattnaik G and Kumar S. (2023). Indigenous Traditional Knowledge and Ethnopharmacology. APRF Publisher, India.

Mety SS, Sahu JK, Malhotra C, Barman H and Kumar S. (2023). Medicinal Climbers. APRF Publisher, India. DOI: https://doi.org/10.5281/zenodo.7592862

Mohanty M, Choudhury R, Kumar S and Maggirwar R. (2021). Phytochemical screening and antibacterial activity of *Dioscorea bulbifera* L. fruits. Plant Archive. 21(9): 862-86.

Nayak S and Kumar S. (2023). Medicinal plants used by tribals of Odisha. APRF & SMPB, Odisha; APRF Publishers, India.

Sethi J, Jena N and Kumar S. (2024). Medicinal Plants of Rourkela Forest Division, Odisha, India. Asian Journal of Environment & Ecology. 23 (12):69-84. https://doi.org/10.9734/ajee/2024/v23i12635.

Sharma BP, Marndi S and Kumar S. (2024). Orchids of India, Food, Medicinal and Ecological Aspects. APRF Publishers, India. DOI: https://doi.org/10.5281/zenodo.11209084

Singh RK and Kumar S. (2024). Medico-Biowealth of India, Volume 18. Ambika Prasad Research Foundation, Odisha, India. DOI: https://doi.org/10.5281/zenodo.14581320.

Sinha MK, Dimri R, Sharma BP, Kumar S and Marndi S. (2023). Medicinal Grasses of India. APRF Publishers, India.

Sinha MK, Panda A, Mishra S and Kumar S. (2024). Medicinal Combretaceae of India, Volume I. Ambika Prasad Research Foundation, Odisha, India. DOI: https://doi.org/10.5281/zenodo.14582202

Swain J, Kumar S, Antunes LLC, Biswal AK and Jena PK. (2022). *Amorpophallus paeoniifolius* (Araceae): A nutraceutical for food disorders, novel bacterial & viral infections. Carpathian Journal of Food Science and Technology. 14(1): 118-136.

Vimala, Das K, Choudhary JR and Kumar S. (2024). Medico Bio-wealth of India. APRF Publishers, India. DOI: https://doi.org/10.5281/zenodo.11180206