

Full Length Research Paper

Woody species composition and structure of Amoro Forest in West Gojjam Zone, North Western Ethiopia

Liyew Birhanu*[†] Tamrat Bekele and Sebsebe Demissew

Department of Plant Biology and Biodiversity Management, Addis Ababa University, P. O. Box 3434,
Addis Ababa, Ethiopia.

Received 16 March, 2018; Accepted 10 May, 2018

Ethiopia is one of the biodiversity rich areas in the world. A study was conducted at Amoro Forest to determine the woody species composition, vegetation and population structure. Vegetation data were collected from 36 plots (900 m²) systematically laid along transects. A hierarchical cluster analysis with R software was used to identify plant communities. Structural analysis of the forest was performed based on frequency, density, diameter at breast height (DBH), basal area and importance value indices (IVI) of woody species. A total of 57 woody species belonging to 38 families were encountered, consisting of 19 tree (33.33%), 31 (54.38%) shrubs, and 7(4.37%) liana species. The vegetation of the forest was classified into four plant communities namely, *Calpurnia aurea-Vernonia myriantha*, *Vernonia myriantha-Euphorbia abyssinica*, *Laggera tomentosa-Solanecio gigas*, *Allophylus abyssinicus-Bersama abyssinica* based on cluster analysis. The total basal area and density of woody plants were 18.5 m²/h and 2860.49 stems/h respectively. The DBH distribution showed a reverse “J” shaped curve, meaning that there is active regeneration and recruitment in the forest. Four representative woody plant population structures were identified, that is, inverted J, Gauss I, Gauss II and J- shaped patterns. There is high anthropogenic effect and high dependence of the local community in Amoro Forest so that forest conservation and restoration measures should be required.

Key words: Ethiopia, Amoro Forest, vegetation communities, anthropogenic activities and population structure.

INTRODUCTION

Ethiopia is an important regional biological diversity hotspot due to wide ranges of altitude, and geographical features, such as high and rugged mountains, flat-topped plateaus, deep gorges, river valleys and rolling plains (Kelbessa et al., 1992; Woldu, 1999). These contributed

to emergence of a variety of habitats suitable for the evolution and survival of various plant and animal species.

The rich biodiversity resources, including forests, are being destroyed at an alarming rate largely due to human

*Corresponding author. E-mail: liyewmtu@gmail.com.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

related disturbances (Aynekulu, 2011). North western highlands of Ethiopia have only fragments of natural forests scattered and confined to inaccessible and sacred places (Wassie et al., 2005), which suggested that the highlands were once covered by high forests. Therefore, appropriate and immediate measures are required to maintain and restore the remaining natural forests. Accelerated deforestation that arise largely due to the conversion of forests to other agricultural land-use types and the overutilization of forest resources to satisfy the food and energy requirements of the increasing population are major environmental concerns in Ethiopia (Friis et al., 2001; Teketay, 2001). In the absence of suitable interventions, current patterns of deforestation will decimate the remaining forests in the near future (Teketay, 1992; Bekele, 1993, 1994).

Amoro Forest is composed of patches of dry evergreen Afromontane forests found in north western part of Ethiopia. It has been continuously exploited by surrounding communities for agricultural land, firewood collection and charcoal production and construction material. *Prunes africana* is highly sought by locals for fuel wood and charcoal production (Personal observation). Detailed description of resource exploitation needed analysis of floristic composition and structural complexity could contribute towards the conservation of vegetation resources. It is additionally important to document the remaining vegetation resources for posterity. Nevertheless, such data on woody plant composition, community and structural complexity are lacking for the Amoro forest in Dega Damot District and hence the current study. This is believed to contribute a lot to the effort being made in the development of a sound management plan for effective conservation of the forest resources in the study area.

MATERIALS AND METHODS

Study area

Amoro Forest is located in Dega Damot District, Amhara Regional State, North western Ethiopia (Figure 1). The forest is located between 10°50'06.53" latitude 37°35'51.94" longitude. The major town nearby is Feresbet and is far from 3 km from the forest. The district is also characterized by good climate for most of the year with annual rainfall between 900 and 1200 ml. Topographically, it consists of 35% mountainous, 30% ups and downs, 20% valleys and 15% plains. The soil of the district is reddish (clay soil), black soil, sand soil, brown soil and white soil types (Dega Damot Woreda Agricultural Office (DWAQ), 2017).

The area is divided into six land use types such as farm land, grazing land, shrub land, settlement, forest and bare land. Different types of crops cultivate in the study area crops including barley (*Hordeum vulgare*), wheat (*Triticum* spp.), faba bean (*Vicia faba*), teff (*Eragrostis tef*), maize (*Zea mays*) and potato (*Solanum tuberosum*) (DWAQ, 2017).

Amoro Forest belongs in the category of dry evergreen Afromontane vegetation (Friis et al., 2011). The vegetation of the

forest is dominated by trees; *Juniperus procera*, *Olea europaea* subsp. *cuspidata*, *Allophylus abyssinicus*, *Apodytes dimidiata*, *Bersama abyssinica*. Shrub and short stature trees such as; *Carissa spinarum*, *Discopodium penninervium*, *Dombeya torrida*, *Lobelia giberroa*, *Myrsine africana* and *Pittosporum viridiflorum*. The most dominant liana is *Ureia hypselodendron*.

Sampling design

Reconnaissance survey was made across the forest in order to get an impression of the site conditions and identify the possible sampling sites. Systematic sampling technique was used for vegetation data. Sampling sites were arranged along transects in two directions (NW and SW) from the top to the base of the forest. The number of plots per transect varies depending on length of the transect and accessibility of the sample plots. A total of 36 sampling plots of each 30 m x 30 m (900 m²) was used for woody species. The distance between two consecutive plots along a line transect was 50 m and the transects were 100 m apart.

Vegetation data collection

Diameter at breast height of each tree and shrub species with a diameter of ≥ 2.5 cm was measured using tree calipers. The number of individuals of each woody species was counted for each plot. Geographical data (altitude, latitude and longitude) were recorded using GPS for each plot. The percentage cover values estimated in each sample plot were converted into cover abundance values using 1-9 modified Braun-Blanquet Scale (Van der Maarel, 1979). Voucher specimens were collected, coded, pressed and dried for subsequent identification and verification at the National Herbarium (ETH), Addis Ababa University, using Volume 8 of Flora of Ethiopia and Eritrea (Hedberg et al., 2009b).

Data analysis

Cluster analysis

Hierarchical cluster analysis was performed using R-free statistical software version 3.4.1 to identify plant communities (R Core Team, 2017). Cluster analysis helps to group together a set of observations (plots or vegetation samples in this study) based on their attributes or floristic similarities (Kent and Coker, 1992). The community types identified from the cluster analysis were further refined in a synoptic table, and species occurrences are summarized as synoptic-cover abundance values. Synoptic values are the product of the species' frequency and average cover abundance value (Bekele, 1993). Dominant species of each community type were identified based on their synoptic values. Finally, the community types were named based on two dominant species. Shannon-Wiener diversity indices and Shannon's evenness were computed to describe species diversity of the plant community types in the vegetation (Kent and Coker, 1992).

$$H' = - \sum_{i=1}^s P_i \ln p_i \quad (1)$$

Where H' = Shannon diversity index; s = number of species, p = proportion of individuals or abundance of the i th species expressed as a proportion of total cover in the sample; and $\ln$ = the natural logarithm.




Figure 1. Map of the study area.

Shannon's evenness index (J) was also calculated using:

$$J = H' / H'_{\max} \quad (2)$$

where, H' = Shannon–Wiener Diversity Index; and $H'_{\max}$ = $\ln s$ where s is the number of species in the sample.

Structural data analysis

Structural characteristics (stem density, basal area, and diameter at breast height (DBH), class distributions) were calculated for each woody plant species. Generally, the following formulas were used to calculate frequency, density and basal area of woody species.

$$\text{Frequency (F)} = \frac{\text{Number of plots in which a species occurs}}{\text{Total number of plots}} \times 100 \quad (3)$$

$$\text{Density} = \frac{\text{Total number of stems of all trees}}{\text{Sample size in hectare}} \quad (4)$$

$$BA = \left(\frac{DBH}{2} \right)^2 \times \pi \text{ with } (\pi = 3.14) \quad (5)$$

where DBH is diameter at breast height.

Importance value indices (IVI) were computed for all woody species based on their relative density (RD), relative dominance (RDO) and relative frequency (RF). This index is used to determine

Table 1. Synoptic table of the four community types (values in bold refer to species used to name community types).

Species	C1	C2	C3	C4
<i>Calpurnia aurea</i>	7.78	3.7	3	0.7
<i>Vernonia myriantha</i>	6.22	6.7	4.2	4.5
<i>Euphorbia abyssinica</i>	2.89	5.7	4	2.5
<i>Clausena anisata</i>	4.8	4.6	1.25	0
<i>Laggera tomentosa</i>	3.1	0	6.5	0
<i>Solanecio gigas</i>	1.5	2.4	6.2	5.7
<i>Urera hypselodendron</i>	1.7	2.7	6.0	2.7
<i>Lobelia giberroa</i>	0	1.9	6.0	1.7
<i>Vernonia amygdalina</i>	0.4	0.3	4.2	0
<i>Allophyllus abyssinicus</i>	1.3	4.7	6.2	7.0
<i>Bersama abyssinica</i>	5.3	1.1	2	6.5
<i>Brucea antidysentrica</i>	5.1	2	2	3.2
<i>Discopodium penninervium</i>	3.4	3.5	3.2	3.0

the overall importance of each species in the forest system (Kent and Coker, 1992).

The Importance Value Index (IVI) for each woody species was computed using the following formula:

$$RDO = \frac{BA}{TBA} \times 100 \quad (6)$$

where

RDO= relative dominance

BA= Basal area

TBA= total basal area

$$Relative\ Density = \frac{Number\ individuals\ of\ each\ species\ per\ ha}{Total\ number\ of\ individuals\ of\ all\ species\ per\ ha} \times 100 \quad (7)$$

$$Relative\ Frequency = \frac{Number\ of\ sample\ plots\ containing\ a\ species}{Sample\ units\ for\ all\ species\ of\ the\ sample} \times 100 \quad (8)$$

$$IVI = Relative\ Density + Relative\ Dominance + Relative\ Frequency, \quad (9)$$

Population structure of woody species was analyzed for the entire forest and each woody species based on defined DBH classes. Diameter was classified into seven classes. DBH classes (cm) were 2.5-10, 10.1-20, 20.1-40, 40.1-60, 60.1-80, 80.1-100, >100 cm).

RESULTS AND DISCUSSION

Forest composition

A total of 57 woody species representing 38 families were recorded from 36 plots (Appendix Table 1). Of the total plant species identified, 19 species (33.33%) are trees while 31 species (54.38%) are shrubs. The remaining 7 species (4.37%) are lianas.

The results of this study show that the woody species composition of Amoro Forest (57 species) were higher than many Afromontane forests in Ethiopia and other tropical forest. For instance, Tadele et al. (2014) recorded a much lower species (50 species) in Zengena Forest in Ethiopia, Berhanu et al. (2016) recorded 66 woody species in Kuandisha Afromontane Forest in Ethiopia; whereas Neelo et al. (2015) recorded 47 woody species in open and exclosed dry woodland sites around *Molapo* farming areas of the Okavango Delta in Botswana. On the other hand, Zegeye et al. (2011) recorded a much higher woody species 143 species in Tara Gedam and Ababay forests in Ethiopia. The reasons for variation in floristic composition at the study sites could be due to excessive anthropogenic disturbances, disparity in conditions for regeneration and exploitation of some species. Besides this, the geographical location of the study area differs from other areas. According to Chen et al. (2004), environmental heterogeneity, regeneration success and competition are also important factors that shape species composition of forests.

Four plant community types were identified from the hierarchical cluster analysis, *Calpurnia aurea*-*Vernonia myriantha*, *Vernonia myriantha*-*Euphorbia abyssinica*, *Laggera tomentosa*-*Solanecio gigas*, *Allophyllus abyssinicus*-*Bersama abyssinica* (Figure 2; Table 1). Unfortunately, this study did not address analyses of a range of possible environmental variables except altitude that could shape the distribution of identified plant communities. The cluster results of four groups of communities in the present study suggest the existence of overlapping altitudes (Körner, 2000).

***Calpurnia aurea*-*Vernonia myriantha* type:** This occurs between altitudinal 2432 and 2615 m a.s.l. This community




Figure 2. Dendrogram of the vegetation community types obtained from hierarchical cluster analysis of Amoro forest. (1= Community type 1, 2= Community type 2, 3= Community type 3 and 4= Community type 4).

type is characterized by *C. aurea* and *V. myriantha* as the dominant shrubs. Other important species in this group include *Clausena anisata*, *Clusia abyssinica*, *B. abyssinica*, *C. anisata*, *Euphorbia abyssinica*, *Maytenus arbutifolia* and *Dovyalis abyssinica*.

***Vernonia myriantha*-*Euphorbia abyssinica* type:** This community type was situated at altitudinal of 2455 to 2622 m a.s.l. This community was dominated by *V. myriantha* and *E. abyssinica*. Some of the species are *C. anisata*, *C. aurea*, *A. abyssinicus*, *D. penninervium* and *U. hypselodendron*.

***Allophyllus abyssinicus*-*Bersama abyssinica* type:** This was encountered at an altitudinal range of 2463–2728 m a.s.l. This community type was dominated by *A. abyssinicus* and *B. abyssinica*. The tree layer consists of tree species such as *P. africana*, *A. dimidiata* and *E. abyssinica*. The shrub layer includes *C. aurea*, *L. giberroa*, *M. arbutifolia*, *D. penninervium* and *Liana* species are *Rubus steudneri* and *Embelia schimperi*.

***Lagera tomentosa*-*Solanecio gigas* type:** This community type extends from 2493 to 2567 m a.s.l. In this community type, *L. tomentosa* and *S. gigas* are the dominant shrub.


Shannon's diversity indices showed that use of the name as described above had the highest species diversity (3.62) followed by Communities 2 and 3 (Table 2). On the other hand, Community 1 had the highest number of species (47) and Communities 3 and 4 the lowest (29). Community 1 followed by communities had the highest evenness value. While Communities 2 and 3 had the least evenness value (Table 2).

Vegetation structure

The density of trees and shrubs with DBH greater than 2.5 cm in the Amoro Forest was 2860.5 stems ha⁻¹ (Appendix Table 2). The forest had relatively high density compared with that of Zengena Forest in North western Ethiopia (Tadele et al., 2014). On the other hand, the density was low compared to some other dry Afromontane Forest in Northwestern Ethiopia such as Tara Gedam and Abebaye Forests (Zegeye et al., 2011), Kuandisha Afromontane Forest (Berhanu et al., 2016) and peninsula of Zegie (Aleign et al., 2007). The density of the woody species varied considerably in the different forests. This could be attributed to variations in topographic gradients and habitat preferences of species forming the forest, and the degree of anthropogenic

Table 2. Species diversity, richness and evenness of the plant community.

S/N	Type of community	Species richness	Diversity index(H)	Shannon_Evenness (J)
1	<i>Calpurni aurea</i> - <i>Vernonia myriantha</i>	47	3.62	0.94
2	<i>Vernonia myriantha</i> - <i>Euphorbia abyssinica</i>	44	3.40	0.898
3	<i>Lagera tomentosa</i> - <i>Solanecio gigas</i>	29	3.00	0.892
4	<i>Vernonia myriantha</i> - <i>Allophyllus abyssinicus</i>	29	3.14	0.93

**Figure 3.** Distribution of woody species density among DBH classes (1=2.5-10 cm, 2=10.1-20 cm, 3=20.1-40 cm, 4=40.1-60 cm, 5=60.1-80 cm, 6=80.100 cm, 7=>100 cm).

disturbances (Whittaker et al., 2003). The density of the desired species reduced due to the devastating effects and lack of afforestation.

The species with the highest frequency value was *V. myriantha* (94.4%) followed by *A. abyssinicus* (88.9%), *P. africana* (83.3%), *Brucea antidyssentrica* and *D. penninervium* (75% each) and *E. abyssinica* (72.2%) (Appendix Table 2). High frequency value represents a wider distribution of the species in the forest. The variation in density and frequency among species may be attributed to differences in site conditions, species characteristics for adaptation, degree of exploitation and conditions for regeneration (Shibru and Balcha, 2004).

The total basal area of woody species in Amoro Forest was 18.5 m² ha⁻¹ and comparable to other Afromontane Forests in Ethiopia like Kuandisha Afromontane Forest (Berhanu et al., 2016) and Zengena Forest (Tadele et al., 2014). However, the basal area of this forest is small compared to other dry Afromontane Forests in Ethiopia like Tara Gedam and Abeyaye Forests (Zegeye et al., 2011), Wof-Washa Forest (Bekele, 1993) Adelle and Boditi Forests (Yineger et al., 2008). The normal basal

area value for virgin tropical forests in Africa is 23-37 m²/ha (Lamprecht, 1989). Thus, the basal area of Amoro Forest is low compared to tropical forests in Africa including dry Afromontane Forests found in Ethiopia. This may be due to Amoro Forest being dominated by shrubs. Besides this, cutting down trees and other factors influenced species by reducing the number of stems desired, affected species diversity and their size.

The distribution of woody species showed in decrease from lower to higher among DBH classes (Figure 3). The forest showed a reverse "J" distribution. These results were similar for those of Savadogo et al. (2007) in Tiogo Forest. The density was most abundant at DBH classes less than 10 cm (1318 stems ha⁻¹), 10 -20 cm (440 stems ha⁻¹), 20-40 cm (205.556 stems ha⁻¹), 40-60 cm (60.802 stems ha⁻¹), 60-80 cm (28.086 stems ha⁻¹) and >100 cm (17.901 stems ha⁻¹). This pattern indicates that the majority of the species had the highest number of individuals in lower DBH which in turn shows that the forest vegetation has good reproduction and recruitment potential.

Schefflera abyssinica had the highest IVI (41.0) followed




Figure 4. Size structure composition of a), *Bersama abyssinica* , b) *Ekebergia capensis* , c) *Schefflera abyssinica*, d), *Prunus africana* tree and shrub species from Amoro forest, Ethiopia. (A=2.5-10 cm, B=10.1-20 cm, C=20.1-40 cm, D=40.1-60 cm, E=60.1-80 cm, F=80.1-100 cm, G=> 100 cm).

by *P. africana* (30.7), *A. abyssinicus* (25.4), *A. dimidiata* (23.3), *V. myriantha* (17.8), *S. gigas* (12.3), *C. aurea* (10.7), *E. abyssinica* (10.6), *B. abyssinica* (9.4) (Appendix Table 2). IVI value is an important parameter that reveals the ecological significance of species in a given ecosystem (Lamprecht, 1989).

Population structure

The woody tree and shrub species in Amoro Forest were analyzed in four representative size structure (Figure 4a-d). The first pattern was an inverted J-shaped distribution

exhibited by species with high number of individuals in the first and second DBH classes and with gradual decrease of larger sized trees (Figure 4a). This pattern was recorded on *B. abyssinica*, suggesting adequate seedling reproduction and regeneration (Bekele, 1993). Gause II (Figure 4b) lacked individuals at intermediate DBH classes and indicates the sizes missing and individuals present at the high and lower DBH classes. The species *Ekebergia capensis* shows this type of pattern. This kind of distribution is observed when there is selective logging of medium sized individual. In my observation, the main reasons for the absence of medium DBH classes were selective cutting for construction,

charcoal production timber and firewood.

The J shaped was characterized by a higher proportion of larger individuals with over 40 cm DBH (15.12 stem/ha). The trend decreased towards lower DBH classes and absence of smaller individuals below 40 cm DBH (0 stem/ha). This regeneration pattern was observed in *S. abyssinica*. Such pattern shows poor reproduction (Bekele, 1993) due to the fact that it is possible most trees are not producing seeds due to age. It also indicates the presence of selective cutting of preferred size classes on juvenile individuals by local. The species with this type of pattern have large individuals that are less competent to reproduce and in a weak position of regeneration status selective cutting or grazing effects on juvenile individuals.

Gause I was represented by *P. africana* (Figure 4d). The first and second DBH classes having low number of individuals, a gradual increase in the number of individuals towards the medium classes, and subsequently a decrease in number towards the higher DBH classes. This pattern indicates a poor reproduction (Bekele, 1993) and recruitment of species which may be associated with human use selectively harvesting branches of stem for construction material and charcoal production.

CONCLUSION AND RECOMMENDATION

The woody plant of Amoro Forest is dominated by shrubs (54.38%). Euphorbaceae was the dominant family with six species (10.52%) followed by Asteraceae family with four species (7.01%). Amoro Forest is mainly a dry evergreen Afromontane Forest which is dominated by small sized tree and shrub species. The general size composition of trees in the forest was inverted J, however there were differences among tree species. *B. abyssinica* species showed typical healthy inverted, whereas *S. abyssinica* species J with predominance of adult and absence of juveniles. *P. africana* and *Ekebergia capensis* showed Gaussian form with absence of intermediately sized individuals attributed to selective harvest. The density of woody species decreases with increasing DBH indicating predominance of small sized individuals in the forest. This implies that the forest is in good state of reproduction.

Therefore, to improve the natural diversity and structure of the forest, to minimize the influence of the surrounding communities and utilize the forest resources sustainably for present and future generation, it is necessary to take measures for protection of the forest human interference and other anthropogenic influences.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

The authors express their deepest thanks to the Agriculture and Rural Development Office of the Dega Damot District as well as to the respective chairpersons. We also thank Addis Ababa University for financial support.

REFERENCES

- Aleign A, Teketay D, Yemshaw Y, Edwards S. (2007). Diversity and status of regeneration of woody plants on the Peninsula of Zegie, Northwestern Ethiopia. *Tropical Ecology* 48(1):37-49.
- Aynekulu E (2011). Forest diversity in fragmented landscapes of northern Ethiopia and implications for conservation. PhD Dissertation. P 142.
- Bekele T (1993). Vegetation ecology of remnant Afromontane forests on the central plateau of Shewa, Ethiopia. Ph.D. Dissertation, Opulus press, Uppsala, pp.1-64.
- Bekele T (1994). Phytosociology and ecology of a humid Afromontane forest on the central plateau of Ethiopia. *Journal of Vegetation Science* 5(1):87-98.
- Berhanu A, Demissew S, Woldu Z, Didita M (2016). Woody species composition and structure of Kuandisha Afromontane forest fragment in Northwestern Ethiopia. *Journal Forestry Research* 28:343-355.
- Chen J, Song B, Rudnicki M, Moeur M, Bible K, North M, Shaw DC, Franklin JF, Braun DM (2004). Spatial relationship of biomass and species distribution in an old-growth Pseudotsuga - Tsuga Forest. *Forest Science* 50(3):364-375.
- Dega Damot Woreda Agricultural office (2017). Description of the study area.
- Friis I (1992). Forest and forest trees of North East tropical Africa. Kew Bulletin Additional Series 15:1-396.
- Friis I, Demissew S (2001) Vegetation Maps of Ethiopia and Eritrea In: Biodiversity Research in the Horn of Africa Region, pp. 399-439 (Friis, I. and Ryding, O., eds). Proceedings of the 3rd International symposium on the Flora of Ethiopia and Eritrea at the Carlsberg Academy, Copenhagen.
- Friis I, Demissew S, Van Bruegel P (2011). Atlas of the Potential Vegetation of Ethiopia. The Royal Danish Academy of Sciences and Letters, Copenhagen, Denmark, P 306.
- Hedberg I, Friis I, Persson E (2009b). Flora of Ethiopia and Eritrea. General part and index to volumes 1-7 (vol. 8). The National Herbarium, Addis Ababa, pp. 1-331.
- Kelbessa E, Demissew S, Woldu Z, Edwards S (1992). Some threatened Endemic plants of Ethiopia. In: (Edwards, S. and Asfaw Z eds.) The status of some plants in parts of tropical Africa NAPRECA, No.2 Botany 2000: East and Central Africa. pp. 35-55.
- Kent M, Coker P (1992). Vegetation description and analysis. A practical approach. Wiley, New York, pp. 1-363.
- Korner C (2000). Why are there global gradients in species richness? Mountains may hold the answer. *Trends in Ecology and Evolution* 15:513- 514.
- Lamprecht H (1989). Silviculture in Tropics. Tropical Forest Ecosystems and their Tree Species- Possibilities and Methods for their Long-term Utilization. TZVerlagsgesellschaftGmbH, Rossdort, Germany.
- Neelo J, Teketay D, Kashe K, Masamba W (2015). Stand Structure, Diversity and Regeneration Status of Woody Species in Open and Exclosed Dry Woodland Sites around Molapo Farming Areas of the Okavango Delta, Northeastern Botswana. *Open Journal of Forestry* 5:313-328.
- R Core Team (2017) R: a language and environment for statistical computing. R foundation for Statistical Computing, Vienna.
- Savadogo P, Tigabu M, Sawadogo L, Odén PC (2007) Woody species composition, structure and diversity of vegetation patches of a

- Sudanian savanna in Burkina Faso. *Bois et Forest Des Tropiques* 294(4):5-20.
- Shibru S, Balcha G (2004). Composition, structure and regeneration status of woody species in Dindin natural forest, Southeast Ethiopia: An implication for conservation. *Ethiopian Journal of Biological Sciences* 3(1):15-35.
- Tadele D, Lulekal E, Damtie D, Assefa A (2014). Floristic diversity and regeneration status of woody plants in Zengena forest, a remnant montane forest patch in northwestern Ethiopia. *Journal Forestry Research* 25(2):329-336.
- Teketay D (1992). Human impacts on a natural Montane forest in southeastern Ethiopia. *Mountain Research Development*, 12:393-400.
- Teketay D (2001). Deforestation, wood famine and environmental degradation in highland ecosystems of Ethiopia: urgent need for actions. *Northeast African Studies* 8:53-76.
- Van der Maarel E (1979). Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetation* 39(2):97-114.
- Whittaker RJ, Willis KJ, Field R (2003). Climatic–energetic explanations of diversity: a macroscopic perspective. In: *Macroecology: concepts and consequences*, Blackburn, T.M. and Gaston, K.J. (eds.), Cambridge University Press; Cambridge. pp. 107-129.
- Wassie A, Teketay D, Powell N (2005). Church forests in north Gonder administrative zone, northern Ethiopia. *Forests, Trees and Livelihoods* 15:349-373.
- Woldu Z (1999). Forests in the vegetation types of Ethiopia and their status in the geographical context. In: Edwards S, Demissie A, Bekele T, Haase G (eds) forest genetic resources conservation: principles, strategies and actions. Proceedings of the National Forest Genetic Resources Conservation Strategy Development Workshop. Institute of Biodiversity Conservation and Research, Addis Ababa, pp. 1-38.
- Yineger H, Kelbessa E, Bekele T, Lulekal E (2008). Floristic composition and structure of the dry Afromontane forest at Bale Mountains National Park, Ethiopia. *Ethiopian Journal of Science* 31(2):103-120.
- Zegeye H, Teketay D, Kelbessa E (2011). Diversity and regeneration status of woody species in Tara Gedam and Ababay forests, northwestern Ethiopia. *Journal Forestry Research* 22(3):315-328.

APPENDIX

Appendix Table 1. List of woody plants recorded at Amoro Forest.

S/N	Scientific name	Family	Vernacular name/Amharic	Habit
1	<i>Acalypha psilostachya</i> Hochst.	Euphorbiaceae	Nacha	Shrub
2	<i>Acanthus sennii</i> Chiov.	Acanthaceae	Kosheshila	Shrub
3	<i>Albizia himperiana</i> Oliv.	Fabaceae	Sesa	Tree
4	<i>Allophylus abyssinicus</i> (Hochst.) Radlk.	Sapindaceae	Abalat	Tree
5	<i>Apodytes dimidiata</i> E.Mey.ex Arn	Icacinaceae	Dong	Tree
6	<i>Arundinaria alpina</i> K. Schum.	Poaceae	Kerkeha	Shrub
7	<i>Asparagus africanus</i> Lam.	Asparagaceae	Yeset Kesit	Liana
8	<i>Bersama abyssinica</i> Fresen.	Melanthaceae	Azamira	Shrub
9	<i>Bridelia micrantha</i>	Euphorbiaceae	Yenebir Tifir	Shrub
10	<i>Brucea antidysenterica</i>	Simaroubaceae	Abalo	Shrub
11	<i>Buddleja polystachya</i> Fresen.	Loganiaceae	Nech Anifar	Shrub
12	<i>Buddleja davidii</i> Franch	Loganiaceae	Tikuir Anfar	Shrub
13	<i>Calpurnia aurea</i> (Ait.) Benth.	Fabaceae	Ligita	Shrub
14	<i>Carissa spinarum</i> L.	Apocynaceae	Agam	Shrub
15	<i>Clausena anisata</i> (Willd.) Benth.	Rutaceae	Limich	Shrub
16	<i>Clematis longicauda</i> Steud. ex A.Rich.	Ranunculaceae	Azo harg	Liana
17	<i>Clusia abyssinica</i> Jaub. & Spach	Euphorbiaceae	Fiyelefej	Shrub
18	<i>Croton macrostachyus</i> Del.	Euphorbiaceae	Bisana	Tree
19	<i>Discopodium penninervium</i> Hochst	Solanaceae	Aluma	Shrub
20	<i>Dodonaea angustifolia</i> L.f	Sapindaceae	Kitkita	Shrub
21	<i>Dombeya torrida</i> (J.F.Gmel) P. Bamps	Sterculiaceae	Wulkifa	Tree
22	<i>Dovyalis abyssinica</i> (A. Rich) Warb	Flacourtiaceae	Koshim	Shrub
23	<i>Ekebergia capensis</i> Spamn.	Meliaceae	Lol	Tree
24	<i>Embelia schimperi</i> Vatke	Myrsinaceae	Enkoko	Liana
25	<i>Erythrina brucei</i> Schweinf.	Fabaceae	Korch	Tree
26	<i>Euphorbia abyssinica</i> Gmel.	Euphorbiaceae	Kulkual	Tree
27	<i>Ficus sur</i> Forssk.	Moraceae	Shola	Tree
28	<i>Juniperus procera</i> Hochst. ex Endl.	Cupressaceae	Yeabesha Tsid	Tree
29	<i>Laggera tomentosa</i> Sch.Bip	Asteraceae	Gimane	Shrub
30	<i>Lippia adoensis</i> Hochst.ex Walp.	Verbenaceae	Kesy	Shrub
31	<i>Lobelia giberroa</i> Hemsl.	Lobeliaceae	Gibera	Shrub
32	<i>Maesa lanceolata</i> Forssk	Myrsinaceae	kilaba	Shrub
33	<i>Maytenus arbutifolia</i> (A.Rich.) Wilczek	Celastraceae	Atat	Shrub
34	<i>Maytenus obscura</i> (A.Rich.) Cuf.	Celastraceae	Qoba	Tree
35	<i>Myrica salicifolia</i> A.Rich	Myricaceae	Shihnet	Tree
36	<i>Myrsine africana</i> L.	Myrsinaceae	Kecho	Shrub
37	<i>Olea capensis</i> L. subsp. <i>macrocarpa</i> (C.H. Wright) Verdc.	Oleaceae	wegeda	Tree
38	<i>Olea europaea</i> L. subsp. <i>cuspidata</i> (Wall. ex G.Don) Cif	Oleaceae	Woyra	Tree
39	<i>Olinia rochetiana</i> A. Juss	Oliniaceae	Tife	Tree
40	<i>Osyris quadripartita</i> Decn.	Santalaceae	Qeret	Shrub
41	<i>Phytolacca dodecandra</i> L'Herit.	Phytolaccaceae	Endod	Liana
42	<i>Pittosporium viridiflorum</i> Sims	Pittosporaceae	weylwiha	Tree
43	<i>Prunus Africana</i> (Hook.f.) Kalkm	Rosaceae	Koma	Tree
44	<i>Rhus glutinosav</i> A.Rich. subsp. <i>glutinosa</i>	Anacardiaceae	Qamo	Tree
45	<i>Ricinus communis</i> L.	Euphorbiaceae	Chakima	Shrub
46	<i>Ritchiea albersii</i> Gilg.	Capparidaceae	Yetota Kolet	Shrub
47	<i>Rosa abyssinica</i> Lindley	Rosaceae	Qega	Shrub
48	<i>Rubus steudneri</i> Schwienf.	Rosaceae	Enjori	Liana

Appendix Table 1. Contd.

49	<i>Rumex nervosus</i> Vahl	Polygonaceae	Ambacho	Shrub
50	<i>Rytigynia neglecta</i> (Hiern) Robyns	Rubiaceae	Dingayseber	Shrub
51	<i>Schefflera abyssinica</i> (Hochst. ex A. Rich.) Harms	Araliaceae	Getem	Tree
52	<i>Solanecio gigas</i> (Vatke) C. Jeffrey	Asteraceae	Yeshikoko gomen	Shrub
53	<i>Solanum giganteum</i> Jacq.	Solanaceae	Tikuir Emby	Shrub
54	<i>Urera hypselodendron</i> (A. Rich) Wedd.	Urticaceae	Lankuso	Liana
55	<i>Vernonia amygdalina</i> Del	Asteraceae	Girawa	Shrub
56	<i>Vernonia myriantha</i> Hook.f.	Asteraceae	Dengorita	Shrub
57	<i>Zehneria Scabra</i> (Linn.f.) Sond	Cucurbitaceae	Nech Hareg	Liana

Appendix Table 2. Density, frequency, basal area and IVI of woody plants at Amoro Forest.

S/N	Species	Density (stem/h)	Frequency (%)	Basal area (m ² /ha)	Relative density (%)	Relative frequency (%)	Relative dominance (%)	IVI
1	<i>Acalypha psilostachya</i>	14.815	27.778	-	0.518	1.618	-	2.136
2	<i>Acanthus sennii</i>	13.580	11.111	-	0.475	0.647	-	1.122
3	<i>Albizia schimperiana</i>	6.790	13.889	0.018	0.237	0.809	0.100	1.146
4	<i>Allophylus abyssinicus</i>	425.926	88.889	0.992	14.890	5.178	5.367	25.435
5	<i>Apodytes dimidiata</i>	18.827	61.111	3.527	0.658	3.560	19.089	23.307
6	<i>Arundinaria alpina</i>	117.284	5.556	0.001	4.100	0.324	0.005	4.429
7	<i>Asparagus africanus</i>	0.617	2.778	-	0.022	0.162	-	0.183
8	<i>Bersama abyssinica</i>	104.630	69.444	0.327	3.658	4.045	1.771	9.474
9	<i>Bridelia micrantha</i>	29.938	36.111	0.091	1.047	2.104	0.494	3.644
10	<i>Brucea antidyssentrica</i>	73.765	75.000	0.020	2.579	4.369	0.109	7.057
11	<i>Buddleja polystachya</i>	10.494	13.889	0.025	0.367	0.809	0.136	1.312
12	<i>Buddleja davidii</i>	8.333	13.889	0.013	0.291	0.809	0.068	1.169
13	<i>Calpurnia aurea</i>	207.716	58.333	0.014	7.262	3.398	0.074	10.734
14	<i>Carissa spinarum</i>	128.704	11.111	0.002	4.499	0.647	0.009	5.155
15	<i>Clausena anisata</i>	152.160	44.444	-	5.319	2.589	-	7.908
16	<i>Clematis longicauda</i>	0.309	2.778	-	0.011	0.162	-	0.173
17	<i>Clutia abyssinica</i>	41.358	11.111	-	1.446	0.647	-	2.093
18	<i>Croton macrostachyus</i>	16.358	27.778	0.030	0.572	1.618	0.161	2.351
19	<i>Discopodium penninervium</i>	62.037	75.000	0.072	2.169	4.369	0.387	6.925
20	<i>Dodonaea angustifolia</i>	0.926	2.778	0.002	0.032	0.162	0.008	0.202
21	<i>Dombeya torrida</i>	10.802	38.889	0.032	0.378	2.265	0.171	2.814
22	<i>Dovyalis abyssinica</i>	54.630	44.444	0.064	1.910	2.589	0.348	4.846
23	<i>Ekebergia capensis</i>	7.099	33.333	0.339	0.248	1.942	1.837	4.027
24	<i>Embelia schimperi</i>	22.840	25.000	-	0.798	1.456	-	2.255
25	<i>Erythrina brucei</i>	0.617	5.556	0.005	0.022	0.324	0.028	0.373
26	<i>Euphorbia abyssinica</i>	124.074	72.222	0.385	4.338	4.207	2.085	10.630
27	<i>Ficus sur</i>	3.704	13.889	0.162	0.129	0.809	0.876	1.814
28	<i>Juniperus procera</i>	0.309	2.778	0.001	0.011	0.162	0.006	0.179
29	<i>Laggera tomentosa</i>	66.667	25.000	-	2.331	1.456	-	3.787
30	<i>Lippia adoensis</i>	0.926	2.778	-	0.032	0.162	-	0.194
31	<i>Lobelia giberroa</i>	71.914	38.889	0.015	2.514	2.265	0.079	4.858
32	<i>Maesa lanceolata</i>	6.790	13.889	0.015	0.237	0.809	0.081	1.128
33	<i>Maytenus arbutifolia</i>	85.494	41.667	0.013	2.989	2.427	0.069	5.485
34	<i>Maytenus obscura</i>	3.395	8.333	0.079	0.119	0.485	0.427	1.032

Appendix Table 2. Contd.

35	<i>Myrsine africana</i>	23.457	5.556	-	0.820	0.324	-	1.144
36	<i>Olea capensis</i>	1.543	11.111	0.548	0.054	0.647	2.968	3.669
37	<i>Olea europaea</i>	0.926	5.556	0.044	0.032	0.324	0.240	0.596
38	<i>Olinia rochetiana</i>	12.963	25.000	0.029	0.453	1.456	0.156	2.065
39	<i>Osyris quadripartita</i>	16.358	25.000	0.006	0.572	1.456	0.031	2.060
40	<i>Phytolacca dodecandra</i>	25.000	27.778	-	0.874	1.618	-	2.492
41	<i>Pittosporum viridiflorum</i>	0.617	5.556	0.013	0.022	0.324	0.072	0.417
42	<i>Prunus africana</i>	55.247	83.333	4.429	1.931	4.854	23.971	30.756
43	<i>Rhus glutinosa</i>	28.395	22.222	0.011	0.993	1.294	0.061	2.348
44	<i>Ricinus communis</i>	1.235	5.556	0.008	0.043	0.324	0.043	0.409
45	<i>Ritchiea albersii</i>	32.099	50.000	0.050	1.122	2.913	0.273	4.308
46	<i>Rosa abyssinica</i>	9.568	19.444	0.005	0.334	1.133	0.027	1.494
47	<i>Rubus steudneri</i>	40.432	36.111	-	1.413	2.104	-	3.517
48	<i>Rumex nervosus</i>	8.951	16.667	-	0.313	0.971	-	1.284
49	<i>Rytigynia neglecta</i>	33.025	52.778	0.023	1.155	3.074	0.123	4.352
50	<i>Schefflera abyssinica</i>	15.123	52.778	6.917	0.529	3.074	37.438	41.041
51	<i>Solanecio gigas</i>	242.284	63.889	0.025	8.470	3.722	0.133	12.324
52	<i>Solanum giganteum</i>	0.309	2.778	-	0.011	0.162	0.001	0.174
53	<i>Urera hypselodendron</i>	70.062	66.667	-	2.449	3.883	-	6.333
54	<i>Vernonia amygdalina</i>	9.259	19.444	0.040	0.324	1.133	0.215	1.672
55	<i>Vernonia myriantha</i>	337.963	94.444	0.086	11.815	5.502	0.464	17.781
56	<i>Zehneria Scabra</i>	1.543	2.778	-	0.054	0.162	-	0.216
Total		2860.185	1713.889	18.477	100	100	100	299.827