PhytoKeys

PhytoKeys 255: 189–201 (2025)
DOI: 10.3897/phytokeys.255.147252

Research Article

Yushania rubrovaginata (Poaceae, Bambusoideae), a new combination for *Sasa rubrovaginata*

Xing Li[1,2,3], Jing-Bo Ni[1,2], Mian Liang[4], Yi-Hua Tong[1,2], Nian-He Xia[1,2]

1 *State Key Laboratory of Plant Diversity and Specialty Crops/Key Laboratory of Digital Botanical Garden of Guangdong Province, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China*
2 *South China National Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China*
3 *Key Laboratory of State Forestry Administration on Tropical Forestry Research, Research Institute of Tropical Forestry, Chinese Academy of Forestry, Guangzhou 510520, China*
4 *The Administration Center of Guangxi Cenwanglaoshan National Nature Reserve, Baise 530000, China*

Corresponding author: Yi-Hua Tong (yh-tong@scbg.ac.cn)

Abstract

Sasa rubrovaginata is transferred to *Yushania* based on morphological and molecular evidence. The lectotype for *S. rubrovaginata* is designated. Besides, a revised description of this species and a key to the four *Yushania* species distributed in Cenwanlaoshan Mountain are also provided.

Key words: Bamboo, Cenwanlaoshan Mountain, *Sasa*, taxonomy, *Yushania*

OPEN ACCESS

Academic editor: Wei Lim Goh
Received: 20 January 2025
Accepted: 16 March 2025
Published: 24 April 2025

Citation: Li X, Ni J-B, Liang M, Tong Y-H, Xia N-H (2025) *Yushania rubrovaginata* (Poaceae, Bambusoideae), a new combination for *Sasa rubrovaginata*. PhytoKeys 255: 189–201. https://doi.org/10.3897/phytokeys.255.147252

Copyright: © Xing Li et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Sasa Makino & Shibata (1901) is a medium genus of Arundinarieae, Bambusoideae (Zhang et al. 2020a), and characterized by having a shrubby habit, leptomorph rhizomes, single branch per culm node, spikelets arranged into a panicle-like inflorescence, and six stamens and three stigmas per floret (Hu 1996; Wang and Stapleton 2006; Clayton et al. 2016; Yi et al. 2008; Shi et al. 2022). About 43 species are recognized in *Sasa* genus at present, which are mainly distributed in East Asia (Vorontsova et al. 2016; Soreng et al. 2022; Li et al. 2023a; POWO 2025). Previous phylogenetic studies indicated that *Sasa* is polyphyletic (Peng et al. 2008; Sungkaew et al. 2009; Triplett and Clark 2010; Zeng et al. 2010; Zhang et al. 2012; Qin et al. 2021), and most Chinese *Sasa* species were successively revised and transferred to *Sinosasa* L. C. Chia ex N. H. Xia et al., *Yushania* Keng f., or *Pseudosasa* Makino ex Nakai in recent studies (Qin et al. 2021; Li et al. 2022, 2023a, 2023b, 2023c). For now, there are only three accepted names under *Sasa* from China (POWO 2025), viz., *S. subglabra* McClure (1940), *S. hainanensis* C. D. Chu & C. S. Chao (Chao et al. 1980) and *S. rubrovaginata* C. H. Hu (1985).

Sasa rubrovaginata was described based on the only collection, *Nanzhidi* (Phytogeography expeditions of South China Institute of Botany) 5102, from Cenwanglaoshan Mountain in Guangxi, China. Only two duplicates of the type were found in N (Fig. 1). Both duplicates (N019023159 and N019023168) only

constitute a leafy branch. No other specimens can be located in the Chinese herbaria. Without any information of rhizome and branch complement, which are very important for the placement of generic position, it was designated as a member of *Sasa*. Thus, in order to ascertain the taxonomic position of *S. rubrovaginata*, it is necessary to recollect complete specimens with rhizome and branch complement from the type locality.

Materials and methods

The complete specimens of *Sasa rubrovaginata* were collected from the type locality, i.e., Cenwanlaoshan Mountain, Langping Town, Tianlin County, Guangxi Zhuang Autonomous Region, China, during a field trip in September 2022. Fresh leaf samples were stored in sealed bags with silica gel for DNA extraction. Magnifier and ruler (0.5 mm scale) were used for observations and measurements. Some minor characters were observed with a stereo microscope (Mshot-MZ101, Guangzhou Micro-shot Technology Co., Ltd, Guangzhou, China). Photos of type specimens of *S. rubrovaginata* were downloaded from the Chinese Virtual Herbarium (https://www.cvh.ac.cn/) for comparison. The description was conducted based on both living and dried materials, and the descriptive terms follow McClure (1966) and Beentje (2016). Herbarium acronyms follow Thiers (2025).

To study the phylogenetic position of *S. rubrovaginata* within the tribe Arundinarieae, phylogenetic analyses were carried out by using the complete chloroplast genome data. A total of 30 representatives from all the five subtribes of the tribe Arundinarieae (Zhang et al. 2020a) were sampled with *Dendrocalamus strictus* (Roxburgh) Nees from the tribe Bambuseae as an outgroup. All the sampled taxa, information of voucher specimens and GenBank accession numbers were listed in Table 1.

DNA extraction, sequencing, assembly and annotation

Total genomic DNA was extracted from dried leaves using the modified CTAB method (Li et al. 2013) and sent to Novo Gene Company (Beijing, China) for DNA assessment. The genomic DNA was then cut up to 350 bp-sized fragments for the construction of the library, and paired-end sequencing was performed on the Illumina Hiseq 4000 platform. A total of 20 G clean data (150 bp read length) were generated from each sample. These clean data were utilized to assemble the plastome by GetOrganelle v.1.7.7 pipeline (Jin et al. 2020) using the plastome of *Phyllostachys edulis* (Carriere) J. Houzeau (accession number: HQ337796) as the reference, with k-mer values of 45, 65, 85, 105, 125, word size of 102, and extension rounds of 20. Bandage software (Wick et al. 2015) was used to visually check if the final result of the assembled genome was circular or not. Finally, the assembled plastome sequence with the same direction as the reference sequence was kept and manually operated in Geneious v. 9.1.4 (Kearse et al. 2012) with the structure of LSC-IRa-SSC-IRb.

Phylogenetic analysis

By using MAFFT v. 7.490, all of the complete chloroplast genomes were concatenated into a data matrix after being aligned. Maximum Likelihood (ML) and

Table 1. A list of the vouchers and GenBank accession numbers for the sampled species in this study.

Taxon	Voucher information	Accession number
Ingroup		
Acidosasa glauca B.M.Yang	CZY56 (IBSC)	OP850353
Ampelocalamus actinotrichus (Merr. & Chun) S.L.Chen, T.H.Wen & G.Y.Sheng	MPF10003 (KUN)	MF066245
Chimonobambusa tumidissinoda Ohrnb.	MPF10083 (KUN)	MF066244
Fargesia albocerea J.R.Xue & T.P.Yi	ZhangYu-QuD588 (SANU)	NC_043891
Fargesia communis T.P.Yi	ZhangYu-QuD540 (SANU)	NC_043934
Fargesia daminiu T.P.Yi & J.Y.Shi	ZhangYu-QuG660 (SANU)	NC_043942
Fargesia hygrophila J.R.Xue & T.P.Yi	ZhangYu-QuD552 (SANU)	NC_043938
Fargesia setosa T.P.Yi	ZhangYu-QuG695 (SANU)	NC_043939
Gaoligongshania megalothyrsa (Hand.-Mazz.) D.Z.Li, Hsueh & N.H.Xia	MPF10056 (KUN)	JX513419
Gelidocalamus stellatus T.H.Wen	BH102 (IBSC)	OP850347
Hsuehochloa calcareus (C.D.Chu & C.S.Chao) D.Z.Li & Y.X.Zhang	MPF10050 (KUN)	KJ496369
Indocalamus longiauritus Hand.-Mazz.	MPF10168 (KUN)	HQ337795
Indocalamus sinicus (Hance) Nakai	ZMY037 (KUN)	MF066250
Indosasa crassiflora McClure	BH58 (IBSC)	OK558536
Oligostachyum sulcatum Z.P.Wang & G.H.Ye	Not provided by the author	MW190089
Phyllostachys edulis (Carriere) J.Houzeau	MPF10163 (KUN)	HQ337796
Pleioblastus maculatus (McClure) C.D.Chu & C.S.Chao	CZY56 (IBSC)	JX513424
Pseudosasa cantorii (Munro) Keng f.	MPF10006 (KUN)	MF066255
Pseudosasa japonica (Siebold & Zucc. ex Steud.) Makino ex Nakai	Pjc-1 (ZJFC)	KT428377
Ravenochloa wilsonii (Rendle) D.Z.Li & Y.X.Zhang	MPF10146 (KUN)	JX513421
Sasa rubrovaginata C.H.Hu	LX178 (IBSC)	PQ010623
Sasa veitchii (Carriere) Rehder	LC1325 (ISC)	KU569975
Shibataea chiangshanensis T.H.Wen	ZLN-2011080 (KUN)	MF066257
Sinosasa fanjingshanensis N.H.Xia, Q.M.Qin & J.B.Ni	BH124 (IBSC)	OP850348
Sinosasa longiligulata (McClure) N.H.Xia, Q.M.Qin & J.B.Ni	CZY163 (IBSC)	OP850351
Sinobambusa tootsik (Makino) Makino ex Nakai	NH031 (IBSC)	OP850357
Yushania confusa (McClure) Z.P.Wang & G.H.Ye	ZhangYu-QuF642 (SANU)	NC_043893
Yushania maculata T.P.Yi	Not provided by the author	OR750784
Yushania niitakayamensis (Hayata) Keng f.	Not provided by the author	MN310560
Yushania shuichengensis T.P.Yi & L.Yang	Not provided by the author	OR750781
Outgroup		
Dendrocalamus strictus (Roxburgh) Nees	zmy018 (KUN)	MK679802

Bayesian Inference (BI) tools in the PhyloSuite v.1.2.3 platform (Xiang et al. 2023; Zhang et al. 2020b) were utilized for phylogenetic reconstructions. The Bayesian Information Criterion (BIC) in ModelFinder (Kalyaanamoorthy et al. 2017) was used to identify the optimal substitution model for ML and BI methods. Maximum likelihood phylogenies were inferred by using IQ-TREE v 2.2.0 (Nguyen et al. 2015) under the best-fit K81u+R4+F model. We then performed 1000 ultrafast bootstrap replicates and 1000 approximate likelihood ratio (SH-aLRT) tests to assess branch supports (Guindon et al. 2010). Bayesian Inference phylogenies were inferred using MrBayes v 3.2.7 (Ronquist et al. 2012) under the GTR+I+G+F model (2 parallel runs, 40,000,000 generations), in which the initial 25% of sampled data were discarded as burn-in. ML and BI trees were visualized in FigTree v. 1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Results

The real identity of *Sasa rubrovaginata*

According to the protologue and field note of the type specimens of *Sasa rubrovaginata*, the gathering was collected at an elevation of 2000 m at Cenwangling (岑王岭) (the main peak of Cenwanlaoshan), Langping (浪平), Tianlin County (田林县). During our field trip to Cenwanlaoshan Mountain, only one bamboo species that has a long-necked pachymorph rhizome and two or three branches on the mid or upper culm nodes were found at the elevation from 1968 m to 2062.5 m (the peak's elevation). The young and unbranched culms with foliage leaves at the apex (Fig. 2) fully matched with the protologue and the type specimens (Fig. 1) of *S. rubrovaginata* as well as the glabrous internodes, white powdery infranodal region, purple-red culm leaf sheath with a hispid base and ciliate margins, caducous culm leaf blades, truncate culm leaf ligules, 5–6 foliage leaves per ultimate branch with the glabrous sheath being ciliate on margins, falcate foliage leaf auricles with developed oral setae, truncate foliage leaf inner ligules, and glabrous foliage leaf blades with conspicuous transverse veins and 5–6 pairs secondary veins. Thus, we are very sure that the specimens we collected are *S. rubrovaginata*. Hu (1985) should take the young and unbranched culm with foliage leaves at the apex as a leafy branch, and supposed that the bamboo possessed solitary branch complement and leptomorph rhizome, just like the case of *S. tomentosa* C. D. Chu & C. S. Chao (≡ *Yushania tomentosa* (C. D. Chao & C. S. Chao) N. H. Xia et al.; see Li et al. 2023a for nomenclatural revision). Owing to the long-necked rhizome (Fig. 3G) and branch complement with mostly solitary branch at lower culm nodes and two to three (Fig. 3C) branches at mid and upper culm nodes, it should be a member of *Yushania* rather than *Sasa*.

After examining the type specimens and referring to the related literature (Yi et al. 2008; Shi et al. 2022; Li et al. 2023a), *S. rubrovaginata* is mostly similar to *Y. tomentosa* by sharing the branch complement with usually solitary branch at lower culm nodes and two to three branches at mid and upper culm nodes (Fig. 3C), the glabrous internodes with purple spots, the culm leaf with sheath being ciliate on margins and leaving persistent remains on internode when falling off, truncate ligule, falcate auricles with radiate oral setae (Fig. 3F), and foliage leaf with sheath being ciliate on margins, falcate auricles with radiate oral setae (Fig. 3D), truncate ligule and glabrous blades, but differs in having purple-red (vs. green to brown) and densely brown hispid (Fig. 3E) (vs. white to yellowish-brown hirsute) culm leaf sheath, and glabrous culm leaf ligules margin (vs. white ciliolate), foliage leaf sheath (vs. densely white hirsute), outer ligule margin (vs. white ciliate) and pseudopetiole (vs. white puberulous). A more detailed comparison of the two species is presented in Table 2.

Phylogenetic analysis

The chloroplast genomes of the sampled species vary from 139,404 bp (*Dendrocalamus strictus*) to 140,064 bp (*Gaoligongshania megalothyrsa* (Hand.-Mazz.) D. Z. Li, Hsueh & N. H. Xia) with an alignment of 144,047 bp. Sequence divergence was observed in this data matrix with 3,903 variable sites (2.71%) comprising 3,087 singleton variable sites (2.14%) and 816 parsimony informative

Table 2. Comparison of *Sasa rubrovaginata* and *Yushania tomentosa*.

Characters	*Sasa rubrovaginata*	*Yushania tomentosa*
Culm leaf		
Sheath	Purple-red when fresh, densely brown hispid	Green to brown when fresh, densely white to yellowish-brown hirsute
Ligule	Glabrous	White ciliolate on the margin
Foliage leaf		
Sheath	Glabrous, initially white powdery	Densely white hirsute
Pseudopetiole	Glabrous	White puberulous
Outer ligule	Glabrous	White ciliate on the margin

sites (0.57%). Only the ML tree was displayed (Fig. 4) with nodal support values from both ML and BI methods labeled on each node. As shown in the phylogenetic tree, *S. rubrovaginata* is distantly related to *S. veitchii* (Carriere) Rehder (= *S. albomarginata* (Miq.) Makino & Shibata, the type of *Sasa*) but forms a monophyletic clade with four *Yushania* species with strong nodal support (BS = 99.6% & PP = 1.00), which also supports that *S. rubrovaginata* should be a member of *Yushania*, rather than *Sasa*.

Discussion

Based on the morphological and molecular evidence, it is concluded that *Sasa rubrovaginata* represents a distinct species of *Yushania*. Accordingly, a new combination of *S. rubrovaginata* under *Yushania* should be made. Hu (1985) designated the collection *Nanzhidi 5102* (N) as the type of *S. rubrovaginata*, but there are two duplicates in N, which should be recognized as syntypes. We designated N019023168 (Fig. 1A) which bears some pencil annotations in Chinese and "Typus" in the identification slip by the author and has relatively better developed culm and foliage leaves as the lectotype of *S. rubrovaginata*.

Taxonomic treatment

Yushania rubrovaginata **(C. H. Hu) N. H. Xia, Y. H. Tong, J. B. Ni & X. Li, comb. nov.**
urn:lsid:ipni.org:names:77360579-1
Figs 1–3

Basionym. *Sasa rubrovaginata* C. H. Hu, Bamboo Research 2(2): 59 (1985)

Type. China. • Guangxi: Tianlin County, Langping Town, Cenwanlaoshan Mountain, elev. 2000 m, 26 Nov. 1957, *Nanzhidi 5102* (lectotype N019023168!, Fig. 1A, designated here; isolectotype N019023159!, Fig. 1B)

Description. Shrubby bamboo. Rhizomes pachymorph, necks 15–30 cm long, 3–5 mm in diameter, solid. Culms 1–3.5 m tall, 5–8 mm in diameter, diffuse; branches usually solitary at lower culm nodes, 2–3 at mid and upper culm nodes; internodes terete, 10–30 cm long, glabrous, densely purple-spotted, thickly white powdery below nodes, hollow; supranodal ridges raised. Culm buds solitary, long-ovate, yellow to light green, ciliate on the margin. Culm leaf sheaths persistent or tardily deciduous, ca. 1/2 as long as internodes, purple-red, densely brown hispid




Figure 1. Lectotype (**A**) and isolectotype (**B**) of *Sasa rubrovaginata* C. H. Hu (*Nanzhi 5102*, N **A** N019023168 **B** N019023159). Photos downloaded from Chinese Virtual Herbarium (https://www.cvh.ac.cn/).

abaxially, densely ciliate on the margin; sheath scar prominent, with persistent remains of sheath base; auricles falcate, 3–5 × 1–2 mm; oral setae developed, radiate; ligule truncate, ca. 0.5 mm high; blades linear-lanceolate to lanceolate, reflexed, easily deciduous, margin serrulate. Foliage leaves 5–13 per ultimate branch; sheath glabrous, margin ciliate; auricles falcate, 1–3 × 0.5–1 mm; oral setae radiate, ca. 1 cm long; inner ligule truncate, ca. 1 mm high; outer ligule and pseudopetioles glabrous; blades broad-lanceolate to lanceolate, 17–26 × 3.5–6 cm, wavy when dry, glabrous, apex acuminate, base cuneate to obtuse; secondary veins 9–10 pairs, transverse veins conspicuous. Inflorescence unknown.

Distribution and habitat. It is only found in Cenwang Mountain, Tianlin County, Guangxi, China. It grows on top of mountains at an altitude of 1968 to 2062.5 meters (the peak's elevation).

Phenology. New shoots from August to September.

Chinese name. 红壳玉山竹 (Chinese pronunciation: hóng ké yù shān zhú).

Notes. Hu (1985) described that the culm leaf auricles and oral setae of *Yushania rubrovaginata* are undeveloped (see Fig. 1). In fact, this bamboo does have developed culm leaf auricles and oral setae (Fig. 3E, F). But the culm leaf auricles and oral setae are easy to fall off, which exactly happens to the type specimens of *Y. rubrovaginata*. Similarly, the culm leaf sheaths of the type specimens are glabrous on the mid and upper parts, and only the basal part is sparsely hispid. Actually, *Y. rubrovaginata* has a thoroughly hispid culm leaf sheath, and pits are left after the trichomes fall off (Fig. 3E).



Figure 2. A specimen of *Sasa rubrovaginata* C. H. Hu collected from the type locality, *X.Li & J.B.Ni LX178* (IBSC). Photo by Xing Li.

Wang and Stapleton (2006) treated *Sasa duplicata* W. T. Lin & Z. J. Feng (1992) as a synonym of *Y. rubrovaginata*. However, the former has a leptomorph rhizome, rather than pachymorph rhizome. Actually, *S. duplicata* owns some characters that are the same as *Pseudosasa cantorii* (Munro) P. C. Keng ex S. L. Chen et al. (Zhu et al. 2006), such as glabrous internodes, the white powdery infranodal region, branch complement with one to three branches at each culm node, culm leaf sheath with ciliate margin, arcuate to truncate culm leaf ligules and falcate auricles with developed oral setae, foliage leaf sheath with ciliate margin, truncate ligules, developed oral setae and lanceolate to oblong-lanceolate blades with conspicuous transverse veins. Thus, *S. duplicata* is probably a synonym of *P. cantorii*, but more work needs to be done to ascertain this.

There are another three *Yushania* species, viz., *Y. cartilaginea* T. H. Wen (1984), *Y. chingii* T. P. Yi and *Y. rugosa* T. P. Yi (1986), distributed in Cenwanglaoshan Mountain. *Yushania rubrovaginata* can be distinguishable from them by having branch




Figure 3. *Yushania rubrovaginata* **A** foliage leafy branch **B** culm bud **C** three branches at an upper culm node **D** partial foliage leafy branch, showing auricles and oral setae **E** culm leaf and white powdery infranodal region **F** apex of culm leaf sheath, showing ligules, auricles and oral setae **G** pachymorph rhizome with long neck. All photos by Xing Li.




Figure 4. Phylogenetic tree derived from MI and BI methods based on the complete chloroplast sequences, showing the phylogenetic position of *Sasa rubrovaginata*. Bootstrap values and posterior probabilities are indicated at each node, and the type of *Sasa* is highlighted in blue.

complement with the solitary branch at lower culm nodes and two to three branches at mid and upper culm nodes, densely purple-spotted internodes, purple-red and densely brown hispid culm leaf sheath, and foliage leaf with developed and radiate oral setae. A key to these *Yushania* species is provided as follows.

Key to *Yushania* species distributed in Cenwanglaoshan Mountain

1a Branches solitary at each culm node.....**2**
2a Culm leaf sheath glabrous; auricles developed and falcate; oral setae radiate with trichomes 5–8 mm long.....***Y. chingii***
2b Culm leaf sheath glabrous or sparsely purple-brown setose at base; auricles absent or small; oral setae absent or weak with trichomes 2–3 mm long.....***Y. rugosa***
1b Branches 1–3 at each culm node.....**3**
3a Internodes densely purple-spotted; culm leaf sheath purple-red, densely brown hispid; foliage leaf sheath with radiate oral setae.....***Y. rubrovaginata***
3b Internodes green, without purple spots; culm leaf sheath green to brown, glabrous; foliage leaf sheath with straight oral setae.....***Y. cartilaginea***

Additional specimens examined

Yushania rubrovaginata: China. • Guangxi: Tianlin County, Langping Town, Cenwanglaoshan Mountain, Cenwangling, 25 September 2022, 24°29'22.4"N, 106°24'5.3"E, elev. 2062 m, *X. Li & J. B. Ni LX178* (IBSC).

Yushania tomentosa: China. • Guangxi: Rongshui County, Jiuwan Mountain, elev. 1400 m, 25 August 1958, *S. H. Chun 15320* (isotypes: NAS00070361, image; WUK0211330, image; N019023167, image; IFP15899999w0005, image); • Rongshui County, Wangdong Township, Jiuwan Mountain, Weilinjiang, 23 September 2022, 25°18'39.3"N, 108°38'13.2"E, elev. 1358 m, *X. Li & J. B. Ni LX168* (IBSC).

Yushania cartilaginea: China. • Guangxi: Baise City, [Tianlin County], Kashan [Laoshan = Cenwanlaoshan] forestry station, elev. 1700 m, 9 April 1982, *W. W. Chou L82433* (holotype: ZJFI).

Yushania chingii: China. • Guangxi: Tianlin County, Laoshan forestry station, elev. 1400 m, 14 January 1990, *J. P. Ruan 90005* (N 019025132, image; N 019025139, image; N 019025140, image).

Yushania rugosa: China. • Guizhou: Wangmo County, Maoping community, elev. 1500–1556 m, 26 August 1981, *T. P. Yi 81118* (holotype: SIFS).

Acknowledgments

We thank Mr. Fu-Hong Long from Langping protection station, Cenwanglaoshan Mountain National Nature Reserve for his assistance in the fieldwork.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was funded by Chinese Academy of Sciences (Biotaxonomic Scientist Post Grant No. CAS-TAX-24-049) and the National Natural Science Foundation of China (Grant No. 32270227).

Author contributions

Funding acquisition: NHX, YHT. Investigation: XL, ML, JBN. Methodology: XL. Supervision: YHT, NHX. Writing - original draft: XL. Writing - review and editing: NHX, YHT.

Author ORCIDs

Xing Li https://orcid.org/0000-0001-7146-0125
Yi-Hua Tong https://orcid.org/0000-0002-5034-005X
Nian-He Xia https://orcid.org/0000-0001-9852-7393

Data availability

All of the data that support the findings of this study are available in the main text.

References

Beentje H (2016) The Kew Plant Glossary, Second Edition. Royal Botanic Gardens, Kew, London, 1–184.

Chao CS, Chu CD, Hsiung WY (1980) A revision of some genera and species of Chinese bamboos. Acta Phytotaxonomica Sinica 18(1): 20–36.

Clayton WD, Vorontsova M, Harman KT, Williamson H (2016) GrassBase-The online world grass flora. https://www.kew.org/data/grassbase/index.html [Accessed 18 January 2025]

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. Systematic Biology 59(3): 307–321. https://doi.org/10.1093/sysbio/syq010

Hu CH (1985) A revision of the genus *Sasa* from China. Bamboo Research 2(2): 56–62.

Hu CH (1996) *Sasa* Makino et Shibata. In: Keng PC, Wang ZP (Eds) Flora Reipublicae Popularis Sinicae (Vol. 9). Science Press, Beijing, 662–675.

Jin JJ, Yu WB, Yang JB, Song Y, dePamphilis CW, Yi TS, Li DZ (2020) GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of organelle genomes. Genome Biology 21(1): 241. https://doi.org/10.1186/s13059-020-02154-5

Kalyaanamoorthy S, Minh BQ, Wong T, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14(6): 587–589. https://doi.org/10.1038/nmeth.4285

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics (Oxford, England) 28(12): 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Li JL, Wang S, Yi J, Wang L, Zhou SL (2013) A modified CTAB protocol for plant DNA extraction. Chinese Bulletin of Botany 48(1): 72–78. https://doi.org/10.3724/SP.J.1259.2013.00072

Li X, Tong YH, Xia NH (2022) *Sasa guangdongensis*, a new synonym of *Acidosasa carinata* (Poaceae: Bambusoideae). Phytotaxa 568(2): 204–212. https://doi.org/10.11646/phytotaxa.568.2.5

Li X, Ni JB, Tan F, Tong YH, Xia NH (2023a) *Yushania tomentosa* (Poaceae, Bambusoideae), a new combination from Guangxi. PhytoKeys 218: 11–18. https://doi.org/10.3897/phytokeys.218.97312

Li X, Ni JB, Xia NH (2023b) The identity of *Sasa oblongula* C. H. Hu (Poaceae, Bambusoideae, Arundinarieae): Evidence from morphology and molecular data. PhytoKeys 226: 17–32. https://doi.org/10.3897/phytokeys.226.101221

Li X, Ni JB, Cai ZY, Tong YH, Xia NH (2023c) *Sinosasa gracilis* (Poaceae: Bambusoideae), a new combination supported by morphological and phylogenetic evidence. PhytoKeys 226: 53–63. https://doi.org/10.3897/phytokeys.226.101164

Lin WT, Feng ZJ (1992) Six new bamboo species from Guangdong. Acta Phytotaxonomica Sinica 30(6): 561–562.

Makino T, Shibata K (1901) On *Sasa*, a new genus of Bambuseae, and its affinities. Botanical Magazine Tokyo 15(168): 18–31. https://doi.org/10.15281/jplantres1887.15.168_18

McClure FA (1940) New genera and species of Bambusaceae from eastern Asia. Lingnan University Science Bulletin 9: 1–67.

McClure FA (1966) The bamboos: A fresh perspective. Harvard University Press, Cambridge, 1–347. https://doi.org/10.4159/harvard.9780674428713

Nguyen LT, Schmidt HA, Haeseler V, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating Maximum-Likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. https://doi.org/10.1093/molbev/msu300

Peng S, Yang HQ, Li DZ (2008) Highly heterogeneous generic delimitation within the temperate bamboo clade (Poaceae: Bambusoideae): evidence from GBSSI and ITS sequences. Taxon 57(3): 799–810. https://doi.org/10.1002/tax.573011

POWO (2025) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:18942-1#children [accessed 9 Jan 2025]

Qin QM, Tong YH, Zheng XR, Ni JB, Xia NH (2021) *Sinosasa* (Poaceae: Bambusoideae), a new genus from China. Taxon 70(1): 27–47. https://doi.org/10.1002/tax.12422

Ronquist F, Teslenko M, Mark VDP, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. https://doi.org/10.1093/sysbio/sys029

Shi JY, Che BY, Zhang YX, Zhou DQ, Ma LS, Yao J, Zhang XZ (2022) *Yushania* Keng f. In: Shi JY, Zhang YX, Zhou DQ, Ma LS, Yao J (Eds) Illustrated Flora of Bambusoideae in China (Vol. 2). Science Press, Beijing & Springer Nature Singapore Pte Ltd., Singapore, 409–481. https://doi.org/10.1007/978-981-16-2758-3_43

Soreng RJ, Peterson PM, Zuloaga FO, Romaschenko K, Clark LG, Teisher JK, Gillespie LJ, Barberá P, Welker CAD, Kellogg EA, Li DZ, Davidse G (2022) A worldwide phylogenetic classification of the Poaceae (Gramineae) III: An update. Journal of Systematics and Evolution 60(3): 476–521. https://doi.org/10.1111/jse.12847

Sungkaew S, Stapleton C, Salamin N, Hodkinson TR (2009) Non-monophyly of the woody bamboos (Bambuseae; Poaceae): A multi-gene region phylogenetic analysis of Bambusoideae s.s. Journal of Plant Research 122(1): 95–108. https://doi.org/10.1007/s10265-008-0192-6

Thiers B (2025) [continuously updated] Index Herbariorum: A global directory of public herbaria and associated staff. http://sweetgum.nybg.org/ih [Accessed 8 January 2024]

Triplett JK, Clark LG (2010) Phylogeny of the temperate bamboos (Poaceae: Bambusoideae: Bambuseae) with an emphasis on Arundinaria and allies. Systematic Botany 35(1): 102–120. https://doi.org/10.1600/036364410790862678

Vorontsova MS, Clark LG, Dransfield J, Govaerts R, Baker WJ (2016) World checklist of bamboos and rattans. INBAR Technical Report No. 37. International Network of Bamboo & Rattan, Beijing, 1–454.

Wang ZP, Stapleton CMA (2006) *Sasa* Makino & Shibata. In: Wu ZY, Raven P (Eds) Flora of China (Vol. 22). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 109–112.

Wen TH (1984) New taxa of Bambusoideae in China (I). Journal of Bamboo Research 3(2): 28–29.

Wick RR, Schultz MB, Justin Z, Holt KE (2015) Bandage: Interactive visualization of de novo genome assemblies. Bioinformatics (Oxford, England) 31(20): 3350–3352. https://doi.org/10.1093/bioinformatics/btv383

Xiang CY, Gao FL, Jakovlić I, Lei HP, Hu Y, Zhang H, Zou H, Wang GT, Zhang D (2023) Using phyloSuite for molecular phylogeny and tree-based analyses. iMeta 2(1): e87. https://doi.org/10.1002/imt2.87

Yi TP (1986) Studies on the genus *Yushania*. Journal of Bamboo research 5(1): 45–64.

Yi TP, Shi JY, Ma LS, Wang HT, Yang L (2008) Iconographia Bambusoidearum Sinicarum. Science Press, Beijing, 1–766.

Zeng CX, Zhang YX, Triplettd JK, Yang JB, Li DZ (2010) Large multi-locus plastid phylogeny of the tribe Arundinarieae (Poaceae: Bambusoideae) reveals ten major lineages and low rate of molecular divergence. Molecular Phylogenetics and Evolution 56(2): 821–839. https://doi.org/10.1016/j.ympev.2010.03.041

Zhang YX, Zeng CX, Li DZ (2012) Complex evolution in Arundinarieae (Poaceae: Bambusoideae): Incongruence between plastid and nuclear GBSSI gene phylogenies. Molecular Phylogenetics and Evolution 63(3): 777–797. https://doi.org/10.1016/j.ympev.2012.02.023

Zhang YX, Guo C, Li DZ (2020a) A new subtribal classification of Arundinarieae (Poaceae, Bambusoideae) with the description of a new genus. Plant Diversity 42(3): 127–134. https://doi.org/10.1016/j.pld.2020.03.004

Zhang D, Gao FL, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020b) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20(1): 348–355. https://doi.org/10.1111/1755-0998.13096

Zhu ZD, Li DZ, Stapleton CMA (2006) *Pseudosasa* Makino ex Nakai. In: Wu ZY, Raven P (Eds) Flora of China (Vol. 22). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 115–121.