



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Paper

A brief study on phytochemical and pharmacological profile of calotropis gigantea

Harshali Ghorpade*, Prafull Chavan, Prerana Bagul, Momin Sanaurrehman

Dr. Uttamrao Mahajan College Of B.pharmacy, Chalisgaon

ARTICLE INFO

Published: 18 Feb. 2025

Keywords:

gigantea Calotropis,
Milkweed phytochemistry,
pharmacological activity,
antioxidant, and anti-
inflammatory properties

DOI:

10.5281/zenodo.14888978

ABSTRACT

The main point is that we may access a variety of life-saving medications, either in modified synthetic form or directly in extract form. Calotropis gigantea is a huge, gregarious shrub with many branches and young branches covered in white, cottony hairs. It has been used as a medicinal plant and a key source of pharmaceuticals since ancient times. It has milky latex in it. Commonly referred to as "milkweed," this plant is native to China, Malaysia, and India and is found practically everywhere in the world. It has long been utilized in the Siddha, Ayurvedic, and Unani medical systems. A variety of illnesses may be treated with the help of this plant's parts, including its roots, bark, latex, leaves, and flowers. Among the many isolated compounds found in this medicinal plant are alkaloids, tannins, resins, flavonoids, terpenoids, cardiac glycosides, and many chemical compounds such as giganteol, α and β calotropeol, β -amyrin, giganteol, and isogiganteol. In addition to these chemical compounds, C. gigantea has some pharmacological activity, including anti-inflammatory, anti-diarrheal, hepatoprotective, anti-asthmatic, antioxidant, antibacterial, antiviral, wound healing, and hypoglycemic properties.

INTRODUCTION

Animals, plants, and other natural items have had a profound influence on human culture and civilization from prehistoric times to the present in various regions of the world, including India. Since the dawn of civilization, people have revered plants, which are preserved as genetic resources and utilized for fuel, food, fiber, fodder, fertilizer,

febrifuge, and other purposes. Calotropis gigantea, a member of the Apocynaceae family, is a well-known medicinal herb that has long been utilized in Ayurvedic, Unani, and Siddha medicine systems. It is also referred to as milkweed or crown flower weed. This is a latex-bearing plant that releases latex when tissue is damaged.

***Corresponding Author:** Harshali Ghorpade

Address: Dr. Uttamrao Mahajan College Of B.pharmacy, Chalisgaon

Email ✉: harshalighorpade1@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



Combinations of tannins, alkaloids, sugar, starch, resins, protein, and gum make up plant latex.



Fig: - Calotropis gigantea plant.

Vernacular Classification: -

India	(Sanskrit) Arka, Ganarupa, Mandara, Vasuka, Svetapushpa, Sadapushpa, Alarka, pratapass, (Hindi) Aak, Madar, (Kannada) Ekka, (Tamil and Malayalam) Erukku, (Telugu) Jilledi puvvu.
English	Crown flower, giant indian milkweed.
Vietnam	B[oot]ng b[oot]ng, 1[as]hen, nam t[it] b[at].
Indonesia	Bidhuri (sundanese, Madurese), sidaguri (Javanese), rubik (Aceh)
Malaysia	Remiga, rembega, kemengu.
Laos	Remiga, rembega, kemengu.
Thailand	Po thuean, paan thuean (northern), rak (central).
Philippines	Kapal-kapal (Tagalog).

Scientific Classification: -

Sr. No	Kingdom	Plantea	Sr. No	Kingdom	Plantea
1	clade	Tracheophytes	1	Order	Gentianales
2	clade	Angiosperms	2	Family	Apocynaceae
3	clade	Eudicots	3	Genus	Caloterpis
4	clade	Asterids	4	Species	C.gigantea

Taxonomical Classification:

Kingdom	Plantae
Order	Gentianales
Family	Apocynaceae
Subfamily	Asclepiadaceae
Genus	Calotropis
Species	C.gigantea

Morphology: -

Root: Simple, branching, woody at the base, with a fissured bark; branches are heavily white tomentose and rather tasty; it is glabrescent in the

early stages. When the plant is cut or broken, white latex is released from every portion.

Leaves: Opposite-decussate, simple, subsessile, exstipulate; oval obovate to fully obovate edge, 5-30X 2.5-15.5 cm, base cordate, apex abruptly and

quickly acuminate to apiculate, margins whole, delectable, white tomentose when young, later glabrescent and glaucous.

Fruit: A simple, fat, enlarged, subglobose to laterally ovoid follicle with a diameter of at least 10 cm. Numerous tiny, flat, oblong seeds that are 6 × 5 mm in size and packed with smooth white pappus that are at least 3 cm long.

Flowers: Actinomorphic, pentamerous, hypogynous, bracteate, complete, sexually unbiased, pedicellate, pedicel 1-3 cm long.

Calyx: Sepal 5, glabrescent, quincuncial aestivation, polysepalous, five lobed, connected at the base in a couple of seconds Androecium The anther is ditheous, gynandrous, and has five stamens.

Inflorescence: Thick, multibloomed, umbellate, peduncled cymes that appear axillary or terminal emerge from the hubs.

Gynoecium: The peltate disgrace has five parallel stigmatic surfaces and is bicarpellary and apocarpous, with styles united at the apex. The embarrassment of framing a gynostegium is akin to anthers.

Natural habitual, geological, or biological sources:

Throughout the country, *Calotropis* grows wild up to 900 meters (msl) and can withstand drought and salt to a moderate degree. It prefers sandy, disturbed soils with 300–400 mm of mean annual rainfall. Because its seeds are dispersed by wind and animals, it quickly becomes a weed along abandoned roads, lagoon shores, and in natural grasslands that have been overgrazed. It usually predominates in areas with disturbed sandy soils and low rainfall, and it loves abandoned agricultural sites. It is said to be an indication of over cultivation. For *C. gigantea*, a broad habitat with little competition is ideal. This species' plants grow in dry environments with rainfall limited to 150–1000 mm per year and in regions with overly drained soil where annual precipitation can reach

up to 2000 mm. Additionally, it may be found in common settings such as densely populated metropolitan areas, seaside dunes, and roadside sand dunes. *C. gigantea* can also be found at elevations of up to 1,000 meters above sea level. The plant is sometimes planted as an ornamental in arid or coastal areas since it is easy to manage, reproduces, and can survive in xerophytic environments. In addition to Madagascar, the Arabian Peninsula, West Africa, North and East Africa, Macaronesia, and South Asia, *C. gigantea* is indigenous to Southern Asia and Indo-China. The plant is a natural species in Australia, Central America, North America, South America, and the West Indies. Many countries now tolerate and grow the plant, including those in Mexico, Central and South America, the Pacific Islands, Australia, and the Caribbean.

The Chief Features: -

- The plant grows well in a range of climate conditions and soil types.
- Thus, it can be found in tropical and
- It grows best on poor soils, especially in areas where overgrazing has reduced competition from indigenous grasses.
- All the subtropical areas of the world, encompassing all
- It doesn't need cultivating methods.
- One of the few plants that grazing animals avoid eating is this one.
- India

Therapeutic Uses:

In addition to treating leprosy, leucoderma, ulcers, tumors, piles, and diseases of the spleen, liver, and abdomen, the plant is purgative, anthelmintic, alexipharmic, and its juice prevents leucoderma, tumors, ascites, and disorders of the abdomen. Wounds, painful or paralyzed joints, and swellings are all treated with the leaves. The leaves are tinctured and used as an antiperiodic to treat intermittent fevers. Ascites, inflammation, rat bites, and tumors. The milk relieves piles, is bitter,



purgative, and laxative. The diaphoretic root bark cures asthma and syphilis. The flower is astringent, analgesic, anthelmintic, sweet, bitter, and medicinal


Chemical Constituent: -

Various chemical constituent isolated from *C.gigantea* Linn:


Class of chemical constituent	Name of chemical constituent	Plant used	Part	Extract Takes
Triterpenoids	Di-(2-ethylhexyl) Phthalate	Flowers		Ethyl acetate extract
	Anhydrosophoradiol-3-acetate			
	Lupeol	Aerial parts		Latex
	α -Taraxerol	Root bark		Ethyl acetate extract
Triterpene esters	γ -Taraxasterol	Aerial parts		Hexane and methanol soluble extract
	Lupenyl-1-acetate	Root bark		Petroleum ether extract
Flavonol	Isorhamnetin	Aerial parts		Methanol extract
Cardiac glycosides	Calotropone	Root bark		Ethanol extarct
	Gofruside			
Steroids	Stigmasterol	Root bark		Methanol extract
	β -Sitosterolacetate			
	β -Sitosterolacetate			Ethyl acetate extract
Resin	β -Amyrin	Root bark		95% Alcohol extract
	β -Amyrin acetate			
Fatty acids	Isovaleric acid	Root bark		95%Alcohol extract
Miscellaneous	Asclepin	Roots		Latex




Lupeol



β - Sitosterol acetate



Calotropone



Isorhamnetin

Chemical structures of various chemical constituents isolated from *C. gigantea*.




Fig. Medicinal properties of *C.gigantea* Linn.

Pharmacological Activities: -

1. Anti-inflammatory activity:

Several experimental animal models were used to assess *Calotropis gigantea*'s anti-inflammatory properties. The anti-

inflammatory properties of chloroform, n-butanol, ethanol, and distilled water extract of *Calotropis gigantea* leaves were examined. Using the carrageenan-induced rat paw oedema method, this activity was contrasted

with that of the common medication, paracetamol. Either a stronger malic enzyme or the filarial worm *Setaria digitata*: specific drug-and herbal extract properties and effects are indicated by the *calotropis gigantea* alkaloid fraction's residual anti-inflammatory activity. Conversely, the anti-inflammatory action is assessed using a model of chronic inflammation caused by adjuvant-induced arthritis and a model of acute inflammation caused by carrageenin-induced kaolin-induced rat paw oedema and cotton-pellets granuloma. Additionally, the anti-inflammatory properties were demonstrated against the albumin denaturation approach.

2. **Anticancer activity:**

Anhydrosophoradiol-3-acetate (A3A), an isolated chemical found in *Calotropis gigantea* flowers, kills viable tumor cells and reduces body weight growth. A3A's ability to effectively suppress EAC growth in vivo has been shown to improve cancer-induced complications.

3. **Antitussive activity:**

Calotropis gigantea leaf extract has antitussive properties since it contains alkaloids and glycosides.

4. **Antidiarrheal activity :**

The anti-diarrheal properties of *Calotropis gigantea* aerial component hydroalcoholic (50:50) extract were evaluated in a rat model of castor oil-induced diarrhea. The gastrointestinal transit rate was determined by dividing the proportion of the longest distance traveled by the charcoal by the total length of the small intestine. The weight and volume of intestinal content caused by castor oil were measure during the enterpoling method. The extract significantly reduced the frequency of fecal dropping and fecal output at dosages of 200 and 400 mg/kg body weight (interaperitoneal dose). Additionally, the

extract greatly decreased the weight and volume of intestinal material.

5. **Analgesic activity:**

Alcohol extract from *Calotropis gigantea* flowers was administered orally, and its analgesic effects were evaluated in mice utilizing thermal and chemical methods. In the acetic acid-induced writhing test, a 20.97% reduction in the number of writhes was seen at doses of 250 mg/kg, respectively. The hot plate method extended the time spent licking the paws. Thirty minutes after the dose was given, the analgesic effect became apparent, and ninety minutes later, it peaked. The analgesic potential of dry latex (DL) derived from *C. gigantea* was evaluated in this study. The effect of DL at a dose of 415 mg/kg against acetic acid-induced writhing was more noticeable than that of an oral dose of aspirin (100 mg/kg). DL (830 mg/kg) produced a modest analgesic effect in the tail-flick assay that was similar to that of aspirin.

6. **Wound healing activity:**

Studies on the activity of wound healing using animal models are many. Utilizing an excision and incision wound model, *Calotropis gigantea* latex was first applied to albino rats. When comparing the latex-treated animal to the control, which was 76.22% with framycetin sulphate cream (1%) as a standard, the latter showed an 83.42% reduction in wound area.

7. **Hair growing Activity:**

The purpose of *Calotropis gigantea* with *Hibiscus rosa sinensis* (HRSF) and polyherbal formulation (HCF) was to determine the effects of the two plants on the commencement and encouragement of hair development in albino rats. The findings and observations of the study were contrasted with those of Minoxidil. Compared to other treatments, *Calotropis gigantea* had reduced potential hair growth activity.



8. CNS Activity:

Calotropis gigantea extract exhibits a variety of neuropharmacological activities, including sedative and hypnotic, antidepressant, antianxiety, analgesic, and more, because it contains the flavonoids chrysin and apigenin. The C. gigantea extract enhances collateral branching, axonal and dendritic length, quantity, and branching orders, even in this kind of activity.

9. Anti-venom Activity:

Antivenom action against viper russell snake venom is provided by Calotropis gigantea plant extract. Calotropis gigantea methanolic extract has beneficial effects that counteract the venom's different actions, including mortality, oedema formation, necrotizing, and hemorrhagic activity. The deadly effects of 2LD50 and 3LD50 venom were substantially mitigated in mice by oral administration of extract at 200 and 400 mg/kg. The plant extract effectively neutralized the 2LD50 and 3LD50 of venom in in vitro experiments at doses of 100, 200, and 400 mg/kg. Methanolic extract from viper venom had an effect on oedema at 60, 120, 180, and 240 minutes.

10. Anti-malaria:

Calotropis gigantea extract exhibits antimalarial action against Plasmodium falciparum and Plasmodium berghei. This plant has outstanding anti-malarial action because it exhibits anti-plasmodial activity against the chloroquine-sensitive Plasmodium falciparum.

Traditional Uses of Calotropis Gigantea:-

1. In Ayurveda: Paralysis, swellings, and intermittent fevers are all treated with the leaves of the C. gigantea plant. Flowers can be used to treat fever, inflammation, helminthic infections, anorexia, asthma, and catarrh. The root bark of the plant is used to treat skin diseases, intestinal worms, helminth infections, and ascites.

2. In Siddha: - The leaves of C. gigantea are utilized for the treatment of dangerous snake bites, periodic fever, vatha illnesses, intestinal worms and ulcers. The roots of this plant are crushed completely and then firmly rubbed over the bite area. The latex of this plant can be used to treat gonococcal arthritis, swellings, dental problems, rat bites, and other rheumatic conditions. Using flowers to treat bronchial asthma

3. In Unani: In the Unani system, root bark powder has traditionally been used to treat dysentery and diarrhea. The plant's root helps with dyspepsia and has carminative properties.

CONCLUSION:

Calotropis gigantea is a potential plant with various healing principles and economic values. It is generally assigned through all over the India, which is use as traditional medicine, ornamental, fuel, fibre, auxiliary plant, mosquito controlling etc. and the various part of this plant viz. root, leaves, root bark, milk, flower are used ethnomedicinally as a remedy for various diseases for human beings. Though Calotropis gigantea has various medicinal applications, but still in future the phytochemicals and unknown compound of this plant need to be standardized and characterization of new potent molecule that suppress various pathological disorders and development of new class of drug therapies for the betterment of health of human beings.

REFERENCES

1. Sureshkumar P, Chezian A, Senthil Raja P and Sathiyapriya J; Computational selections of terpenes present in the plant Calotropis gigantea as mosquito larvicide's by blocking the sterol carrying protein, Bangladesh J Pharmacol, 2012, 7: 1-5
2. Singh U, Wadhwani AM, Johri BM. Dictionary of Economic Plants of India. New Delhi: Indian council of Agricultural Research; 1996:38-39.



3. Aarti C. A review on pharmacological and biological properties of *Calotropis gigantea*. International Journal of Recent Scientific Research. 2014;5(4):716-719.
4. https://en.wikipedia.org/wiki/Calotropis_gigantea
5. Divya N, Singh AB. A Review on Brief Study of *Calotropis gigantea* Linn. Journal of Drug Delivery and Therapeutics. 2021; 11(5):224-228.
6. Amutha A., Jeyalalitha T and Kohila M. *Calotropis gigantea* a review paper. International Journal of Recent Scientific Research. 2018; 9(10):29386-29390.
7. Chatterjee A, Pakrashi SC. The Treatise on Indian Medicinal Plants. V-IV. New Delhi: National Institute of Science Communication and Information Resources. 2003: 128-129.
8. Gamble J S; Flora of the Presidency of Madras, Vol. I, II, III, Botanical survey of India, Calcutta, 1935.
9. Singh N, Gupta P, Pathak AK, Patel AV. *Calotropis gigantea*: a review on its phytochemical & pharmacological profile. International Journal of Pharmacognosy. 2014;1(1): 1-8.
10. Mushir A, Jahan N, Ahmed A. A review on phytochemical and biological properties of *Calotropis gigantea* (Linn.) R.Br. Discovery Phytomedicine 2016; 3 (2): 15-21
11. Rahman MS, Moly NN, Hossen MJ. Antimicrobial, Cytotoxic and Antioxidant Activity of The Exudate of *Calotropis gigantea*. International Journal of Pharmaceutical Sciences and Research. 2013 Feb 1; 4(2):745.
12. Kumar PS, Suresh E, Kalavathy S. Review on a potential herb *Calotropis gigantea* (L.) R. Br. Scholars Academic Journal of Pharmacy. 2013; 2(2):135-43.
13. Kori P, Alawa P. Antimicrobial activity and phytochemical analysis of *Calotropis gigantea* root, latex extracts. IOSR J. Pharm. 2014; 4(6):7-11. <https://doi.org/10.9790/3013-04060207011>
14. Kumar G, Karthik L, Rao KV. A review on pharmacological and phytochemical profile of *Calotropis gigantea* Linn. Pharmacologyonline. 2011; 1:1-8.
15. Amutha A., Jeyalalitha T and Kohila M. 2018, *Calotropis Gigantea* A Review Paper. Int J Recent Sci Res. 9(10), pp. 29386- 29390.
16. Sarkar S, Chakraverty R, Ghosh A. *Calotropis Gigantea* Linn. -a complete busket of Indian traditional medicine. Int. J. Pharm. Res. Sci. 2014; 2(1):7-17.
17. Deshpande S, Deshpande K, Tomar E. *Calotropis Gigantea*: A Phytochemical Potential. (2018)
18. Gharge VG, Ghadge DM, Shelar PA, Yadav AV. Importance of Pharmacognostic study of medicinal plants *Calotropis gigantea* (Linn.): A review. Int J Pharmacognosy. 2017; 4(11):363-71.
19. Kumar D, Kumar S. *Calotropis gigantea* (L.) Dryand-A review update. Indian Journal of
20. Gamble J S; Flora of the Presidency of Madras, Vol. I,II, III, Botanical survey of India, Calcutta, 1935. (sajp).
21. Singh, U., A.M. Wadhwani, and B.M. Johri,1996. Dictionary of Economic Plants of India. Indian Council of Agricultural Research, NewDelhi. p. 38-39. Rastogi, Ram, 1991.
22. Gamble J S; Flora of the Presidency of Madras, Vol. I,II, III, Botanical survey of India, Calcutta, 1935.
23. Vaidya A: Pharm Res. India (Pharma Pulse – Suppl), 1998; 44-45.
24. Haslam EJ and Nat: Prod. 59, 1996; 205-215.
25. Yelne MB, Sharma PC, Dennis TJ. Database onmedicinal plants used in ayurveda, central council for research in ayurveda and siddha,New Delhi; Vol. 2,69-73(2000).



26. Chitme HR, Chandra R, Kaushik S, Studies on anti-diarrhoeal activity of *Calotropis*.
27. Sharma AP and Tripathi BD; Assessment of atmospheric PAHs profile through *Calotropis gigantea* R.Br. leaves in the vicinity of an Indian coal-fired power plant, *Environ Monit Assess.*, 2009, 149: 477 –482.
28. Gamble J S; *Flora of the Presidency of Madras*, Vol. I, II, III, Botanical survey of India, Calcutta, 1935.
29. Ahmed KK, Rana AC, Dixit VK. *Calotropis* species (Asclepiaceae): A comprehensive review. *Pharmacogn Mag* 2005; 1:48-52.
30. Parrotta JA. *Healing Plants of Peninsular India*. Wallingford, UK and New York: CAB International; 2001. p. 944 *Research in Pharmacy and Biotechnology*. 2015 May 1; 3(3):218.

HOW TO CITE: Harshali Ghorpade*, Prafull Chavan, Prerana Bagul, Momin Sanaurrehman, A brief study on phytochemical and pharmacological profile of *calotropis gigantea*, *Int. J. of Pharm. Sci.*, 2025, Vol 3, Issue 2, 1554-1562. [https://doi.org/ 10.5281/zenodo.14888978](https://doi.org/10.5281/zenodo.14888978)

