

A new species of *Sedum* (Crassulaceae, Saxifragineae) from Guangxi, China

Chun-Yu Zou^{1,2}, Tao Meng¹, Shi-Yong Meng³, Yan Liu¹ 

¹ Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin, 541006, Guangxi, China

² Laboratory of Systematic Evolution and Biogeography of Woody Plants, College of Nature Conservation, Beijing Forestry University, Beijing, 100871, China

³ College of Life Sciences, Peking University, Beijing, 100871, China

Corresponding author: Yan Liu (gxibly@163.com)

Abstract

A new species, *Sedum guangxiense* **sp. nov.**, discovered in Guangxi Province, China, is described and illustrated. Molecular phylogenetic analyses and morphological differences indicate that this species is well separated from its close relatives in *Sedum*, justifying its recognition as a distinct new species. Morphologically, it closely resembles *Sedum tosaense* and *S. emarginatum* in its leaf shape, inflorescence type and life form, but it can be easily distinguished in its erect stems when young, much larger leaves, narrowly triangular petals, ovate nectar scales and oblong anthers.

Key words: Phylogeny, sect. *Sedum*, *Sedum guangxiense*, taxonomy



Academic editor: Joachim Thiede

Received: 21 May 2024

Accepted: 27 January 2025

Published: 20 February 2025

Citation: Zou C-Y, Meng T, Meng S-Y, Liu Y (2025) A new species of *Sedum* (Crassulaceae, Saxifragineae) from Guangxi, China. *PhytoKeys* 252: 287–299. <https://doi.org/10.3897/phytokeys.252.128011>

Copyright: © Chun-Yu Zou et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The genus *Sedum* Linnaeus (1753: 430) (stonecrops) comprises ca. 470 species mainly distributed in the Northern Hemisphere, but some extend to the Southern Hemisphere in Africa and South America (Fu and Ohba 2001). It typically thrives in open, climatically (semi-)arid environments and comprises herbs to (sub-)shrubs with succulent leaves (Stephenson 1994). The molecular study by Messerschmid et al. (2020) showed that *Sedum* is highly polyphyletic and, in order to render *Sedum* monophyletic, about 14 genera derived from within *Sedum* would have to be merged with it. This broader concept of *Sedum* would encompass approximately 755 species, mainly distributed in the Mediterranean, Central America, the Himalayas and East Asia (Stephenson 1994; Thiede and Eggli 2007).

In China, there are approximately 120 species of *Sedum* including 91 endemics which are particularly abundant in the south-western region. *Sedum* is classified into sect. *Filipes* (Fröd.) S.H. Fu (1965: 115), sect. *Oreades* (Fröd.) K.T. Fu (1974: 52) and sect. *Sedum* (Fu and Ohba 2001) which is by far the largest. China stands out as one of the primary centres of biodiversity for *Sedum*, with several studies undertaken (Fu 1965, 1974; Rao 1996; Fu and Ohba 2001). However, ongoing exploration reveals new species within the *Sedum* genus in

China, indicating that the full extent of its diversity is yet to be uncovered (Zou et al. 2020; Ying 2022a, 2022b; Huang et al. 2023; Meng et al. 2023).

Guangxi Province, located in southern China, boasts a significant biodiversity. The challenge of accurately identifying *Sedum* species within this region is difficult due to the presence of over 17 species. Notably, during the taxonomic revision of *Sedum* in Guangxi, anomalous specimens were collected in Yongfu County, Guangxi Province, China, in 2013. Subsequently, a field survey was conducted in May 2020 to investigate these unidentified specimens. Field observations suggested similarities to *S. tosaense* Makino (1892: 52). However, a thorough examination of both living and herbarium specimens revealed distinctions from *S. tosaense*, particularly in the erect stems during early growth stages, petal morphology, nectar scale shape and style length.

During a field survey of plant resources in Fangcheng City, Guangxi, China, in April 2021, populations of an unknown *Sedum* species were discovered in forests. Through field observations and detailed comparisons, we concluded that this plant is conspecific with the one discovered in Yongfu County, establishing it as a putatively undescribed species. This study intends to determine the phylogenetic position of these plants and to provide a comparative morphological analyses with its close relatives within the genus *Sedum*. To achieve this, phylogenetic analyses were conducted using molecular sequence data from nuclear markers (nrITS) within Eastern Asian *Sedum* sect. *Sedum*. In this study, we clarify the status of the putatively new species by determining its phylogenetic position and comparing its morphology with that of its close relatives.

Material and methods

To examine its phylogenetic relationships and confirm the morphological distinctiveness of this new species from other taxa, voucher specimens and living individuals of *S. guangxiense* were collected from Yongfu County and Fangcheng City. We consulted the relevant literature (Fu 1965, 1974; Rao 1996; Fu and Ohba 2001; Wu et al. 2012; Yang et al. 2012; Xie et al. 2014; Ito et al. 2014a, 2014b, 2017a, 2017b, 2020b; Lu et al. 2019; Zou et al. 2020; Ying 2022a, 2022b; Huang et al. 2023; Meng et al. 2023) and studied herbarium specimens at the Herbaria IBK, IBSC, KUN and PE, therewith comparing our new species with all described Eastern Asian species of *Sedum* sect. *Sedum*.

To determine the phylogenetic position of the new species, nuclear internal transcribed spacer (ITS) sequences of 62 accessions were utilised representing 61 species (Mort et al. 2002; Mayuzumi and Ohba 2004; Ito et al. 2014a, 2017a, 2017b, 2018, 2020a; Xie et al. 2014; Zou et al. 2020; Kim et al. 2023; Huang et al. 2023), including the putatively new species. Fifty-six species belong to *Sedum* sect. *Sedum*, while two species belong to *S.* sect. *Oreades*. These were obtained from GenBank and included in the ITS analysis (Table 1). Based on a phylogenetic study of Crassulaceae (Mayuzumi and Ohba 2004), *Aeonium castello-paivae* Bolle (1859: 240), *A. gomerense* Praeger (1929: 473), *A. lancerottense* Praeger (1932: 190), *A. viscatum* Bolle (1859: 241), *Greenovia aizoon* Bolle (1859: 242) and *Phedimus takesimensis* (Nakai) 't Hart (1995: 165) were selected as outgroup members (Table 1).

Table 1. Taxon, locality, accession number and reference for internal transcribed spacer (ITS) sequences of *Sedum*, *Aeonium*, *Greenovia* and *Phedimus* species registered in the GenBank database. The sequences were used for molecular analyses.

Taxon	Origin	Accession number	References
Ingroup			
Sedum section Oreades			
<i>S. trullipetalum</i>	Nepal	AB088630	Mayuzumi and Ohba (2004)
<i>S. oreades</i>	Nepal	AB088632	Mayuzumi and Ohba (2004)
Sedum section Sedum			
<i>S. actinocarpum</i>	China: Taiwan	LC229265	Ito et al. (2017a)
<i>S. alfredii</i>	China: Guangdong	AB930259	Ito et al. (2014a)
<i>S. arisanense</i>	Japan	LC229273	Ito et al. (2017b)
<i>S. baileyi</i>	China	FJ919935	Unpublished
<i>S. brachyrinchum</i>	China: Taiwan	LC229276	Ito et al. (2017b)
<i>S. boninense</i>	Japan	LC530821	Ito et al. (2020a)
<i>S. bulbiferum</i>	Japan	KM111165	Xie et al. (2014)
<i>S. danjoense</i>	Japan	LC260127	Ito et al. (2017a)
<i>S. emarginatum</i>	China: Anhui	EU592006	Unpublished
<i>S. erythrospermum</i>	China: Taiwan	AB906473	Ito et al. (2014a)
<i>S. formosanum</i>	China: Taiwan	AB930271	Ito et al. (2014a)
<i>S. guangxiense</i> -FC	China: Guangxi	OL693034	This study
<i>S. guangxiense</i> -YF	China: Guangxi	OL693035	This study
<i>S. hakonense</i>	Japan	AB930278	Ito et al. (2014a)
<i>S. hangzhouense</i>	China: Zhejiang	LC260130	Ito et al. (2017b)
<i>S. japonicum</i>	Japan	LC229237	Ito et al. (2017a)
<i>S. jinglanii</i>	China: Guangdong	OQ162326	Huang et al. (2023)
<i>S. jiulongshanense</i>	China: Zhejiang	LC229243	Ito et al. (2017a)
<i>S. kiangnanense</i>	China: Zhejiang	LC229244	Ito et al. (2017a)
<i>S. kwanwuense</i>	China: Taiwan	LC229293	Ito et al. (2017a)
<i>S. lineare</i>	Japan	AB088623	Mayuzumi and Ohba (2004)
<i>S. lipingense</i>	China: Guizhou	MN150061	Zhang et al. (2019)
<i>S. lungtsuanense</i>	China: Zhejiang	LC260131	Ito et al. (2017b)
<i>S. mexicanum</i>	Japan	LC229247	Ito et al. (2017b)
<i>S. makinoi</i>	Japan	AB906476	Ito et al. (2014a)
<i>S. microsepalum</i>	China: Taiwan	LC229281	Ito et al. (2017a)
<i>S. morrisonense</i>	China: Taiwan	AB906477	Ito et al. (2014a)
<i>S. mukojimense</i>	Japan	LC530823	Ito et al. (2020a)
<i>S. multicaule</i>	Nepal	AB088631	Mayuzumi and Ohba (2004)
<i>S. nagasakianum</i>	Japan	LC229249	Ito et al. (2017a)
<i>S. nanlingense</i>	China: Guangxi	MN105949	Zou et al. (2020)
<i>S. nokoense</i>	China: Taiwan	AB906478	Ito et al. (2014a)
<i>S. oligospermum</i>	China	LC229250	Ito et al. (2017a)
<i>S. onychopetalum</i>	China: Nanjin	KM111148	Xie et al. (2014)
<i>S. oryzifolium</i>	South Korea	KF954525	Unpublished
<i>S. polytrichoides</i>	China: Anhui	KM111143	Xie et al. (2014)
<i>S. rupifragum</i>	Japan	LC229254	Ito et al. (2017b)
<i>S. sarmentosum</i>	Japan	LC229255	Ito et al. (2017a)

Taxon	Origin	Accession number	References
<i>S. satumense</i>	Japan	LC229256	Ito et al. (2017a)
<i>S. sekiteiense</i>	China: Taiwan	LC229295	Ito et al. (2017a)
<i>S. subtile</i>	Japan	LC229257	Ito et al. (2017a)
<i>S. taiwanalpinum</i>	China: Taiwan	LC229278	Ito et al. (2017a)
<i>S. taiwanianum</i>	China: Taiwan	LC229296	Ito et al. (2017a)
<i>S. tarokoense</i>	China: Taiwan	LC229298	Ito et al. (2017b)
<i>S. tetractinum</i>	China: Zhejiang	LC260135	Ito et al. (2017b)
<i>S. tianmushanense</i>	China: Zhejiang	LC229261	Ito et al. (2017a)
<i>S. tosaense</i>	Japan	AB088620	Mayuzumi and Ohba (2004)
<i>S. triactina</i>	Japan	AB088629	Mayuzumi and Ohba (2004)
<i>S. tricarpum</i>	Japan	LC229259	Ito et al. (2017a)
<i>S. truncatistigmum</i>	China: Taiwan	LC229304	Ito et al. (2017a)
<i>S. uniflorum</i>	Japan	LC530832	Ito et al. (2018)
<i>S. yabeanum</i>	Japan	AB906490	Ito et al. (2014a)
<i>S. zentaro-tashiroi</i>	Japan	AB088619	Mayuzumi and Ohba (2004)
<i>S. wenchuanense</i>	China	ON707681	Unpublished
Outgroups			
<i>Aeonium castello-paivae</i>	Spain: Canary Islands	AY082127	Mort et al. (2002)
<i>Aeonium gomerense</i>	Spain: Canary Islands	AY082133	Mort et al. (2002)
<i>Aeonium viscatum</i>	Spain: Canary Islands	AY082154	Mort et al. (2002)
<i>Aeonium lancerottense</i>	Spain: Canary Islands	AY082143	Mort et al. (2002)
<i>Greenovia aizoon</i>	Spain: Canary Islands	AY082112	Mort et al. (2002)
<i>Phedimus takesimensis</i>	Korea	OP346962	Kim et al. (2023)

The total genomic DNA was extracted from silica gel-dried leaf materials using the CTAB protocol (Doyle and Doyle 1987). DNA sequences of nuclear ribosomal ITS were selected as a marker for molecular phylogenetic studies. The nrITS region was amplified using polymerase chain reactions (PCR) with universal primers, following the protocol of Huang et al. (2017). The complete DNA sequences were submitted to GenBank (Table 1). The ITS sequences obtained by PCR were aligned using MUSCLE version 3.8.31 (Edgar 2004) and were adjusted manually in Bioedit 5.0.9 (Hall 1999). A phylogeny was constructed using Maximum Likelihood (ML) and Bayesian Inference (BI). ML analyses were performed using RAxML v.7.0.4 (Stamatakis et al. 2008) and the substitution model was estimated with ModelTest (Posada and Crandall 1998). BI was conducted in MrBayes 3.2.6 (Ronquist et al. 2012) with the optimal substitution model selected by ModelTest (Posada and Crandall 1998) according to the Akaike Information Criterion. All BI analyses were run for 1,000,000 generations, with four chains in two parallel runs and one tree sampled every 5,000 generations. The convergence of two parallel runs was guaranteed by a splitting frequency of less than 0.005. All other parameters were set to default. The first 25% of the sampled trees corresponding to the burn-in period were discarded and the remaining trees were used to construct a majority rule consensus tree.

Results

Morphological comparison

In morphological terms, two species closely resemble the undescribed species: *S. tosaense* and *S. emarginatum* Migo (1937: 224). It shares similarities with *S. tosaense* in leaf shape, inflorescence type with three branches (each 2-forked) and the basal leaves forming a rosette. However, the newly-discovered species is distinguished by its erect stems when young, narrowly triangular petals, ovate nectar scales, oblong anthers and longer styles. In contrast, the leaves of sterile shoots of *S. tosaense* are rounder, the leaf base is long-attenuate, but shorter, the nectar scales are subquadrangular at the base and the anthers are oblong-ovoid. It is similar to *S. emarginatum* in having spatulate to obovate leaves with a notch or shallow indentation, but differs in having erect stems when young and alternate leaves.

Molecular analysis

ML and BI phylogenetic analyses were performed, based on the ITS sequences of 54 *Sedum* species. This set included two ITS sequences from our undescribed species and 53 ITS sequences from GenBank. A 50% majority rule consensus tree of all post-burn-in trees using the optimal substitution model GTR and Bayesian posterior probabilities was generated (PP, Fig. 1). ML analyses were performed with the substitution model GTRGAMMA and 1,000 rapid bootstrap searches. The topology of the ML tree was highly similar to that of the Bayesian tree and the bootstrap support values for this tree were depicted on Fig. 1 (BS). BI and ML analyses confirmed that the putative new species constituted a well resolved clade, sister to a clade with *S. oligospermum*, *S. sekiteiense* and *S. actinocarpum* and 13 further *Sedum* species (PP = 0.99, BS = 99).

Discussion

The molecular phylogenetic analysis carried out in this study indicates that *S. guangxiense* constitutes a well-supported distinct clade, which is, with high support, sister to a well-supported clade (named clade A in Fig. 1) which includes *S. oligospermum*, *S. sekiteiense*, *S. actinocarpum* and 13 further species (Fig. 1). However, morphologically, *S. guangxiense* can be readily distinguished by its biennial life form (Table 2), erect stems in youth, larger leaves with a retuse apex, usually 5-merous, but sometimes 4- or 6-merous flowers and longer styles. Although it bears resemblance to *S. tosaense* and *S. emarginatum*, these species are distantly related to *S. guangxiense* in the molecular tree (Fig. 1). Specifically, *S. tosaense* and *S. emarginatum* differ in having a perennial life form, prostrate stems when young, smaller leaves with a rounded and emarginate apex, smaller anthers and shorter styles (Table 2). In addition, *S. guangxiense* (Table 2) differs from *S. tosaense* in its ovate (vs. subquadrangular) nectar scales and from *S. emarginatum* in its alternate leaves (vs. opposite and loosely arranged leaves), strongly unequal (vs. subequal) sepals and ovate (vs. oblong) nectar scales.

In recent years, more than 25 new species of *Sedum* have been reported in China, especially in southwest China. This high level of species richness indicates that this area should be further explored to fully unravel the rich biodiversity there.




Figure 1. Bayesian phylogenetic tree, based on ITS sequences of Eastern Asian *Sedum*. Bootstrap percentages from the ML analysis (left) and Bayesian posterior probabilities (right) are shown at the nodes.

Table 2. Morphological comparisons between *Sedum guangxiense*, *S. tosaense* and *S. emarginatum*, based on our own measurements of herbarium and living specimens of all three species as well as literature data (Fu and Ohba 2001).

Characters	<i>S. guangxiense</i>	<i>S. tosaense</i>	<i>S. emarginatum</i>
Life form	Biennial	Perennial	Perennial
Flowering stems	Erect when young	Prostrate when young, later erect	Prostrate when young, later erect
Leaf arrangement	Basal leaves forming a rosette	Basal leaves forming a rosette	Basal leaves opposite, loosely arranged
Leaves	Alternate, spatulate to obovate, 1.5–3.5 × 0.5–1 cm, apex usually retuse, rarely round	Alternate, linear-spatulate, 1.2–2 × 0.5–1 cm, apex rounded and emarginate	Opposite, spatulate-obovate to broadly obovate, 1–2 × 0.5–1 cm, apex rounded and emarginate
Inflorescence type	Cyme often 3-branched, each 2-forked	Cyme often 3 (sometimes 4)-branched, each 2-forked	Cymes usually 3-branched, each 2-forked
Flowers	Usually pentamerous, sometimes 4- or 6-merous	Pentamerous	Pentamerous
Carpels	Usually 5, sometimes 4 or 6	5	5
Sepals (colour, shape and size)	Green, spatulate to obovate, strongly unequal, 3–8 × 1–3 mm	Green, linear-spatulate, strongly unequal, 4.5–10 × 2–4.5 mm	Green, spatulate-obovate to broadly obovate, 4–5 × 1–1.5 mm
Petals (colour, shape and size)	Yellow, narrowly triangular, 5–6 × 1.1–1.5 mm	Yellow, narrowly elliptic-lanceolate, 5–6 × 1.5–2.5 mm	Yellow, linear-lanceolate to lanceolate, 6–8 × 1.5–2 mm
Nectar scales (colour, shape and size)	Yellow, ovate, 0.3–0.4 mm	Yellowish, subquadrangular, 0.2–0.4 mm	Yellow, oblong, ca. 0.3–0.6 mm
Anthers (colour, shape and size)	Reddish-brown, oblong, ca. 0.8 mm	Reddish-brown, oblong-ovoid, 0.5–0.6	Reddish-brown, oblong-ovoid, 0.4–0.5
Style length	1.6–2 mm	1–1.2 mm	1.1–1.3 mm
Primary flowering season	April to May	April to May	May to June

Taxonomic treatment

Sedum guangxiense Yan Liu & C.Y.Zou, sp. nov.

[urn:lsid:ipni.org:names:77356943-1](https://nomenclature.ipni.org/names/77356943-1)

Figs 2, 3

Type. CHINA • Guangxi: Fangcheng District, Na-suo Town, top of Nanshan. Elevation, 830 m; 21°44'49"N, 108°6'37"E, 24 April 2021. *H. L. Chen* ZCY1978 (holotype: IBK!, isotype: PE!).

Diagnosis. *Sedum guangxiense* is similar to *S. tosaense* and *S. emarginatum* in its leaf shape and inflorescence type, but can be distinguished from the latter two by its erect stems in youth (vs. prostrate when young), much larger leaves (1.5–3.5 cm long vs. 1–2 cm), narrowly triangular petals (vs. narrowly elliptic-lanceolate and linear-lanceolate to lanceolate), ovate nectar scales (vs. subquadrangular and oblong) and larger, oblong anthers (rather than oblong-ovoid).

Description. Biennial herb, fleshy, glabrous. Roots fibrous, with several adventitious roots in the leaf axils of the basal nodes. Stems stout, erect when young, with irregular branching, to 10–25 cm tall. Leaves alternate, base spurred, sessile, thick, spatulate to obovate, 1.5–3.5 cm long, 0.5–1 cm wide, apex usually retuse, rarely round, base long-attenuate. Inflorescences terminal, many flowered cymes, usually with three branches, each 2-forked; bracts leaf-like, 5–10 mm long, 2–5 mm wide; flowers usually sessile, usually 5-, sometimes 4- or 6-merous; Sepals usually 5, sometimes 4 or 6, free, green, fleshy, strongly unequal in size, spatulate to obovate, 3–8 mm long, 1–3 mm wide, apex usually retuse, rarely round. Petals usually 5, sometimes 4 or 6, yellow, narrowly




Figure 2. *Sedum guangxiense* **A** plant in flower **B** sterile plant **C** inflorescence from below **D** flowering branch **E** strongly unequal sepals **F** side view of a flower with carpels removed, showing the sepals, petals, stamens and nectar scales **G** adaxial leaf surface **H** unripe follicles **I** carpels connate at the base in longitudinal section **J** ovules **K** abaxial leaf surface. A, photographed by Hai-ling Chen in Fangcheng District (corresponding to the holotype *H. L. Chen* ZCY1978); B, C, photographed by Chun-Yu Zou in Yongfu County (corresponding to the paratype *C. Y. Zou & J. Q. Huang*, ZCY1977).




Figure 3. Holotype of *Sedum guangxiense*, H. L. Chen ZCY1978 (IBK).

triangular, 5–6 mm long, 1.1–1.5 mm wide, apex acuminate, slightly connate at base. Stamens usually 10, sometimes 8 or 12, shorter than petals, erect at anthesis, arranged in two whorls; antesealous ones 3.5–4 mm, antepetalous ones 3–3.5 mm; anthers oblong, ca. 0.8 mm long, reddish-brown before dehiscence. Nectar scales ovate, 0.3–0.4 mm. Carpels usually 5, sometimes 4 or 6, connate at the base for ca. 0.3–0.5 mm, gibbous ventrally, 5–6 mm long; styles 1.6–2 mm long. Fruits star-shaped, many seeded follicles, spreading. Flowering April to May, fruiting May to July.

Distribution and habitat. The species is known from Bai-shou Town in Yongfu County and Na-suo Town in Fangcheng District, Guangxi, China. It grows on mossy rocks in secondary broadleaf forests at elevations of 200–550 m (Fig. 4).

Etymology. The specific epithet refers to the distribution in Guangxi Province, China.

Additional specimen examined (paratypes). CHINA • Guangxi: Yongfu County, Bai-shou Town, on mossy rocks along streams, elev. 263 m, 25°11'43.29"N, 110°49'42.89"E, 10 May 2021, C. Y. Zou & J. Q. Huang, ZCY1977 (IBK!); • Fangcheng District, Na-suo Town, the top of Nanshan, 21°44'49"N, 108°6'37"E, 19 April 2021, Y. G. Liu, Q. G. Yang & H. L. Cheng 1208 (IBK!).




Figure 4. Distribution of *Sedum guangxiense* Yan Liu & C. Y. Zou (pentagram stars) in Guangxi, China. The red pentagram star indicates the locality of the holotype (type locality).

Acknowledgments

We are very thankful to Hai-ling Chen for her help in collecting herbarium specimens. We also thank Mr. Jin-quan Huang for helping us during the field survey.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by Fundamental Research Fund of Guangxi Institute of Botany (Grant no. GZY 21011) and the National Scientific Foundation of China (Grant no. 41661012).

Author contributions

All authors have contributed equally.

Author ORCIDs

Yan Liu  <https://orcid.org/0000-0001-8931-9656>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bolle C (1859) Addenda ad floram Atlantidis, praecipue insularum Canariensium Gorgadumque. *Bonplandia* 7: 238–246.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Edgar RC (2004) MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32(5): 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Fu S (1965) Species et combinationes novae Crassulaceae Sinicarum. *Acta Phytotaxonomica Sinica Additamentum* 1: 111–128.
- Fu K (1974) Revision of the section *Oreades* on Chinese *Sedum*. *Acta Phytotaxonomica Sinica* 12(1): 51–73. <https://www.jse.ac.cn/EN/Y1974/V12/I1/51>
- Fu KJ, Ohba H (2001) Crassulaceae. In: Wu ZY, Raven PH (Eds) *Flora of China*, vol. 8. Science Press, Beijing and Missouri Botanical Garden, St. Louis, 202–268.
- Hall T (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hart H (1995) Intrafamilial and generic classification of the Crassulaceae. In: 't Hart H, Egli U (Eds.) *Evolution and systematics of the Crassulaceae*. Backhuys Publishers, Leiden, 159–172.
- Huang YS, Guo J, Zhang Q, Lu ZC, Liu Y (2017) *Semiaquilegia guangxiensis* (Ranunculaceae), a new species from the limestone areas of Guangxi, China, based on morphological and molecular evidence. *Phytotaxa* 292(2): 180–188. <https://doi.org/10.11646/phytotaxa.292.2.5>
- Huang YS, Meng KK, Sun YY, Chen ZX, Fan Q (2023) A new species of *Sedum* (Crassulaceae) from Mount Danxia in Guangdong, China. *PhytoKeys* 221: 117–129. <https://doi.org/10.3897/phytokeys.221.97495>
- Ito T, Chen R, Yang Q, Saito Y, Yokota M, Kokubugata G (2014a) Taxonomic reexamination of *Sedum formosanum* (Crassulaceae) in Japan, Taiwan, and the Philippines based on molecular data. *Journal of Phytogeography and Taxonomy* 62: 1–9. <http://id.ndl.go.jp/bib/025974235>
- Ito T, Nakamura K, Park CH, Song GP, Maeda A, Watanabe Y, Kokubugata G (2014b) Nuclear and plastid DNA data confirm that *Sedum tosaense* (Crassulaceae) has a disjunct distribution between Pacific mainland Japan and Jeju Island, Korea. *Phytotaxa* 177(4): 221–230. <https://doi.org/10.11646/phytotaxa.177.4.3>
- Ito T, Nakanishi H, Chichibu Y, Minoda K, Kokubugata G (2017a) *Sedum danjoense* (Crassulaceae), a new species of succulent plants from the Danjo Islands in Japan. *Phytotaxa* 309(1): 23–34. <https://doi.org/10.11646/phytotaxa.309.1.2>
- Ito T, Yu CC, Nakamura K, Chung KF, Yang QE, Fu CX, Qi ZC, Kokubugata G (2017b) Unique parallel radiations of highmountainous species of the genus *Sedum* (Crassulaceae)

- on the continental island of Taiwan. *Molecular Phylogenetics and Evolution* 113: 9–22. <https://doi.org/10.1016/j.ympev.2017.03.028>
- Ito T, Yu CC, Kokubugata G (2018) Reconsiderations of distribution and taxonomic status of infraspecific taxa in *Sedum japonicum* (Crassulaceae) based on morphological and molecular data. *Bulletin of the National Museum of Nature and Science. Series B, Botany* 44(2): 73–83. <https://ndlsearch.ndl.go.jp/books/R000000004-I029071511>
- Ito T, Goto M, Nakano H, Minoda K, Kokubugata G (2020a) A new species of succulent plants from the Muko-jima group of the Bonin Islands, Japan: *Sedum mukojimense* (Crassulaceae). *Phytotaxa* 450(2): 188–198. <https://doi.org/10.11646/phytotaxa.450.2.4>
- Ito T, Yu CC, Yokota M, Kokubugata G (2020b) *Sedum formosanum* subsp. *miyakojimense* (Crassulaceae), a new subspecies from Miyako-jima Island of the Ryukyu Islands, Japan. *PhytoKeys* 148: 51–70. <https://doi.org/10.3897/phytokeys.148.48957>
- Kim Y, Kim SH, Yang J, Cho MS, Koldaeva M, Ito T, Kim SC (2023) Plastome-based backbone phylogeny of East Asian *Phedimus* (Subgenus *Aizoon*: Crassulaceae), with special emphasis on Korean endemics. *Frontiers in Plant Science* 14: 1089165. <https://doi.org/10.3389/fpls.2023.1089165>
- Linnaeus C (1753) *Species Plantarum* 1. Impensis Laurentii Salvii, Stockholm, 560 pp.
- Lu CT, Lin HW, Wang JC (2019) Two new species of *Sedum* (Crassulaceae) from Taiwan. *Taiwania* 64(4): 426–431. <https://doi.org/10.6165/tai.2019.64.426>
- Makino T (1892) Notes on Japanese plants, XV. *Botanical Magazine Tokyo* 6: 52. <https://doi.org/10.15281/jplantres1887.6.45>
- Mayuzumi S, Ohba H (2004) The phylogenetic position of Eastern Asian Sedoideae (Crassulaceae) inferred from chloroplast and nuclear DNA sequences. *Systematic Botany* 29(3): 587–598. <https://doi.org/10.1600/0363644041744329>
- Meng SY, Chen B, Sodmergen (2023) *Sedum danxiacola* (Crassulaceae), an endemic new species from the Danxia landform, China. *Kew Bulletin* 78(4): 673–681. <https://doi.org/10.1007/s12225-023-10122-y>
- Messerschmid TFE, Klein JT, Kadereit G, Kadereit JW (2020) Linnaeus's folly—phylogeny, evolution and classification of *Sedum* (Crassulaceae) and Crassulaceae subfamily Sempervivoideae. *Taxon* 69(5): 892–926. <https://doi.org/10.1002/tax.12316>
- Migo H (1937) Notes on the flora of south-eastern China. (Section III). *The Journal of the Shanghai Science Institute* 3: 221–228.
- Mort ME, Soltis DE, Soltis PS, Francisco-Ortega J, Santos-Guerra A (2002) Phylogenetics and evolution of the Macaronesian clade of Crassulaceae inferred from nuclear and chloroplast sequence data. *Systematic Botany* 27: 271–288.
- Posada D, Crandall KA (1998) Modeltest: Testing the model of DNA substitution. *Bioinformatics (Oxford, England)* 14(9): 817–818. <https://doi.org/10.1093/bioinformatics/14.9.817>
- Praeger RL (1929) *Semperviva* of the Canary Islands area. *Proceedings of the Royal Irish Academy* 38B: 454–499.
- Praeger LR (1932) An account of the *Sempervivum* group (Reprint, 2012). *Plant Monograph reprints*. J. Cramer & H. K. Swann, Stuttgart.
- Rao G (1996) Notes on some species of *Sedum* (Crassulaceae) in China. *Zhiwu Fenlei Xuebao* 34(6): 621–626. <https://www.jse.ac.cn/EN/Y1996/V34/I6/621>
- Ronquist F, Teslenko M, Mark PVD, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>

- Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology* 57(5): 758–771. <https://doi.org/10.1080/10635150802429642>
- Stephenson R (1994) *Sedum*. cultivated stonecrops. Timber Press, Portland, Oregon, 355 pp.
- Thiede J, Eggli U (2007) Crassulaceae. In: Kubitzki K (Ed.) *Flowering Plants Eudicots. The Families and Genera of Vascular Plants* (Vol. 9). Springer, Berlin/Heidelberg, 83–118. https://doi.org/10.1007/978-3-540-32219-1_12
- Wu LH, Liu YJ, Zhou SB, Guo FG, Bi D, Guo XH, Baker AJM, Smith JAC, Luo YM (2012) *Sedum plumbizincicola* X.H. Guo et S.B. Zhou ex L.H. Wu (Crassulaceae): A new species from Zhejiang Province, China. *Plant Systematics and Evolution* 299(3): 487–498. <https://doi.org/10.1007/s00606-012-0738-x>
- Xie DM, Peng DY, Fang CW, Qin MJ, Wang DQ, Huang LQ (2014) *Sedum spiralifolium* (Crassulaceae): A new species from Anhui Province, China. *Phytotaxa* 183(3): 171–182. <https://doi.org/10.11646/phytotaxa.183.3.4>
- Yang CD, Wang XY, Gou GQ (2012) *Sedum fanjingshanensis* C. D. Yang et X. Y. Wang, a new species of *Sedum* L. *Bulletin of Botanical Research* 32(4): 389–391.
- Ying SS (2022a) Four New species of genus *Sedum* (Crassulaceae) from Taiwan. *New Taxa and New Names* 4: 87–112.
- Ying SS (2022b) Two New species of genus *Sedum* (Crassulaceae) from Taiwan. *New Taxa and New Names* 5: 39–49.
- Zhang RB, Deng T, Dou QL, He L, Lv XY, Jiang H (2019) *Sedum lipingense* (Crassulaceae) identifying a new stonecrop species in SE Guizhou, China, based on morphological and molecular evidence. *PhytoKeys* 134(3): 125–133. <https://doi.org/10.3897/phytokeys.134.38287>
- Zou CY, Meng SY, Lu ZC, Liu Y (2020) *Sedum nanlingense* (Crassulaceae), a new species from Guangxi, China. *Phytotaxa* 447(3): 176–184. <https://doi.org/10.11646/phytotaxa.447.3.3>