



OPEN ACCESS

EDITED AND REVIEWED BY
S.Reddy Palli,
University of Kentucky, United States

*CORRESPONDENCE
Kamala Jayanthi Pagadala Damodharam
✉ kamalajayanthi.pd@icar.gov.in

†These authors have contributed
equally to this work and share
first authorship

RECEIVED 01 April 2026
REVISED 09 April 2026
ACCEPTED 11 May 2026
PUBLISHED 10 June 2026

CITATION
Palanivelu K, Balakrishnan U,
Damodharam KJP, Krishnasamy S,
Singh S and Dhayalan A (2026)
Correction: Morphometric and
molecular insights into *Bactrocera*
dorsalis (Hendel, 1912) (Diptera:
Tephritidae) infestation on *Ziziphus*
mauritiana Lamk. (Indian Jujube).
Front. Insect Sci. 6:1845367.
doi: 10.3389/finsc.2026.1845367

COPYRIGHT
© 2026 Palanivelu, Balakrishnan,
Damodharam, Krishnasamy, Singh and
Dhayalan. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/)
(CC BY). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication
in this journal is cited, in accordance
with accepted academic practice. No
use, distribution or reproduction is
permitted which does not comply with
these terms.

Correction: Morphometric and molecular insights into *Bactrocera dorsalis* (Hendel, 1912) (Diptera: Tephritidae) infestation on *Ziziphus* *mauritiana* Lamk. (Indian Jujube)

Kavin Palanivelu ^{1†}, Usharani Balakrishnan ^{2†},
Kamala Jayanthi Pagadala Damodharam ^{3*},
Suresh Krishnasamy ⁴, Sandeep Singh ⁵
and Arul Dhayalan ⁶

¹Department of Agricultural Entomology, Agricultural College and Research Institute, Tamil Nadu
Agricultural University (TNAU), Madurai, Tamil Nadu, India, ²Indian Council of Agricultural Research
(ICAR) - Krishi Vigyan Kendra, TNAU, Aruppukottai, Tamil Nadu, India, ³Indian Council of Agricultural
Research (ICAR) - National Professor, Division of Crop Protection, Indian Institute of Horticultural
Research, Bengaluru, India, ⁴Indian Council of Agricultural Research (ICAR) - Krishi Vigyan Kendra,
Madurai, Tamil Nadu, India, ⁵Indian Council of Agricultural Research (ICAR) - All India Coordinated
Research Project (AICRP) on Fruits, Department of Fruit Science, Punjab Agricultural University,
Ludhiana, Punjab, India, ⁶Division of Crop Protection, Indian Institute of Horticultural Research,
Bengaluru, India

KEYWORDS

Bactrocera dorsalis, ber, molecular characterization, morphometrics, phylogeny tree,
Principal Component Analysis (PCA)

A Correction on

Morphometric and molecular insights into *Bactrocera dorsalis* (Hendel,
1912) (Diptera: Tephritidae) infestation on *Ziziphus mauritiana* Lamk.
(Indian Jujube)

By Palanivelu K, Balakrishnan U, Damodharam KJP, Krishnasamy S, Singh S and Dhayalan A (2026)
Front. Insect Sci. 6:1716183. doi: 10.3389/finsc.2026.1716183

There was a mistake in [Figure 1c](#) as published. The figure and its legend were incorrect.
The original figure legend was:

“Figure 1 Morphological features of adult characters oriental fruit fly, *B. dorsalis*
(Tephritidae, Diptera). (a) Head and Ommatidia, the compound eyes are large and oval of
optical units of ommatidium. (b) Thorax with anepisternum and mediotergite (below the
scutellum) yellow with black spots are present on the margins of the thorax. (c) Wings with four
yellow parallel crossband, one band occurs in region of sub basal, second one is on discal region,
third origins on subapical and third and fourth bands extend toward apical margins.”

The corrected [Figure 1](#) and its caption appear below.

This correction does not change the scientific conclusions of the article in any way.

The original version of this article has been updated.




FIGURE 1
Morphological features of adult characters oriental fruit fly, *B. dorsalis* (Tephritidae, Diptera). **(a)** Head and ommatidia; the compound eyes are large and oval, with distinct optical units (ommatidia). **(b)** Thorax; the scutum is reddish brown to black, with yellow lateral vittae. **(c)** Wings largely hyaline, with a narrow, continuous costal band confluent with vein R2 + 3.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.