

## Full Length Research Paper

# Chemical compositions and antimicrobial activity of twig essential oils from three *Xylopi*a (Annonaceae) species

Siti Humeirah Abdul Ghani<sup>1\*</sup>, Nor Azah Mohamad Ali<sup>1</sup>, Mailina Jamil<sup>1</sup>, Mastura Mohtar<sup>2</sup>, Saiful Azmi Johari<sup>2</sup>, Mazurah Mohamad Isa<sup>2</sup> and Mohd Faridz Zoll Patah<sup>1</sup>

<sup>1</sup>Herbal Product Development Programme, Natural Products Division, Forest Research Institute Malaysia (FRIM), 52109 Kepong, Selangor Darul Ehsan, Malaysia.

<sup>2</sup>Bioactivity Programme, Natural Products Division, Forest Research Institute Malaysia (FRIM), 52109 Kepong, Selangor Darul Ehsan, Malaysia.

Received 4 June, 2015; Accepted 26 October, 2015

The chemical composition of twig essential oils of *Xylopi*a *malayana*, *Xylopi*a *elliptica* and *Xylopi*a *fusca* were analyzed using GC and GCMS. Both *X. malayana* (12.9%) and *X. fusca* (11.8%) twig oils contained spathulenol as the major constituent while *X. elliptica* twig oil was found to be rich in terpinen-4-ol (11.9%). The antimicrobial activities of the essential oils were evaluated against six gram positive bacteria, two gram negative bacteria and yeast by using the broth microdilution method. The twig oil of *X. elliptica* at a concentration ranging from 156 to 625 µg/ml was found to be active against all bacterial and yeast strains tested. Both *X. malayana* and *X. fusca* twig oils demonstrated weak activity toward all microorganisms tested with the MIC value of 5000 µg/ml. Gram negative bacteria's, *Pseudomonas aeruginosa* and *Escherichia coli* seemed to be resistant to the essential oils tested. α-Pinene showed moderate to strong inhibitory effect against all the microorganism strains tested with the MIC value ranging from 325 to 2500 µg/ml, as compared to the other standard compounds which showed weak inhibitory even at the highest concentrations used. This finding demonstrated that the twig oil of *X. elliptica* possesses antimicrobial activity which may be useful and potential ingredient in the production of health care products.

**Key words:** *Xylopi*a twigs, essential oils, GC & GCMS analysis, antimicrobial activity.

## INTRODUCTION

The family Annonaceae comprises about 120 genera and more than 2000 species (Leboeuf et al., 1982) and is distributed mainly in South America, Paleotropical Africa

and Southeast Asia. The plants of the family Annonaceae can be divided into three categories which are trees, shrubs, woody climbers and commonly found in the

\*Corresponding author. E-mail: [norazah@frim.gov.my](mailto:norazah@frim.gov.my). Tel: +603- 62797344 . Fax: +603-62729805.

lowland forest (Kochummen, 1997). The shape of the plant's leaf is simple and alternate. The bark is smooth, brown or dark green in colors and fragrant. The flowers are solitary or clusters and possessing a strong aromatic smell while the fruits are normally oblong in shape, stalked or sessile. Many members of Annonaceae family are used by local native in traditional medicine for the treatment of skin diseases, fever and stomachaches (Wiart, 2006). Phytochemical studies on the Annonaceae plants resulted in the isolation of several compounds for examples isoquinoline alkaline alkaloids (Mukhtar et al., 2000) and acetogenins (Alali et al., 1999) which have been found to possess antimicrobial, insecticidal, antimalarial and antioxidant properties (Iwu et al., 1999). The genus *Xylopi*a comprises about 170 species and they are widely distributed throughout Southeast Asia, Australia and Melanesia (Hyland and Whiffin, 1993). In Malaysia, eight species namely *Xylopi*a *malayana*, *Xylopi*a *caudata*, *Xylopi*a *ferrugineae*, *Xylopi*a *magna*, *Xylopi*a *fusca*, *Xylopi*a *elliptica*, *Xylopi*a *stenopetala* and *Xylopi*a *subdehiscens* have been documented by Kochummen (1972). The plants are usually shrubs or small trees in nature and commonly found in lowland, peat swamp and mountain forests forest. The secondary metabolites from *Xylopi*a species have been known for their great usage in folk medicine, food flavoring and medical industries (Leboeuf et al., 1982). Besides, the essential oils of some *Xylopi*a sp. are also used in traditional medicine as a remedy in the treatment of stomach-ache and intestines. *Xylopi*a species are sources of volatile components, diterpenes and alkaloid, which have been reported to possess antimicrobial, anti-inflamantory and antimalarial properties. Volatile constituents and bioactivity studies are available in the literature on *Xylopi*a *aethiopica* (Issakou et al., 2014; Sylvain et al, 2014; Vvry et al, 2014), *Xylopi*a *longifolia* (Fourier et al, 1993), *Xylopi*a *parviflora* (Woguem et al, 2014; Samuel et al., 2015), *Xylopi*a *aromatica* (Lago et al, 2003) and *Xylopi*a *langsдорфiana* (Correia et al., 2015).

*X. malayana* or locally known as Jangkang is a small tree up to 20 m high and occurs in the northern part of East Malaysia. The morphology of leaves are elliptic-oblong whilst flowers are normally small and aromatic. In Peninsular Malaysia, the leaves of *X. malayana* are traditionally be used for treatment after childbirth (Kamarudin, 1988). The chemical constituent and antimicrobial activity of essential oils from the leaf of *X. malayana* have been previously reported by Nor Azah et al. (1996). *X. elliptica* or lilan is a medium sized tree which could grow up to 8 to 10 m tall and is one of the endemic plants. The leaves are membranous and blade elliptic while the flowers are axillary with oblong fruits. *X. fusca* is widely distributed in Southeast Asia. It is a medium to large trees with 125 m in height and commonly found in peat swamp forest. *X. fusca* is known as jangkang paya. In traditional medicine practice, the plant used as an abortifacient and as an antidote for

insect-bites. The wood of *X. fusca* is used in making a pineapple box whilst the barks of this plant are also used for walls of huts (Whitmore, 1972). The chemical composition of the leaf essential oils from *X. malayana*, *X. fusca* and *X. elliptica* have been examined by Siti Humeirah et al. (2010) but no information is available on the twig oil of these plants. Therefore, in this study, the chemical compositions of essential oils of twig part from *X. malayana*, *X. elliptica* and *X. fusca* as well as their antimicrobial properties were explored.

## MATERIALS AND METHODS

### Plant

The twig parts of *X. malayana* (FRI 54729), *X. elliptica* (FRI 54736) and *X. fusca* (FRI 54733) were randomly collected from Pasoh Forest Reserve, Negeri Sembilan, Malaysia on November 2007. All the plant materials were identified by Mr. Kamarudin Salleh from Forest Research Institute Malaysia (FRIM) and the voucher specimens were deposited at the FRIM herbarium

### Extraction of essential oils

The plant material were cut into small pieces, weighed (200 to 400 g) and hydrodistilled separately in clavenger type apparatus for 6 h. The oily layer obtained from each plant material were separated, dried over anhydrous sodium sulfate and stored at 4°C until used. The yield (%) of essential oil was also determined by Azeotropic methods (British Pharmacopoeia, 2004).

### GC and GCMS analysis

The chemical constituents of essential oils were analysed by GC and GCMS. GC analysis was carried out using Shimadzu GC2010 (FID) chromatography with a CBP-5 column (25 mm × 0.25 µm × 0.25 mm film thickness). GCMS analysis was performed using a Agilent GCMS 7890A/5975C series MS with a HP-5MS column (30 mm × 0.25 µm × 0.25 mm film thickness). The components of essential oils were identified by comparison of their retention indices with literature values (Adams, 2001) and confirmed by comparison of their mass spectral data with those from the Wiley (1990), HPCH 2205.L and NIST05a.L libraries. The Kovats indices (Kovats, 1965) of each component were determined relative to the retention time of a series of n alkanes (C<sub>8</sub>-C<sub>20</sub>). The relative proportions of the chemical compounds were expressed as percentages obtained by peak-area normalization, all relative response factors being taken as one.

### Antimicrobial assay

The twig oils of *X. malayana*, *X. elliptica* and *X. fusca* were screened for their antimicrobial activity against nine strains of microorganisms including six gram positive bacteria *Staphylococcus aureus* ATCC 25923 (MSSA); *S. aureus* ATCC 33591 (MRSA); *S. aureus* ATCC 70069 (VISA); *Vancomycin Resistant S. aureus* (VRSA156), *Vancomycin Intermediate S. aureus* (VISA24) and *S. epidermidis* ATCC 12228. The gram negative bacteria including *Pseudomonas aeruginosa* ATCC 25668, *Escherichia coli* ATCC 10536 and yeast *Candida albicans* ATCC 10231 using the Minimum Inhibitory Concentration (MIC)

method as described previously by Saiful et al. (2008). The gram positive and negative bacteria and yeast were purchased from American Type Culture Collection (ATCC) while VRSA156 and VISA24 were lab passage derived mutants from clinical MRSA isolate.

#### Minimum inhibitory concentration (MIC)

The minimum inhibitory concentration (MIC) values of twig oils were determined using the double-broth microdilution method. Briefly, the well A was loaded with essential oils stock solution (100 mg/ml) and topped up with respectively broth. The mixture in well A was mixed thoroughly and transferring to well B to H which are already contained with respectively broth. The same procedure (well A) was repeated for well B to H to create a serial doubling dilution of the oils with concentration ranging from 10 to 0.078 mg/ml. Finally, a suspension containing  $10^8$  colony-forming units (CFU)/ml which are equivalent to McFarland standard tube no. 0.5 used (Vandepitte et al., 1991) were inoculated into the mixture to produce a final volume of 100  $\mu$ l. Oxacillin and cyclohexamide were used as positive control whiles DMSO-broth mixture (5%) was used as a negative control. The experiments were performed in triplicates and repeated twice. The lowest concentration which produces no visible growth or remains clear after observed through the macroscopic evaluation was taken as the MIC. The activity was categorized as strong (MIC  $\leq$  1000  $\mu$ g/ml), moderate (999  $\mu$ g/ml < MIC < 4900  $\mu$ g/ml) and weak (MIC  $\geq$  5000  $\mu$ g/ml) (Ibrahim et al., 2009).

## RESULTS AND DISCUSSION

### Essential oil compositions

The twig parts of *X. malayana* (354.3 g), *X. fusca* (252.2 g) and *X. elliptica* (400.0 g) were extracted by hydrodistillation and yielded 0.08, 0.08 and 0.05% (v/w) of oils respectively. Both twig oils from *X. elliptica* and *X. malayana* were yellowish in color while the twig oil from *X. fusca* was colorless. The yield percentages of essential oils from fresh plant materials of *X. malayana* varied greatly from the previous study by Nor Azah et al. (1996). The twigs of *X. malayana* on the earlier study yielded 0.40% of essential oils. According to Yang et al. (2002), the variation factor of essential oil contents could be due to extraction time and the ratio of water and twig used in the study. The chemical composition of the twig oils are shown in Table 1. GC/MS analysis resulted in the identification of 29 (61.2% of the total oil), 17 (45.3% of the total oil) and 33 (79.7% of the total oil) components from the twig oils of *X. malayana*, *X. fusca* and *X. elliptica*, respectively. Components were mainly made up of monoterpenes hydrocarbon, oxygenated monoterpenes, sesquiterpenes hydrocarbon and oxygenated sesquiterpenes. The major constituents of the twig oil from *X. malayana* were spathulenol (12.9%), cyperotundone (4.9%), elemol (8.3%) and  $\alpha$ -eudesmol (5.9%). Other components were  $\gamma$ -gurjunene (2.0%),  $\alpha$ -terpineol (1.9%),  $\beta$ -pinene (1.5%) and  $\beta$ -phellandrene (0.9%). Spathulenol (36.0%) are identified as major component in the twig oil of *X. malayana*, previously

mentioned by Nor Azah et al. (1996). However, in the present study, spathulenol was constituted only 12.9% of the total *X. malayana* twig oil. Differences in the chemical composition of *X. malayana* twig oils could be attributed to different collecting time, climate effect on the plants which are growing from the different habitat, method of extraction used and geographic origin of plant studied (Cimanga et al., 2002). On the other hand, *X. elliptica* twig oil was found to be rich in terpinen-4-ol (11.9%), *p*-cymene (7.4%), spathulenol (6.3%),  $\alpha$ -cadinol (6.8%), guaialol (4.5%) and linalool (4.2%). The twig oil was also made up of  $\alpha$ -pinene (3.2%), 1,8-cineole (2.7%),  $\alpha$ -terpineol (1.8%), sabinene (2.5%) and  $\beta$ -elemene (0.8%). The twig oil of *X. fusca* was mainly made up of spathulenol (11.8%) and  $\alpha$ -cadinol (8.2%) as the major components. Other components present in appreciable amount were 1-epi cubenol (3.6%), vulgarone B (4.8%) and caryophyllene oxide (2.0%).

Several studies on the chemical composition of the essential oils of other *Xylopi*a species have shown that monoterpenes and sesquiterpenes are the major constituents of essential oils. Spathulenol, which was detected in the twig oils of the *X. elliptica*, *X. fusca* and *X. malayana* was identified as the major component in the *X. aromatica* (Founier et al., 1994) and *Xylopi*a *emarginata* (Lago et al., 2003) essential oils. Both monoterpenoid, *p*-cymene and  $\alpha$ -pinene were detected as—major constituents in the essential oil of *Xylopi*a *sericea* (Lemos et al., 1992). Other chemical constituents for example terpinen-4-ol,  $\beta$ -pinene,  $\delta$ -cadinene,  $\delta$ -cymene and  $\alpha$ -copaene were also found in other part of *Xylopi*a species Nor Azah et al. (1996); Lemos et al. (1992). However, the essential oils obtained from *X. fusca* contained certain compounds which are not observed in the other oils namely; isopulegyl acetate, gymnomitrol, 14-hydroxy- $\delta$ -cadinene and 14-hydroxy- $\alpha$ -muurolene which could be considered unique to the species. The presence of some terpene type compounds in appreciable amounts supported the use of these compounds as chemical marker in the species identification purposes. Phytochemical analysis of the oils indicated that the chemical constituents represent a valuable chemotaxonomic characterization for supporting the identification of the genus *Xylopi*a (Siti Humeirah et al., 2010).

### Antimicrobial activity

The antimicrobial assays of the twig oils of *X. malayana*, *X. fusca* and *X. elliptica*, together with eight reference standards which are  $\alpha$ -pinene, geranyl acetate, linalool, limonene, geraniol, terpinen-4-ol, 1,8-cineole and  $\beta$ -pinene were carried out by broth microdilution method against nine different strains of microorganisms. According to the results in Table 2, the twig oil from *X. elliptica* have been shown to possess the strongest

**Table 1.** Chemical compositions (%) of the twig essential oil from *X. elliptica*, *X. fusca* and *X. malayana*..

| Compound                         | Retention index (RI) | Twig oils concentration (%) |                 |                    | Method identification |
|----------------------------------|----------------------|-----------------------------|-----------------|--------------------|-----------------------|
|                                  |                      | <i>X. elliptica</i>         | <i>X. fusca</i> | <i>X. malayana</i> |                       |
| $\alpha$ -pinene                 | 935                  | 3.2                         | -               | 0.4                | RI,MS, CO             |
| sabinene                         | 977                  | 2.5                         | -               | -                  | RI,MS                 |
| $\beta$ -pinene                  | 980                  | 3.2                         | -               | 1.5                | RI,MS,CO              |
| myrcene                          | 989                  | 0.5                         | -               | -                  | RI,MS,CO              |
| p-cymene                         | 1023                 | 7.4                         | -               | 0.2                | RI,MS                 |
| $\beta$ -phellandrene            | 1029                 | 1.6                         | -               | 0.9                | RI,MS                 |
| 1,8-cineole                      | 1037                 | 2.7                         | -               | -                  | RI,MS,CO              |
| (Z)- $\beta$ -ocimene            | 1038                 | 0.7                         | -               | -                  | RI,MS                 |
| $\gamma$ -terpinene              | 1059                 | -                           | -               | 0.3                | RI,MS                 |
| terpinolene                      | 1085                 | -                           | -               | 0.2                | RI,MS                 |
| linalool                         | 1099                 | 4.2                         | -               | 0.8                | RI,MS,CO              |
| cis-p-menth-2-en-1-ol            | 1121                 | 0.4                         | -               | -                  | RI,MS                 |
| trans-p-menth-2-en-1-ol          | 1141                 | 0.6                         | -               | -                  | RI,MS                 |
| trans-pinocarveol                | 1144                 | -                           | -               | 0.8                | RI,MS                 |
| pinocarvone                      | 1163                 | -                           | -               | 0.5                | RI,MS                 |
| para-metha-1,5-diene-8-ol        | 1175                 | -                           | -               | 0.3                | RI,MS                 |
| terpinen-4-ol                    | 1177                 | 11.9                        | 0.1             | 1.4                | RI,MS,CO              |
| $\alpha$ -terpineol              | 1192                 | 1.8                         | -               | 1.9                | RI,MS                 |
| cis-piperitol                    | 1194                 | 0.3                         | -               | -                  | RI,MS                 |
| myrtenal                         | 1201                 | -                           | -               | 0.8                | RI,MS                 |
| verbenone                        | 1210                 | -                           | 0.1             | 0.1                | RI,MS                 |
| trans-carveol                    | 1232                 | -                           | 0.1             | 0.1                | RI,MS                 |
| 5-hydroxy-isobornyl isobutanoate | 1252                 | 0.3                         | -               | -                  | RI,MS                 |
| bornyl acetate                   | 1280                 | 0.4                         | -               | 1                  | RI,MS,CO              |
| $\delta$ -elemene $\epsilon$     | 1333                 | 0.3                         | -               | -                  | RI,MS                 |
| $\alpha$ -terpinyl acetate       | 1344                 | 0.8                         | -               | -                  | RI,MS                 |
| $\alpha$ -copaene                | 1367                 | 0.7                         | 0.1             | 0.3                | RI,MS                 |
| $\beta$ -elemene                 | 1386                 | 0.8                         | 0.1             | 0.5                | RI,MS                 |
| cyperene                         | 1396                 | 1.2                         | -               | -                  | RI,MS                 |
| $\beta$ -caryophyllene           | 1415                 | -                           | 0.1             | 0.2                | RI,MS                 |
| $\delta$ -copaene                | 1435                 | 0.3                         | -               | -                  | RI,MS                 |
| $\gamma$ -cadinene               | 1494                 | 0.4                         | -               | -                  | RI,MS                 |
| bicyclogermacrene                | 1498                 | -                           | -               | 1.2                | RI,MS                 |
| 4-epi-cis-dihydroagarofuran      | 1502                 | -                           | -               | 0.9                | RI,MS                 |
| $\alpha$ -murolene               | 1512                 | -                           | 2               | -                  | RI,MS                 |

**Table 1.** Contd.

|                                 |      |     |      |      |       |
|---------------------------------|------|-----|------|------|-------|
| $\delta$ -cadinene              | 1518 | 1.1 | 2.1  | 0.9  | RI,MS |
| maaliol                         | 1555 | 0.3 | -    | -    | RI,MS |
| elemol                          | 1561 | 2   | -    | 8.3  | RI,MS |
| spathulenol                     | 1584 | 6.3 | 11.8 | 12.9 | RI,MS |
| viridiflorol                    | 1590 | 4.3 | -    | -    | RI,MS |
| caryophyllene oxide             | 1597 | -   | 2    | -    | RI,MS |
| guaial                          | 1601 | 4.5 | -    | -    | RI,MS |
| $\gamma$ -gurjunene             | 1607 | -   | -    | 2    | RI,MS |
| 14-hydroxy- $\alpha$ -muurolene | 1621 | -   | 3.1  | -    | RI,MS |
| 1-epi-cubenol                   | 1634 | -   | 3.6  | -    | RI,MS |
| cyperotundone                   | 1638 | 1   | -    | 4.9  | RI,MS |
| $\beta$ -cedrene                | 1644 | 6.5 | -    | -    | RI,MS |
| $\alpha$ -cadinol               | 1661 | 6.8 | 8.2  | -    | RI,MS |
| $\alpha$ -eudesmol              | 1663 | -   | -    | 5.9  | RI,MS |
| bulnesol                        | 1671 | 0.7 | -    | -    | RI,MS |
| isopulegyl acetate              | 1674 | -   | 3.1  | -    | RI,MS |
| vulgarone B                     | 1692 | -   | 4.8  | 2.6  | RI,MS |
| gymnomitrol                     | 1855 | -   | 2.2  | -    | RI,MS |
| 14-hydroxy- $\delta$ -cadinene  | 1875 | -   | 1.8  | -    | RI,MS |

Percentages were calculated based on results obtained from gas chromatography on CBP-5 column; RI = retention indices; MS= mass fragmentation; CO= co-chromatography with authentic sample.

antibacterial and antifungal activity against all gram positive bacteria and yeast strains tested with the MIC values ranging from 156  $\mu\text{g/ml}$  to 625  $\mu\text{g/ml}$ . Gram negative bacteria's, *P. aeruginosa* and *E.coli* seemed to be resistant to the essential oils tested. Both *X. fusca* and *X. malayana* twig oils however showed less inhibitory effect on the growth of bacteria and yeast strains even at the highest doses (5000  $\mu\text{g/ml}$ ) used. The results indicated the gram positive bacteria are more sensitive to plant oil than gram negative bacteria due to their restrictive outer membrane barrier which are generally less susceptible and destabilization. The inhibitory effects of *X. elliptica*

twig oil on bacterial strains tested could be related to the present of  $\alpha$ -pinene (Magwa et al., 2006), terpinen-4-ol (Barel et al., 1991) and  $p$ -cymene which have been reported their high level of antimicrobial activity. In addition, it is also possible that the other minor or trace compounds might be involved in some types of antimicrobial synergism with other active components of essential oil. Various studies have reported antimicrobial activities for essential oils from *Xylopi*a species. Lemos et al. (1992) reported the essential oil of *X. sericea* which contains 1,8-cineole as the major constituents showed effective antifungal activity against *Candida* strains. The essential oils of

*Xylopi*a *longifolia* from France were also found to possess antibacterial and antifungal activity (Fourier et al., 1993). The antimicrobial activities of essential oil from *Xylopi*a *parviflora* and *Xylopi*a *aethiopica* from Africa have been reported by Bakarnga et al. (2014) and Woguem et al. (2014).

Beside essential oil, the antimicrobial properties of selected standard compounds were also conducted in the present study. Based on the results summarized in Table 2, the antibacterial and antifungal activity of the five reference standards indicated that  $\alpha$ -pinene had the moderate to strong inhibitory effect against all

**Table 2.** Minimum Inhibitory Concentration ( $\mu\text{g/ml}$ ) of twig oils from three *Xylopi*a species and standard compounds.

| Species/ Standards        | Parts | sa2   | sa3   | sa7   | VISA  | VRSA  | Se    | Pa    | Ec    | Ca    |
|---------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <i>X. elliptica</i>       | twig  | 625   | 625   | 625   | 312   | 312   | 156   | >5000 | >5000 | 312   |
| <i>X. fusca</i>           | twig  | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 |
| <i>X. malayana</i>        | twig  | 5000  | 5000  | 5000  | 5000  | 5000  | >5000 | >5000 | >5000 | >5000 |
| <b>Standard compounds</b> |       |       |       |       |       |       |       |       |       |       |
| $\alpha$ -pinene          |       | 625   | 2500  | 1250  | 1250  | 1250  | 625   | 2500  | 325   | 325   |
| $\beta$ -pinene           |       | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 |
| linalool                  |       | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 2500  | 2500  |
| limonene                  |       | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  |
| Geranyl acetate           |       | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  | 5000  |
| 1,8-cineole               |       | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 | >5000 |
| Terpinen-4-ol             |       | 5000  | 5000  | 5000  | 5000  | 5000  | 2500  | 2500  | 625   | 2500  |
| Geraniol                  |       | 5000  | 5000  | 5000  | 2500  | 5000  | 1250  | 2500  | 1250  | 2500  |
| <b>Positive control</b>   |       |       |       |       |       |       |       |       |       |       |
| Streptomycin sulfate      |       | NT    | NT    | NT    | NT    | NT    | 156   | 2500  | 2500  | NT    |
| Oxacillin                 |       | 312   | 312   | 625   | 2500  | 1250  | NT    | NT    | NT    | NT    |
| Cycloheximide             |       | NT    | NT    | NT    | NT    | NT    | NT    | NT    | NT    | 1250  |

Sa2, *Staphylococcus aureus* ATCC 25923 (MSSA); Sa3, *S. aureus* ATCC 33591 (MRSA/VSSA); Sa7, *S. aureus* ATCC 70069 (MRSA); VRSA156, Vancomycin Resistant *S. aureus*; VISA24, Vancomycin Intermediate *S. aureus*; Se, *S. epidermidis* ATCC 12228; Pse, *Pseudomonas aeruginosa* ATCC 25668; Ec, *Escherichia coli* ATCC 10536; Ca, *Candida albicans* ATCC 10231; NT, Not Test.

bacterial and yeast growth at the concentration of 325 to 2500  $\mu\text{g/ml}$ . *Xylopi*a *epidermidis*, *E. coli* and *C. albicans* were most susceptible to the compound investigated and hence support the earlier finding on  $\alpha$ -pinene's significant antimicrobial potential (Dorman and Deans, 2000). The reference standards of  $\beta$ -pinene and 1,8-cineole however inhibited weak (MIC values of 5000  $\mu\text{g/ml}$ ) antibacterial and antifungal activity against all microorganisms tested. Terpinen-4-ol, the most abundant component in the twig oil of *X. elliptica*, has been shown to have a moderate to strong antibacterial activity against *P. aeruginosa* and *E. coli* strains with the MIC values of 2500 and 625  $\mu\text{g/ml}$ . Terpinen-4-ol however showed weak inhibitory effect against MRSA strains tested. The results however was not in agreement with a previous study reported by Barel et al. (1991) which found the bacteriostatic effects of tested compound, terpinen-4-ol against several microorganisms. Linalool was found to be less effective to all microorganisms tested with the MIC values ranging from 1250 to 5000  $\mu\text{g/ml}$ .

## Conclusion

The characteristics of the essential oils from three *Xylopi*a species are represent a valuable chemotaxonomic tool for the identification of the genus *Xylopi*a. Considering the relatively high yield of monoterpenoids and bioactive component in the oils,

*Xylopi*a could be a potential new source of fragrance ingredient for food, pharmaceutical and cosmetic industries.

## Conflict of interests

The author(s) did not declare any conflict of interest.

## ACKNOWLEDGEMENTS

The authors appreciate the effort of Mr. Kamarudin Salleh from Herbarium unit, FRIM in the identification of the plant sample and highly grateful to all the researchers from Natural Products Division for their technical assistance and encouragements during the work. This research was supported by the Ministry of Science, Technology and Innovation (MOSTI) fund no: 02-03-10SF0047.

## REFERENCES

- Adams RP (2001). Identification of essential oils components by GC/ quadrupole mass spectrometry. Allured Publishing Corporation: Carol Stream, IL.
- Alali FQ, Rogers L, Zhang Y, McLaughlin JL (1999). Goniatriocin and (2,4-*cis*- and - *trans*-) xylomaticinones, bioactive annonaceous acetogenins from *Goniathalamus giganteus*. J. Nat. Prod. 62:31-34.
- Barel S, Segal R, Yashphe J (1991). The antimicrobial activity of the

- essential oil from *Achillea fragrantissima*. J. Ethnopharmacol. 33:187-191.
- British Pharmacopeia (2004). Edition I and II. London, HMSO.
- Cimanga K, Kambu K, Tona L, Apers S, De Bruyne T, Hermans N, Totté J, Pieters L, Vlietinck AJ (2002). Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo. J. Ethnopharmacol. 2(79): 213-220.
- Correia AC, Ferreira TF, Martins IR, Macedo CL, Monteiro FS, Costa VC, Tavares JF, Silva MS, Paredes-Gamero EJ, Buri MV, Rigoni VL, Nouailhetas VL, Silva BA (2015). Essential oil from the leaves of *Xylopia langsdorffiana* (Annonaceae) as a possible spasmolytic agent. Nat. Prod. Res. 29(10):980-984.
- Dorman HJ, Deans SG (2000). Antimicrobial agents from plants: antibacterial activity of plant volatile oils. J. Appl. Microbiol. 88:308-316.
- Fourier G, Hadjiakoondi A, Lebouef M, Cave A, Fourniat J and Charles B (1993). Chemical and biological studies of *Xylopia longifolia* A.D.C. essential oils. J. Essent. Oil Res. 5:403-410.
- Ibrahim H, Syamsir DR, Aziz AN, Awang K, Nor Azah MA, Mastura M, Rasadah MA (2009). Essential oils of *Elettariopsis curtisii* (Zingiberaceae) and their antimicrobial activities. J. Essent. Oil Res. 21:464-466.
- Hyland BPM, Whiffin T (1993). *Australian Tropical Rain Forest Trees*. An Interactive identification system. Volume 2. CSIRO Australia, Melbourne.
- Issakou BV, Jean BH, Patrick VT, Lauve RY, Magali GB, Audrey G, Marcel G, Lucain W, Youssouf S, Pierre MJ, Fabrice FB and Chantal M (2014). Composition and cytotoxic activity of essential oils from *Xylopia aethiopica* (Dunal) A. Rich., *Xylopia parviflora* (A. Rich.) Benth.) and *Monodora myristica* (Gaertn) growing in Chad and Cameroon. BMC Complement. Altern. Med. 14:125
- Iwu M, Duncan AR, Okunji CO (1999). *New antimicrobials of plants origin*. In: Janick, J. (Ed.). Perspectives on new crops and new uses. ASHS Press, Alexandria, VA. pp. 457-462.
- Kamarudin MS (1988). Ethnobotanical Importance of the Asiatic Annonaceae. In: Soepadmo E, SH Goh, WH Wong, Laily, BD & CH Chuah (eds.). Proceedings of the Malaysian Traditional Medicine Seminar, Kuala Lumpur. pp. 80-87.
- Kochummen KM (1972). Family: Annonaceae. Pp. 61-99 in Whitmore, T.C. (Ed.) *Tree Flora of Malaya*. Vol. 1. Malayan Forest Records. No.26. Forest Department, Kuala Lumpur.
- Kochummen KM (1997). Tree flora of Pasoh Forest. Forest Research Institute Malaysia, Kuala Lumpur.
- Kovats E (1965). Gas chromatographic characterization of organic substances in the retention index system. Adv. Chromatogr.1:229-247.
- Lago JH, Moreira IC, Tanizaki TM, Moreno PR, Roque NF (2003). Mono and sesquiterpenes from the leaf essential oil of *Xylopia brasiliensis* Spreng. (Annonaceae). J. Essent. Oil Res. 15:406-407.
- Leboeuf M, Cave A, Bhaumik PK, Mukherjee B, Mukherjee R (1982). The phytochemistry of the Annonacea. Phytochemistry 21(12):2783-2813.
- Lemos TG, Monte FQ, Matos FA, Alencar JW, Craveiro A, Barbosa RB, Lima EO (1992). Chemical composition and antimicrobial activity of essential oils from Brazilian plants. Fitoterapia 3:266-268.
- Magwa ML, Gundidza M, Gweru N, Humphrey G (2006). Chemical composition and biological activities of essential oil from the leaves of *Sesuvium portulacastrum*. J. Ethnopharmacol. 103:85-89.
- Mukhtar MR, Awang K, Mustafa MR, Norhanom AW, Ashril Y and Hadi AH (2000). Chemical constituents and bioactive compounds of *Goniothalamus tortilipetalus* Hend. (Annonaceae). Malay. J. Sci. 19:7-12.
- Nor Azah MA (1996). Kajian terhadap minyak pati daripada Genus *Xylopia*. Master thesis. Universiti Kebangsaan Malaysia.
- Saiful AJ, Mastura M, Zarizal S, Shuhaimi M, Ali AM (2008). Efflux genes and active efflux activity detection in Malaysian clinical isolates of Methicillin-Resistant *Staphylococcus aureus* (MRSA). J. Basic Microbiol. 48:245-251.
- Samuel AB, Olufemi OR, Ganiyu O, Michael OA (2015). *First report of toxicity of Xylopia parviflora* (A. Rich.) Benth (Annonaceae) root bark's essential oil against cowpea seed bruchid, *Callosobruchus maculatus* Fabricius (Coleoptera: Chrysomelidae: Bruchinae). Nat. Prod. Res. 29(4): 349-352.
- Siti Humeirah AG, Nor Azah MA, Mailina J, Muhajir H, Puad A (2010). Chemical composition of three *Xylopia* leaf essential oils from Pasoh Forest Reserve, Negeri Sembilan, Malaysia. J. Trop. For. Sci. 22(1):1-4
- Sylvain LS, Nicoletta B, Francois T, Jean JE, Chiara MG, Rosalba L and Fausto G (2014). Effect of mild heat treatments on the antimicrobial activity of essential oils of *Curcuma longa*, *Xylopia aethiopica*, *Zanthoxylum xanthoxyloides* and *Zanthoxylum lepreurii* against *Salmonella enteritidis*. J. Essent. Oil Res. 27(1): 52-60.
- Vandepitte J, Andepitte J, Engback K, Piot P, Heuck CC (1991). Basic microbiology procedure in clinical bacteriology. World Health Organization, Geneva. 88.
- Vyry WN, Misra L and Venkatesh KR (2014). Antibacterial Activity of Essential Oils of Edible spices, *Ocimum canum* and *Xylopia aethiopica*. J. Food Sci. 79(5): 972-977.
- Whitmore TC (1972). *Tree Flora of Malaysia*. A manual for foresters. Vol 1. Forest Department, Ministry of Agriculture and Land Malaysia.
- Wiat C (2006). Medicinal plants of the Asia Pacific: drugs for the future? World Scientific Publishing Co. Pte. Ltd.
- Woguem V, Fogang HP, Maggi F, Tapondjou LA, Womeni HM, Quassinti L, Bramucci M, Vitali LA, Petrelli D, Lupidi G, Papa F, Vittori S, Barboni L (2014). Volatile oil from striped African pepper (*Xylopia parviflora*, Annonaceae) possesses notable chemopreventive, anti-inflammatory and antimicrobial potential. Food Chem. 149: 183-189.