



# Cassava by-products in broiler diets: effects on growth performance based on a systematic review

Wanessa Daróz Matte<sup>1</sup> · Cynthia Pieri Zeferino<sup>1</sup> · Henrique Melo da Silva<sup>2</sup> · Jean Kaique Valentim<sup>3</sup> · Melody Martins Cavalcante Pereira<sup>4</sup> · Maria Eduarda Vignoli Nabhan<sup>4</sup> · Livia Costa Silva<sup>4</sup> · Sarah Sgavioli<sup>4</sup>

Received: 10 May 2026 / Accepted: 31 July 2026  
© The Author(s) 2026

## Abstract

The use of cassava (*Manihot esculenta* Crantz) by-products as alternative energy sources in broiler diets has increased, but their effects on growth performance remain inconsistent across studies. This systematic review evaluated the effects of replacing corn with cassava by-products (cassava meal, pulp, and bagasse) on broiler performance using a qualitative direction-of-effect synthesis. A total of 36 studies published between 2005 and 2025 were included, comprising 187 dietary comparisons. Most studies were conducted in Brazil (36.1%), Nigeria (25.0%), and Indonesia (18.2%). Because measures of variability were frequently unavailable, responses in feed intake (FI), body weight gain (BWG), and feed conversion ratio (FCR) were classified as positive, negative, or neutral relative to the reference treatment. Feed intake was predominantly unaffected (54.4% neutral responses). For BWG, negative responses were most frequent (49.2%), followed by neutral (44.4%) and positive responses (6.4%). For FCR, neutral (45.5%) and negative (44.4%) responses occurred at similar frequencies, whereas positive responses accounted for 10.2% of comparisons. Negative responses became more frequent as dietary inclusion levels increased, particularly for BWG and FCR, and were most evident at inclusion levels above 25%. Most studies (86.1%) were classified as having a high risk of bias or some concerns, and measures of variability were frequently not reported, limiting quantitative comparison of effect estimates and the strength of inference. Overall, cassava by-products were more frequently associated with reduced growth performance at higher dietary inclusion levels, although responses varied according to by-product type, processing method, inclusion level, and study conditions. These findings support cautious dietary formulation and highlight the need for standardized experimental reporting to better define safe and effective inclusion levels for individual cassava by-products in broiler nutrition.

**Keywords** Alternative feed ingredients · Broiler chickens · Cassava by-products · Corn replacement · Feed efficiency · Growth performance · Systematic review

## Introduction

In traditional poultry production systems, corn and soybean meal account for approximately 90% of the nutritional composition of diets. However, in least-cost feed formulations, alternative ingredients may be incorporated to optimize production costs depending on their local availability (Uchegbu et al. 2011). Among these alternatives, cassava (*Manihot esculenta* Crantz) is particularly viable in tropical countries because of its high starch content (Latif and Müller, 2015) and broad availability. Brazil produced 20.57 million tons of cassava harvested from 1.3 million hectares as of September 2025 (IBGE, 2025).

Starch-processing industries generate large quantities of by-products, such as bagasse and bran, during cassava

---

✉ Jean Kaique Valentim  
jean.valentim@ufrj.br

<sup>1</sup> Postgraduate Program in Animal Production, Universidade Brasil, Descalvado, São Paulo 13690-000, Brazil

<sup>2</sup> Dairy Cattle Research Lab, Universidade Federal de Mato Grosso, Sinop, Mato Grosso 78557-267, Brazil

<sup>3</sup> Department of Animal Production, Universidade Federal Rural do Rio de Janeiro, Seropédica, Rio de Janeiro 23890-000, Brazil

<sup>4</sup> Graduate Program in Animal Science, Pontifícia Universidade Católica do Paraná, Curitiba, Paraná 80215-901, Brazil

industrialization. These by-products consist largely of fibrous root material containing residual starch that is not extracted during processing (Leonel and Cereda 2000). Driven by sustainability goals, there is increasing interest in incorporating these by-products into animal nutrition, particularly due to their reduced environmental contamination associated with hydrogen cyanide (HCN) (Cereda 2001; Nambisan 2011; Oliveira et al. 2012). The main limitations to the use of cassava in poultry diets are its high crude fiber content, the presence of HCN, and its low crude protein concentration (Diarra and Devi 2015).

The metabolizable energy value of cassava residues is approximately 2973 kcal ME/kg, making them a feasible substitute for corn in poultry feeding programs (Rostagno et al. 2024). The solid residue resulting from cassava starch extraction contains fibrous material (approximately 11% crude fiber) and non-extracted starch (60–70%), with minimal amounts of crude protein, lipids, vitamins, and minerals (Aro et al. 2010). Nutrient composition varies according to cassava cultivar, plant maturity, and processing method, among other factors (Pandey et al. 2000), and these factors must be considered and properly adjusted during diet formulation.

According to Ogbuewu and Mbajiorgu (2023), up to 25% cassava can be included in layer diets without adverse effects on feed intake, feed conversion ratio, or egg production, indicating a promising future for cassava as an energy source for laying hens. However, although scientific literature on cassava by-products in broiler nutrition exists, the available evidence remains fragmented, with considerable variation in by-product type, dietary inclusion level, rearing phase, and reported performance outcomes. Consequently, a systematic review using a qualitative direction-of-effect synthesis was conducted to identify overall response patterns and knowledge gaps in the current literature. The objective of the present systematic review was to evaluate the effects of replacing corn with cassava by-products in broiler diets on growth performance across different rearing phases using a qualitative direction-of-effect synthesis.

## Materials and methods

### Literature search strategy

A systematic literature search was conducted in the databases Web of Science, Scopus, and PubMed to identify studies evaluating the use of cassava by-products in broiler diets. The search was performed in March 2025 and included studies published between 2005 and 2025. The search strategy was developed based on a Population–Intervention–Outcome (PIO) framework, where Population was

defined as broiler chickens, Intervention as dietary inclusion of cassava by-products, and Outcome as growth performance parameters. This systematic review was conducted according to the PRISMA guidelines.

The following search string was used: (cassava OR “cassava by-product\*” OR “cassava residue\*” OR “cassava pulp” OR “cassava flour” OR “cassava bagasse”) AND (broiler\* OR “broiler chicken\*” OR poultry) AND (performance OR “growth performance” OR “body weight gain” OR “feed intake” OR “feed conversion ratio”) AND (diet\* OR feed OR nutrition OR “corn replacement”). Search terms were adapted as needed for each database. In addition, the reference lists of all eligible studies were manually screened to identify potentially relevant articles not retrieved through the electronic database search. Only studies published in English in peer-reviewed journals were considered. This language restriction may have introduced language bias and should be considered when interpreting the findings. This systematic review was not prospectively registered in an international review registry.

### Study selection and eligibility criteria

All retrieved records were imported into Rayyan for screening and duplicate removal (Ouzzani et al. 2016). Two independent reviewers screened titles and abstracts, followed by full-text assessment. Disagreements were resolved through discussion or consultation with a third reviewer.

Studies were included if they met the following criteria:

1. In vivo experimental studies involving broiler chickens;
2. Evaluation of diets containing cassava by-products (pulp, bagasse, or meal);
3. Inclusion of a control or reference group;
4. Reporting at least one performance outcome (feed intake, body weight gain, or feed conversion ratio);
5. Cassava by-products used as partial or total replacement of corn without confounding inclusion of other major energy sources.

Studies were excluded if they: (i) involved other poultry species; (ii) did not report performance outcomes; (iii) included cassava in combination with other major dietary energy replacements that prevented isolated evaluation; (iv) were not published in English.

### Data extraction and database construction

Data were extracted independently by two reviewers using a standardized spreadsheet. Extracted information included: study identification (authors, year, country); experimental design (number of animals, replicates, duration); type of

cassava by-product (pulp, bagasse, meal); dietary inclusion level (%); production phase (starter, grower–finisher, total period); performance outcomes (FI, BWG, FCR); nutritional characteristics of diets when available. When discrepancies occurred, consensus was reached through discussion. Each dietary treatment within a study was recorded as a separate comparison for qualitative synthesis.

### Risk of bias assessment

Risk of bias was assessed using the Cochrane Collaboration Risk of Bias 2 (RoB 2) tool (Sterne et al. 2019), adapted for animal nutrition studies according to the recommendations of Sargeant and O'Connor (2014), as no universally accepted risk-of-bias assessment tool is currently available for controlled animal nutrition experiments. Five domains were evaluated: D1: randomization process; D2: deviations from intended interventions; D3: missing outcome data; D4: measurement of outcomes; D5: selection of reported results.

Each domain was classified as low risk, some concerns, or high risk. Assessments were conducted independently by two reviewers, with disagreements resolved by consensus. Results were visualized using the “robvis” package (McGuinness & Higgins 2021) in R.

### Data synthesis and direction-of-effect analysis

A qualitative synthesis based on a direction-of-effect approach was conducted to summarize the effects of cassava by-products on broiler growth performance. This approach was selected because many of the included studies did not consistently report measures of variability (e.g., standard deviation or standard error), precluding quantitative comparison of effect estimates across studies.

Each comparison was classified as positive (+), negative (−), or neutral (0) relative to the control group (0% inclusion), or to the lowest inclusion level when a control group was not available. For feed intake (FI) and body weight gain (BWG), increases relative to the reference were classified as positive and decreases as negative. For feed conversion ratio (FCR), reductions (improved efficiency) were considered positive, whereas increases were considered negative.

Classification was based on reported mean values regardless of statistical significance because variability measures were inconsistently reported across studies. When regression models were presented, classification was based on the biological response pattern.

Quadratic responses were considered positive when performance improved up to an apparent optimal inclusion level. Each dietary inclusion level within a study was treated as a separate comparison for descriptive purposes; however, this approach may introduce dependency among

observations and is acknowledged as a limitation of the qualitative synthesis. To facilitate interpretation, inclusion levels were grouped into predefined categories low (2–8%), moderate (10–25%), and high (> 25%), allowing identification of overall response patterns across studies.

## Results

### Study selection and characteristics

The literature search identified 376 records, of which 202 duplicates were removed. After title and abstract screening, 96 studies were excluded for not meeting the eligibility criteria, resulting in 78 articles for full-text assessment. Of these, 42 were excluded due to absence of relevant performance outcomes, inclusion of confounding dietary ingredients, or language restrictions. A total of 36 studies met the eligibility criteria and were included in the qualitative synthesis (Fig. 1). The included studies were published between 2005 and 2025 and were predominantly conducted in Brazil (36.1%), Nigeria (25.0%), and Indonesia (18.2%). Cassava pulp was evaluated in 24 studies (66.7%), followed by cassava flour in seven studies (19.4%) and cassava bagasse in five studies (13.9%) (Table 1).


Most studies evaluated the grower–finisher phase (33.3%) or the total rearing period (27.8%), with fewer studies focusing exclusively on the starter phase or combined production phases. Feed intake (FI), body weight gain (BWG), and feed conversion ratio (FCR) were reported in 97.2% of the included studies (Table 1).

### Risk of bias assessment

Overall, 13.9% of the studies were classified as having low risk of bias, whereas 47.2% were classified as high risk and 38.9% as presenting some concerns. The most frequent sources of bias were related to missing outcome data (Domain 3) and measurement of outcomes (Domain 4) (Fig. 2). Domains related to deviations from intended interventions (Domain 2) and selection of reported results (Domain 5) showed the highest proportion of low-risk classifications.

### Direction-of-effect synthesis

A total of 187 comparisons were included in the qualitative synthesis. For feed intake (FI), neutral responses were the most frequent (54.4%), followed by negative (26.4%) and positive responses (19.2%). For body weight gain (BWG), negative responses were the most frequent (49.2%), followed by neutral (44.4%) and positive responses (6.4%). For



**Fig. 1** PRISMA flow diagram of the study selection process for the systematic review

feed conversion ratio (FCR), neutral (45.5%) and negative (44.4%) responses occurred at similar frequencies, whereas positive responses accounted for 10.2% of the comparisons (Fig. 3).

### Influence of inclusion level

When comparisons were stratified according to dietary inclusion level (2–8%, 10–25%, and >25%), negative responses became more frequent with increasing inclusion level, particularly for body weight gain (BWG) and feed conversion ratio (FCR) (Fig. 3). In contrast, neutral responses were

more frequent at lower dietary inclusion levels for all evaluated outcomes. Feed intake (FI) responses were distributed across all inclusion categories, with neutral responses remaining the most frequent regardless of dietary inclusion level (Fig. 3).

### Relationship between risk of bias and direction of effect

Most comparisons were derived from studies classified as having high risk of bias or some concerns. No clear

**Table 1** Characteristics of the studies included in the systematic review, including author, year of publication, country, cassava by-product, production phase, and performance outcomes evaluated

| Code | Author                | Year        | Country                | By-product  | Phases                                    | Performance*        |
|------|-----------------------|-------------|------------------------|-------------|-------------------------------------------|---------------------|
| 1    | Do Nascimento et al.  | 2005        | Brazil                 | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 2    | Khempaka et al.       | 2009        | Thailand               | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 3    | Aderemi et al.        | 2010        | Nigeria                | Flour       | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 4    | Ali-Mursyid et al.    | 2010        | Indonesia              | Bagasse     | Total                                     | FI, BWG, FCR        |
| 5    | Ferreira et al.       | 2012        | Brazil                 | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 6    | Sousa et al.          | 2012        | Brazil                 | Bagasse     | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 7    | Tang et al.           | 2012        | Australia              | Pulp        | Starter; Grower/Finisher and Total        | FI, BWG, FCR        |
| 8    | Oso et al.            | 2014        | Nigeria                | Pulp        | Total                                     | FI, BWG, FCR        |
| 9    | Tesfaye et al.        | 2013        | Ethiopia               | Pulp        | Starter and Finisher                      | FI, BWG, FCR        |
| 10   | Akapo et al.          | 2014        | Nigeria                | Pulp        | Total                                     | FI, BWG, FCR        |
| 11   | Khempaka et al.       | 2014        | Japan                  | Pulp        | Total                                     | FI, BWG, FCR        |
| 12   | Lubis et al.          | 2007        | Indonesia              | Bagasse     | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 13   | Oladunjoye et al.     | 2014        | Nigeria                | Flour       | Total                                     | FI, BWG, FCR        |
| 14   | Picoli et al.         | 2014        | Brazil                 | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 15   | Geron et al.          | 2015        | Brazil                 | Pulp        | Total                                     | FI, BWG, FCR        |
| 16   | Holanda et al.        | 2015        | Brazil                 | Flour       | Grower/Finisher                           | FI, BWG, FCR        |
| 17   | Sugiharto et al.      | 2016        | Indonesia              | Pulp        | Grower/Finisher                           | BWG, FCR            |
| 18   | Broch et al.          | 2017        | Brazil                 | Bagasse     | Starter and Total                         | FI, BWG, FCR        |
| 19   | Ojewola et al.        | 2017        | Nigeria                | Flour       | Total                                     | FI, BWG, FCR        |
| 20   | Dayal et al.          | 2018        | Nigeria                | Flour       | Grower and Grower/Finisher                | FI, BWG, FCR        |
| 21   | Parente et al.        | 2018        | Brazil                 | Bagasse     | Grower/Finisher                           | FI, BWG, FCR        |
| 22   | Da Silva et al.       | 2019        | Brazil                 | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 23   | Sugiharto et al.      | 2019        | Indonesia              | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 24   | Yadav et al.          | 2019        | United States (Hawaii) | Pulp        | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 25   | Chang'a et al.        | 2019        | Australia              | Pulp        | Starter; Grower/Finisher and Total        | FI, BWG, FCR        |
| 26   | Oyewole et al.        | 2020        | Nigeria                | Pulp        | Total                                     | FI, BWG, FCR        |
| 27   | Trautenmüller et al.  | 2020        | Brazil                 | Pulp        | Starter; Grower/Finisher and Total        | FI, BWG, FCR        |
| 28   | <b>Almeida et al.</b> | <b>2020</b> | <b>Brazil</b>          | <b>Pulp</b> | <b>Starter; Grower/Finisher and Total</b> | <b>FI, BWG, FCR</b> |
| 29   | Ferreira et al.       | 2021        | Brazil                 | Pulp        | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 30   | Ojedian et al.        | 2022        | Nigeria                | Pulp        | Total                                     | FI, BWG, FCR        |
| 31   | Okpara et al.         | 2022        | Nigeria                | Pulp        | Total                                     | FI, BWG, FCR        |
| 32   | Vieira et al.         | 2022        | Brazil                 | Flour       | Grower/Finisher                           | FI, BWG, FCR        |
| 33   | Puspita et al.        | 2023        | Indonesia              | Pulp        | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 34   | Ramadhani et al.      | 2025        | Indonesia              | Pulp        | Grower/Finisher                           | FI, BWG, FCR        |
| 35   | Yang et al.           | 2024        | China                  | Pulp        | Starter and Grower/Finisher               | FI, BWG, FCR        |
| 36   | Osei-Adjei et al.     | 2025        | Ghana                  | Flour       | Grower/Finisher                           | FI, BWG, FCR        |

FI: feed intake; BWG: body weight gain; FCR: feed conversion ratio.

relationship was observed between the risk-of-bias category and the direction of effect across the evaluated outcomes.

## Discussion

The present systematic review provides a comprehensive synthesis of the available evidence on the use of cassava by-products as partial or total replacements for corn in broiler diets. The results indicate that these ingredients were more frequently associated with reduced growth performance, particularly at dietary inclusion levels above 25%, although considerable variability was observed among studies.

A plausible explanation for these findings is the high fiber concentration present in cassava by-products. With increased inclusion in the diet, the amount of carbohydrates with glycosidic bonds ( $\alpha$ -1,6  $\beta$ -1,4, and  $\beta$ -1,6), which are resistant to hydrolysis in the digestive tract, also increases, potentially affecting the digestion of other nutrients, leading to more significant nutrient loss (Raupp et al. 2002) and reduced nutrient availability.

Physiologically, birds have limited utilization of fibrous compounds due to their digestive system characteristics and nutritional interactions, which affect nutrient digestion (Costa et al. 2009; Macari and Maiorka 2017).

Additionally, the high fiber content in food may increase digesta viscosity in the chicken small intestine, decreasing

**Fig. 2** Risk-of-bias assessment of the included studies across the five evaluated domains using the adapted RoB 2 tool

|                            | Risk of bias domains |    |    |    |    |         |
|----------------------------|----------------------|----|----|----|----|---------|
|                            | D1                   | D2 | D3 | D4 | D5 | Overall |
| Do Nascimento et al., 2005 | +                    | +  | -  | +  | +  | -       |
| Khempaka et al., 2009      | +                    | +  | ×  | -  | +  | ×       |
| Aderemi et al., 2010       | ×                    | +  | +  | -  | +  | ×       |
| Ali-Mursyid et al., 2010   | +                    | +  | ×  | +  | +  | ×       |
| Ferreira et al., 2012      | +                    | +  | ×  | -  | +  | ×       |
| Sousa et al., 2012         | +                    | +  | -  | -  | +  | -       |
| Tang et al., 2012          | +                    | +  | ×  | -  | +  | ×       |
| Oso et al., 2013           | +                    | +  | ×  | -  | +  | ×       |
| Tesfaye et al., 2013       | +                    | +  | +  | +  | +  | +       |
| Akapo et al., 2014         | +                    | +  | ×  | +  | +  | ×       |
| Khempaka et al., 2014      | +                    | +  | ×  | +  | +  | ×       |
| Lubis et al., 2014         | +                    | +  | +  | -  | +  | -       |
| Oladunjoye et al., 2014    | ×                    | +  | +  | -  | +  | ×       |
| Picoli et al., 2014        | +                    | +  | -  | -  | +  | -       |
| Geron et al., 2015         | -                    | -  | -  | -  | +  | -       |
| Holanda et al., 2015       | +                    | +  | -  | -  | +  | -       |
| Sugiharto et al., 2016     | +                    | +  | +  | -  | +  | -       |
| Broch et al., 2017         | +                    | +  | -  | -  | +  | -       |
| Ojewola et al., 2017       | +                    | +  | ×  | +  | +  | ×       |
| Dayal et al., 2018         | +                    | +  | -  | +  | +  | -       |
| Parente et al., 2018       | +                    | +  | ×  | +  | +  | ×       |
| Da Silva et al., 2019      | +                    | +  | -  | -  | +  | -       |
| Sugiharto et al., 2019     | ×                    | +  | ×  | +  | +  | ×       |
| Yadav et al., 2019         | +                    | +  | +  | +  | +  | +       |
| Chang'a et al., 2020       | +                    | +  | -  | -  | +  | -       |
| Oyewole et al., 2020       | +                    | +  | ×  | +  | +  | ×       |
| Trautenmüller et al., 2020 | +                    | +  | -  | +  | +  | -       |
| Ferreira et al., 2021      | +                    | +  | -  | +  | +  | -       |
| Ojediran et al., 2022      | +                    | +  | ×  | +  | +  | ×       |
| Okpara et al., 2022        | +                    | +  | ×  | +  | +  | ×       |
| Vieira et al., 2022        | +                    | +  | -  | +  | +  | -       |
| Puspita et al., 2023       | +                    | +  | ×  | +  | +  | ×       |
| Almeida et al., 2024       | +                    | +  | +  | +  | +  | +       |
| Ramadhanti et al., 2024    | +                    | +  | +  | +  | +  | +       |
| Yang et al., 2024          | +                    | +  | ×  | -  | +  | ×       |
| Osei-Adjei et al., 2025    | +                    | +  | +  | +  | +  | +       |


**Domains:**  
D1: Bias arising from the randomization process.  
D2: Bias due to deviations from intended intervention.  
D3: Bias due to missing outcome data.  
D4: Bias in measurement of the outcome.  
D5: Bias in selection of the reported result.

**Judgement**  
 High  
 Some concerns  
 Low

the digestibility and absorption of other nutrients (Hetland et al. 2004). When this fiber-rich content comes into contact with water, gums are formed that increase bolus size, and through gastric distension, FI is reduced due to a feeling of satiety (Macari and Maiorka 2017). This physiological

mechanism explains why fiber-rich foods contribute to reduced FI in chickens.

The low protein content of cassava by-products, along with the challenges of high fiber content and anti-nutritional factors, limits their inclusion (Valdivié et al. 2008). Consequently, one key to the successful use of cassava by-products as a



**Fig. 3** Direction-of-effect responses for feed intake (FI), body weight gain (BWG), and feed conversion ratio (FCR) according to dietary inclusion level (%) in broilers fed cassava by-products

corn substitute in poultry diets is determining the appropriate replacement level and product origin to ensure adequate nutritional composition concerning lysine and methionine (Okereke 2012).

Moderate levels of dietary fiber have been shown to improve nutrient digestibility and performance in poultry, primarily due to their role in the development of the gastrointestinal tract and in modulating intestinal content characteristics (Röhe and Zentek 2021). In addition, studies indicate that supplementation with soluble or insoluble fiber can be beneficial to poultry health, with recommended inclusion levels ranging from 2 to 2.4% in the diet (Jha and Mishra 2021).

Despite the recognized importance of fiber for intestinal histomorphology and nutrient digestibility, only 28% of the studies included in the present review (10 out of 36) evaluated these parameters.

Measures of variability, such as standard deviation (SD) or standard error of the mean (SEM), were not consistently reported across the included studies, preventing reliable quantitative comparison of effect estimates. Consequently, the evidence was synthesized using a qualitative direction-of-effect approach (Higgins and Green, 2023).

The omission of SD was the most prominent issue, affecting 85.7% (30 of 35 studies) for feed intake and 86.1% (31 of 36 studies) for both feed conversion ratio and body weight gain. The absence of SEM was slightly lower, occurring in 42.9% (15 of 35 studies) for feed intake and in 38.9% (14 of 36 studies) for feed conversion ratio and weight gain. These omissions limited the assessment of estimate precision and within-study variability.

The direction-of-effect assessment showed a predominance of neutral responses for feed intake (FI), whereas negative responses were more frequently observed for body

weight gain (BWG) and, to a lesser extent, feed conversion ratio (FCR). These findings suggest that cassava by-products were more frequently associated with changes in growth performance than with voluntary feed intake.

In broiler chickens, feed intake is primarily regulated by dietary energy density; thus, birds tend to increase intake to compensate for diets with lower metabolizable energy (Macari and Maiorka 2017). However, this compensatory response may be constrained by physical and physiological factors, such as increased dietary fiber, gut fill, and digesta viscosity, which can limit intake capacity (Rostagno et al. 2024).

As a result, even when feed intake remains unchanged, reduced nutrient digestibility and energy availability may impair growth, leading to lower BWG and poorer FCR. The greater frequency of neutral responses for FI compared with BWG and FCR suggests that growth performance was more sensitive to increasing dietary inclusion levels of cassava by-products than voluntary feed intake.

These findings should be interpreted considering the methodological limitations and overall quality of the included studies.

In a meta-regression conducted by Ogbuewu and Mbajiorgu (2023), evaluating cassava inclusion in diets for laying hens, the authors observed that substitution levels above 25% increased feed intake and reduced eggshell thickness, egg production, and egg weight. These adverse effects may be associated with antinutritional factors (Omede et al. 2017) found in cassava-based diets that exceeded the tolerance threshold of laying hens (Chauynarong et al. 2009). Similarly, the wide variation in dietary inclusion levels observed in the present review may have contributed to the heterogeneous performance responses reported among studies. Dietary inclusion levels ranged from 2 to 100% for cassava pulp, 2.5 to 30% for cassava bagasse, and 6.8 to 100% for cassava meal, highlighting the lack of standardization across experiments.

**Cassava pulp.** Cassava pulp is the by-product with the least industrial processing, resulting in minimal alteration to its nutritional composition and a higher fiber concentration than other cassava by-products. The pulp typically contains 50–70% starch and 20–30% fiber, mainly cellulose and non-starch polysaccharides (Rattanachomsri et al. 2009).

**Cassava bagasse.** Cassava bagasse is a by-product of starch extraction (Leonel and Cereda 2000). During processing, cassava roots are peeled, crushed, and washed to extract starch, resulting in a coarse residue. This material typically contains approximately 1.5% crude protein, 11% crude fiber, and 2,450 kcal/kg of metabolizable energy on a dry matter basis (Rodrigues and Campos 2001), and may contain up to 80% residual starch.

**Cassava meal.** According to Rostagno et al. (2024), whole cassava root meal contains 2.64% crude protein, 4.21%

crude fiber, 73.7% starch, and 2,973 kcal/kg of metabolizable energy for poultry.

An economic analysis by Sousa et al. (2012) examined the inclusion of cassava bagasse in two rearing phases (from d 1–21 and d 22–40) in broiler chicken production. The results indicated that the lowest feed cost per kilogram of live weight gain and the best productive efficiency index were achieved with dietary inclusion levels of up to 5% cassava bagasse, indicating increased profitability. Above 5%, feed costs increased because of the need to supplement amino acids to maintain a balanced diet.

In regions with high cassava production, safe levels of corn substitution that do not impair broiler performance could promote sustainable production by utilizing these by-products and reducing diet costs (Uchegbu et al. 2011). Cassava by-products can be included in monogastric diets as a strategy to reduce corn use and, consequently, feed costs (Ferreira et al. 2012). However, their main limitation is the high fiber content, which often reduces nutrient utilization. High levels of crude fiber, particularly from plant cell wall components, can negatively affect diet digestibility (Brito et al., 2008).

The risk-of-bias assessment showed that only 13.9% of the studies were classified as having a low risk of bias (Tsfaye et al. 2013; Yadav et al. 2019; Almeida et al., 2020; Ramadhanti et al., 2025; Osei-Adjei et al. 2025), whereas the remaining studies were classified as presenting either some concerns or a high risk of bias. Missing outcome data were identified in 15 studies (41.6%), mainly due to bird mortality, without appropriate methodological adjustments to account for its effects on feed intake and feed conversion ratio. In addition, 16 studies (44.4%) presented subjective methodologies or lacked sufficient detail regarding the measured variables, likely due to the assumption of standard practices in performance evaluation.

Overall, 86.1% of the studies were classified as having high risk of bias or some concerns, indicating important methodological limitations across the evidence base. Although these studies provide useful quantitative data, their results should be interpreted with caution, and improvements in experimental design and reporting are needed in future research.

The geographic distribution of the included studies showed that Brazil, Nigeria, and Indonesia accounted for the majority of studies, with the cassava species *Manihot esculenta* being the most commonly used across these regions.

The processing methods of cassava by-products varied widely among studies. Most involved dehydration, either sun-drying or industrial drying, while others included grinding, washing, peeling, and, in fewer cases, fermentation. In general, studies used residues from pulp or bagasse processing with relatively simple processing methods, regardless of geographic origin. A temporal analysis indicated a

progression from traditional sun-drying and simple grinding techniques to more advanced methods, such as oven-drying, evaluation of cyanogenic compound volatilization, and controlled fermentation using microorganisms, which may contribute to improved utilization of cassava by-products.

Regarding hydrogen cyanide (HCN) content, only seven studies reported its concentration, with values ranging from 3.97 to 43.40 mg/kg. Cassava is considered one of the most important cyanogenic species in tropical regions due to the natural presence of linamarin and lotaustralin (Cereda and Mattos 1996). Other economically important plants, such as sorghum, plum, bamboo, and rubber tree, also contain varying levels of HCN (Haque and Bradbury 2002), which can lead to endogenous cyanide production upon ingestion. Identifying cassava varieties with low cyanogenic potential is essential to improve safety and reduce the risk of toxicity.

Overall, the findings of this study indicate that cassava by-products were more frequently associated with variable effects on broiler performance, with negative responses occurring more frequently at dietary inclusion levels above 25%. However, the considerable heterogeneity among studies, differences in processing methods and dietary inclusion levels, and the predominance of studies with methodological limitations highlight the need for cautious interpretation.

In this context, the effective use of cassava by-products in poultry nutrition depends on appropriate processing, careful formulation, and appropriate dietary inclusion levels. Future research should focus on standardized experimental designs, comprehensive reporting of nutritional and toxicological parameters, and consistent reporting of measures of variability (e.g., SD and SEM) to better define safe and optimal inclusion strategies.

## Conclusion

Cassava by-products may represent alternative feed ingredients in broiler diets. However, they were more frequently associated with reduced growth performance at dietary inclusion levels above 25%, although responses varied according to the type of cassava by-product, processing method, and experimental conditions. Their effective use depends on appropriate processing, careful dietary formulation, and the adoption of suitable dietary inclusion levels to minimize nutritional limitations and maximize their potential as a partial substitute for corn. Future studies should adopt standardized experimental protocols and report measures of variability (e.g., SD and SEM) to strengthen the available evidence and better define safe inclusion levels for each cassava by-product.

**Acknowledgements** The authors gratefully acknowledge the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES),

Brazil – Finance Code 001, the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Universidade Brasil, Universidade Federal de Mato Grosso (UFMT), Universidade Federal Rural do Rio de Janeiro (UFRRJ), and Pontifícia Universidade Católica do Paraná (PUCPR) for their institutional and academic support.

**Author Contributions** Conceptualization, W. D. M., H. M. S., and C. P. Z.; methodology, W. D. M., S. S., H. M. S., M. E. V. N., L. C. S., and C. P. Z.; validation, S. S. and J. K. V.; investigation, S. S., M. M. C. P., L. C. S., and M. E. V. N.; resources, S. S., L. C. S., and M. E. V. N.; formal analysis, M. M. C. P., L. C. S., and M. E. V. N.; software, M. M. C. P.; data curation, M. M. C. P., L. C. S., and M. E. V. N.; writing—original draft preparation, S. S. and J. K. V.; writing—review and editing, S. S., M. M. C. P., and J. K. V.; visualization, S. S. and J. K. V.; supervision, S. S., J. K. V., and C. P. Z.; project administration, W. D. M., S. S., H. M. S., and C. P. Z.; funding acquisition, W. D. M., H. M. S., and C. P. Z. All authors have read and agreed to the published version of the manuscript.

**Funding** The Article Processing Charge (APC) for the publication of this research was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) (ROR identifier: 00x0ma614). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001, and by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

**Data availability** The data presented in this study will be made available in supplementary materials in Excel (.xlsx) format.

## Declarations

**Ethics approval** This article does not contain any studies with human participants or animals performed by any of the authors.

**Conflict of interest** The authors report there are no competing interests to declare.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Aderemi F (2010) Utilization of graded levels of biodegraded cassava peels in broiler ration. *Electron J Environ Agric Food Chem* 9:672–678. <https://www.ajol.info/index.php/tjas/article/view/202144>
- Akapo AO, Oso AO, Bamgbose AM, Sanwo KA, Jegede AV, Sobayo RA, Olorunsola RA (2014) Effect of feeding cassava (*Manihot esculenta* Crantz) root meal on growth performance, hydrocyanide intake and haematological parameters of broiler chicks. *Trop*

- Anim Health Prod 46:1167–1172. <https://doi.org/10.1007/s1125-0-014-0622-5>
- Ali-Mursyid WM, Zuprizal Z, Nur-Cahyanto M (2010) Corn substitution using fermented solid cassava-waste on broiler chicken. J Indones Trop Anim Agric 35:9–15. <https://eprints.undip.ac.id/17157/>
- Almeida AZ, Eyng C, Nunes RV, Broch J, Souza C, Sangalli GG, Tenório KI (2020) Carcass characteristics and meat quality of slow-growing broilers fed diets containing dry residue of cassava with or without the addition of carbohydrases. Trop Anim Health Prod 52:2677–2685. <https://doi.org/10.1007/s11250-020-02308-4>
- Aro SO, Aletor VA, Tewe OO, Agbede JO (2010) Nutritional potentials of cassava tuber wastes: a case study of a cassava starch processing factory in southwestern Nigeria. Livest Res Rural Dev 22:11. <https://www.lrrd.cipav.org.co/lrrd22/11/aro22213.htm>
- Broch J, Nunes RV, Oliveira V, Silva IM, Souza C, Wachholz L (2017) Dry residue of cassava as a supplementation in broiler feed with or without addition of carbohydrases. Semina Ciênc Agrár 38:2641–2658. <https://doi.org/10.5433/1679-0359.2017v38n4Supl1p2641>
- Cereda MP (2001) Caracterização dos subprodutos da industrialização da mandioca. Manejo, uso e tratamento de subprodutos da industrialização da mandioca. São Paulo: Fundação Cargill 4:31–35
- Cereda MP, Mattos MCY (1996) Linamarin: the toxic compound of cassava. J Venom Anim Toxins 2(1):6–12. <https://doi.org/10.1590/S0104-79301996000100002>
- Chang'a EP, Abdallah ME, Ahiwe EU, Mbaga S, Zhu ZY, Fru-Nji F, Iji PA (2019) Replacement value of cassava for maize in broiler chicken diets supplemented with enzymes. Asian-Australas J Anim Sci 33:1126
- Chauynarong N, Elangovan AV, Iji PA (2009) The potential of cassava products in diets for poultry. World Poult Sci J 65:23–36. <https://doi.org/10.1017/S0043933909000026>
- Costa FGP, Goulart CC, Costa JS, Souza CJ, Dourado LRB, Silva JHV (2009) Desempenho, qualidade de ovos e análise econômica da produção de poedeiras semipesadas alimentadas com diferentes níveis de raspa de mandioca. Acta Sci 31:13–18. <https://doi.org/10.4025/actascianimsci.v31i1.457>
- da Silva IM, Broch J, Wachholz L, Souza C, Dalolio FS, Teixeira LV, Nunes RV (2019) Dry residue of cassava associated with carbohydrases in diets for broiler chickens. J Appl Poult Res 28:1189–1201. <https://doi.org/10.3382/japr/pfz085>
- Dayal AD, Diarra SS, Lameta S, Devi A, Amosa F (2018) High cassava peel meal-based diets with animal fat and enzyme for broilers. Livest Res Rural Dev 30:99. <https://lrrd.cipav.org.co/lrrd30/6/siaka30099.html>
- Diarra SS, Devi A (2015) Feeding value of some cassava by-products meal for poultry: a review. Pakistan J Nutr 14:735–741. <https://repository.usp.ac.fj/id/eprint/8419/1/PJN.pdf>
- Instituto Brasileiro de Geografia e Estatística (IBGE) (2025) Levantamento Sistemático da Produção Agrícola. Brazil. <https://sidra.ibge.gov.br/tabela/6588resultado>. Accessed Aug. 2024
- Ferreira AHC, Lopes JB, Abreu MLT, Merval RR, Ribeiro MN, Silva FES, Andrade FT (2012) Cassava root scrapings for 22 to 42-day-old broilers in high-temperature environments. R Bras Zootec 41:1442–1447. <https://doi.org/10.1590/S1516-3598201200060018>
- Ferreira BL, Brito DAP, Bayão GFV, Serra JL, Sousa EM, Santos WS, Lima V (2021) Utilization of increasing levels of cassava root meal in the diet of broiler. Ciênc Anim Bras 22:e69289. <https://doi.org/10.1590/1809-6891v22e-69289>
- Geron LJV, Costa FG, Moraes KJS, Oliveira EM, Gomes RD, Pereira SR, Pelicia K (2015) Consumo, desempenho e rendimento de carcaça de frangos de corte alimentados com rações contendo raspa de mandioca residual desidratada. Bol Ind Anim 72:304–310. [http://web.archive.org/web/20180605044655id\\_/http://www.iz.sp.gov.br/pdfsbia/1451413465.pdf](http://web.archive.org/web/20180605044655id_/http://www.iz.sp.gov.br/pdfsbia/1451413465.pdf)
- Haque MR, Bradbury JH (2002) Total cyanide determination of plants and foods using the picrate and acid hydrolysis methods. Food Chem 77:107–114. [https://doi.org/10.1016/S0308-8146\(01\)00313-2](https://doi.org/10.1016/S0308-8146(01)00313-2)
- Hetland H, Choct M, Svihus B (2004) Role of insoluble nonstarch polysaccharides in poultry nutrition. Worlds Poult Sci J 60:415–422. <https://doi.org/10.1079/WPS200325>
- Higgins JPT, Green S (eds) (2023) Cochrane handbook for systematic reviews of interventions. Version 6.4. Cochrane. <https://training.cochrane.org/handbook>
- Holanda MAC, Holanda MCR, Vigoderes RB, Dutra WM, Albino LFT (2015) Desempenho de frangos caipiras alimentados com farelo integral de mandioca. Rev Bras Saúde Prod Anim 16:106–117. <https://doi.org/10.1590/S1519-99402015000100012>
- Jha R, Mishra P (2021) Dietary fiber in poultry nutrition and their effects on nutrient utilization, performance, gut health, and on the environment: a review. J Anim Sci Biotechnol 12:51. <https://doi.org/10.1186/s40104-021-00576-0>
- Khempaka S, Molee W, Guillaume M (2009) Dried cassava pulp as an alternative feedstuff for broilers: Effect on growth performance, carcass traits, digestive organs, and nutrient digestibility. J Appl Poult Res 18:487–493. <https://doi.org/10.3382/japr.2008-00124>
- Khempaka S, Thongkratok R, Okrathok S, Molee W (2014) An evaluation of cassava pulp feedstuff fermented with *A. oryzae* on growth performance, nutrient digestibility and carcass quality of broilers. J Poult Sci 51:71–79. <https://doi.org/10.2141/jpsa.0130022>
- Latif S, Muller J (2015) Potential of cassava leaves in human nutrition: a review. Trends Food Sci Technol 44:147–158. <https://doi.org/10.1016/j.tifs.2015.04.006>
- Leonel M, Cereda MP (2000) Extração da fécula retida no resíduo fibroso do processo de produção de fécula de mandioca. Food Sci Technol 20:122–127. <https://doi.org/10.1590/S0101-2061200000100023>
- Lubis AD, Darmawan B, Ningrum H, Noormasari IY, Nakagoshi N (2007) Evaluation of fermented cassava (*Manihot esculenta* Crantz) pulp as feed ingredient for broiler. Tropics 17:73–80. <https://doi.org/10.3759/tropics.17.73>
- Macari M, Maiorka A (2017) Fisiologia das aves comerciais. 2nd ed. Funep, Jaboticabal, Brazil 53–63
- McGuinness LA, Higgins JPT (2021) robvis: Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. Res Synth Methods 12:55–61. <https://doi.org/10.1002/jrsm.1411>
- Nambisan B (2011) Strategies for elimination of cyanogens from cassava for reducing toxicity and improving food safety. Food Chem Toxicol 49:690–693. <https://doi.org/10.1016/j.fct.2010.10.035>
- Nascimento GAJD, Costa FGP, Amarante Júnior VDS, Barros LR (2005) Efeitos da substituição do milho pela raspa de mandioca na alimentação de frangos de corte, durante as fases de engorda e final. Ciência e Agrotecnol 29:200–207. <https://doi.org/10.1590/S1413-70542005000100025>
- Ogbuewu IP, Mbajioru CA (2023) Meta-analysis of the influence of dietary cassava on productive indices and egg quality of laying hens. Heliyon 9(3). <https://linkinghub.elsevier.com/retrieve/pii/S2405844023012057>
- Ojediran T, Aroyehun B, Emiola A (2022) Evaluation of cassava distillers' waste meal in the diet of broiler chicken. Anim Sci Genet 18:41–55. <https://doi.org/10.5604/01.3001.0015.8961>
- Ojewola GS, Okeke UL, Olojede AO (2017) Effects of enzyme supplementation of cassava flour-based diets on performance and economic efficiency of broiler chicken. [https://managementjournal.usamv.ro/pdf/vol.17\\_1/Art43.pdf](https://managementjournal.usamv.ro/pdf/vol.17_1/Art43.pdf)
- Okereke CO (2012) Utilization of cassava, sweet potato, and cocoyam meals as dietary sources for poultry. World J Eng Pure Appl Sci 2:63. <https://www.proquest.com/openview/c937b4441122dcb42a1842870a98a2e>

- Okpara O, Obakanurhe O, Onowhakpor CN, Sorhue UG, Gbayisomore OS (2022) Effect of replacing maize with processed cassava on the growth performance and haematological characteristics of broiler chickens. *Anim Nutr Feed Technol* 22:155–166. <https://doi.org/10.5958/0974-181X.2022.00013.0>
- Oladunjoye IO, Ojebiyi OO, Rafiu TA (2014) High methionine supplementation improves the nutritional value of cassava peel meal for broiler chicken. *Livest Res Rural Dev* 27:26–30. <https://www.lrrd.org/lrrd26/4/olad26063.htm>
- Oliveira NTD, Uchôa SCP, Alves JMA, Sediya T, Albuquerque JDAAD, Souza ED, Melville CC (2012) Ácido cianídrico em tecidos de mandioca em função da idade da planta e adubação nitrogenada. *Pesqui Agropecu Bras* 47:1436–1442. <https://doi.org/10.1590/S0100-204X2012001000004>
- Omede AA, Ahiwe EU, Zhu ZY, Fru-Nji F, Iji PA (2017) Improving cassava quality for poultry feeding through application of biotechnology. In: Waisundara V (ed) *Cassava*. <https://doi.org/10.5772/intechopen.72236>
- Osei-Adjei A, Hamidu JA, Adjei-Mensah B, Hanson PNA, Sarfo GK, Donkoh A (2025) Evaluating cassava starch residue as a maize substitute in broiler chicken diets. *Food Sci Nutr* 13:e70360. <https://doi.org/10.1002/fsn3.70360>
- Oso AO, Akapo O, Sanwo KA, Bamgbose AM (2014) Utilization of unpeeled cassava (*Manihot esculenta* Crantz) root meal supplemented with or without charcoal by broiler chickens. *J Anim Physiol Anim Nutr* 98:431–438. <https://doi.org/10.1111/jpn.12088>
- Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A (2016) Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 5(1):210. <https://doi.org/10.1186/s13643-016-0384-4> PMID: 27919275; PMCID: PMC5139140
- Oyewole NO, Akinyele SA, Ogunsipe MH (2020) Effect of cassava pulp substituting maize on the growth performance and haematobiochemical attributes of broiler chickens. *Bone* 4:4–50
- Pandey A, Soccol CR, Nigam P, Soccol VT, Vandenbergue LPS, Mohan R (2000) Biotechnological potential of agro-industrial residue II: Cassava bagasse. *Bioresour Technol* 74:81–87. [https://doi.org/10.1016/S0960-8524\(99\)00143-1](https://doi.org/10.1016/S0960-8524(99)00143-1)
- Parente IP, Albino LFT, Rodrigues KF, Vaz RGMV, Sousa LF, Fonseca FLR, Noleto RA (2018) Cassava bagasse and annatto colorific (*Bixa orellana* L.) in diets for slow-growing broilers from 30 to 90 days of age. *Rev Bras Saúde Prod Anim* 19:58–68. <https://doi.org/10.1590/S1519-99402018000100006>
- Picoli KP, Murakami AE, Nunes RV, Amaral Duarte CR, Eyng C, Ospina-Rojas IC (2014) Cassava starch factory residues in the diet of slow-growing broilers. *Trop Anim Health Prod* 46:1371–1381. <https://doi.org/10.1007/s11250-014-0649-7>
- Puspita PS, Hermana W, Pramujo M, Nahrowi N (2023) The effect of isoamylase application in cassava root meal on broiler growth performance, feed retention time and metabolite profile. *BIO Web Conf* 81:00044. <https://doi.org/10.1051/bioconf/20238100044>
- Ramadhani SAH, Ridla M, Mutia R, Hermana W (2025) Performance and health profile of IPB-D1 chickens supplemented with cassava (*Manihot esculenta* Crantz) leaf extract via drinking water. *J Ilmu Nutr Teknol Pakan* 23:74–81. <https://doi.org/10.29244/jin tp.23.1.74-81>
- Rattanachomsri U, Tanapongpipat S, Eurwilaichitr L, Champred V (2009) Simultaneous nonthermal saccharification of cassava pulp by multienzyme activity and ethanol fermentation by *Candida tropicalis*. *J Biosci Bioeng* 107:488–493. <https://doi.org/10.1016/j.jbiosc.2008.12.024>
- Raupp DDS, Marques SHDP, Rosa DA, Caldi CM, Cremasco ACV, Banzatto DA (2002) Arrastre via fecal de nutrientes da ingestão produzido por bagaço de mandioca hidrolisada. *Sci Agric* 59:235–242. <https://doi.org/10.1590/S0103-90162002000200005>
- Röhe I, Zentek J (2021) Lignocellulose as an insoluble fiber source in poultry nutrition: a review. *Journal of Animal Science and Biotechnology* 12:82. <https://doi.org/10.1186/s40104-021-00594-y>
- Rodrigues AA, Campos OF (2001) Resíduos industriais da raiz da mandioca na alimentação de bovinos. In: Cereda, M. P. (Coord.), *Manejo, uso e tratamento de subprodutos da industrialização da mandioca* (pp. 240–258). São Paulo, Brazil: Fundação Cargill. <http://www.infoteca.cnptia.embrapa.br/infoteca/handle/doc/44869>
- Rostagno HS, Teixeira AO, Calderano AA, Hannas MI, Sakomura NK, Perazzo FG, Rocha GS, Saraiva A, Abreu MLT, Genova JL, Tavernari FC, et al. (2024) Tabelas Brasileiras para Aves e Suínos: Composição de Alimentos e Exigências Nutricionais. 5th ed. Suprema, Viçosa. <https://doi.org/10.26626/978-85-8179-206-4.2024.B001>
- Sargeant JM, O'Connor AM (2014) Introduction to systematic reviews in animal agriculture and veterinary medicine. *Zoonoses Public Health* 61(Suppl 1):3–9. <https://doi.org/10.1111/zph.12128>
- Sousa JPL, Rodrigues KF, Albino LFT, Santos Neta ER, Vaz RGMV, Parente IP, Amorim AF (2012) Bagaço de mandioca em dietas de frangos de corte. *Rev Bras Saúde Prod Anim* 13:1044–1053. <https://www.scielo.br/j/rbspa/a/PhV3tBLkGFzm4mLhrTybQkC/>
- Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng HY, Corbett MS, Eldridge SM, et al. (2019) RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 366:14898. <https://doi.org/10.1136/bmj.14898>
- Sugiharto S, Yudiarti T, Isroli I (2016) Performances and haematological profile of broilers fed fermented dried cassava (*Manihot esculenta* Crantz). *Trop Anim Health Prod* 48:1337–1341. <https://cyberleninka.ru/article/n/a-review-on-fungal-fermented-cassava-pulp-as-a-cheap-alternative-feedstuff-in-poultry-ration>
- Sugiharto S, Yudiarti T, Isroli I (2019) Growth performance, haematological parameters, intestinal microbiology and carcass characteristics of broiler chickens fed two-stage fermented cassava pulp during finishing phase. *Trop Anim Sci J* 42:113–120. <https://journal.ipb.ac.id/index.php/tasj/article/view/24431>
- Tang DF, Ru YJ, Song SY, Choct M, Iji PA (2012) The effect of cassava chips, pellets, pulp and maize based diets on performance, digestion and metabolism of nutrients for broilers. <http://docsdrive.com/pdfs/medwelljournals/javaa/2012/1332-1337.pdf>
- Tesfaye E, Animut G, Urge M, Dessie T (2013) Cassava root chips as an alternative energy feed ingredient in broiler ration. *Int J Poult Sci* 12:298–306. <https://web.archive.org/web/20141005233041id/http://www.pjbs.org:80/ijps/fin2371.pdf>
- Trautenmüller H, Broch J, Pires Filho IC, Silva IM, Frank R, Schöne RA, Nunes RV (2020) Energetic values and inclusion levels of the dry residue of cassava in broiler diet. *Braz J Poult Sci* 22:eRBCA–2019. <https://doi.org/10.1590/1806-9061-2019-1183>
- Uchegbu MC, Ogbuwu IP, Obimara SO, Obua BE (2011) Effect of replacement of maize with cassava root meal fortified with palm oil on performance of starter broilers. *Online J Anim Feed Res* 1:171–175. <https://www.ojafr.com/main/attachments/article/78/OJAfr-%20A27.pdf>
- Valdivié M, Leyva C, Cobo R, Ortiz A, Dieppa O, Febles M (2008) Total substitution of corn by cassava (*Manihot esculenta*) meal in broiler chicken diets. *J Agric Sci* 42:61. <https://www.cabidigitallibrary.org/doi/full/https://doi.org/10.5555/20123007207>
- Vieira SS, Vieira ES, Barbosa FRS, Lima ADCS, Marinho AM, Reis CP, Neta ERS (2022) Cassava residues in the diet of slow-growing broilers. *Acta Amazon* 52:189–198. <https://doi.org/10.1590/1809-4392202100843>
- Yadav S, Mishra B, Jha R (2019) Cassava (*Manihot esculenta*) root chips inclusion in the diets of broiler chickens: effects on growth performance, ileal histomorphology and cecal volatile fatty acid production. *Poult Sci* 98:4008–4015. <https://doi.org/10.3382/ps/pez143>

Yang Y, Lei F, Zhang Z, Liu L, Li Q, Guo A (2024) Effects of cassava root meal on growth performance, apparent nutrient digestibility, organ and intestinal indices and slaughter performance of yellow-feathered broiler chickens. *Trop Anim Health Prod* 56:274. <https://doi.org/10.1007/s11250-024-04135-3>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.