

RESEARCH

Open Access



The study on interacting factors and functions of *GASA6* in *Jatropha curcas* L.

Xue Li^{1,2,3}, Ming-sheng Zhang³, Liang-qing Zhao⁴, Qian-qian Ling-hu³ and Gang Xu^{2*}

Abstract

Background The gibberellic acid-stimulated Arabidopsis (*GASA*) gene encodes a class of cysteine-rich functional proteins and is ubiquitous in plants. Most *GASA* proteins influence the signal transmission of plant hormones and regulate plant growth and development, however, their function in *Jatropha curcas* is still unknown.

Results In this study, we cloned *JcGASA6*, a member of the *GASA* family, from *J. curcas*. The *JcGASA6* protein has a *GASA*-conserved domain and is located in the tonoplast. The three-dimensional structure of the *JcGASA6* protein is highly consistent with the antibacterial protein Snakin-1. Additionally, the results of the yeast one-hybrid (Y1H) assay showed that *JcGASA6* was activated by *JcERF1*, *JcPYL9*, and *JcFLX*. The results of the Y2H assay showed that both *JcCNR8* and *JcSIZ1* could interact with *JcGASA6* in the nucleus. The expression of *JcGASA6* increased continuously during male flower development, and the overexpression of *JcGASA6* was associated with filament elongation of the stamens in tobacco.

Conclusion *JcGASA6*, a member of the *GASA* family in *J. curcas*, play an important role in growth regulation and floral development (especially in male flower). It is also involved in the signal transduction of hormones, such as ABA, ET, GA, BR, and SA. Also, *JcGASA6* is a potential antimicrobial protein determined by its three-dimensional structure.

Keywords Floral development, *GASA* family, *JcGASA6*

Background

The gibberellic acid-stimulated Arabidopsis (*GASA*) gene family is a plant-specific group of genes. This family comprises many genes, and most members are regulated by gibberellin (GA). By far, a great number of *GASA* genes have been isolated and identified from *Populus trichocarpa*, *Glycine max*, *Arabidopsis*, and *Solanum*

tuberosum [1–4]. The structure and function of this kind of proteins are deeply understood by the analysis of different *GASA* members identified from a variety of plant species. *GASA* protein family is a class of cysteine-rich functional proteins. They all have a highly conserved *GASA* domain (marked by 12 cysteines), which is essential for their normal function [1, 2].

The *GASA* proteins regulate plant growth and development, including seed germination, lateral root formation, stem elongation, flowering, flower and fruit development, biotic and abiotic stress responses, and signal transduction of hormones [5–9]. The functions of the genes in the *GASA* family have been mostly studied in *Arabidopsis*. Most *GASA* proteins are involved in hormone signal transduction. *GAST1*, *GASA4*, *GASA6*, *GASA9* and *GASA14* are involved in gibberellin signal transduction and are located downstream of DELLA protein [5, 10]. The *GASA* proteins are also involved abscisic acid

*Correspondence:

Gang Xu
xg335300@aliyun.com

¹ School of Chinese Ethnic Medicine, Guizhou Minzu University, Guiyang 550025, Guizhou, China

² School of Chinese Medicinal Resource, Guangdong Pharmaceutical University, Guangzhou 510006, Guangdong, China

³ School of Life Sciences/Key Laboratory of Plant Resource Conservation and Germplasm Innovation in Mountainous Region (Ministry of Education), Guizhou University, Guiyang 550025, Guizhou, China

⁴ Guizhou Botanical Garden, Guiyang 550005, China



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(ABA) signal transduction. The expression of *NtGASA1*, *NtGASA2*, *NtGASA9* and *AtGASA14* can be induced by ABA [5, 11]. However, no study has shown that proteins in the GASA family can perform ethylene (ET) signal transduction, although crosstalk between ET and gibberellin is common in seed germination, stem elongation, and flowering [12, 13]. Several GASA proteins also regulate flower development. For example, the overexpression of *AtGASA5* causes delayed flowering, whereas the mutation of *AtGASA5* leads to early flowering [14]. Functional deletions in *AtGASA6* and *AtGASA4* were found to cause late flowering, but early flowering was caused by only the overexpression of *AtGASA6* [6].

Jatropha curcas (Euphorbiaceae) is widely distributed in tropical and subtropical areas and is an ideal bioenergy crop for its oil-rich seeds that are especially rich in unsaturated fatty acids [15, 16]. However, due to low seed yield, the economic benefits of further expanding the *Jatropha*-based biodiesel industry are limited. The low ratio of female to male flowers (1:10–1:30) is a critical factor attributed to the low seed yield of *J. curcas* [15, 17]. The development of the flower of *J. curcas* has received much attention. In another study, using transcriptome data, we found that *JcGASA6*, a member of the GASA family, is differentially expressed in the flower buds during the development of the flowers of *J. curcas* [18]. However, the *JcGASA6* functions have not been further investigated.

In this study, *JcGASA6* was cloned from *J. curcas* and expressed in vitro. Upstream regulators of *JcGASA6* and interactive proteins with *JcGASA6* protein have been evaluated and screened. Moreover, we observed the expression pattern of *JcGASA6* at several key stages during flower development and further analyzed the phenotypic of overexpression *JcGASA6* in tobacco. Our results clarified the function of *JcGASA6* from the protein and gene levels, and also enriched the signal network of GASA protein in hormone crosstalk.

Results

Bioinformatics analysis and subcellular localization of *JcGASA6* protein

We obtained the coding sequence (CDS) of *JcGASA6* from cDNA library of flower bud. The CDS of *JcGASA6* contains 327bp and encodes 108 amino acids. The start codon is ATG and the stop codon is TAG (Fig. S1). Amino acid sequence alignment between *JcGASA6* and 14 members of the GASA family from *Arabidopsis* show that *JcGASA6* protein has a special sequence composed of 12 cysteines (C-3X-C-2X-RC-8X-C-3X-C-2X-2C-2X-C-X/2X-CV-2X-G-2X-G-4X-C-X/2X-CY-10X-KCP). This special sequence is a conserved domain of GASA family (Fig. 1A). The phylogenetic tree of protein sequences between *JcGASA6* protein and *Arabidopsis*

GASA family indicated that *JcGASA6* protein is most closely related to *AtGASA4* protein sequences (Fig. 1B). The first 24 amino acids at the N-terminal of *JcGASA6* protein were the putative signal peptides according to UniProt (Fig. S2). This suggests that *JcGASA6* may transfer after synthesis. Further study of the localization of *JcGASA6* protein. Under the excitation of 480nm, the results of laser confocal microscopy showed that the control protein (pBWA(V)HS-GFP) could emit light normally and located in the cell membrane, while the fusion protein (pBWA(V)HS-GASA6-GFP) could emit light in the tonoplast besides the cell membrane. Co-localization of the marker protein in tonoplast and the fusion protein showed that green and red fluorescence could be detected at tonoplast (Fig. 1C). This indicated that *JcGASA6* protein was located in the tonoplast.

Expression and identification of *JcGASA6* in vitro

To further study *JcGASA6* from the protein level, *JcGASA6* was cloned into pColdII vector, then was overexpressed in BL21 and ESLA to get *JcGASA6* protein. *JcGASA6* encodes a total of 108 amino acids and the signal peptide consists of the first 24 amino acids. Therefore, the molecular weight of the recombinant protein *JcGASA6* was 10.97kDa after adding the tag sequence of vector and removing the signal peptide. SDS-PAGE analysis showed that the expression products from BL21 (with pColdII-*JcGASA6*) or ESLA (with pColdII-*JcGASA6*) contain a 10.97kDa protein compared with the control (Fig. 2A). This indicates that *JcGASA6* was successfully expressed in BL21 and ESLA. Additionally, the 10.97kDa protein was distributed in the precipitates released from *E.coli* rupture (Fig. 2B) and was not detected in the supernatants of *E.coli* rupture (Fig. 2C). This indicates that *JcGASA6* was expressed in inclusion bodies way. The 10.97kDa protein was further identified by ESI-LC-MS/MS. The results showed that a total of 17 peptides from the 10.97kDa protein matched the amino acid sequence of *JcGASA6* protein (Table 1) and the coverage of amino acids match with *JcGASA6* protein reached 83% (Fig. 2D). Therefore, the 10.97kDa protein is *JcGASA6* protein.

JcGASA6 protein and Snakin-1 protein (crystal 5e5t.1A) have the highest homology according to the results from the SWISS-MODEL database. Therefore, taking the crystal of 5e5t.1A as the template, we get the speculative three-dimensional structure of *JcGASA6* protein after homologous modeling. Both *JcGASA6* protein and Snakin-1 protein have two important structures—short helices. The first short helix is composed of $\alpha 1$ and $\alpha 2$, and the second short helices is composed of $\alpha 3$ and $\alpha 4$ (Fig. 3A). These two structures are essential for the function of GASA family protein [19]. Moreover, like




Fig. 1 Amino acid sequence alignment analysis and subcellular localization of JcGASA6 protein. **A** Multiple amino acid sequence alignment of JcGASA6 with GASA family. **B** Phylogenetic analysis of the JcGASA6 protein. **C** The subcellular localization of JcGASA6 protein. Red arrow indicates the cell membrane, and white arrow indicates the tonoplast

Snakin-1 protein, JcGASA6 protein contains 12 cysteines. The 12 cysteine residues form 6 disulfide bonds (Fig. 3B, C), which is of great significance to maintaining the stability of protein three-dimensional structure [20].

Promoter sequence characteristics and upstream regulatory factors of JcGASA6

To further reveal the potential signal transduction pathway involved by JcGASA6, the promoter of JcGASA6 was isolated from *J. curcas* genome DNA. Except for conserved motifs (AT-TATATA-box, CAAT-box, and TATA-box), JcGASA6 promoter included several motifs with unknown function. Additionally, JcGASA6 promoter also included an ethylene-responsive motif which suggested the expression of JcGASA6 may be regulated by ET (Fig. S3). Then, the upstream regulatory factors

of JcGASA6 were screened by yeast one-hybrid (Y1H) system using JcGASA6 promoter as bait. Three genes, which may bind to the promoter of JcGASA6, were screened from cDNA library of *J. curcas* flower. The three genes included JcFLX-like (ID: XM_012232748.2), JcERF1 (ID: XM_012213416.2), and JcPYL9 (ID: XM_012227842.2). To confirm that the three genes can bind to the promoter of JcGASA6, the three genes were tested by Y1H individually. Self-activation of promoters is common, so 3-amino-1,2,4-triazole (3-AT) with suitable concentration is used to inhibit the self-activation of promoters [21]. In our study, 3-AT with a concentration of 130 mM completely inhibit the self-activation of JcGASA6 promoter (Fig. 4A). Additionally, the yeast cells, which contain pGADT7-JcFLX and JcGASA6-Pro-pHis2, pGADT7-JcPYL9 and JcGASA6-Pro-pHis2,




Fig. 2 Prokaryotic expression and identification of *JcGASA6*. **A** Whole bacteria SDS-PAGE analysis of *JcGASA6* expression. **B** SDS-PAGE analysis of precipitation from bacteria. **C** SDS-PAGE analysis of supernatant from bacteria. M, marker; lane 0, control (empty pColdII); lane 1–3, expression of *JcGASA6* in BL21 (DE3); lane 4–6, expression of *JcGASA6* in ES*L*a; lane 1 and 4, induction with 0.10 mM IPTG for 16 hours; lane 2 and 5, induction with 0.25 mM IPTG for 16 hours; lane 3 and 6, induction with 0.25 mM IPTG for 4 hours; red arrow indicates *JcGASA6* protein. **D** Mass spectrometry identification results of expression product from *JcGASA6*

Table 1 Seventeen peptides matched with amino acid sequence of *JcGASA6*

No.	position	Matched peptide sequence	MW(Da)		Error rate/Da
			Calculation	Observation	
1	1–26	MNHKVHHHHHHQVMANDTQYHLDSGR	3165.4383	3166.4219	−0.0237
2	27–43	YGPGLKSYQCPSECTR	1988.8720	1989.8728	−0.0066
3	34–43	SYQCPSECTR	1230.4878	1230.4884	0.0006
4	34–43	SYQCPSECTR	1287.5093	1287.5027	−0.0066
5	34–43	SYQCPSECTR	1287.5093	1287.5027	−0.0066
6	34–44	SYQCPSECTRR	1386.5889	1386.574	−0.0149
7	34–44	SYQCPSECTRR	1443.6104	1443.6012	−0.0092
8	34–44	SYQCPSECTRR	1443.6104	1443.6012	−0.0092
9	34–44	SYQCPSECTRR	1329.5675	1329.5389	−0.0286
10	48–60	TQYHKPCMFFCQK	1717.7648	1717.7571	−0.0077
11	48–60	TQYHKPCMFFCQK	1774.7863	1774.7654	−0.0209
12	48–60	TQYHKPCMFFCQK	1774.7863	1774.7654	−0.0209
13	48–60	TQYHKPCMFFCQK	1790.7812	1790.7656	−0.0156
14	65–76	CLCVPPPGFYGNK	1410.6424	1411.6293	−0.0205
15	77–86	SVCPCYNNWK	1270.5344	1270.5083	−0.0261
16	77–86	SVCPCYNNWK	1327.5559	1327.5404	−0.0155
17	77–86	SVCPCYNNWK	1327.5559	1327.5404	−0.0155

or pGADT7-*JcAP2* and *JcGASA6*-Pro-pHis2, can grow normally on SD/−His-Leu-Trp medium with 130 mM 3-AT (Fig. 4A). These results indicated that all of *JcFLX*, *JcPYL9*, and *JcERF1* could interact with *JcGASA6* promoter to regulate the expression of *JcGASA6*. Furthermore, the regulation of three genes on *JcGASA6* was




Fig. 3 Comparison of JcGASA6 protein and Snakin-1 protein. **A** Amino acid sequence alignment of JcGASA6 protein and Snakin-1 protein. **B** Speculative three-dimensional structure of JcGASA6 protein. **C** Three-dimensional structure of Snakin-1 protein

determined by dual-luciferase assays. The results of dual-luciferase assays showed that all of JcFLX, JcPYL9, and JcERF1 could enhance the activity of Luc driven by *JcGASA6* promoter and the activity of LUC was the strongest in *JcERF1/JcGASA6* group (Fig. 4B). This suggested that all of the three genes (*JcFLX*, *JcPYL9* and *JcERF1*) could interact with *JcGASA6* promoter to activate the expression of *JcGASA6*, and *JcERF1* has the strongest activation effect on *JcGASA6*.

Proteins interacting with JcGASA6 protein

In addition to screening the upstream regulators of *JcGASA6*, we also screened the interacting proteins of *JcGASA6* protein by using yeast two-hybrid (Y2H) system. On SD/−Ade/−His/−Leu/−Trp medium, the yeast cells with pGBKT7-53 and pGADT7-T plasmids can grow normally, while the yeast cells with pGBKT7 and pGADT7 plasmids (negative control) and yeast cells with pGBKT7-*JcGASA6* and pGADT7 plasmids can not grow normally (Fig. S4). These results indicated that *JcGASA6* protein without transcriptional self-activation activity.

JcGASA6 was used as the bait to screen its interacting proteins from flower bud cDNA library, and five proteins interacting with *JcGASA6* were obtained. The five proteins include JcCNR8 (*J. curcas* cell number regulator 8, ID: XM_012226926.2), JcAMs (*J. curcas* transcription factor ABORTED MICROSPORES, ID: XM_020682349.1), JcAPRR2 (*J. curcas* two-component response regulator-like APRR2, ID: XM_012218861.2), JcFRI (*J. curcas* FRIGIDA-like protein 4a FRI, ID: XM_012214239.2) and JcSIZ1 (*J. curcas* E3 SUMO-protein ligase SIZ1, ID: XM_012209470.2). To verify the interaction between the five proteins with *JcGASA6*, we used Y2H assay. Yeast cells, which containing JcCNR8 and *JcGASA6*, JcAMs and *JcGASA6*, JcAPRR2 and *JcGASA6* or JcSIZ1 and *JcGASA6*, grew normally on SD/−Trp/−Leu/−Ade/−His and SD/−Trp/−Leu/−Ade/−His (X-a-gal) medium, while the yeast cells with JcFRI and *JcGASA6* could not grow (Fig. 5A). These results suggested four proteins JcCNR8, JcAMs, JcAPRR2, and JcSIZ1 may interact with *JcGASA6* protein. BiFC assay was carried out to further confirm these results. The BiFC assay show in Fig. 5B, the yellow fluorescence could be detected only when




Fig. 4 All of JcFLX, JcPYL9 and JcERF1 binds to the promoter of *JcGASA6* and activate its expression. **A** Yeast one-hybrid assay shows all of JcFLX, JcPYL9 and JcERF1 bind to *JcGASA6* promoter. **B** Dual-luciferase assays showed all of JcFLX, JcPYL9 and JcERF1 can activate the expression of *JcGASA6*. Diverse lowercase letters indicate significant differences ($P < 0.05$). The significance of difference was analyzed by Tukey's test

the protein pair of JcCNR8 and JcGASA6 or JcSIZ1 and JcGASA6 were transiently co-expressed in leaf epidermal cells, and the yellow fluorescence was detected in the nucleus (Fig. 5B). These results indicated that both JcCNR8 and JcSIZ1 could interact with JcGASA6 in nucleus.

Expression pattern of *JcGASA6* in the flower development of *J. curcas*

Our previous studies suggest that *JcGASA6* may also be involved in flower development [18], we further

analyzed the expression pattern of *JcGASA6* during flower development. In order to accurately judge the expression pattern of *JcGASA6* during flower development of *J. curcas*, the flower of *J. curcas* was observed at the morphological and histological. When the bud length of *J. curcas* is about 0.15 mm, the primordium of petal has appeared, but the sexual differentiation of flower has not yet begun (Fig. 6St0). During the development of female flowers, the carpel did not fully heal when the ovary length is about 0.80 mm (Fig. 6ST1). Histological




Fig. 5 Identification of proteins interacted with JcGASA6. **A** Y2H assay to verify the interactions of JcGASA6 with JcCNR8, JcAMs, JcAPRR2, JcFRI and JcSIZ1. **B** BiFC assay of the interactions of JcGASA6 with JcAMs, JcAPRR2, JcCNR8 and JcSIZ1; Bright: bright field without fluorescence signal; NLS: blue fluorescent signal located in the nucleus; YFP: yellow fluorescent signal of interacting proteins; Merged: Overlap of yellow and blue fluorescence signals in bright field; Bars: 48 μm




Fig. 6 Morphological and histological observation on flower of *J. curcas*. ST1 to ST4 represent the morphology of female flower and the corresponding histological structures are St1 to St4. ST5 to ST8 represent the morphology of male flower, and the corresponding histological structures are St5 to St8. St0, represent the morphology of undifferentiated flower. **A** Expression profile of *JcGASA6* during the development of flower, asterisks indicate a significant difference, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Ca, carpel; Pp, petal primordium; S, stamen; Me, megaspore; Fm, functional megaspore; Mes, mononuclear embryo sac; Mi, microsporocyte; Snp, single nucleus pollen; Mp, mature pollen. Bars = 50 μ m for histological structures of flower. Bars = 270 μ m for morphology of female flower. Bars = 20 μ m in St0

analysis showed that the female flower was at the stage of megasporocyte (Fig. 6St1). After that, the megasporocyte undergoes two meiosis to form functional megaspore (Fig. 6ST2), at this time, the carpel has basically healed (Fig. 6St2), and the ovary length was about 1.10 mm. The functional megaspore develops further and enters the mononuclear embryo sac stage (Fig. 6St3), the ovary length was about 1.50 mm (Fig. 6ST3). Finally, functional megaspores formed mature embryo sac with 8-core 7 cells through three mitoses (Fig. 6St4), and the ovary length was about 3.20 mm (Fig. 6ST4). During the development of male flower, the male flower was in the stage of microspore mother cell when the flower bud length is about 0.50 mm (Fig. 6St5, ST5). After two times of meiosis, the microsporocyte entered the tetrad stage (Fig. 6St6), the length of bud was 1.20 mm (Fig. 6ST6). Microsporocyte was released to the anther chambers to form single nucleus pollen (Fig. 6St7), the length of the bud was 2.00 mm (Fig. 6ST7). Finally, single nucleus pollen undergoes nuclear division to form mature pollen with two-cell (Fig. 6St8), the length of bud was about 3.00 mm (Fig. 6ST8). In female flowers, the expression of *JcGASA6* gradually increased from the undifferentiated stage (St0) to the megasporocyte meiosis stage (St2) and reached the highest lever at the megasporocyte meiosis stage. After that, the expression of *JcGASA6* gradually decreased and reached the lowest level at the mature embryo sac stage (St4). These results suggested that *JcGASA6* plays an important role in the early development of female flower. In male flowers, the expression of *JcGASA6* increased from undifferentiated stage (St0) to mature pollen stage (St8) and reached the highest level at mature pollen stage. These results indicate that *JcGASA6* plays an important role in the whole development of male flower. Moreover, the expression of *JcGASA6* was not significantly different in the early development between female (St1-St2) and male flower (St5-St6), while in the late development of female (St3-St4) and male flowers (St7-St8), the expression of *JcGASA6* increased significantly in male flowers. This suggested that *JcGASA6* was more important

for the development of male flower than female flower (Fig. 6A).

Overexpression of *JcGASA6* promoted the elongation of stamen filament in tobacco

The expression pattern of *JcGASA6* during flower development suggests that *JcGASA6* plays an important role in flower development, especially in male flower development. To further reveal the role of *JcGASA6* in flower development, overexpression of *JcGASA6* in tobacco (wild-type tobacco were controls, WT). There was no significant difference in plant height, ground diameter and style length between wild-type tobacco and transgenic tobacco (Table 2). However, the stamen of transgenic tobacco (TR) was higher than the stigma of ovary, and the filament length of TR tobacco was longer than WT tobacco (Table 2 and Fig. 7C, D). This result suggested that *JcGASA6* could promote the elongation of stamen filament. Moreover, the leaves of TR tobacco are longer and wider than those of WT tobacco (Table 2 and Fig. 7A, B). This result indicated that *JcGASA6* was also involved in plant growth.

Discussion

Gibberellic Acid-stimulated Arabidopsis (GASA) protein, also known as Snakin protein, is a kind of CRP (cysteine-rich peptides) protein [8]. GASA family proteins have been found in many plants and have many members, characterize by a conserved domain containing 12 cysteines [11]. In present study, *JcGASA6* encoded a protein identified as *JcGASA6* protein. *JcGASA6* is a member of GASA family, suggested by its typical domain containing 12 cysteines (Fig. 1A). It was with a length of 108 amino acids and a molecular weight of 10.97 KD, containing a signal peptide at N-terminal. It showed the closest genetic relationship with *AtGASA4* (Fig. 1B).

Antimicrobial peptides (AMPs) are excellent candidate drugs against drug-resistant pathogens. The structure of AMPs, especially an α -helical hairpin structure, plays an important role in killing pathogens [20]. Snakin is the only GASA member with antibacterial effect [3, 22]. Two short helices structures (dbHTH) in Snakin-1 protein are similar to the α -helical hairpin in antimicrobial peptide EcAMP1 [23]. The short helices and six disulfide bonds

Table 2 Comparison of growth and development between wild-type tobacco and transgenic tobacco

Sample	Plant height (cm)	Ground diameter (mm)	Leaf length (cm)	Leaf width (cm)	Style length (cm)	Filament length (cm)
WT	29.18 ± 1.52a	9.79 ± 0.38a	24.00 ± 0.36a	12.59 ± 0.06a	5.31 ± 0.03a	4.93 ± 0.06a
TR	29.28 ± 1.64a	10.00 ± 0.51a	31.23 ± 0.37b	15.40 ± 0.38b	5.31 ± 0.03a	5.61 ± 0.07b

Different lowercase letters in each column indicate the significance between wild-type tobacco and transgenic tobacco ($P < 0.05$)




Fig. 7 Phenotype of tobacco overexpressing *JcGASA6*. **A** and **B** plant shape of wild-type tobacco and transgenic tobacco. **C** The flower of wild-type tobacco. **D** The flower of transgenic tobacco. Sf, stamen filament, S, stigma. Bars = 1.8 cm in **B**, **C**, **D**. Bars = 7.5 cm in **A**

can form a large positive electrostatic surface (Fig. 3C). The positive electrostatic surface makes Snakin-1 protein play an antibacterial role [20]. Similar to the Snakin-1 protein, the speculative three-dimensional structure of JcGASA6 protein have two important structures—short helices and six disulfide bonds (Fig. 3), which might endow it with an antibacterial role.

Protein localization is often closely related to its function. Most GASA proteins have signal peptides at the N-terminal, and they are located in the plasma membrane or cell wall. AtGASA14 protein is located in the plasma membrane which will make it easier to regulate the balance of reactive oxygen [5]. GhGASA10 (a GASA protein in Cotton) was localized in the plasma membrane, which conducive to synthesize proteins in the cell wall to promotes the cell elongation in fiber [24]. AtGASA6 protein is located in the cell wall, which helps to regulate the elongation of hypocotyl and seed germination [13]. In our study, The N-terminal of JcGASA6 protein also has a signal peptide (Fig. S3). But unlike other members of the GASA protein, JcGASA6 is located at the tonoplast (Fig. 1C). This suggests that the function

of JcGASA6 protein may be different from other members of GASA family.

GASA family is firstly thought to play important role in gibberellin signal pathway. AtGASA6 and AtGASA4 are located the downstream of DELLA protein in GA signal pathway [10]. To data, most members of the GASA family are found to be widely involved in hormone signal transduction, such as ABA, ET, GA, BR (brassinolide), SA (salicylic acid). OsGSR1 control BR biosynthesis to regulate salt stress [25]. Snakin-1 is involved in hormonal balance and Snakin-1 silencing enhanced GA and SA levels [26]. AtGASA6 is down-regulated by ABA and can integrate GA and ABA to promote cell elongation and seed germination [13]. Our study also found that ABA receptor JcPYL9 protein could directly activate the expression of JcGASA6 (Fig. 4), supporting that the members of GASA family would be involved in ABA signal transduction. A few evidence showed that members of GASA protein family were involved in ET signal transduction. FaGAST2, a member of GASA family in strawberry, was up-regulated by ethephon [7]. ERF1 is the target gene of the core transcription factor EIN3 of the ET signaling

pathway. EIN3 protein can activate the expression of *ERF1* through binding to the promoter of *ERF1* [27]. Our results showed that the expression of *JcGASA6* could be activated by ERF1 protein in *J. curcas* (Fig. 4), which further confirmed that GASA protein family is involved in ET signal transduction.

AtGASA4 (closeted genetic relationship with *JcGASA6* protein) and *AtGASA6* (the homologous protein of *JcGASA6*) are involved in flower development. The *gasa4* mutant showed more leaf buds before generating flowers, but overexpression of *AtGASA4* could not promote flowering transformation. This indicates that *AtGASA4* is not enough to induce flowering transformation [28]. *AtGASA4* and *AtGASA6* also affect flowering time, but *AtGASA6* plays a predominant role in causing early flowering [6]. *FLX* is reported to be involved in the regulation of early flowering [29]. In present study, *FLX* could directly activate the expression of *JcGASA6*. Although we did not observe abnormal flowering in transgenic tobacco, overexpression of *JcGASA6* caused elongated filaments in tobacco (Figs. 6 and 7C). Furthermore, the expression of *JcGASA6* increased continuously during male flower development. All these results suggested that *JcGASA6* would contribute to the development of male flowers, especially the filaments of stamens. This affection of *JcGASA6* on flower is different from other *GASA* family (such as *AtGASA4* and *AtGASA6*). On the other hand, *JcGASA6* protein could interact with *JcCNR8* protein. *CNR* can control organ size by regulating cell number [30]. These would explain that the leaf size of transgenic tobacco was larger than wild-type tobacco (Fig. 7A, B) (Table 3). Therefore, *JcGASA6* is also involved in growth regulation. Also, this is the first time to find that the *GASA* family can interact with *CNR* protein, which may support *AtGASA10* facilitate wall growth [4].

Additionally, the *JcGASA6* protein also could interact with SUMO E3 ligase (*JcSIZ1*). SUMO (Small

Ubiquitin-like Modifier) E3 ligase is an important member involved in protein SUMO modification. SUMO E3 ligase recruits target proteins and promote the binding of SUMO to the target protein, then the activity of the target protein changes reversibly [31, 32]. Therefore, the activity of *JcGASA6* protein may be modified by SUMO.

Conclusions

JcGASA6, a member of the *GASA* family, has an antibacterial structure similar to the antibacterial peptide Snakin-1, and is located in the tonoplast. In addition, *JcGASA6* can be activated by the ABA receptor *JcPYL9* and the core transcription factor *JcERF1*, which supports that *JcGASA6* can participate in ABA and ET signal transduction. Meanwhile, *JcGASA6* is activated by the early flowering gene *FLX*, and its expression increases continuously during male flower development. Abnormal flowering was not observed after overexpressing *JcGASA6*, but the elongation of stamen filaments was caused by the overexpression of *JcGASA6*. These results confirmed that *JcGASA6* is involved in flower development. Additionally, overexpression of *JcGASA6* increased the leaf size in transgenic plants, and *JcGASA6* can interact with *CNR* (a protein with the regulation cell number). These results indicate that *JcGASA6* is also involved in growth regulation. Moreover, *JcGASA6* interacts with ubiquitin ligase *SIZ1* also implies that its activity may be modified by SUMO. All in all, our results have revealed the functions of *JcGASA6* from multiple perspectives.

Methods

Plant materials and flower collection

The flower buds of *J. curcas* L. were collected from Zhenfeng, Guizhou Province, China (36°14,050.2'N, 87°51,047.8'E). Flower buds for morphological and microscopic observation were temporarily stored in the mixture of acetaldehyde acetic acid 50% alcohol (4: 6: 90, v/v), and for RNA extraction were temporarily stored in RNAlocker (Tiandz, Inc., Beijing China). All samples were placed on ice.

JcGASA6 isolation and sequence analysis

The full-length cDNA of *JcGASA6* was cloned from flower bud by RACE-pcr, then the obtained cDNA sequences were aligned in NCBI database (Accession number: KU500008).

Sequence of the *GASA* family of *Arabidopsis thaliana* was obtained from NCBI database (Table S1). Multiple sequence alignment used DNAMAN, phylogenetic tree constructed by MEGA 6.0, and signal peptide predicted by UniProt. (<https://www.uniprot.org/peptidesearch/>).

Table 3 Parameters of developmental stages of J. curcas flower

Developmental stage	Flower organ size/mm
Undifferentiated stage (St0)	0.15 ± 0.01
♀ Megasporocyte stage (St1)	0.80 ± 0.03
♀ Megasporocyte meiosis stage (St2)	1.00 ± 0.06
♀ Mononuclear embryo sac stage (St3)	1.50 ± 0.03
♀ Mature embryo sac stage (St4)	3.20 ± 0.04
♂ Microsporocyte stage (St5)	0.50 ± 0.01
♂ Tetrad stage (St6)	1.20 ± 0.03
♂ Single nucleus pollen stage (St7)	2.00 ± 0.05
♂ Mature pollen stage (St8)	3.00 ± 0.05

Subcellular localization, expression and identification of JcGASA6 protein

The full-length cDNA of *JcGASA6* was fused with the pBWA(V)HS-osGFP then recombinant plasmid pBWA(V)HS-*JcGASA6*-osGFP was transfected transformed into rice protoplasts by PEG (polyethylene glycol). The protoplasts were observed by confocal laser microscope under the excitation of 480 nm wavelength after dark culture at 28 °C for 48 hours (FV10-ASWO-LYMPUS, Japan) [33]. Gamatip protein located in the tonoplast was used as a marker [34]. All primers used for subcellular localization are listed in Table S2.

The coding sequence of *JcGASA6* was amplified by specific primers (*JcGASA6*ex-F and *JcGASA6*ex-R), then the coding sequence was connected to the expression vector pCold II after digesting with Nde I and Hind III. The sequence of recombinant plasmid *JcGASA6*-pCold II was examined using vector primers (pCold II-F and pCold II-R) (Table S2). The recombinant plasmid was transformed into *E. coli* competent cells (BL21 or ESLA) to overexpress *JcGASA6*, and the empty vector pCold II was transformed into BL21 or ESLA as the control. Then the BL21 or ESLA were cultured at 16 °C, and isopropyl β-D-thiogalactoside (IPTG) was used as an expression inducer. The details of method same as previously reported [35].

The expression products of *JcGASA6* in *E. coli* were analyzed by SDS-PAGE electrophoresis. Cut off the SDS-PAGE glue containing *JcGASA6* protein, and wash the SDS-PAGE glue with ultrapure water and decolorization with acetonitrile mixture. The decolorized SDS-PAGE glue was digested by trypsin overnight at 37 °C to form enzymolysis solution, then the enzymolysis solution was identified by LC-MS/MS method. The method was consistent with the previously reported [36]. The three-dimensional structure of protein was analyzed by the SWISS-MODEL database (<https://www.swissmodel.expasy.org/interactive>).

Promoter isolation and analysis, construction of cDNA library for yeast-hybrid system

The promoter sequence of *JcGASA6* was obtained from the published genomic database (ID: 105640538), and PlantCARE was used for promoter sequence analysis (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>). The cDNA library of Yeast-Hybrid System was constructed by mixed RNA from flowers at different developmental stages. The primary library was constructed using attB2 as linker and ATTB-A1, ATTB1-B and ATTB1-C as primers. The clone number of primary library was 8.04×10^6 cfu. The plasmid of primary library was extracted and transferred into DH10B by the

electrotransfer method, and then the secondary library was obtained. The clone number of secondary library was 1.31×10^7 cfu. The method is consistent with that previously reported [37] (Table S3).

Yeast-one hybrid (Y1H) and dual-luciferase assay

The promoter of *JcGASA6* was amplified using specific primer pro-*JcGASA6*-F/R, then fusion to the pHIS2 (Table S4). Co-transferred fusion plasmid and secondary library to Y187 yeast system, then screen the upstream regulator of *JcGASA6*. The screening process refers to the previous method [38]. Three upstream regulators (*JcFLX*, *JcERF1* and *JcPYL9*) were screened from the secondary library. The interaction between *JcGASA6* promoter and these three regulators was verified one by one with Y187 yeast system [38]. pGADT7 and p53-pHis2 were co-transformed into Y187 as negative controls. pGADT7-53 and p53-pHis2 were co-transformed into Y187 as positive controls. The promoter self-activation of *JcGASA6* was detected using Y187 with plasmids pGADT7 and *JcGASA6*-Pro-pHis2.

The primer *JcGASA6*-luc-F/R was used to construct pGreenII 0800-*JcGASA6*-luc, and three primers (AP2-F/R, FLX-F/R and PYL9-F/R) was used to construct regulators (pGreenII 62-*JcFLX*-SK, pGreenII 62-*JcERF1*-SK and pGreenII 62-PYL9-SK). Co-transferred pGreenII 0800-pro-*JcGASA6*-luc and regulator into tobacco leaves, then detected the fluorescence value (Dual-Luciferase Assay System, Promega) (Table S4) [38]. pGreenII 62-SK and pGreenII 0800-Luc were co-transformed into tobacco leaves as negative controls.

Yeast-two hybrid (Y2H) and bimolecular fluorescence complementation (BiFC) assay

Full-cDNA *JcGASA6* fusion with pGBKT7 by using *GASA6*-GBK-F/R primer, then both of pGBKT7-*JcGASA6* and pGADT7-AD were transferred into AH109 by LiAc method, then the AH109 was cultured for the detection of self-activation activity. Co-transferred pGBKT7-*JcGASA6* and secondary library to AH109 for screening the interaction proteins of *JcGASA6*. Five proteins were screened, including *JcCNR8*, *JcAMs*, *JcAPRR2*, *JcFRI* and *JcSIZ1*. The interaction between the five proteins and *JcGASA6* were verified by one-to-one in AH109 [39] (Table S5). pGADT7 and pGBKT7 were co-transformed into AH109 as negative controls. pGADT7-53 and pGBKT7-T were co-transformed into AH109 as positive controls.

Full-cDNA *JcCNR8*, *JcSIZ1*, *JcAMs*, and *JcAPRR2* were fused with PSPYCE-35S respectively by using specific primer, and Full-cDNA *JcGASA6* was fused with PSPYNE-35S. Co-transferred PSPYNE-35S-*JcGASA6* and

PSPYCE-35S-JcCNR8/PSPYCE-35S-JcSIZ1/PSPYCE-35S-JcAMs/PSPYCE-35S-*JcAPRR2* into EHA105. The five types of EHA105 infected tobacco leaves respectively, then the fluorescence signal was detected 72 hours after infection [39] (Table S5). PSPYCE-35S and PSPYNE-35S were co-transformed into tobacco leaves as negative controls. PSPYCE-35S-bZIP63 and PSPYNE-35S-bZIP63 were co-transformed into tobacco leaves as positive control.

Morphology and microscopic observation of flower

Flower buds were classified according to their length, dissected under stereoscope, observed and photographed. Since the flower bud of undifferentiated stage is too small to be observed clearly under stereoscope, it was observed under the scanning electron microscope. The undifferentiated flower were fixed in the mixture of formaldehyde-acetic acid-50% ethanol, and then dissected and observed by electron microscope [18]. According to the classification, paraffin sections of these flower buds were made [40] (Table 3).

Expression of *JcGASA6* during flower development determined by qRT-PCR

The total RNA of flower was extracted by RNA isolation kit (Omega Bio-Tek, Beijing, China), qualified RNA was used to synthesize the first strand cDNA (Takara, Beijing, China). PCR amplification was used Bio-Rad CFX system (Bio-Rad, USA). *Beta-tubulin* and *actin* as internal control. The $2^{-\Delta\Delta CT}$ method was used to calculate the relative expression of *JcGASA6* [41]. All primers used for PCR amplification are listed in Table S6 and each sample reaction was repeated three times.

Overexpression of *JcGASA6* in *Nicotiana tabacum* L.

Fusion full-length cDNA of *JcGASA6* with the pBWA(V)KS-GUS. The recombinant plasmid (pBWA(V)KS-*JcGASA6-GUS*) was transformed into agrobacterium tumefaciens (GV3101), then positive agrobacterium tumefaciens were screened. Tobacco (K326) leaves infected by GV3101 were cultured in dark at 25°C for 2 days. After tobacco leaves were differentiated into seedlings, positive tobacco was detected (Fig. S5), and wild-type tobacco was used as control [42] (Table S6).

Abbreviations

GASA	Gibberellic acid-stimulated Arabidopsis
GA	Gibberellin
ABA	Abscissic Acid
ET	Ethylene
SA	Salicylic Acid
BR	Brassinolide
CDS	Coding Sequence
Y1H	Yeast One-Hybrid

3-AT	3-Amino-1,2,4-Triazole
Y2H	Yeast Two-Hybrid
WT	Wild-Type
TR	Transgenic
AMPs	Antimicrobial peptides
SUMO	Small Ubiquitin-like Modifier

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12870-023-04067-4>.

Additional file 1.

Additional file 2.

Acknowledgements

We are grateful to XIAN CHUNFENG BIOTECHNOLOGY CO., LTD for their excellent technical assistance.

Authors' contributions

Xue Li, Ming-sheng Zhang, Xu Gang conceived and designed the research, Xue Li, Liang-qing Zhao, Qian-qian Ling-hu performed the experiments and analyzed the data. Xue Li wrote the paper. Xu Gang, supervision, funding acquisition. All authors read and approved the manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (No. 31760198), and the Scientific Research Project of Ordinary Undergraduate Colleges and Universities of Guizhou Provincial Department of Education (Qian Jiao Ji[2022]No. 145).

Availability of data and materials

All data generated or analyzed in our study are available in this article and its supplementary information files. Gene sequences can be down-loaded at NCBI database (<https://www.ncbi.nlm.nih.gov/>). The GenBank accession number of *JcGASA6* is KU500008, *AtGASA1* is P46689.2, *AtGASA2* is P46688.1, *AtGASA3* is P46687.1, *AtGASA4* is P46690.2, *AtGASA5* is Q84J95.1, *AtGASA6* is Q6NMQ7.1, *AtGASA7* is O82328.1, *AtGASA8* is O80641.1, *AtGASA9* is Q8GWK5.1, *AtGASA10* is Q8LFM2.1, *AtGASA11* is F4IQJ4.1, *AtGASA12* is Q6GKX7.1, *AtGASA13* is A8MR46.1, and *AtGASA14* is Q9LFR3.1.

Declarations

Ethics approval and consent to participate

Before conducting the research, the permission from research institute of forest resources and environment of Guizhou University to collect and analyze the *J. curcas* L samples documented in this work was obtained. All the methods complied with Chinese regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Received: 13 July 2022 Accepted: 16 January 2023

Published online: 18 February 2023

References

1. Wu K, Qu Y, Rong H, Han X, Tian Y, Xu L. Identification and Expression Analysis of the *Populus trichocarpa* GASA-Gene Family. Int J Mol Sci. 2022;23(3):1507.

2. Ahmad MZ, Sana A, Jamil A, Nasir JA, Ahmed S, Hameed MU, Abdullah: a genome-wide approach to the comprehensive analysis of GASA gene family in *Glycine max*. *Plant Mol Biol*. 2019;100(6):607–20.
3. Almasia NI, Molinari MP, Maroniche GA, Nahirnak V, Barrios Baron MP, Taboga OA, et al. Successful production of the potato antimicrobial peptide Snakin-1 in baculovirus-infected insect cells and development of specific antibodies. *BMC Biotechnol*. 2017;17(1):75.
4. Trapalis M, Li SF, Parish RW. The *Arabidopsis* GASA10 gene encodes a cell wall protein strongly expressed in developing anthers and seeds. *Plant Sci*. 2017;260:71–9.
5. Sun SL, Wang HX, Yu HM, Zhong CM, Zhang XX, Peng JZ, et al. GASA14 regulates leaf expansion and abiotic stress resistance by modulating reactive oxygen species accumulation. *J Exp Bot*. 2013;64(6):1637–47.
6. Qu J, Kang SG, Hah C, Jang JC. Molecular and cellular characterization of GA-stimulated transcripts GASA4 and GASA6 in *Arabidopsis thaliana*. *Plant Sci*. 2016;246:1–10.
7. Moyano-Canete E, Bellido ML, Garcia-Caparrós N, Medina-Puche L, Amil-Ruiz F, Gonzalez-Reyes JA, et al. FaGAST2, a strawberry ripening-related gene, acts together with FaGAST1 to determine cell size of the fruit receptacle. *Plant Cell Physiol*. 2013;54(2):218–36.
8. Zhang S, Wang X. One new kind of phytohormonal signaling integrator: up-and-coming GASA family genes. *Plant Signal Behav*. 2017;12(2):e1226453.
9. Fan S, Zhang D, Zhang L, Gao C, Xin M, Tahir MM, et al. Comprehensive analysis of GASA family members in the *Malus domestica* genome: identification, characterization, and their expressions in response to apple flower induction. *BMC Genomics*. 2017;18(1):827.
10. Zhang SC, Wang XJ. Expression pattern of GASA, downstream genes of DELLA, in *Arabidopsis*. *Chin Sci Bull*. 2008;53(24):3839–46.
11. Li Z, Gao J, Wang G, Wang S, Chen K, Pu W, et al. Genome-wide identification and characterization of GASA gene family in *Nicotiana tabacum*. *Front Genet*. 2021;12:768942.
12. Fu X, Gao X, Liu X. Integration of ethylene and gibberellin signaling. Springer Netherlands. 2015;1:153–73.
13. Zhong CM, Xu H, Ye ST, Wang SY, Li LF, Zhang SC, et al. Gibberellic acid-stimulated *Arabidopsis*6 serves as an integrator of gibberellin, Absciscic acid, and glucose signaling during seed germination in *Arabidopsis*(1[OPEN]). *Plant Physiol*. 2015;169(3):2288–303.
14. Zhang S, Yang C, Peng J, Sun S, Wang X. GASA5, a regulator of flowering time and stem growth in *Arabidopsis thaliana*. *Plant Mol Biol*. 2009;69(6):745–59.
15. Albuquerque N, Garcia-Almodovar RC, Valverde JM, Burgos L, Martinez-Romero D. Characterization of *Jatropha curcas* accessions based in plant growth traits and oil quality. *Ind Crop Prod*. 2017;109:693–8.
16. Qian JF, Shi HX, Yun Z. Preparation of biodiesel from *Jatropha curcas* L. oil produced by two-phase solvent extraction. *Bioresour Technol*. 2010;101(18):7025–31.
17. Gangwar M, Shankar J. Molecular mechanisms of the floral biology of *Jatropha curcas*: opportunities and challenges as an energy crop. *Front Plant Sci*. 2020;11:609.
18. Xu G, Huang J, Yang Y, Yao YA. Transcriptome Analysis of Flower Sex Differentiation in *Jatropha curcas* L. Using RNA Sequencing. *PLoS One*. 2016;11(2):e0145613.
19. Zhang M, Cheng W, Wang J, Cheng T, Zhang Q. Genome-Wide Identification, Evolution, and Expression Analysis of GASA Gene Family in *Prunus mume*. *Int J Mol Sci*. 2022;23(18):10923.
20. Yeung H, Squire CJ, Yosaatmadja Y, Panjikar S, Lopez G, Molina A, et al. Radiation damage and racemic protein crystallography reveal the unique structure of the GASA/Snakin protein superfamily. *Angew Chem Int Ed Engl*. 2016;55(28):7930–3.
21. Deng CY, Wang JY, Lu CF, Li YF, Kong DY, Hong Y, et al. CcMYB6-1 and CcbHLH1, two novel transcription factors synergistically involved in regulating anthocyanin biosynthesis in cornflower. *Plant Physiol Bioch*. 2020;151:271–83.
22. Oliveira-Lima M, Benko-Iseppon AM, Neto J, Rodriguez-Decuadro S, Kido EA, Crovella S, et al. Snakin: structure, roles and applications of a plant antimicrobial peptide. *Curr Protein Pept Sci*. 2017;18(4):368–74.
23. Barashkova AS, Ryazantsev DY, Rogozhin EA. Rational Design of Plant Hairpin-like Peptide EcAMP1: Structural-Functional Correlations to Reveal Antibacterial and Antifungal Activity. *Mol*. 2022;27(11):3554.
24. Chen B, Sun Y, Tian Z, Fu G, Pei X, Pan Z, et al. GhGASA10-1 promotes the cell elongation in fiber development through the phytohormones IAA-induced. *BMC Plant Biol*. 2021;21(1):448.
25. Li Q, Xu F, Chen Z, Teng Z, Sun K, Li X, et al. Synergistic interplay of ABA and BR signal in regulating plant growth and adaptation. *Nat Plants*. 2021;7(8):1108–18.
26. Nahirnak V, Rivarola M, Almasia NI, Barrios Baron MP, Hopp HE, Vile D, et al. Snakin-1 affects reactive oxygen species and ascorbic acid levels and hormone balance in potato. *PLoS One*. 2019;14(3):e0214165.
27. Binder BM. Ethylene signaling in plants. *J Biol Chem*. 2020;295(22):7710–25.
28. Roxrud I, Lid SE, Fletcher JC, Schmidt ED, Opsahl-Sorteborg HG. GASA4, one of the 14-member *Arabidopsis* GASA family of small polypeptides, regulates flowering and seed development. *Plant Cell Physiol*. 2007;48(3):471–83.
29. Shen L, Zhang Y, Sawettalake N. A molecular switch for FLOWERING LOCUS C activation determines flowering time in *Arabidopsis*. *Plant Cell*. 2022;34(2):818–33.
30. Wang GL, Zhang CL, Huo HQ, Sun XS, Zhang YL, Hao YJ, et al. The SUMO E3 ligase MdSIZ1 Sumoylates a cell number regulator MdCNR8 to control organ size. *Front Plant Sci*. 2022;13:836935.
31. Streich FC, Lima CD. Capturing a substrate in an activated RING E3/E2-SUMO complex. *Nature*. 2016;536(7616):304.
32. Benlloch R, Lois LM. Sumoylation in plants: mechanistic insights and its role in drought stress. *J Exp Bot*. 2018;69(19):4539–54.
33. Yarra R, Xue Y. Ectopic expression of nucleolar DEAD-box RNA helicase OsTOGR1 confers improved heat stress tolerance in transgenic Chinese cabbage. *Plant Cell Rep*. 2020;39(12):1803–14.
34. Besse M, Knipfer T, Miller AJ, Verdeil JL, Jahn TP, Fricke W. Developmental pattern of aquaporin expression in barley (*Hordeum vulgare* L.) leaves. *J Exp Bot*. 2011;62(12):4127–42.
35. Li Q, Zhu TT, Zhang R, Bu QT, Yin J, Zhang L, et al. Molecular cloning and functional analysis of hyoscyamine 6 betahydroxylase (H6H) in the poisonous and medicinal plant *Datura innoxia* mill. *Plant Physiol Bioch*. 2020;153:11–9.
36. Liu G, Tian H, Huang YQ, Hu J, Ji YX, Li SQ, et al. Alterations of Mitochondrial Protein Assembly and Jasmonic Acid Biosynthesis Pathway in Honglian (HL)-type Cytoplasmic Male Sterility Rice. *J Biol Chem*. 2012;287(47):40051–60.
37. Mitsuda N, Ikeda M, Takada S, Takiguchi Y, Kondou Y, Yoshizumi T, et al. Efficient yeast one-/two-hybrid screening using a library composed only of transcription factors in *Arabidopsis thaliana*. *Plant Cell Physiol*. 2010;51(12):2145–51.
38. Shi SL, Liu Y, He YJ, Li LZ, Li DL, Chen HY. R2R3-MYB transcription factor SmMYB75 promotes anthocyanin biosynthesis in eggplant (*Solanum melongena* L.). *Sci Hortic-Amsterdam*. 2021;255(1):12.
39. Liu YD, Wen L, Shi Y, Su DD, Lu W, Cheng YL, et al. Stress-responsive tomato gene SIGRASA4 function in drought stress and abscisic acid signaling. *Plant Sci*. 2021;304:110804.
40. Chen DH, Molitor AM, Xu L, Shen WH. *Arabidopsis* PRC1 core component AtRING1 regulates stem cell-determining carpel development mainly through repression of class I KNOX genes. *BMC Biol*. 2016;14(1):112.
41. Shen XP, Xu LA, Liu YH, Dong H, Zhou D, Zhang YZ, et al. Comparative transcriptome analysis and ChIP-sequencing reveals stage-specific gene expression and regulation profiles associated with pollen wall formation in *Brassica rapa*. *BMC Genomics*. 2019;20(1):264.
42. Gomez LM, Teixeira-Silva NS, Caserta R, Takita MA, Marques MOM, de Souza AA. Overexpression of *Citrus reticulata* SAMT in *Nicotiana tabacum* increases MeSA volatilization and decreases *Xylella fastidiosa* symptoms. *Planta*. 2020;252(6):103.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.