



SALTISH: The SALt-affected vegeTation dataset of Tuscany coaStal Habitats, central Italy*

Hamid Gholizadeh¹, Gianmaria Bonari^{1,2}, Emilia Pafumi^{1,2}, Andrea Bertacchi³, Mariasole Calbi⁴, Paolo Castagnini¹, Daniela Ciccarelli⁵, Emanuele Fanfarillo^{1,2}, Giulio Ferretti⁶, Tiberio Fiaschi¹, Bruno Foggi⁴, Matilde Gennai⁴, Lorenzo Lazzaro⁴, Michele Mugnai⁴, Simona Sarmati^{2,7}, Daniele Viciani⁴, Claudia Angiolini^{1,2**}, Simona Maccherini^{1,2**}

¹ Department of Life Sciences, University of Siena, Siena, Italy

² NBFC, National Biodiversity Future Center, Palermo, Italy

³ Department of Agriculture, Food and Agro-Environmental Sciences, University of Pisa, Pisa, Italy

⁴ Department of Biology, University of Florence, Florence, Italy

⁵ Department of Biology, University of Pisa, Pisa, Italy

⁶ Botanical Garden “Giardino dei Semplici”, Florence, Italy

⁷ Department of Sciences, University of Rome Tre, Rome, Italy

Corresponding author: Gianmaria Bonari (gianmaria.bonari@unisi.it)

Academic editor: Pauline Delbosc ♦ Received 18 December 2024 ♦ Accepted 3 February 2025 ♦ Published 17 April 2025

Abstract

Surveying vegetation is essential for documenting plant diversity, especially for coastal vegetation that results among the most threatened ecosystems globally. To support conservation and management programs, we developed the SALt-affected vegeTation dataset of Tuscany coaStal Habitats (SALTISH). This dataset comprises 734 newly sampled vegetation plots of 4 m² (2 m × 2 m) from the Tuscany region in central Italy, including 569 sand dune plots and 165 salt marsh plots, recorded between 2018 and 2023. In total, the dataset contains 4,541 occurrences of vascular plant taxa. Overall, it comprehends 257 vascular plant taxa belonging to 165 genera and 56 families. The Poaceae family is the most diverse, represented by 50 taxa, while the most represented genus is *Juncus*, with seven species. Species richness within individual plots ranges from one to 55 species, with 622 plots (84%) containing fewer than 10 species. *Juniperus macrocarpa* emerges as the most frequent and dominant species in the dataset. *Helichrysum stoechas*, *Festuca fasciculata*, and *Medicago littoralis* are present in over 20% of the plots, whereas 157 taxa are recorded in fewer than 1% of plots. The dataset includes noteworthy taxa: four Italian endemics (*Centaurea aplolepa* subsp. *subciliata*, *Limonium etruscum*, *L. multifforme*, and *Solidago virgaurea* subsp. *littoralis*), eight taxa listed as threatened in the Italian Red List, and 18 archaeophyte and neophyte alien species. SALTISH provides critical data for monitoring and conserving threatened coastal habitats in Tuscany. This resource will facilitate comparisons of biodiversity status and vegetation changes over time and will aid in identifying habitats harboring rare and endangered plant species.

Keywords

Coastal dunes, Italy, Tuscany, salt marshes, vegetation dataset, vegetation plot

* Topical Collection: “Vegetation databases: enhancing data integration and accessibility for ecological research”. Edited by Adrian Indreica, Kiril Vassilev, Pauline Delbosc, Federico Fernández-González, Irena Axmanová, Borja Jiménez-Alfaro, Gianmaria Bonari.

** These authors contributed equally to this work.

Introduction

Coastal habitats are increasingly threatened by global changes. These areas face direct and indirect impacts from sea-level rise (Jones et al. 2013), as well as anthropogenic activities such as coastal development, habitat degradation, and agricultural practices (Seitz et al. 2014; Gennai et al. 2022). Coastal sand dunes and salt marshes support numerous rare and endangered plant species (Acosta et al. 2009), and they serve as important nurseries for juvenile fish and invertebrates (Lefcheck et al. 2019). However, approximately 86% of European coastal habitats are at high or moderate risk from unsustainable coastal construction and development (Seitz et al. 2014). The 21% reduction in the quality of coastal dunes in Italy over the last 50 years affected negatively about half of the current area, and they can be categorised as vulnerable (Gigante et al. 2018; Sarmati et al. 2025). To address these threats, the EU Natura 2000 network of protected areas aims to conserve the most endangered species and habitats.

In recent decades, vegetation archives containing ecological plot data have significantly expanded. Advances in software and techniques for data storage and analysis have enabled researchers to create new vegetation databases and cover under-sampled regions (Hennekens and Schaminée 2001; Tichý 2002; Peterka et al. 2015; Marcenò and Jiménez-Alfaro 2017). Vegetation plot datasets are fundamental resources for vegetation syntheses (Bonari et al. 2021; Kavğacı et al. 2021; Jiroušek et al. 2022; Novák et al. 2023b; Peterka et al. 2023) and have proven invaluable for biodiversity and macroecology research (Wüest et al. 2020; Cai et al. 2023). The European Vegetation Archive (EVA; Chytrý et al. 2016) is the primary repository for vegetation data in Europe, containing many national and regional datasets. Nevertheless, efforts to survey under-sampled vegetation types, digitize data, and develop new datasets continue at various scales across Europe and other continents (Bonari et al. 2019; Gholizadeh et al. 2019; Alessi et al. 2022; Novák et al. 2023a; Bagella et al. 2024).

Italy has over 7,900 km of coastline, and the coastal habitats of the Tuscany region support a significant portion of the human population as well as numerous tourism-related activities (Cipriani et al. 2013; Cappucci et al. 2024). This long and accessible coastline has attracted botanists for decades. Although many vegetation plots from this part of the Mediterranean coast are available, they vary in size and do not adequately cover the inner zones of the coastal gradient (Prisco et al. 2012; Marcenò and Jiménez-Alfaro 2017; Alessi et al. 2022).

Coastal areas play a critical role in the provision of habitat for biodiversity, carbon sequestration, water supply, and enabling socio-economic activities (Everard et al. 2010; Ford et al. 2012). To ensure the preservation of these functions, it is crucial to create and maintain updated vegetation datasets. Such data help monitor the distribution of plant communities and habitats, as well as endangered and invasive alien plant species, track their invasiveness, and assess the potential transformations of vegetation

types. In Tuscany, human pressure on these habitats has led to a steady reduction in their number and area since the early 20th century (Bertacchi et al. 2021).

Tuscany's coastal zone includes a variety of protected areas, such as regional parks, state nature reserves, and special areas for conservation. Generally, the southern part of the region is better preserved than the north, where human activities are more intensive (Ciccarelli et al. 2014). Nevertheless, recent studies have identified sites with high conservation value that remain unprotected (Pafumi et al. 2024).

The MONITORARE project (2017–2018) and the NATNET project (NATura Network Toscana 2019–2024) were launched to develop a dataset for monitoring and conserving plant and animal species and their habitats in the Tuscany region of central Italy. These initiatives, conducted in collaboration with the Tuscan Regional Administration and researchers from the Universities of Siena, Florence, and Pisa, aim to fill gaps in knowledge and improve conservation efforts (Gennai et al. 2022). This paper presents the structure and content of the SALT-affected vegetation dataset of Tuscany coastal Habitats (SALTISH), which contains recently collected vegetation plots as part of these projects.

Study area

The study area is about 450 km long and includes protected areas and unprotected regions throughout the coastline of Tuscany, central Italy. The map of the study area is shown in Suppl. material 1. The macrobioclimate of the area is Mediterranean with upper meso-mediterranean thermotype and ombrotype ranging from lower humid to upper dry in the North and South, respectively (Pesaresi et al. 2017). Geologically, Late Quaternary sands, pebbles, and muds cover most of the coastal areas in the Tuscany region. However, some small patches with ocean-derived sand rocks and other substrates exist in these areas (Carmignani et al. 2013).

Different habitat and vegetation types occur along the coastal areas of Tuscany. Following the sea-inland gradient, the annual vegetation of drift lines, stable coastal grasslands, coastal dune scrubs and woods, salt meadows and steppes, and heliophilous scrubs can be found in these areas (Bertacchi et al. 2021; Gennai et al. 2022; Pafumi et al. 2024).

Data collection and preparation

Following a stratified random sampling design and several field excursions to cover most parts of the region, all vegetation plots in our dataset were surveyed between April 2018 and October 2023 during the coastal vegetation season. Plot data were gathered with a positional accuracy ranging from 0 to 79 meters (mean = 4.7 meters) using a hand-held GPS unit in the field for the MONITORARE and NATNET projects. These projects focused

on habitat monitoring and developing a comprehensive vegetation dataset for various coastal ecosystems and habitats (Gennai et al. 2022). The dataset includes two main habitat types in coastal areas: 569 plots were sampled in sand dunes, while 165 plots were sampled in salt marshes.

To minimize the influence of plot size on species richness and other vegetation parameters, all plots were sampled within a standardized 4 m² area (2 m × 2 m) (Dengler et al. 2008; Acosta et al. 2009), which is also recommended in the Monitoring Manual of Italy for coastal areas (Angelini et al. 2016). Vascular plant taxa were recorded based on the percentage cover of each species within the plots.

Scientific plant names and taxonomical concepts were standardized following the Portal to the Flora of Italy (2024). Life forms, chorotypes, alien species categories, and conservation status for each taxon were assigned based on Pignatti et al. (2017–2019) and Orsenigo et al. (2018; 2020). To assess species dependency on salinity in both habitats, salinity relationships and salinity indicator values for taxa were extracted from the FloraVeg.EU database (Chytrý et al. 2024; FloraVeg 2024).

Furthermore, each plot's habitat type was classified according to the EUNIS habitat classification at level 3, applying the EUNIS Expert System (Chytrý et al. 2020), the 92/43/EEC habitat classification adapted for Italy (Biondi et al. 2009), and expert-based classification for the remaining 76 unclassified plots. The EUNIS classification was performed in R 4.3.2 (R Core Team 2023) using R code implemented by Bruehlheide et al. (2021).

Structure and content of the dataset

The SALTISH dataset is based on the standard header data fields of TURBOVEG 2.166 (Hennekens and Schaminée 2001). It encompasses 28 data fields, including elevation, geographic coordinates, slope, and aspect as topographic variables, the percentage cover of total vegetation, tree, shrub, and herb layers, and other plot-level information, including the protection status (Suppl. material 2). The dataset includes 734 georeferenced vegetation plots, including 257 vascular plant taxa with 4,541 occurrence data.

The species richness of plots varies between one and 55, with 622 (85%) plots having less than 10 species and three plots having 20 or more taxa. The average species richness per plot in sand dunes and salt marshes is 6.9 and 3.8, respectively (Fig. 1A).

A total of 257 vascular plant taxa belonging to 165 genera and 56 families were identified. Among these, 53 families belong to the Angiosperms, two families (Cupressaceae with four taxa and Pinaceae with two taxa) are Gymnosperms, and one family (Equisetaceae, represented by a single species found exclusively in sand dunes) is a fern. While 219 taxa of 53 families were recorded in sand dunes, only 72 taxa of 22 families were found in salt marshes. Poaceae is the largest family in both habitats with 37 taxa in sand dunes and 25 taxa in salt marshes. Asteraceae and Amaranthaceae are the second-largest families in sand dunes and salt marshes, respectively. Fabaceae, the third-largest family with 19 taxa in sand dunes, has no species in salt marshes (Fig. 1B).

Juncus with seven taxa is the most diverse genus in the dataset. However, the most abundant genera are different in sand dunes and salt marshes. *Juncus* with six taxa in sand dunes and *Sporobolus* with four taxa in salt marshes are the most abundant genera. *Medicago* and *Trifolium*, with six and five taxa respectively, are other abundant genera in sand dunes, and they are missing in salt marshes. However, *Euphorbia* with six taxa is one of the second-abundant genera in the dataset with almost similar abundance in sand dunes and salt marshes (Fig. 1C).

The dominant life forms are Therophytes (40%) and Hemicryptophytes (27%) in sand dunes and conversely Hemicryptophytes (37%) and Therophytes (31%) in salt marshes. Geophytes and Phanerophytes are the less frequent life forms in these habitats, respectively. However, Phanerophytes with 16% have a relatively good proportion of life forms in sand dunes (Fig. 1D).

The phytogeographical distribution of taxa shows Mediterranean (44%) and Eurasian (18%) elements are the most common chorotypes in the dataset, as well as in sand dunes and salt marshes, separately (Fig. 1E). Moreover, the dataset contains four Italian endemics (*Centaurea aplolepa* subsp. *subciliata*, *Limonium etruscum*, *L. multifforme*, and *Solidago virgaurea* subsp. *litoralis*), and a total of eight threatened taxa of Italian Red List (Table 1).

Table 1. The endemic and threatened taxa and their categories from the Italian Red List (Orsenigo et al. 2018, 2020), as well as their distribution and frequency in each habitat. Plant taxa were ordered alphabetically according to chorotype (Pignatti et al. 2017–2019). SD: sand dunes; SM: salt marshes.

Plant taxa	Chorotype	IUCN threat category	Distribution (Species frequency)
<i>Centaurea aplolepa</i> subsp. <i>subciliata</i>	Endemic	Endangered (EN)	SD (4)
<i>Limonium etruscum</i>	Endemic	Critically Endangered (CR)	SM (9)
<i>Limonium multifforme</i>	Endemic	Vulnerable (VU)	SD (1), SM (1)
<i>Solidago virgaurea</i> subsp. <i>litoralis</i>	Endemic	Endangered (EN)	SD (5)
<i>Chamaerops humilis</i> subsp. <i>humilis</i>	Mediterranean	Near Threatened (NT)	SD (2)
<i>Euphorbia barrelieri</i>	Mediterranean	Endangered (EN)	SD (1)
<i>Triglochin barrelieri</i>	Mediterranean	Endangered (EN)	SM (4)
<i>Samolus valerandi</i>	Wide distribution	Least Concern (LC)	SD (1)




Figure 1. Species richness per plot (A), the contribution of the eight most frequent families (B), the eight most abundant genera (C), the proportion of different life forms (D), chorological spectrum (E), and the eight most frequent taxa (F). The subspecies designations for *Cakile maritima* subsp. *maritima* and *Calamagrostis arenaria* subsp. *arundinacea* are not reported in the figure for graphical reasons.

There are also 18 archaeophyte and neophyte alien species that are categorised as invasive and naturalised in Italy (Table 2).

The most frequent taxa (present in at least 30% of plots) including *Juniperus macrocarpa*, *Helichrysum stoechas*, and *Festuca fasciculata* in sand dunes, and *Salicornia perennis*, *Puccinellia festuciformis*, and *Halimione portulacoides* in salt marshes only have been recorded in one habitat (Fig. 1F). Moreover, the five most dominant taxa in the dataset containing *J. macrocarpa* and *Calamagrostis arenaria* subsp. *arundinacea* in sand dunes, and *S. perennis*, *Arthrocnemum macrostachyum*, and *H. portulacoides* in salt marshes were distributed only in one of them.

According to salinity relationship classification, plants adapted to growing in non-saline areas are the most common in both habitats. However, approximately 60% of the taxa in salt marshes are classified as those adapted to saline and slightly saline or brackish areas. On the other hand, for 12 taxa in our dataset, salinity indicator values are recorded as seven or more, and most of these are present only in salt marshes. Among these species with a high tolerance to salinity, four species including *Limonium multifforme*, *L. narbonense*, *Galatella tripolium* (one occurrence), and *Sporobolus pungens* (63 occurrences) are recorded in sand dunes (Fig. 2A, B).

Table 2. The alien taxa and their associated status (Portal to the Flora of Italy 2024) in Italy and Tuscany, as well as their distribution and frequency in each habitat. Plant taxa were ordered alphabetically. SD: sand dunes; SM: salt marshes.

Plant taxa	Alien species status (in Italy / Tuscany)	Distribution (Species frequency)
<i>Ambrosia psilostachya</i>	Neophyte species (Invasive / Naturalized)	SD (12)
<i>Arundo donax</i>	Archaeophyte species (Invasive / Invasive)	SD (1)
<i>Avena fatua</i>	Archaeophyte species (Naturalized / Alien)	SD (1)
<i>Avena sterilis</i>	Archaeophyte species (Naturalized / Naturalized)	SD (9)
<i>Crepis sancta</i> subsp. <i>nemausensis</i>	Neophyte species (Invasive / Invasive)	SD (1)
<i>Erigeron canadensis</i>	Neophyte species (Invasive / Invasive)	SD (9), SM (3)
<i>Erigeron sumatrensis</i>	Neophyte species (Invasive / Invasive)	SD (2)
<i>Euphorbia serpens</i> subsp. <i>serpens</i>	Neophyte species (Naturalized / Naturalized)	SM (5)
<i>Oxalis stricta</i>	Neophyte species (Invasive / Casual)	SM (1)
<i>Paspalum vaginatum</i>	Neophyte species (Naturalized / Naturalized)	SD (1)
<i>Phalaris canariensis</i>	Neophyte species (Invasive / Naturalized)	SD (1)
<i>Pinus pinea</i>	Archaeophyte species (Naturalized / Naturalized)	SD (11)
<i>Pittosporum tobira</i>	Neophyte species (Naturalized / Naturalized)	SD (3)
<i>Sporobolus indicus</i>	Neophyte species (Naturalized / Naturalized)	SM (1)
<i>Sporobolus pumilus</i>	Neophyte species (Invasive / Invasive)	SD (12), SM (1)
<i>Symphyotrichum squamatum</i>	Neophyte species (Invasive / Invasive)	SM (24)
<i>Xanthium orientale</i>	Neophyte species (Invasive / Invasive)	SD (61), SM (3)
<i>Yucca gloriosa</i>	Neophyte species (Invasive / Invasive)	SD (5)




Figure 2. Salinity dependency status of plant taxa: **A)** Occurrence in different areas (64 taxa missed information), **B)** Salinity indicator value (40 taxa missed information).

The vegetation plots are mainly included in protected areas (549 protected vs 185 non-protected plots). Those protected areas are distributed across different designation types of protected areas. Specifically, five designation types are present in the study area: Special Protection Area, Special Area for Conservation, State Nature Reserve, Regional Nature Reserve, and Regional Park. Moreover, 482 plots of 549 plots in protected areas, are protected by two (81 plots), three (398 plots), or four (3 plots) types of protection (Fig. 3).

The classification of habitat types for each vegetation plot identified 14 92/43/EEC habitat types and seven EUNIS habitat types. 370 plots classified under 92/43/EEC codes were sampled from areas designated as Special Protection Areas, Special Areas for Conservation, or both. However, 27.4% of the plots could not be classified into any 92/43/EEC habitat type, and 3% could not be assigned to any EUNIS habitat type (Table 3).

Table 3. Classification of the vegetation plots according to 92/43/EEC and EUNIS habitat typologies. Habitats are sorted by decreasing the number of assigned vegetation plots.

	92/43/EEC	No. plots (%)	EUNIS	No. plots (%)
Sand dunes	2250	135 (18.4)	N16	319 (43.5)
	2110	79 (10.8)	N14	109 (14.9)
	2120	67 (9.1)	N1B	64 (8.7)
	2210	38 (5.2)	N12	48 (6.5)
	1210	21 (2.9)	N1J	7 (1)
	2230	13 (1.8)	N1G	6 (0.8)
	2260	12 (1.6)	MA25	4 (0.5)
	2270	8 (1.1)	Not-classified	12 (1.6)
	Not-classified	196 (26.7)		
Salt marshes	1420	88 (12)	MA25	147 (20)
	1410	52 (7.1)	N1J	8 (1.1)
	1310	8 (1.1)	Not-classified	10 (1.4)
	6420	6 (0.8)		
	1510	6 (0.8)		
	Not-classified	5 (0.7)		




Figure 3. Number of plots in protected and non-protected areas and types of protection. The protected areas are shown on the map.

Conclusions

The information present in the SALTISH dataset strengthens the knowledge of habitat diversity in the Tuscany coastal habitats. This dataset, which compiles comprehensive data on plant species, community composition, and habitat types, is crucial for understanding the diverse ecosystems that characterize Tuscany region's coasts. It also helps defining all vegetation units of Tuscany's sand dunes and salt marshes. It can be used to document changes over time and monitor shifts in vegetation patterns, which can be influenced by factors such as climate change, human activity, and invasive species. Moreover, the SALTISH dataset facilitates biodiversity assessments and helps identify areas of ecological importance that require conservation efforts. Additionally, this dataset serves as a reference for conservation programs of threatened coastal habitats in central Italy. It can support decisions regarding land use, coastal protection, and the restoration of degraded areas. It also provides a basis for future studies on the effects of environmental stressors on plant communities and their

resilience. In conclusion, the SALTISH dataset is not only an essential resource for scientific research but also an effective tool for promoting nature conservation. However, data collection and storage in SALTISH is still ongoing process. Future efforts should be directed towards rush communities to assess the entire sea-inland gradient fully.

Data availability

Data will be available by contacting the authors.

Author contributions

HG led the writing with contributions from GB, EP, CA, and SM. The data was collected by AB, CA, PC, DC, LDD, GF, SS, BF, TF, MM, and DV. TF, BF, SS, and EF identified plants in the lab. HG and EP did the analysis. HG, MG, BF, SS, and TF prepared the dataset. TF classified plots remained unclassified. EP prepared the figures. CA and

SM provided funding. All authors reviewed and approved the manuscript before submission.

Competing interest

The authors have declared that no competing interests exist.

Acknowledgements

We thank Letizia Di Domizio for her help in the vegetation sampling.

We acknowledge financial support under the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.1, Call for tender No. 104 published on 2.2.2022 by the Italian Ministry of University and Research (MUR), funded by the European Union – NextGenerationEU – Project Title Prioritisation of coastal areas for plant diversity conservation through a multidisciplinary approach – CUP B53D23012040006 – Grant Assignment Decree No 1015 of 7 July 2023 adopted by the Italian Ministry of University and Research (MUR).

We acknowledge the financial support of Regione Toscana through the MonitoRare and Nat-Net projects.

References

- Acosta A, Carranza ML, Izzi CF (2009) Are there habitats that contribute best to plant species diversity in coastal dunes? *Biodiversity and Conservation* 18: 1087–1098. <https://doi.org/10.1007/s10531-008-9454-9>
- Alessi N, Bruzzaniti V, Buldrini F, Centomo E, Cervellini M, ... Chiarucci A (2022) AMS-VegBank: a new database of vegetation plots for the Italian territory. *Vegetation Classification and Survey* 3: 177–185. <https://doi.org/10.3897/VCS.85083>
- Angelini P, Casella L, Grignetti A, Genovesi P (2016) Manuali per il monitoraggio di specie habitat di interesse comunitario (Direttiva 92/43/CEE) in Italia. ISPRA, Serie Manuali e linee guida. <http://is-praambiente.gov.it>
- Bagella S, Caria MC, Bonari G, Malavasi M, Melis R, ... Riviaccio G (2024) A thematic vegetation dataset of Sardinian GRasslands (SAGRA). *Plant Sociology* 61(2): 41–47. <https://doi.org/10.3897/pls2024612/03>
- Bertacchi A, Lombardi T, Saggese A, Lazzeri V (2021) The vegetation of a relict salt marsh area in the Pisan coast in the context of brackish wetlands of Tuscany. *Plant Sociology* 58(1): 41–53. <https://doi.org/10.3897/pls2021581/03>
- Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, ... Zivkovic L (2009) Manuale italiano di interpretazione degli Habitat (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. <http://vnr.unipg.it/habitat/> [Accessed on 15 November 2024]
- Bonari G, Knollová I, Vlčková P, Xystrakis F, Çoban S, ... Chytrý M (2019) CircumMed Pine Forest Database: an electronic archive for Mediterranean and Submediterranean pine forest vegetation data. *Phytocoenologia* 49(3): 311–318. <https://doi.org/10.1127/phyto/2019/0311>
- Bonari G, Fernández-González F, Çoban S, Monteiro-Henriques T, Bergmeier E, ... Chytrý M (2021) Classification of the Mediterranean lowland to submontane pine forest vegetation. *Applied Vegetation Science* 24(1): e12544. <https://doi.org/10.1111/avsc.12544>
- Bruehlheide H, Tichý L, Chytrý M, Jansen F (2021) Implementing the formal language of the vegetation classification expert systems (ESy) in the statistical computing environment R. *Applied Vegetation Science* 24: e12562. <https://doi.org/10.1111/avsc.12562>
- Cai L, Kreft H, Taylor A, Denelle P, Schrader J, ... Weigelt P (2023) Global models and predictions of plant diversity based on advanced machine learning techniques. *New Phytologist* 237(4): 1432–1445. <https://doi.org/10.1111/nph.18533>
- Cappucci S, Carillo A, Iacono R, Moretti L, Palma M, ... Sannino G (2024) Evolution of coastal environments under inundation scenarios using an oceanographic model and remote sensing data. *Remote Sensing* 16(14): 2599. <https://doi.org/10.3390/rs16142599>
- Carmignani L, Conti P, Cornamusi G, Pirro A (2013) Geological map of Tuscany (Italy). *Journal of Maps* 9: 487–497. <https://doi.org/10.1080/17445647.2013.820154>
- Chytrý M, Hennekens SM, Jiménez-Alfaro B, Knollová I, Dengler J, ... Yamalov S (2016) European Vegetation archive (EVA): an integrated database of European vegetation plots. *Applied Vegetation Science* 19: 173–180. <https://doi.org/10.1111/avsc.12191>
- Chytrý M, Tichý L, Hennekens SM, Knollová I, Janssen JAM, ... Schaminée JHJ (2020) EUNIS Habitat Classification: Expert system, characteristic species combinations and distribution maps of European habitats. *Applied Vegetation Science* 23: 648–675. <https://doi.org/10.1111/avsc.12519>
- Chytrý M, Řezníčková M, Novotný P, Holubová D, Preislerová Z, ... Axmanová I (2024) FloraVeg.EU – an online database of European vegetation, habitats and flora. *Applied Vegetation Science* 27: e12798. <https://doi.org/10.1111/avsc.12798>
- Ciccarelli D, Di Bugno C, Peruzzi L (2014) Checklist della flora vascolare psammofila della Toscana. *Atti della Società Toscana di Scienze Naturali, Memorie, Serie B* 121[2014]: 37–88. <https://doi.org/10.2424/ASTSN.M.2014.05>
- Cipriani LE, Pranzini E, Vitale G (2013) Coastal erosion in Tuscany: Short vs. medium term evolution. In: Cipriani LE (Ed.) *Coastal Erosion Monitoring. A Network of Regional Observatories – Results from RESMAR Project*; Nuova Grafica Fiorentina, 135–155.
- Dengler J, Chytrý M, Ewald J (2008) Phytosociology. *Encyclopedia of Ecology* 4(5): 2767–2779. <https://doi.org/10.1016/B978-008045405-4.00533-4>
- Everard M, Jones MLM, Watts B (2010) Have we neglected the societal importance of sand dunes? An ecosystem services perspective. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 476–487. <https://doi.org/10.1002/aqc.1114>
- FloraVeg.EU (2024) Database of European Vegetation, Habitats and Flora. <https://www.floraveg.eu> [Accessed on 15 November 2024]
- Ford H, Garbutt A, Jones DL, Jones L (2012) Impacts of grazing abandonment on ecosystem service provision: Coastal grassland as a model system. *Agriculture, Ecosystems and Environment* 162: 108–115. <https://doi.org/10.1016/j.agee.2012.09.003>
- Gennai M, Angiolini C, Bertacchi A, Gabellini A, Sarmati S, ... Foggi B (2022) Studying local species assemblages of salt-affected vegetation for monitoring Natura 2000 habitats. *Plant Sociology* 59(1): 1–10. <https://doi.org/10.3897/pls2022591/01>
- Gholizadeh H, Naqinezhad A, Chytrý M (2019) Hyrcanian forest vegetation database. *Phytocoenologia* 49: 209–210. <https://doi.org/10.1127/phyto/2018/0315>
- Gigante D, Acosta ATR, Agrillo E, Armiraglio S, Assini S, ... Viciani D (2018) Habitat conservation in Italy: the state of the art in the light of the first European Red List of Terrestrial and Freshwater Habitats.

- Rendiconti Lincei. Scienze Fisiche e Naturali 29: 251–265. <https://doi.org/10.1007/s12210-018-0688-5>
- Hennekens SM, Schaminée JHJ (2001) TURBOVEG, a comprehensive database management system for vegetation data. *Journal of Vegetation Science* 12: 589–591. <https://doi.org/10.2307/3237010>
- Jiroušek M, Peterka T, Chytrý M, Jiménez-Alfaro B, Kuznetsov OL, ... Hájek M (2022) Classification of European bog vegetation of the *Oxycocco-Sphagnetum* class. *Applied Vegetation Science* 25(1): e12646. <https://doi.org/10.1111/avsc.12646>
- Jones L, Garbutt A, Hansom J, Angus S (2013) Impacts of climate change on coastal habitats. *MCCIP Science Review* 2013: 167–179. <https://doi.org/10.14465/2013.arc18.167-179>
- Kavgacı A, Balpınar N, Öner HH, Arslan M, Bonari G, ... Čarni A (2021) Classification of forest and shrubland vegetation in Mediterranean Turkey. *Applied Vegetation Science* 24(2): e12589. <https://doi.org/10.1111/avsc.12589>
- Lefcheck JS, Hughes BB, Johnson AJ, Pfirrmann BW, Rasher DB, ... Orth RJ (2019) Are coastal habitats important nurseries? A meta-analysis. *Conservation Letters* 12: e12645. <https://doi.org/10.1111/conl.12645>
- Marcenò C, Jiménez-Alfaro B (2017) The Mediterranean *Ammophila*-*tea* Database: a comprehensive dataset of coastal dune vegetation. *Phytocoenologia* 47: 95–105. <https://doi.org/10.1127/phyto/2016/0133>
- Novák P, Kalníková V, Szokala D, Aleksanyan A, Batsatsashvili K, ... Zúkal D (2023a) Transcaucasian Vegetation Database – a phytosociological database of the Southern Caucasus. *Vegetation Classification and Survey* 4: 231–240. <https://doi.org/10.3897/VCS.105521>
- Novák P, Willner W, Biurrun I, Gholizadeh H, Heinken T, ... Chytrý M (2023b) Classification of European oak–hornbeam forests and related vegetation types. *Applied Vegetation Science* 26(1): e12712. <https://doi.org/10.1111/avsc.12712>
- Orsenigo S, Montagnani C, Fenu G, Gargano D, Peruzzi L, ... Rossi G (2018) Red Listing plants under full national responsibility: Extinction risk and threats in the vascular flora endemic to Italy. *Biological Conservation* 224: 213–222. <https://doi.org/10.1016/j.biocon.2018.05.030>
- Orsenigo S, Fenu G, Gargano D, Montagnani C, Abeli T, ... Rossi G (2020) Red list of threatened vascular plants in Italy. *Plant Biosystems* 155(2): 310–335. <https://doi.org/10.1080/11263504.2020.1739165>
- Pafumi E, Angiolini C, Sarmati S, Bacaro G, Fanfarillo E, ... Maccherini S (2024) Spatial patterns of coastal dune plant diversity reveal conservation priority hotspots in and out a network of protected areas. *Global Ecology and Conservation* 54: e03085. <https://doi.org/10.1016/j.gecco.2024.e03085>
- Pesaresi S, Biondi E, Casavecchia S (2017) Bioclimates of Italy. *Journal of Maps* 13(2): 955–960. <https://doi.org/10.1080/17445647.2017.1413017>
- Peterka T, Jiroušek M, Hájek M, Jiménez-Alfaro B (2015) European Mire Vegetation Database: a gap-oriented database for European fens and bogs. *Phytocoenologia* 45: 291–298. <https://doi.org/10.1127/phyto/2015/0054>
- Peterka T, Hájková P, Jiroušek M, Hinterlang D, Chytrý M, ... Hájek M (2023) Formalized classification of the class *Montio-Cardaminetum* in Europe: towards a consistent typology of spring vegetation. *Preslia* 95(3): 347–383. <https://doi.org/10.23855/preslia.2023.347>
- Pignatti S, Guarino R, La Rosa M (2017–2019) *Flora d'Italia* (2nd edn., Vols 1–4). Edagricole di New Business Media, Milano.
- Portal to the Flora of Italy (2024) Portal to the Flora of Italy. <http://dryades.units.it/floritaly> [Accessed on 15 November 2024]
- Prisco I, Carboni M, Acosta ATR (2012) VegDunes – a coastal dune vegetation database for the analysis of Italian EU habitats. *Biodiversity and Ecology* 4: 191–200. <https://doi.org/10.7809/b-e.00076>
- R Core Team (2023) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Sarmati S, Angiolini C, Sperandii MG, Barták V, Gennai M, ... Bazzichetto M (2025) A complex interplay between natural and anthropogenic factors shapes plant diversity patterns in Mediterranean coastal dunes. *Landscape Ecology* 40: 20. <https://doi.org/10.32942/X20044>
- Seitz RD, Wennhage H, Bergström ULF, Lipcius RN, Ysebaert T (2014) Ecological value of coastal habitats for commercially and ecologically important species. *ICES Journal of Marine Science* 71(3): 648–665. <https://doi.org/10.1093/icesjms/fst152>
- Tichý L (2002) JUICE, software for vegetation classification. *Journal of Vegetation Science* 13: 451–453. <https://doi.org/10.1111/j.1654-1103.2002.tb02069.x>
- Wüest RO, Zimmermann NE, Zurell D, Alexander JM, Fritz SA, ... Karger DN (2020) Macroecology in the age of Big Data—Where to go from here? *Journal of Biogeography* 47(1): 1–12. <https://doi.org/10.1111/jbi.13633>

Supplementary material 1

Location of the 734 vegetation plots along the Tuscan coast (central Italy)

Authors: Hamid Gholizadeh, Gianmaria Bonari, Emilia Pafumi, Andrea Bertacchi, Mariasole Calbi, Paolo Castagnini, Daniela Ciccarelli, Emanuele Fanfarillo, Giulio Ferretti, Tiberio Fiaschi, Bruno Foggi, Matilde Gennai, Lorenzo Lazzaro, Michele Mugnai, Simona Sarmati, Daniele Viciani, Claudia Angiolini, Simona Maccherini

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/ved.144362.suppl1>

Supplementary material 2

Data stored in the SALTISH dataset

Authors: Hamid Gholizadeh, Gianmaria Bonari, Emilia Pafumi, Andrea Bertacchi, Mariasole Calbi, Paolo Castagnini, Daniela Ciccarelli, Emanuele Fanfarillo, Giulio Ferretti, Tiberio Fiaschi, Bruno Foggi, Matilde Gennai, Lorenzo Lazzaro, Michele Mugnai, Simona Sarmati, Daniele Viciani, Claudia Angiolini, Simona Maccherini

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/ved.144362.suppl2>