

REFERENCES

- ANON. (1998): Biogeography of *Scylla tranquebarica*. Bioinformatics Centre, National Institute of Oceanography, Goa.
- CHHAPGAR, B.F. (1957): On the marine crabs (Decapoda: Brachyura) of Bombay State. *J. Bombay Nat. Hist. Soc.* 54(2 & 3): 399-439.
- CHHAPGAR, B.F. (1962): Crab fishing at Bombay. *J. Bombay Nat. Hist. Soc.* 59(1): 306-309.
- ESTAMPADOR, E.P. (1949): Studies on *Scylla* (Crustacea: Portunidae). Revision of the genus. *Philip. J. Sci.* 78(1): 95-108.
- FUSEYA, R. (1998): Studies on the species identification of the genus *Scylla*. Ph.D. Thesis of the Tokyo University of Fisheries, 170 pp. [In Japanese].
- FUSEYA, R. & S. WATANABE (1995): Notes on the taxonomy of the genus *Scylla*. *Cancer* (4): 5-8. [In Japanese].
- FUSEYA, R. & S. WATANABE (1996): Genetic variability in the mud crab genus *Scylla*. (Brachyura: Portunidae). *Fish. Sci.* 62(5): 705-709.
- FUSHIMI, H. & S. WATANABE (2001): "Problems in species identification of the mud crab genus *Scylla* (Brachyura: Portunidae)". Research article on Web. 5 pp.
- JOEL, D.R. & P.J. SANJEEVA RAJ (1983): Taxonomic remarks on two species of the genus *Scylla* De Haan (Portunidae: Brachyura) from Pulicat Lake. *Indian J. Fish.* 30: 13-26.
- KATHIRVAL, M. & S. SRINIVASAGAM (1992): Taxonomy of the mud crab, *Scylla serrata* (Forsk.) from India. The mud crab. Pp. 132-172. In: A report on the seminar convened in Surat Thani, Thailand. (Ed: Angell, C.A.). November 5-8, 1991. Bay of Bengal Programme, Madras, India.
- KEENAN, C.P., P.J.F. DAVIE & D.L. MANN (1998): A revision of the genus *Scylla* de Haan, 1833 (Crustacea: Decapoda: Brachyura: Portunidae). *The Raffles Bulletin of Zoology* 46(1): 217-245.
- OSHIRO, N. (1988). Mangrove crabs (*Scylla* spp.). Aquaculture in tropical areas (S. Syokita, ed.), Midorishobo, Tokyo: 198-209. [In Japanese]
- OVERTON, J.L., D.J. MACINTOSH, & R.S. THORPE (1997): Multivariate analysis of mud crab *Scylla serrata* (Brachyura: Portunidae) from four locations in South-east Asia. *Mar. Biol.* 128: 55-62.
- SAKAI, T. (1976): Crabs of Japan and the adjacent seas. *Kodansha*, Tokyo. Pp. 335-336.
- SERENE, R. (1952): Les especes du genre *Scylla* a Nhatrang (Vietnam). *Proc. Indo-Pacific Fish. Council* 3(2): 133-137.
- STEPHENSON, W. & B. CAMPBELL (1960): The Australian Portunids (Crustacea: Portunidae). IV. Remaining genera. *Austr. J. Mar. & Freshwater Res.* 11: 73-122.
- WATANABE, S. & R. FUSEYA (1997): Notes on the identification of the species in genus *Scylla*. *Cancer* 6: 33-36.

25. NEW SITES OF *NEPENTHES KHASIANA* FROM MEGHALAYA WITH NEW EASTERN AND WESTERN RANGE EXTENSIONS

The pitcher plant *Nepenthes khasiana* Hk. f. is the only *Nepenthes* species of carnivorous herbs, found in the Indian subcontinent. It is endemic to Meghalaya. *Nepenthes khasiana* was known only from the southern faces of Meghalaya plateau. Its reported western limit is Baghmara (90° 40' E) in South Garo Hills district, while the eastern limit

was at Jowai in Jaintia Hills district (92° 12' E) (Rodgers and Gupta 1989). The eastern range extended recently when a site was discovered near Umtra, also in Jaintia Hills (Choudhury 2000).

I report some more sites, which were hitherto unrecorded or overlooked including new eastern and western




Fig. 1: Map of Meghalaya showing the localities of *N. khasiana*

range extensions of *Nepenthes khasiana*. In October 2002, a new site was found more than a kilometre northwest of the site near Umtra. This site is smaller than the earlier one and is on the left side of National Highway 44, while coming from Khliehriat. On December 14, 2002, I found many small sites scattered widely from Rymbai village (25° 20' N, 92° 20' E) (Fig. 1) towards Lakadong. I drove for another 2 km (25° 19' N, 92° 19' E) and found scattered plants. The plants reportedly occur farther down towards Lakadong, but may not be present at Lakadong. The elevation of these patches was 1100 m above msl.

I then drove east of Khliehriat. The first patch, a large one along a *nullah* was near Sutnga village (25° 22' N, 92° 26' E) at 950 m elevation. The plant is reported from all around the village, but I could not visit all the sites. Farther east, up to about 4 km east of Sutnga I found some more pitcher plants. This site (25° 21' N, 92° 27' E) at 900 m elevation is the new eastern range of *khasiana*. There was a report from a site at 92° 28' E, which I could not visit.

Towards west, I received reports of occurrence at Chokpot (25° 18' N, 90° 25' E) (Surajit Roy, pers. comm.) and Deku area (25° 20' N, 90° 21' E) (Chagla Sangma, Awal Marak pers. comm.), both in South Garo Hills district. The known west-east extent of *Nepenthes khasiana*, which was 90° 40' E to 92° 25' E, has now extended to 90° 21' E to 92° 28' E with a number of new sites (Fig. 1).

In Jaintia Hills, coal mines using crude methods and its allied activities such as clearance of forest and levelling for storage of coal, setting up of labourers' camps, truck parking, shops and hotels have become a major threat. The labourers, truck drivers, shops and hotels also need wood for fuel and heating (in winter). It is high time that the coal mining, which is the main economy of the area is modernised so that its impact on the environment can be minimised. Some of the larger patches, such as the one along the *nullah* near Sutnga village, should be protected by erecting fence and banning coal mining and felling.

ACKNOWLEDGEMENTS

I thank T. Deb Roy, DFO (Wildlife), Jowai who informed me of possible new sites in the area and also helped during my visits. I also thank N. Sutnga, L. Nongkhla, G. Ringkhlem and Hakim for helping in the field survey.

March 19, 2003

ANWARUDDIN CHOUDHURY

The Rhino Foundation for nature in NE India,

C/o Assam Co., Ltd., Bamunimaidam,

Guwahati 781 021, Assam,

India.

Email: badrul@sancharnet.in

REFERENCES

CHOUDHURY, A.U. (2000): Range extension of *Nepenthes khasiana* in the Jaintia Hills, Meghalaya. *J. Bombay Nat. Hist. Soc.* 97(1): 166-167.

RODGERS, W.A. & S. GUPTA (1989): The Pitcher Plant (*Nepenthes khasiana* Hk. f.) Sanctuary of Jaintia Hills, Meghalaya: lessons for conservation. *J. Bombay Nat. Hist. Soc.* 86: 17-21.

26. *LINDERNIA ESTAMINODIOSA* (BLATT. & HALLB.) MUKHERJEE (SCROPHULARIACEAE): A NEW DISTRIBUTIONAL RECORD TO ANDHRA PRADESH

The genus *Lindernia* All., comprising about 100 species (Sivarajan and Mathew 1983), is described under various generic names, such as *Lindernia* All., *Vandellia* L., *Bonnaya* Link & Otto and *Ilysanthes* Rafin. The genus is represented by 22 species in India (Sivarajan and Mathew 1983; Cook 1996) and 11 species in Andhra Pradesh (Pullaiah and Ali Moulali 1997).

While studying the aquatic and wetland angiosperm diversity in Andhra Pradesh, we came across an interesting taxon of *Lindernia* All., with a sizable population, growing in moist locality, near Tada, Nellore district, Andhra Pradesh. After a critical examination of the material with the help of literature it was identified as *Lindernia estaminodiosa* (Blatt. & Hallb.) Mukherjee. A thorough perusal of regional floras and monographs revealed that this taxon is hitherto not reported from Andhra Pradesh. A detailed description, brief

account on phenology along with illustrations is given to facilitate easy field identification.

Lindernia estaminodiosa (Blatt. & Hallb.) Mukh. In: *J. Ind. Bot. Soc.* 24: 133. 1945.

Bonnaya estaminodiosa Blatt. & Hallb. In: *J. Bombay Nat. Hist. Soc.* 24: 416. 1918.

Annual, erect herb, 6-15 cm, tall; stem slender, sharply quadrangular, branched, glabrous. Leaves opposite, sessile, oblong-ob lanceolate, 10-30 x 3-5 mm, 1-nerved, lateral nerves obscure, base decurrent, margin with distinctly saw-like teeth, apex obtuse. Flowers in lax terminal and lateral racemes; bract subulate, 4 x 0.2 mm, 3-nerved, margin distantly setaceous above the middle, apex acute; pedicel stout, 4 mm long. Sepals deeply 5-lobed almost to the base, 3-4 x 0.2 mm, lobes lanceolate, margin and apex setaceous above the middle, apex subacute. Petals slightly pinkish, 3 x 1.5 mm, tube linear, 2 mm



Choudhury, Anwaruddin. 2005. "New Sites of Nepenthes Khasiana from Meghalaya with new Eastern and Western Range Extensions." *The journal of the Bombay Natural History Society* 102, 252–253.

View This Item Online: <https://www.biodiversitylibrary.org/item/188389>

Permalink: <https://www.biodiversitylibrary.org/partpdf/154835>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.