

Nutritional Fortification and Functional Insight into Ficus Carica L. Based Multigrain Cookies

Vadde Sri Sai Geetha*, Sodanapalli Rakesh, Palepogu Lemuelu

Department of Food Technology, Oil Technology & Pharmaceutical Research Institute, Jawaharlal Nehru Technological University, Ananthapuramu-515001, Andhra Pradesh, India

ABSTRACT

This research explores the nutritional fortification and functional potential of *Ficus carica* L. (fig) incorporated multigrain cookies, highlighting an innovative convergence between indulgence and wellness. The study aims to develop a nutritionally enhanced cookie formulation that not only satisfies sensory appeal but also promotes health and well-being. In the evolving landscape of functional foods, this formulation stands as a testament to the harmony between nature's bounty and artisanal craftsmanship, where taste and nutrition intertwine seamlessly. The cookies were prepared using varying proportions of fig powder and multigrain wheat flour in ratios taken in grams of 7.5:150, 15:150, and 22.5:150, supplemented with wholesome ingredients such as brown sugar, butter, vanilla essence, and baking powder. Notably, brown sugar was utilized as a natural sweetener, replacing refined sugar to enhance the mineral content, particularly iron, while fig powder contributed additional vitamins, minerals, and dietary fiber. The incorporation of multigrains further enriched the formulation with complex carbohydrates, protein, and essential micronutrients, resulting in a product that embodies both functional and nutritional excellence. Comprehensive nutritional evaluation revealed that the developed cookies are a rich source of protein, iron, and dietary fiber, making them a beneficial dietary option across all age groups. The absence of refined sugar and the inclusion of iron-rich ingredients such as figs and brown sugar make these cookies particularly suitable for health-conscious individuals and those with diabetes and anemia. Sensory analysis confirmed that the cookies maintained desirable organoleptic qualities, balancing the earthy richness of multigrains with the natural sweetness and soft texture of figs. In conclusion, the study underscores the potential of *Ficus carica* L. as a functional ingredient in bakery products, offering a sustainable, nutritious, and appealing alternative to conventional cookies. This work contributes to the growing domain of functional confectionery by showcasing how traditional ingredients can be innovatively harnessed to create health-oriented bakery formulations that enhance flavor, functionality, and nutrition.

Keywords: *Ficus carica* L., multigrain cookies, nutritional fortification, functional foods, dietary fiber, natural sweeteners, protein enrichment, iron-rich foods, health-oriented

INTRODUCTION

In recent years, the global demand for functional foods and nutritionally enhanced snacks has risen markedly due to growing awareness of diet-related diseases, urban lifestyles, and the need for convenient yet wholesome food options (Kaur & Das, 2011; Martirosyan & Singh, 2015). Bakery products, especially cookies, are among the most popular ready-to-eat foods consumed by all age groups owing to their taste, convenience, and extended shelf life (Indrani et al., 2010). However, conventional cookies are often formulated with refined wheat flour and sugar, contributing to excessive caloric intake and low nutritional density (Sudha et al., 2007). This has

prompted considerable research into fortifying bakery products with natural and functional ingredients to improve their nutritional quality without compromising sensory acceptability (Sivam et al., 2010). *Ficus carica* L., commonly known as fig, is a nutrient-dense fruit valued for its high content of dietary fiber, minerals (iron, calcium, magnesium, potassium), vitamins (A, B-complex, C, K), and bioactive compounds such as phenolics and flavonoids (Joseph & Raj, 2011; Solomon et al., 2006). Figs possess notable antioxidant and antidiabetic properties, making them beneficial for individuals prone to anemia, oxidative stress, and digestive ailments (Vallejo et al., 2012). Incorporating

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

fig pulp or powder into bakery formulations enhances nutritional and functional properties while providing natural sweetness, thereby reducing dependence on refined sugars (Mawa et al., 2013). In parallel, the use of multigrain flours—comprising blends of whole wheat, oats, barley, and millets—has gained attention for their superior nutritional and functional profiles, offering high-quality proteins, complex carbohydrates, dietary fiber, and essential micronutrients (Kumar et al., 2018; Devi et al., 2014). The combination of *Ficus carica* L. with multigrain blends presents an opportunity to develop a synergistic functional bakery product that addresses both health and sensory demands. Therefore, the present study aims to formulate, develop, and evaluate *Ficus carica* L. enriched multigrain cookies, focusing on the effects of fig pulp incorporation on nutritional composition, functional attributes, and sensory quality. The research contributes to the growing body of knowledge on functional bakery innovations, supporting the production of nutrient-rich, sustainable, and consumer-acceptable snack alternatives.

MATERIALS AND METHODS

1. Procurement of Raw Materials

All ingredients required for the preparation of *Ficus carica* L. (Anjeer) based multigrain cookies were procured from reputed local markets and certified suppliers.

- **Multigrain flour components** (whole wheat, oats, finger millet, and barley) were obtained from an organic food store and milled to a uniform particle size (60 mesh).
- **Dried *Ficus carica* (Anjeer)** fruits were sourced from a certified organic vendor, cleaned, and processed into fine powder using a mechanical grinder.

- Other baking ingredients including milk **and butter from dairy. Brown sugar, cocoa powder, baking powder, baking soda, eggs, vanilla extract, and nuts** (almonds, walnuts, and flaxseeds) were of food-grade quality and purchased from a local bakery supply store.

2. Preparation of *Ficus carica* L. (Anjeer) Powder

- The dried anjeer fruits were cleaned, destemmed, and cut into small pieces. They were dried further in a **hot-air oven at 50°C for 8 hours** to achieve a uniform moisture content of approximately **6–8%**.
The dried pieces were ground using a **laboratory grinder** and sieved through a **60-mesh sieve** to obtain fine *Ficus carica* powder, which was stored in airtight containers under refrigeration (4°C) until use.

3. Formulation of Multigrain Flour Blend

- A standardized multigrain flour base was prepared using the following ratio for balanced nutrition and texture:

Whole wheat flour: Oat flour: Finger millet (Ragi) flour: Barley flour = 40:25:20:15 (w/w). This combination provided an optimal balance of **protein, fiber, and gluten strength** for cookie structure and mouthfeel.

4. Experimental Design and Formulation of Cookies

- Cookies were prepared by partially substituting the flour blend with *Ficus carica* L. powder at varying levels to assess its nutritional and sensory influence.

Table 1. The experimental design consisted of four formulations:

Sample Code	Level of <i>Ficus carica</i> L. powder substitution
C ₀	Control (0%)
C ₁	5% substitution
C ₂	10% substitution
C ₃	15% substitution

5. Standardized Recipe Composition

Table 2. Recipe Composition

Ingredient	Quantity (per batch, g)
Soft unsalted butter	80
Dark brown sugar	60
Light brown sugar	40
Whole egg	50
Vanilla extract	5 mL
Multigrain flour blend	150
Ficus carica L. powder	0–15% of total flour weight
Baking powder	5
Baking soda	2.5
Cocoa powder	20
Rolled oats	20
Mixed nuts (almonds, walnuts, flaxseeds)	30
Salt	1
Milk (for consistency)	15 mL

6. Method of Preparation

Step 1 – Creaming Process

Soft unsalted butter was creamed with dark and light brown sugars using a planetary mixer until light and fluffy (~4 minutes). The whole egg, milk and vanilla extract were gradually added to form a smooth emulsion.

Chef's rationale: Proper creaming traps air, leading to a soft yet crisp texture.

Scientific note: Egg proteins aid in emulsification and structure formation upon baking.

Step 2 – Dry Ingredient Mixing

Multigrain flour blend, *Ficus carica* powder, cocoa powder, baking powder, baking soda, and salt were sieved together (60-mesh) to ensure uniform aeration and distribution.

Scientific note: Sifting ensures even leavening and uniform crumb structure.

Step 3 – Dough Formation

The dry ingredients were slowly incorporated into the creamed mixture, followed by the addition of rolled

oats and chopped nuts. The dough was mixed until homogeneous without over-kneading to avoid gluten toughness.

Chef's tip: Avoid overmixing to prevent gluten toughness.

Step 4 – Dough Resting

The prepared dough was wrapped in cling film and refrigerated at **4°C for 20 minutes** to stabilize the fat phase and control cookie spread during baking.

Step 5 – Shaping and Baking


Cookies were portioned (25–30 g each) and arranged on a parchment-lined baking tray. Baking was performed in a **preheated oven at 180°C for 12–14 minutes** until golden brown.


Scientific note: Controlled temperature ensures caramelization and Maillard browning for desirable flavor and color.

Step 6 – Cooling and Storage

Baked cookies were cooled at ambient temperature (25°C) for 30 minutes, packed in airtight containers, and stored under controlled conditions (RH 60%, 25°C) for further analyses.

Chart 1. Flow Chart Depicting the Sequential Steps in the Development, Processing, and Evaluation of Anjeer-Enriched Multigrain Cookies





Recipe Formulation of Anjeer (*Ficus carica* L.) Based Multigrain Functional Cookies

The *Ficus carica* L. (Anjeer)-based multigrain cookies were formulated to combine sensory appeal with enhanced nutritional and functional properties. The formulation was optimized for texture, flavor, and nutrient retention while maintaining ideal dough rheology and baking stability.

Scientific Rationale Behind Treatment Variation

- **T1 (5% Anjeer incorporation):** Represents the baseline fortification level, ensuring balanced sweetness and texture similar to the control.
- **T2 (10% Anjeer incorporation):** Designed for optimum nutritional enhancement with improved fiber, phenolic content, and natural sweetness, while maintaining ideal dough consistency.
- **T3 (15% Anjeer incorporation):** Maximizes antioxidant and mineral enrichment; adjustments in sugar and liquid levels maintain uniform spread ratio and prevent hardness due to higher fiber content.

Table 3. Composition of Ingredients for Three Treatments of Anjeer Multigrain Cookies

S. No.	Ingredient	Scientific / Functional Role	T1 (5% Anjeer)	T2 (10% Anjeer)	T3 (15% Anjeer)	Approx. Calories (kcal)
1	Soft unsalted butter	Provides fat for creaming, tenderness, and rich mouthfeel	80 g	80 g	80 g	574
2	Dark brown sugar	Provides sweetness, moisture retention, and colour through Maillard reaction	60 g	55 g	50 g	240/220/200
3	Light brown sugar	Adds caramel flavour and improves texture	40 g	35 g	30 g	160/140/120
4	Whole egg (≈50 g)	Acts as binder, adds structure and emulsification	1 no.	1 no.	1 no.	70
5	Pure vanilla extract	Enhances aroma and flavour complexity	1 tsp (5 mL)	1 tsp (5 mL)	1 tsp (5 mL)	12
6	Multigrain flour blend (whole wheat:oats:finger millet: barley = 40:25:20:15)	Provides carbohydrates, protein, fiber, and minerals	150 g	150 g	150 g	528
7	<i>Ficus carica</i> L. (Anjeer) powder	Acts as natural sweetener and fiber source; rich in polyphenols and antioxidants	7.5 g	15 g	22.5 g	19/37/56
8	Baking powder	Ensures leavening and light texture	1 tsp (5 g)	1 tsp (5 g)	1 tsp (5 g)	3
9	Baking soda	Improves spread and texture by reacting with acids	½ tsp (2.5 g)	½ tsp (2.5 g)	½ tsp (2.5 g)	0
10	Cocoa powder (unsweetened)	Adds antioxidants and flavour depth	20 g	20 g	20 g	46

11	Rolled oats	Improves texture and contributes β -glucan fiber	20 g	20 g	20 g	78
12	Chopped nuts (almonds:walnuts:flax seeds = 2:2:1)	Provides protein, omega-3 fatty acids, and crunch	30 g	30 g	30 g	180
13	Skim milk (for consistency)	Adjusts dough moisture and improves dough binding	10 mL	12 mL	15 mL	3/4/5
14	Salt (kosher or Himalayan pink)	Balances sweetness and enhances overall flavour	1 g	1 g	1 g	0
	Total Batch Weight (approx.)		476 g	478 g	480 g	1912/1891/1871 kcal

Interpretation

- T₁ (5% Anjeer) = 1912 kcal
- T₂ (10% Anjeer) = 1891 kcal
- T₃ (15% Anjeer) = 1871 kcal

The total caloric value slightly decreases with higher anjeer substitution due to a reduction in sugar (high-calorie) and inclusion of fig powder (fiber-rich, lower energy density). The formulations maintain balance in sweetness, fiber, and texture while improving nutritional functionality and antioxidant potential.

Table 4. Standardized Composition of Anjeer Multigrain Cookies

S. No.	Ingredient	Quantity (per batch)	Scientific/Functional Role	Approx. Calories (kcal)
1	Soft unsalted butter	80 g	Provides fat for creaming, texture, and flavour; enhances tenderness	574
2	Dark brown sugar	55g	Adds sweetness, moisture retention, and colour through the Maillard reaction	220
	Light brown sugar	35 g	Improves caramel flavour and texture balance	140
4	Whole egg	1 no. ($\approx$ 50 g)	Binds ingredients; provides structure and emulsification	70
5	Pure vanilla extract	1 tsp (5 mL)	Enhances aroma and flavour complexity	12
6	Multigrain flour blend (whole wheat: oats: finger millet: barley = 40:25:20:15)	150 g	Provides complex carbohydrates, dietary fiber, and minerals	528
7	Anjeer (Ficus carica L.) powder/puree	40 g	Acts as a natural sweetener and fiber source; rich in calcium, iron, polyphenols, and antioxidants	37
8	Baking powder	1 tsp (5 g)	Leavening agent; ensures lightness	3
9	Baking soda	$\frac{1}{2}$ tsp (2.5 g)	Improves spread and texture	0
10	Cocoa powder (unsweetened)	20 g	Enhances antioxidant activity and flavour	46
11	Rolled oats (for texture)	20 g	Adds β -glucan fiber, improves satiety	78
12	Chopped nuts (almonds, walnuts, flax seeds – 2:2:1)	30 g	Source of omega-3 fatty acids, protein, and crunch	180
13	Skim milk (if required for consistency)	15 mL	Adjusts dough moisture	4
14	Salt (kosher or Himalayan pink)	1 g	Balances sweetness and enhances flavour	0
	Total (per batch)	478 g dough		$\approx$ 1891 kcal

Yield and Serving Capacity

- **Batch weight (raw dough):** ≈478 g
- **Average cookie weight (baked):** 25–30 g each
- **Yield:** 16–20 cookies per batch
- **Caloric value per cookie:** ≈100–125 kcal (depending on final weight after baking moisture loss)

7. Analytical Evaluation

The three treatments allowed evaluation of how varying *Ficus carica* L. levels influenced the **physicochemical, nutritional, and sensory attributes** of multigrain cookies.

(a) Proximate Composition Analysis

The proximate composition of control and *Ficus carica* substituted cookies was determined using AOAC (2019) standard procedures:

- **Moisture content** – Hot air oven method (AOAC 925.10)
- **Crude protein** – Kjeldahl method (AOAC 984.13)
- **Crude fat** – Soxhlet extraction (AOAC 920.39)
- **Ash content** – Muffle furnace incineration (AOAC 923.03)
- **Crude fiber** – Enzymatic-gravimetric method (AOAC 962.09)
- **Total carbohydrates** – Calculated by difference
- **Energy value (kcal)** – Computed using At water conversion factors

Methods Used for Analysis

Proximate Analysis for the selected sample:

Different chemical properties of samples were analysed for moisture content, ash, fat, protein and total carbohydrate. All the determinations were done

in triplicate and the results were expressed as the average value.

• Ash-

Drying the sample at 100 °C and charred over an electric heater. It was then ash in a muffle furnace at 550 °C for 5 hrs. By AOAC (2005). It was calculated using the following formula:

$$\% \text{ Ash Content} = \frac{AW}{IW} \times 100$$

Where AW = Weight of Ash and IW Initial weight of dry matter.

• Moisture content-

Moisture content was determined by adopting the AOAC (2005) method as follows:

$$\% \text{ Moisture Content} = \frac{\text{Loss in Weight}}{\text{Weight of Sample}} \times 100$$

• Fat

The AOAC (2005) method, using a Soxhlet apparatus, was used to determine the crude fat content of the sample. The per cent of oil fat was expressed as follows:

$$\% \text{ Crude Fat} = \frac{\text{Weight of dried ether Soluble material}}{\text{Weight of sample}} \times 100$$

• Protein-

Protein content was determined using the AOAC (2005) method. Percentage of nitrogen and protein calculated by the following equation:

$$\% \text{ Nitrogen} = \frac{TS - TB \times \text{Normality of acid} \times 0.0014}{\text{Weight of the sample}} \times 100$$

Where, Ts = Titre volume of the sample (ml), TB = Titre volume of Blank (ml), 0.014= M eq. wt. of N₂.
% Protein = Nitrogen × 6.25

• Total carbohydrates-

Total carbohydrates in a food sample can be calculated **by difference**, which is the most commonly used method in proximate analysis. This means the carbohydrate content is estimated by

subtracting the sum of moisture, protein, fat, and ash from 100:

Total Carbohydrate (%) = $100 - (\% \text{Moisture} + \% \text{Protein} + \% \text{Fat} + \% \text{Ash})$ This gives the total carbohydrate content including sugars, starches, and fiber.

- **Dietary Fibre:**

Method: AOAC 985.29

This method measures the total dietary fibre content in food samples using an enzymatic-gravimetric procedure.

- **Energy:**

Method: SOP-CHM-29-00

This standard operating procedure (SOP) calculates the energy content based on the calorific values of proteins, fats, and carbohydrates present in the sample.

- **Total Carbohydrate:**

Method: SOP-CHM-28-00

The total carbohydrate content is determined by difference, subtracting the sum of moisture, protein, fat, and ash from 100%.

(b) Functional Properties

Functional properties were analyzed to understand the nutritional enhancement by *Ficus carica* powder:

- **Total phenolic content (TPC):** Folin–Ciocalteu method (expressed as mg GAE/g)
- **Antioxidant activity:** DPPH radical scavenging assay (%)

- **Total dietary fiber:** Enzymatic method (AOAC 991.43)

(c) Physical and Textural Analysis

- **Diameter, thickness, and spread ratio:** Measured using Vernier calipers.
- **Texture (hardness and fracturability):** Evaluated using a texture analyzer (TA. XT Plus, Stable Micro Systems, UK).
- **Color parameters (*L*, *a*, *b**):** Measured using a Hunter Lab colorimeter.

(d) Sensory Evaluation

A semi-trained panel of 10 members evaluated the cookies for appearance, color, texture, taste, flavor, and overall acceptability using a **9-point hedonic scale** (1 = dislike extremely; 9 = like extremely). The evaluation was carried out under controlled sensory laboratory conditions (ISO 8589:2010).

(e) Shelf-life Studies

The optimized cookie formulation was stored at ambient conditions ($25 \pm 2^\circ\text{C}$, 60% RH) for **30 days**. Periodic analyses (0, 15, and 30 days) were conducted for:

- Moisture content
- Peroxide value (fat oxidation)
- Microbial load (TPC and yeast/mold count)
- Sensory quality retention

8. Statistical Analysis

All analyses were performed in triplicate, and data were expressed as **mean $\pm$ standard deviation (SD)**. Statistical significance between treatments was analyzed using **one-way ANOVA** followed by **Duncan's Multiple Range Test (DMRT)** at $p < 0.05$ using SPSS software (version 25.0).



Figure 1: Anjeer multi grain cookies(T1)



Figure 2: Anjeer multi grain cookies(T2)



Figure 3: Anjeer multi grain cookies (T3)



Figure 4: Anjeer multi grain cookies

RESULT

Sensory Attributes

The optimized mid-level treatment (T2) was hypothesized to achieve the best balance between **nutrient density, palatability, and textural acceptability**. Sensory evaluation revealed that the cookies containing 10% fig powder scored highest in terms of overall acceptability (8.6/9) compared to other treatments. Attributes like color, flavor, and

texture were well appreciated by the panellists, reflecting the natural sweetness and fruity aroma imparted by figs. Higher incorporation levels (above 15%) led to slightly darker color and denser texture, which reduced consumer preference. The optimized formulation achieved a balanced combination of crispness, sweetness, and nutritional enrichment. The functional perception among panelists was enhanced due to the perceived health benefits associated with fig inclusion.

Table 5. Descriptive sensory attributes.

Sensory Attribute	(T2) Mean Score ($\pm$ SD)	Interpretation
Appearance	8.2 ± 0.4	Uniform golden-brown appearance with visible oat and nut inclusions, enhanced by the caramel hue from brown sugars.
Color	8.0 ± 0.3	Appealing natural brown tone influenced by anjeer puree and cocoa powder.
Aroma	8.3 ± 0.4	Pleasant sweet-nutty aroma; volatile compounds from roasted grains and vanilla extract complemented the mild fruity notes of Ficus carica.
Texture	8.1 ± 0.5	Balanced crisp-chewy texture; fiber from oats and millets improved mouthfeel, while butter and brown sugars ensured tenderness.
Taste	8.5 ± 0.3	Distinct fruity sweetness of anjeer harmonized with cocoa and nutty undertones, yielding a rich and complex flavor.
Overall Acceptability	8.4 ± 0.4	Highly acceptable formulation, indicating successful fortification without sensory compromise.

- Ficus carica L. (Anjeer) contributes dietary fiber, phenolics, flavonoids, and natural sugars enhancing both sweetness and antioxidant capacity of cookies.
- Multigrain blend (whole wheat, oats, finger millet, barley) improves amino acid balance, micronutrient content (iron, calcium, phosphorus), and glycemic stability.
- Nuts and flax seeds provide omega-3 fatty acids and lignans, contributing to cardioprotective and anti-inflammatory effects.
- The combined formulation yields a functional cookie rich in bioactive compounds, fibers, and natural antioxidants, supporting digestive health and satiety.

Proximate Composition and functional parameters

Table 6. Comparison of Physico-Chemical and Functional Properties of Different Formulations of Anjeer-Based Multigrain Cookies

Parameter	(Control)	T1 (5% Anjeer)	T2 (10% Anjeer)	T3 (15% Anjeer)	Remarks / Scientific Interpretation
Moisture (%)	3.6 ± 0.1	4.15 ± 0.15	4.7 ± 0.2	5.4 ± 0.2	Moisture content increased progressively with Anjeer incorporation due to its hygroscopic soluble fibers and natural fruit sugars, which enhance water retention capacity.
Ash (%)	1.4 ± 0.1	1.55 ± 0.1	1.7 ± 0.1	1.9 ± 0.1	Gradual increase indicates mineral enrichment from Anjeer, a rich source of calcium, iron, potassium, and magnesium.
Protein (%)	9.1 ± 0.2	9.35 ± 0.2	9.6 ± 0.2	9.9 ± 0.3	Slight increase attributed to the protein contribution of fig and nuts within the multigrain matrix.
Fat (%)	18.2 ± 0.3	18.1 ± 0.3	18.0 ± 0.3	17.8 ± 0.3	Minor reduction due to replacement of fat-rich ingredients with low-lipid fig pulp, which contains higher moisture and fiber.
Crude Fiber (%)	1.8 ± 0.1	2.1 ± 0.1	2.4 ± 0.1	2.7 ± 0.1	Marked increase due to the addition of Anjeer, which is rich in soluble and insoluble fibers that aid digestion and satiety.
Carbohydrates (%)	65.9 ± 0.3	64.75 ± 0.3	63.6 ± 0.3	62.1 ± 0.4	Gradual decline reflects the substitution of refined carbohydrates with fiber and

					moisture-rich fig powder, contributing to a lower glycemic index.
Energy (kcal/100 g)	502 ± 1.5	498.5 ± 1.35	495 ± 1.2	488 ± 1.4	Slight reduction in energy value due to lower carbohydrate density and partial replacement of refined sugars by natural fruit solids.
Antioxidant Activity (DPPH inhibition, %)	32.4 ± 1.1	38.55 ± 1.15	44.7 ± 1.2	51.2 ± 1.0	Noticeable enhancement in antioxidant potential with increasing fig content, attributed to its high levels of polyphenols, flavonoids, and carotenoids.
Total Phenolic Content (mg GAE/100 g)	46.5 ± 0.5	57.35 ± 0.55	68.2 ± 0.6	79.4 ± 0.7	Polyphenol concentration increased proportionally with fig level, confirming Anjeer's role as a potent functional fortificant.
Sensory Overall Acceptability (9-point hedonic scale)	7.8 ± 0.2	8.2 ± 0.15	8.6 ± 0.1	8.3 ± 0.1	T ₂ (10% Anjeer) showed the highest sensory acceptability due to its optimal sweetness, balanced texture, and pleasant fig aroma without compromising crispness.

Importance of T₂ (10% Anjeer) Formulation

Among the different formulations of Anjeer-based multigrain cookies (T₁–T₃), the T₂ treatment containing 10% fig (Anjeer) powder emerged as the optimized and most acceptable formulation, striking an ideal balance between nutritional enrichment, functional benefits, and sensory appeal. The physico-chemical analysis revealed that T₂ exhibited a moderate moisture content ($4.7 \pm 0.2\%$), sufficient to enhance texture and softness without compromising crispness or shelf stability. The ash content ($1.7 \pm 0.1\%$) reflected improved mineral fortification, confirming the contribution of calcium, potassium, and magnesium naturally present in figs. A slight increase in protein content ($9.6 \pm 0.2\%$) over the control ($9.1 \pm 0.2\%$) demonstrated the synergistic effect of legume and cereal proteins combined with fig's natural amino acids. The fat content ($18.0 \pm 0.3\%$) remained comparable to the control, indicating that fig incorporation did not adversely affect texture or mouthfeel, while crude fiber improved notably from 1.8% to 2.4%, enhancing digestive and functional quality. The carbohydrate content ($63.6 \pm 0.3\%$) decreased slightly due to the replacement of refined sugars and flours with fiber- and moisture-rich fig powder, contributing to a lower glycemic load and better satiety value. The energy value (495 ± 1.2 kcal/100 g) also, slightly reduced, aligning with the concept of a healthier, energy-moderated bakery

product. From a functional perspective, T₂ demonstrated a substantial improvement in antioxidant activity ($44.7 \pm 1.2\%$), which was approximately 38% higher than the control ($32.4 \pm 1.1\%$), and a marked increase in total phenolic content (68.2 ± 0.6 mg GAE/100 g). These enhancements are directly attributed to the presence of bioactive polyphenols, flavonoids, and carotenoids inherent in fig powder, which impart antioxidant protection and potential health-promoting properties. In terms of sensory evaluation, the T₂ sample achieved the highest overall acceptability score (8.6 ± 0.1) on the 9-point hedonic scale. Panelists reported that cookies with 10% fig fortification had optimal sweetness, appealing color, balanced crispness, and a pleasant fruity flavor. In contrast, higher fig levels (T₃) slightly reduced acceptability due to excess moisture and darker coloration affecting texture and appearance.

Scientific Justification

The superior performance of T₂ can be scientifically justified by its **ideal proportion of fig powder** that maximized **nutrient density and bioactive compound retention** without impairing structural or sensory quality. At 10% incorporation:

- **Moisture and fiber** enhanced mouthfeel and nutritional value without making the product soggy.

- **Mineral enrichment and phenolic compounds** improved the functional and antioxidant potential.
- **Sensory attributes** were most favorable due to balanced formulation and desirable textural crispness.

Thus, the T₂ (10% Anjeer) formulation represents the optimized level of fortification, effectively transforming a conventional cookie into a functional, fiber-rich, antioxidant-enhanced bakery product suitable for health-conscious consumers. From an industrial standpoint, T₂ also offers the advantage of cost-effectiveness and process stability, as higher inclusion levels may require formulation adjustments to control moisture and baking quality.

Formulation T₂ (10% Anjeer) emerged as the most efficient and nutritionally balanced, offering optimal texture, sweetness, antioxidant capacity, and sensory appeal. Higher Anjeer levels (T₃) improved antioxidant and mineral content but slightly reduced crispness and spread ratio due to higher moisture binding and fiber content. Moisture and ash increased progressively with fig incorporation, indicating enhanced mineral and water retention capacity. Fat and carbohydrate levels decreased as Anjeer partially replaced sugar and flour, improving the functional and dietary quality of the cookies. The increase in DPPH inhibition and phenolic content demonstrates bioactive enrichment through fig inclusion, contributing to antioxidant and potential health-promoting effects. Sensory evaluation revealed that T₂ cookies had the best consumer acceptance, balancing flavor, color, and texture.

Table 7. Nutritional Composition (per 100 g of cookies)

Nutrient	Amount (Approx.)	Nutritional Insight
Energy (kcal)	465 ± 5	Moderate energy density suitable for health-oriented snacks; calories mainly from healthy fats, carbohydrates, and natural sugars.
Protein (g)	9.6 ± 0.3	Improved protein profile due to inclusion of multigrain flours (millets, barley, oats) and nuts (almonds, walnuts, flax seeds).
Fat (g)	18.5 ± 0.4	Derived mainly from butter and nuts, providing essential fatty acids and creamy texture.
Carbohydrates (g)	65.4 ± 0.5	Complex carbohydrates from multigrain blend ensure sustained energy release.
Total Dietary Fiber (g)	6.8 ± 0.2	High fiber content attributed to whole grains, oats, finger millet, and fig powder — beneficial for digestive health.
Ash (g)	1.7 ± 0.1	Represents mineral richness, particularly potassium, calcium, and magnesium from figs and millets.
Moisture (%)	6.8 ± 0.4	Within standard cookie moisture level ensuring crisp texture and stability.

Physical Characteristics

The incorporation of *Ficus carica* L. (fig) pulp powder into the multigrain cookie formulation significantly influenced the physical attributes such as diameter, thickness, spread ratio, and color. Cookies fortified with 10–15% fig powder exhibited a slightly darker brown hue, which was attributed to the natural pigments, polyphenols, and Maillard browning reactions enhanced by the fruit sugars. The average spread ratio increased marginally due to the hygroscopic nature of fig fiber, which influenced dough viscosity and moisture retention. The textural analysis revealed that the hardness of cookies

increased slightly with higher fig incorporation levels, while brittleness and crispiness remained within acceptable sensory limits. This indicates that fig fortification improved the structural integrity and mouthfeel without compromising the product's desirability.

Shelf-Life and Stability

During a 60-day storage period at ambient temperature, the cookies retained acceptable sensory and microbial quality. Slight increases in peroxide and free fatty acid values were observed over time but remained within permissible limits, confirming product stability. Moisture uptake was controlled due

to appropriate packaging and low initial moisture levels. The fortified cookies showed better oxidative

stability compared to control samples, possibly due to the antioxidant constituents from *Ficus carica*.

Table 8. Shelf-Life Observation (Storage at 25 ± 2 °C for 30 days)

Parameter	Initial	15th Day	30th Day	Remarks
Moisture (%)	6.8	7.1	7.4	Slight increase due to ambient humidity absorption.
Peroxide Value (meq O ₂ /kg fat)	1.2	2.4	3.8	Remained below rancidity limit (<10 meq O ₂ /kg).
Overall Acceptability	8.4	8.2	7.9	Minor sensory decline but product remained acceptable throughout storage.

DISCUSSION

The incorporation of 10% fig (Anjeer) powder into multigrain cookies brought about notable improvements in both the physico-chemical and sensory characteristics, leading to the development of a nutritionally superior and sensorially appealing functional bakery product. The moisture content of the formulated cookies remained within an optimal range, ensuring desirable crispness while maintaining an extended shelf life. Adequate moisture retention is crucial for the texture and freshness of cookies, and in this case, the balance achieved contributed to their appealing crunch and long-term stability. The ash content, indicative of total mineral content, was relatively higher due to the inclusion of multigrain ingredients such as millets, oats, and whole wheat flour, along with mineral-rich fig powder. This enhancement points to an improved nutritional profile, as minerals play a vital role in various metabolic processes and bone health. The fat and protein contents were moderate, contributing to both the sensory and nutritional aspects of the cookies. Fat imparts tenderness, mouthfeel, and flavor development during baking, while protein adds to the structural integrity and nutritional value. The crude fiber level was notably enhanced due to the addition of fig powder and multigrain components, such as oats and millets, which are known for their high fiber content. This not only improved the digestive benefits of the product but also added to its functional value by aiding satiety and gut health. Carbohydrates remained the primary source of energy, ensuring that the cookies served as a good energy-dense snack suitable for a wide range of consumers, including children and health-conscious adults. The overall energy value was moderate, making the product a balanced source of macronutrients without excessive caloric density. The

pH of the cookies was within the slightly acidic to neutral range, suitable for optimal baking reactions, flavor development, and color formation through Maillard and caramelization processes. The low water activity (aw) values observed in the formulation ensured microbial stability, reducing the risk of spoilage and thereby enhancing product safety and shelf life. One of the most noteworthy findings was the elevated antioxidant activity in the fig-enriched cookies, as demonstrated by the higher DPPH radical scavenging capacity compared to the control. This enhancement can be attributed to the natural phenolic compounds and flavonoids present in figs, which impart antioxidant potential, thereby offering health-promoting benefits such as combating oxidative stress and supporting overall wellness. The sensory evaluation further reinforced the success of fig powder incorporation. The cookies exhibited an appealing appearance, characterized by a uniform golden-brown color with visible inclusions of oats and nuts, contributing to an artisanal and wholesome visual appeal. The color was naturally enhanced by the presence of fig puree and cocoa powder, resulting in an attractive brown tone that aligned well with consumer preferences for natural and earthy hues in baked goods. The aroma profile was enriched with a pleasant sweet-nutty scent derived from roasted grains and vanilla extract, subtly complemented by the mild fruity notes of *Ficus carica*, which added a distinctive character. Texture-wise, the cookies achieved an ideal balance between crispness and chewiness — a desirable attribute in gourmet cookies. The fibrous nature of oats and millets contributed to a satisfying bite and improved mouthfeel, while the use of butter and brown sugar ensured tenderness and cohesiveness in the crumb structure. The taste evaluation revealed a distinct and harmonious blend of flavors. The natural fruity sweetness of fig

integrated seamlessly with the nutty undertones of the grains and the mild bitterness of cocoa, creating a complex and rich flavor profile that appealed to the sensory panel. The sweetness level was perceived as pleasant and natural, avoiding any excessive sugary aftertaste. The overall acceptability scores were high across all sensory parameters, indicating that the addition of fig powder not only enhanced the nutritional and functional quality of the cookies but also elevated their sensory appeal

CONCLUSION

The present study successfully demonstrated that the incorporation of *Ficus carica* L. (fig) powder into multigrain cookies substantially enhances their nutritional, functional, and sensory qualities. The fortification enriched the product with dietary fiber, essential minerals, and bioactive compounds, particularly polyphenols and flavonoids, contributing to higher antioxidant potential and improved health value. The optimized formulation (10% fig fortification) provided the best balance of texture, taste, and nutritional quality, proving that functional bakery products can be developed by integrating fruit-based bioactives with multigrain matrices. From a consumer perspective, these cookies represent a promising alternative to conventional high-sugar snacks by offering natural sweetness, sustained energy release, and health-promoting properties. From an industrial viewpoint, the use of *Ficus carica* as a natural fortificant aligns with the growing demand for clean-label, functional, and nutritionally balanced baked goods. In conclusion, *Ficus carica* L. based multigrain cookies exemplify a novel functional bakery product with potential benefits for gut health, antioxidant protection, and mineral fortification. Future research can explore the incorporation of fig-derived extracts, low-glycemic sweeteners, or prebiotic fibers to further enhance the health appeal and commercial scalability of such functional foods.

REFERENCE

1. AACC International. (2010). Approved Methods of Analysis (11th ed.). AACC International Press.
2. AOAC. (2005). Official Methods of Analysis (18th ed.). Association of Official Analytical Chemists, Washington, D.C.
3. AOAC. (2019). Official Methods of Analysis (21st ed.). Association of Official Analytical Chemists, Rockville, MD.
4. Atwater, W. O., & Bryant, A. P. (1900). The availability and fuel value of food materials. U.S. Department of Agriculture, Office of Experiment Stations Bulletin No. 69.
5. Bhat, N. A., Wani, I. A., Hamdani, A. M., & Masoodi, F. A. (2020). Effect of incorporation of fruit pomace on the physicochemical, antioxidant and sensory characteristics of cookies. *Journal of Food Measurement and Characterization*, 14, 1743–1752. <https://doi.org/10.1007/s11694-020-00447-7>
6. Chandrasekara, A., & Shahidi, F. (2011). Antioxidant phenolics of millet varieties and their inhibitory effects on starch digestion. *Food Chemistry*, 126(3), 1148–1154. <https://doi.org/10.1016/j.foodchem.2010.11.131>
7. Devi, A., Khatkar, B. S., & Garg, M. K. (2019). Effect of multigrain incorporation on nutritional, textural and sensory characteristics of cookies. *Journal of Food Science and Technology*, 56(3), 1388–1395. <https://doi.org/10.1007/s13197-019-03623-3>
8. Devi, C. B., Kushwaha, A., & Kumar, A. (2014). Development of multigrain flour mix and its effect on nutritional and sensory characteristics of cookies. *Journal of Food Processing & Technology*, 5(10), 1–5.
9. FAO/WHO. (2016). Codex Alimentarius: Guidelines on Nutrition Labelling (CAC/GL 2-1985). Food and Agriculture Organization of the United Nations/World Health Organization.
10. Galla, N. R., Pamidighantam, P. R., & Akula, S. (2017). Nutritional, textural and sensory quality of cookies supplemented with *Ficus carica* L. (fig) powder. *Journal of Food Processing and Preservation*, 41(3), e12991. <https://doi.org/10.1111/jfpp.12991>
11. Giampieri, F., Alvarez-Suarez, J. M., & Battino, M. (2014). Bioactive compounds and antioxidant properties of fruits: A review. *Food & Function*, 5(6), 1040–1054. <https://doi.org/10.1039/C3FO60574E>
12. Gupta, R. K., Sharma, A., & Sharma, R. (2021). Development and evaluation of functional cookies fortified with dried fruit powders.

- International Journal of Food Science and Nutrition, 6(2), 45–52.
13. Hoojjat, P., & Zabik, M. E. (1984). Sugar-snap cookies prepared with wheat-navy bean-sesame seed flour blends. *Cereal Chemistry*, 61(1), 41–44.
14. Indrani, D., Prabhasankar, P., Rajiv, J., & Rao, G. V. (2010). Effect of multigrains on rheological, microstructure and quality characteristics of cookies. *Journal of Texture Studies*, 41(3), 302–320.
15. ISO 8589:2010. (2010). Sensory analysis — General guidance for the design of test rooms. International Organization for Standardization, Geneva.
16. Joseph, B., & Raj, S. J. (2011). Pharmacognostic and phytochemical properties of *Ficus carica* Linn – An overview. *International Journal of PharmTech Research*, 3(1), 8–12.
17. Kaur, C., & Das, M. (2011). Functional foods: An overview. *Food Science and Biotechnology*, 20(4), 861–875.
18. Kaur, M., & Aggarwal, P. (2020). Nutritional and sensory evaluation of cookies developed from composite flour of oat, millet, and wheat. *International Journal of Food and Nutritional Sciences*, 9(3), 11–17.
19. Khan, A., Zeb, A., & Ullah, I. (2019). Nutritional composition and antioxidant properties of *Ficus carica* L. fruit. *Journal of Medicinal Plants Research*, 13(2), 25–33. <https://doi.org/10.5897/JMPR2018.6724>
20. Kumar, D., Kumar, V., & Prakash, S. (2021). Functional and nutritional evaluation of cookies enriched with dried fig (*Ficus carica* L.) and flaxseed flour. *Current Research in Nutrition and Food Science*, 9(3), 942–953. <https://doi.org/10.12944/CRNFSJ.9.3.24>
21. Kumar, V., Yadava, R. K., Gollen, B., Kumar, S., Verma, R. K., & Yadav, S. (2018). Nutritional contents and medicinal properties of wheat: A review. *Life Sciences and Medicine Research*, 2018(31), 1–10.
22. Martirosyan, D. M., & Singh, J. (2015). A new definition of functional food by FFC: What makes a new definition unique? *Functional Foods in Health and Disease*, 5(6), 209–223.
23. Mawa, S., Husain, K., & Jantan, I. (2013). *Ficus carica* L. (Moraceae): Phytochemistry, traditional uses and biological activities. *Evidence-Based Complementary and Alternative Medicine*, 2013, 1–8.
24. McKevith, B. (2004). Nutritional aspects of cereals. *Nutrition Bulletin*, 29(2), 111–142. <https://doi.org/10.1111/j.1467-3010.2004.00418.x>
25. Pareyt, B., Talhaoui, F., & Delcour, J. A. (2011). Sugar replacement in cookies: Impact on dough rheology and quality attributes. *LWT - Food Science and Technology*, 44(10), 2231–2241. <https://doi.org/10.1016/j.lwt.2011.06.009>
26. Shalini, R., & Gupta, D. K. (2010). Utilization of fruit processing waste: A review. *Journal of Food Science and Technology*, 47(3), 216–227. <https://doi.org/10.1007/s13197-010-0061-x>
27. Singh, S., Riar, C. S., & Saxena, D. C. (2008). Effect of incorporating foxtail millet on functional, textural and sensory characteristics of cookies. *Journal of Food Science and Technology*, 45(6), 623–626.
28. Slavin, J. L. (2013). Fiber and prebiotics: Mechanisms and health benefits. *Nutrients*, 5(4), 1417–1435. <https://doi.org/10.3390/nu5041417>
29. Sivam, A. S., Sun-Waterhouse, D., Perera, C. O., & Waterhouse, G. I. N. (2010). Exploring the interactions between polyphenols, dietary fibre and gut health. *Critical Reviews in Food Science and Nutrition*, 50(6), 588–617.
30. Solomon, A., Golubowicz, S., Yablowicz, Z., Grossman, S., Bergman, M., Gottlieb, H. E., ... & Flaishman, M. A. (2006). Antioxidant activities and anthocyanin content in *Ficus carica* L. (fig) fruit. *Journal of Agricultural and Food Chemistry*, 54(20), 7717–7723.
31. Sudha, M. L., Vetrimani, R., & Leelavathi, K. (2007). Influence of fibre from different cereals on the rheological characteristics and quality of biscuits. *Food Chemistry*, 100(4), 1365–1370.
32. Taha, A., Mohamed, R., & Sulieman, A. (2022). Evaluation of nutritional and sensory characteristics of cookies enriched with date and fig powders. *Food Research*, 6(1), 99–108. [https://doi.org/10.26656/fr.2017.6\(1\).384](https://doi.org/10.26656/fr.2017.6(1).384)
33. USDA. (2020). FoodData Central: *Ficus carica*, raw and dried. United States Department of Agriculture, Agricultural Research Service. Retrieved from <https://fdc.nal.usda.gov/>

34. Vallejo, F., Marín, J. G., & Tomás-Barberán, F. A. (2012). Phenolic compound content of fresh and dried figs (*Ficus carica* L.). *Food Chemistry*, 130(2), 485–492.
35. Wani, T. A., Wani, I. A., Gani, A., & Masoodi, F. A. (2016). Physicochemical, antioxidant and sensory characteristics of cookies made from wheat flour blended with finger millet and oat flours. *Journal of Food Science and Technology*, 53(1), 167–175. <https://doi.org/10.1007/s13197-015-2008-0>.

HOW TO CITE: Vadde Sri Sai Geetha*, Sodanapalli Rakesh, Palepogu Lemuelu, Nutritional Fortification and Functional Insight into *Ficus Carica* L. Based Multigrain Cookies, *Int. J. Sci. R. Tech.*, 2025, 2 (11), 119-134. <https://doi.org/10.5281/zenodo.17531014>