Journal of Clinical Practice and Medical Research
ISSN 3083-7146
open access • peer reviewed •

Review article

Reconciling Scientific Evidence with Perception: The Case for Palm Oil's Evidence-Based Health Benefits Across Multiple Physiological Systems

Loso Judijanto
IPOSS Jakarta, Indonesia

Abstract

Palm oil (Elaeis guineensis), the most widely consumed vegetable oil globally, has been the subject of considerable debate over its health implications. This qualitative literature review synthesizes recent evidence (2010-2025) examining the health benefits of palm oil and its derivatives for human health. This review analyzes articles to identify significant health-promoting properties of palm oil's unique bioactive constituents, including tocotrienols, carotenoids, and phytosterols. Evidence demonstrates that palm oil contributes to cardiovascular protection through cholesterol management and anti-atherogenic mechanisms, provides neuroprotection against neurodegenerative diseases, supports metabolic regulation, including glycemic control, exhibits antioxidant and anti-inflammatory effects, and demonstrates potential anti-cancer properties. Red palm oil, particularly rich in carotenoids, addresses vitamin A deficiency in vulnerable populations, while palm oil derivatives find extensive applications in food, pharmaceutical, and cosmetic industries. The findings indicate that when consumed as part of a balanced diet, palm oil does not pose incremental cardiovascular risk and offers measurable health benefits through its bioactive compounds. This review concludes that an evidence-based understanding should replace perception-driven narratives about palm oil consumption, recognizing its nutritional value when responsibly sourced and consumed.

Keywords: *palm oil, tocotrienols, carotenoids, cardiovascular health, neuroprotection, antioxidant, anti-inflammatory, vitamin A deficiency, oleochemicals, bioactive compounds.*

Received: 19.11.2025 **Revised:** 05.12.2025 **Accepted:** 06.12.2025 **Published:** 11.12.2025

Funding: This research did not receive any financial support.
Conflict of interest: The authors declare no conflict of interest.
How to cite: Judijanto L. Reconciling Scientific Evidence with Perception: The Case for Palm Oil's Evidence-Based Health Benefits Across Multiple Physiological Systems. J Clin Pract Med Res, 2025;1(3):61-6. DOI: 10.59324/jcpmr.2025.1(3).11

Introduction

Background and Global Context

Palm oil, extracted from the mesocarp of *Elaeis guineensis* fruit, represents the world's most widely produced and consumed vegetable oil, having surpassed soybean oil in global production volumes over recent decades. Indonesia and Malaysia collectively account for approximately 85% of global palm oil production, with Indonesia alone producing 49.12 million tons of crude palm oil (CPO) in 2020. The oil exists in various forms: crude palm oil (CPO) from the fruit mesocarp, palm kernel oil (PKO) from the seed kernel, and red palm oil (RPO), a minimally refined, carotenoid-rich variant that retains its characteristic reddish-orange color [1,2].

Despite its widespread use in food products, cosmetics, and industrial applications, palm oil remains polarized in public discourse. Environmental concerns related to deforestation and biodiversity loss have dominated discussions, often overshadowing scientific evidence regarding its nutritional properties. Simultaneously, nutritional controversies have emerged, with some sources incorrectly attributing cholesterol content to this plant-based oil or suggesting adverse cardiovascular effects comparable to those of animal fats [3].

Problem Statement and Research Gap

A significant disparity exists between public perception and accumulating scientific evidence regarding palm oil's health effects. While negative health perceptions persist, driven in part by historical campaigns against tropical oils, recent research reveals considerable health benefits from palm oil's unique bioactive constituents. Red palm oil contains 600-750 ppm of carotenoids (primarily α- and β-carotenes), 710-774 ppm of vitamin E (predominantly tocotrienols), 325-365 ppm of phytosterols, and 18-25 ppm of coenzyme Q10 [2,4-7].

From an initial pool of studies on palm oil and health, systematic screening of open-access, peer-reviewed articles published between 2020 and 2025 reveals emerging evidence for cardiovascular protection, antioxidant defense, neuroprotection, and improved lipid profiles. However, comprehensive qualitative reviews integrating these findings across multiple health dimensions remain limited, creating a knowledge gap that this review addresses [4].

This work is licensed under a Creative Commons Attribution 4.0 International License. The license permits unrestricted use, distribution, and reproduction in any medium, on the condition that users give exact credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if they made any changes.

Objectives and Scope of the Review

This review evaluates recent scientific evidence (2010-2025) on the health benefits of palm oil and its derivatives for human health. Primary objectives include: (1) examining bioactive constituents and their mechanisms of action, (2) synthesizing evidence across cardiovascular, neurological, metabolic, dermatological, and immunological health dimensions, and (3) evaluating palm oil derivatives' applications in food, pharmaceutical, and cosmetic industries. The review emphasizes research that provides evidence-based insights to inform health professionals, researchers, and consumers about palm oil's therapeutic potential.

Materials and Methods

Literature Search Strategy

Comprehensive literature searches were conducted across multiple databases, including PubMed, Google Scholar, Web of Science, MEDLINE, Scopus, and specialized journals. Primary search terms included "palm oil," "red palm oil," "tocotrienols," "health benefits," "cardiovascular," and "carotenoids." Secondary terms encompassed specific bioactive compounds ("beta-carotene," "phytosterols," "coenzyme Q10") and health outcomes ("neuroprotection," "antioxidant," "anti-inflammatory," "cholesterol," "diabetes"). Derivative-specific searches incorporated "palm kernel oil," "oleochemicals," "palm derivatives," and "fatty acid methyl esters."

Inclusion criteria specified: (1) publication period 2010-2025 to capture comprehensive evidence, (2) peer-reviewed articles in English-language journals, (3) studies examining health benefits, bioactive constituents, or clinical applications of palm oil and derivatives, and (4) research designs including randomized controlled trials, clinical studies, systematic reviews, and high-quality observational studies [8].

Data Extraction and Analysis Framework

Information extracted from selected studies included: bioactive compound identification and quantification; health outcomes measured; study populations and designs; key findings and effect sizes; and mechanistic insights. Thematic analysis identified recurring health benefit themes, organized findings by health dimension (cardiovascular, neurological, metabolic, dermatological, anti-cancer), and recognized patterns across studies. The quality assessment evaluated the research methodology, the strength of the evidence, and the consistency of findings across independent investigations [9].

Results and Analysis

Bioactive Constituents of Palm Oil and Derivatives

Palm oil's health benefits derive from its unique composition of bioactive compounds. The vitamin E complex in red palm oil comprises both tocopherols (30%) and tocotrienols (70%), with concentrations ranging from 600 to 1000 ppm[2]. Unlike most vegetable oils, in which α-tocopherol predominates, palm oil's vitamin E profile features α-tocopherol and γ-tocotrienol as its major components. Tocotrienols exhibit superior antioxidant activity compared to tocopherols due to their unsaturated isoprenoid side chains, which enable better penetration into saturated fatty tissue layers, such as the brain and liver [10].

Carotenoid content distinguishes red palm oil as the richest natural source of β-carotene, containing 500-700 ppm in crude palm oil. Red palm oil provides 15 times more retinol (provitamin A) equivalents than carrots and 300 times more than tomatoes. The α-, β-, and γ-carotenes in red palm oil exhibit quantifiable provitamin A activity, with bioavailability enhanced when consumed in oil form, as a minimum of 5 grams of fat is required for optimal carotenoid absorption [2,10].

The fatty acid profile comprises approximately 44% palmitic acid (C16:0), 5% stearic acid (C18:0), 39% oleic acid (C18:1), and 10% linoleic acid (C18:2), representing a balanced ratio of saturated to unsaturated fatty acids [11]. Critically, the stereospecific positioning of fatty acids in palm oil triacylglycerols differs from that in animal fats, with unsaturated fatty acids predominantly occupying the sn-2 position, thereby enhancing digestibility and metabolic handling [12].

Secondary bioactive compounds include phytosterols (325-365 ppm), which competitively inhibit cholesterol absorption, and coenzyme Q10 (10-80 ppm), a critical component in cellular energy production and antioxidant defense [7]. Squalene and polyphenolic compounds contribute additional antioxidant capacity [5,6,13].

Cardiovascular and Lipid Health Benefits

Recent clinical trials demonstrate palm oil's beneficial effects on lipid profiles. A double-blind, randomized controlled trial involving 100 hyperlipidemic Indonesian adults consuming plant sterol-enriched palm oil for 8 weeks showed significant reductions in total cholesterol (6.08%) and LDL cholesterol (7.71%) compared with the control groups. The intervention also improved total cholesterol: HDL and LDL:HDL ratios, indicating comprehensive lipid profile enhancement [14].

Tocotrienols exert cholesterol-lowering effects through multiple mechanisms. They inhibit 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG-CoA reductase), the rate-limiting enzyme in cholesterol biosynthesis. Specifically, δ-tocotrienol and γ-tocotrienol promote HMG-CoA reductase degradation, reducing endogenous cholesterol synthesis. Clinical studies report 7-38% reductions in LDL cholesterol with tocotrienol supplementation [15].

A randomized clinical trial examining tocotrienol-rich fraction (TRF) supplementation (300 mg/day for three months) in patients with chronic kidney disease demonstrated significant reductions in LDL cholesterol in hemodialysis patients ($p=0.04$) and decreased C-reactive protein levels in non-dialysis patients ($p=0.05$). These findings suggest that TRF supplementation mitigates dyslipidemia and inflammation, key cardiovascular risk factors [16].

Palm oil consumption shows non-inferiority to other vegetable oils in healthy populations. Meta-analyses indicate that, while palm oil increases LDL cholesterol compared with polyunsaturated fat-rich oils, it significantly increases HDL cholesterol by 0.07 mmol/L without adversely affecting total cholesterol or triglycerides compared with trans-fat-containing partially hydrogenated oils. A study comparing palm olein with coconut oil found that palm olein consumption resulted in more favorable changes in total cholesterol, LDL cholesterol, TC:HDL ratio, and fasting plasma glucose [17].

Tocotrienols regulate thrombosis and atherosclerosis through multiple pathways. They reduce lipid peroxidation, thereby preventing the formation of oxidized LDL, a critical step in atherosclerotic plaque development. Palm-derived tocotrienols inhibit inflammatory biomarkers, including soluble intercellular adhesion molecule (sICAM), soluble vascular cell adhesion molecule (sVCAM), and reduce monocyte binding activity in endothelial cells, demonstrating anti-atherogenic potential. Red palm oil administration reduces hepatic malondialdehyde, conjugated dienes, and pro-inflammatory cytokines (IL-1β, IL-6, TNF-α), while improving antioxidant enzyme activities [18].

Antioxidant and Anti-inflammatory Effects

Palm oil's antioxidant capacity stems primarily from its tocotrienol and carotenoid content. These compounds function as potent free radical scavengers, interrupting oxidation cascades by donating phenolic hydrogen to peroxyl radicals. In a lipopolysaccharide-induced hepatic injury model, red palm oil pretreatment (200 μL/day for 28 days) significantly reduced hepatic oxidative stress markers and increased antioxidant enzyme activities, including superoxide dismutase, glutathione reductase, and glutathione peroxidase[2,19].

Mechanistic studies reveal that red palm oil upregulates nuclear factor erythroid 2-related factor 2 (Nrf2), a redox-sensitive transcription factor controlling antioxidant gene expression. Nrf2 activation increases

expression of downstream targets, including glutamate-cysteine ligase (GCL) and heme oxygenase-1 (HO-1), enhancing cellular antioxidant capacity. Simultaneously, red palm oil suppresses nuclear factor kappa B (NF-κβ) expression and the associated inflammatory protein IκB kinase (IκKβ), attenuating inflammatory responses [2,19].

Clinical evidence supports anti-inflammatory effects. A study in Indonesian families consuming crude palm oil (140 ml weekly for two months) demonstrated significant reductions in C-reactive protein levels from 0.111 to 0.103 ($p<0.05$). Palm-derived tocotrienols inhibit the production of inflammatory mediators, including interleukin-6 and nitric oxide, in LPS-stimulated macrophages [20].

The oxidative stability of palm oil itself contributes to health benefits. Palm olein demonstrates superior oxidative stability compared to more polyunsaturated fats, with only 22-41% change from baseline after 168 hours of storage, compared to 320-500% change for soybean oil. This stability prevents the formation of toxic oxidation products during cooking and storage, thereby maintaining nutritional quality [21,22].

Neuroprotection and Cognitive Benefits

Tocotrienols exhibit preferential distribution to brain tissue due to their lipophilic properties and ability to penetrate polyunsaturated fat-rich neural membranes more efficiently than tocopherols. A study examining red palm oil's neuroprotective effects in a D-galactose-induced Alzheimer's disease rat model found that red palm oil pretreatment (200-400 mg/kg) significantly improved spatial working memory performance in Y-maze tests and increased brain glutathione levels [2].

Red palm oil administration reversed cognitive impairment by counteracting oxidative stress. Glutathione concentrations increased significantly in red palm oil-treated groups, providing defense against reactive oxygen species that cause neuronal damage and memory loss. The antioxidant properties protect neurons from oxidative injury associated with neurodegenerative processes [2].

Tocotrienols demonstrate specific mechanisms for neuroprotection. They reduce β-amyloid plaque accumulation, a hallmark of Alzheimer's disease pathology, and attenuate neuroinflammation. The vitamin E family, particularly tocotrienols, provides neuroprotection against Parkinson's-induced neurodegeneration, with administration improving motor deficits in Parkinson's models [23].

Systematic reviews of essential oils and natural compounds for neurodegenerative diseases consistently identify palm oil-derived tocotrienols as promising neuroprotective agents, particularly for early-stage intervention in Alzheimer's and Parkinson's diseases. The molecular mechanisms include modulation of cholinergic transmission, enhancement of antioxidant enzyme activities, and maintenance of cerebral blood flow [24].

Metabolic and Diabetes Management Benefits

Evidence regarding palm oil and glucose metabolism indicates neutral to beneficial effects. Multiple studies demonstrate that palm oil consumption does not adversely affect insulin secretion or glucose homeostasis. Research by Sundram et al. and subsequent investigators found that palm oil consumption does not influence insulin activity, secretion, or blood glucose content [25].

A study examining the consumption of tropical vegetable oils (palm, coconut, groundnut) in patients with type 2 diabetes found no negative impact on blood glucose levels, with the oils not inducing dyslipidemia or impairing glycemic control when used with metformin therapy. Interestingly, epidemiological data suggest that the reduction of palm oil consumption correlates with increased diabetes cases in some populations [26].

Tocotrienols contribute to metabolic health through multiple pathways. They modulate glucose metabolism, enhance insulin sensitivity, and reduce chronic inflammation associated with metabolic disorders. Tocotrienol-rich fraction supplementation may reduce metabolic endotoxemia by maintaining gut barrier integrity and reducing lipopolysaccharide absorption, thereby modulating inflammatory responses [10].

Critically, interesterified palm oil demonstrates different metabolic effects than natural palm oil. Interesterified palm oil, where chemical modification redistributes fatty acids to increase saturated fat at the sn-2 position, impairs insulin signaling in liver and muscle tissues and increases insulin secretion from pancreatic β-cells, indicating insulin resistance. These findings emphasize the importance of consuming natural, non-interesterified palm oil products [27].

Skin Health and Dermatological Applications

Palm oil and its derivatives are widely used in skincare and cosmetic formulations. The oil's bioactive constituents—vitamins A, C, and E, as well as beta-carotene—provide multiple dermatological benefits. When applied topically, palm oil deeply nourishes the skin, creates a protective barrier against environmental pollutants, and restores hydration by sealing in moisture [28].

For inflammatory skin conditions, palm oil's anti-inflammatory properties provide therapeutic benefits. Its rich nutrient profile helps calm inflamed skin in conditions such as eczema and psoriasis, reducing redness and discomfort. The antioxidant vitamins combat free radicals responsible for cellular damage and premature aging [29,[30].

Palm oil demonstrates anti-aging properties through multiple mechanisms. Vitamins and antioxidants help reduce oxidative stress on skin cells, helping mitigate fine lines and wrinkles. Coenzyme Q10, naturally present in palm oil, protects skin from premature aging, increases elasticity, and supports collagen content. Regular application supports skin regeneration and wound healing through vitamin A-mediated mechanisms [28].

Palm-based oleochemicals dominate the personal care industry, valued at USD 19.94 billion in 2025 and projected to reach USD 28.51 billion by 2034. Palm oil-derived emulsifiers, surfactants, and fatty alcohols provide excellent dispersion characteristics, produce abundant foam, and possess broad-spectrum antimicrobial properties effective against bacteria and viruses while maintaining skin health [31].

Anti-cancer and Cellular Protection Properties

Tocotrienols demonstrate significant anti-cancer activity across multiple cancer types. A systematic review analyzing the mechanisms of action of palm oil tocotrienols in tumorigenesis found that tocotrienols suppress the growth of various malignancies, including breast, lung, prostate, ovarian, liver, brain, colon, pancreatic, and myeloma cancers [32].

Molecular mechanisms include: (1) disruption of cancer cell proliferation signals through cell cycle regulation, (2) enhanced apoptosis (programmed cell death) in malignant cells, (3) inhibition of angiogenesis, preventing tumor blood supply development, and (4) suppression of metastatic potential. Tocotrienols act on intracellular and extracellular targets involved in cell signaling pathways, demonstrating positive effects on cell proliferation, apoptosis, angiogenesis, metastasis, and inflammation [33].

High-dose tocotrienol-rich fraction supplementation (up to 250 mg/kg body weight) demonstrates tolerability in healthy mice and anticancer effects in syngeneic mouse breast cancer models, with reduced tumor size, decreased metastasis rates, and increased populations of cytotoxic T cells and natural killer cells. The immunomodulatory effects enhance the body's natural cancer defense mechanism s[34].

Importantly, tocotrienols selectively target cancer cells while protecting healthy cells from oxidative damage. This selectivity provides therapeutic potential without adverse effects on normal cellular metabolism, making tocotrienols promising candidates for cancer prevention and adjunctive cancer therapy [35].

Vitamin A Deficiency Prevention

Vitamin A deficiency affects approximately 190 million school-aged children and 19 million pregnant women globally, concentrated in Sub-Saharan Africa and Southeast Asia. Red palm oil provides a highly efficacious solution for improving vitamin A status in at-risk populations [36].

As the richest naturally occurring source of β-carotene, red palm oil offers superior bioavailability compared to synthetic vitamin A supplements. Intervention studies from Africa, Asia, and Latin America demonstrate that red palm oil, when used as a dietary supplement, in-home fortificant, or to fortify staple foods, effectively improves plasma levels of β-carotene, retinol, and α-carotene [2].

A four-day supplementation study comparing red palm oil (25 mg/day carotenoids) with synthetic all-trans β-carotene (25 mg/day) in 72 healthy volunteers found comparable increases in plasma β-carotene levels, confirming the effectiveness of natural palm oil carotenoids. An eight-week study in cystic fibrosis patients—who experience pancreatic insufficiency, leading to malabsorption of lipophilic vitamins—found that daily red palm oil supplementation improved plasma levels of β-carotene, retinol, and α-carotene despite these challenges [2].

Red palm oil intervention in school feeding programs in Burkina Faso and provitamin A status studies using β-carotene-fortified biscuits made with red palm oil demonstrate equivalent improvements in vitamin A status compared to synthetic β-carotene fortification. The oil form provides optimal fat content for carotenoid absorption while delivering additional health-promoting compounds [36].

Palm Oil Derivatives Applications

Palm oil-based oleochemicals represent versatile renewable materials with applications across food, pharmaceutical, and industrial sectors. The global market, valued at USD 27.69 billion in 2025, is projected to reach USD 39.94 billion by 2032, exhibiting a 5.37% CAGR.

In food and beverage applications, palm oil derivatives function as emulsifiers, stabilizers, and texture enhancers. RBD (refined, bleached, deodorized) palm olein is used as cooking oil, while RBD stearin is used in margarine, shortening, and bakery fats. Palm kernel oil's high lauric acid content makes it ideal for creamers, chocolate products, and confectionery products. The high oxidative stability, affordability, and trans-fat-free profile make palm derivatives preferred for baking, frying, and processed foods [37].

Pharmaceutical applications leverage glycerin for medicinal formulations, capsules, syrups, and topical creams. Isopropyl myristate and other palm-derived esters provide moisturizing properties in pharmaceutical preparations. The ability to meet stringent pharmaceutical standards establishes palm oleochemicals as essential healthcare ingredients[37].

Industrial applications include fatty acid methyl esters (FAME) for biodiesel production, offering renewable alternatives to fossil fuels. Palm oil's high productivity per hectare (10 times that of rapeseed or soybeans), low water, pesticide, and fertilizer requirements, and its perennial nature make it an optimal feedstock for biodiesel synthesis[38], [39]. Transesterification processes convert palm oil into methyl esters with yields exceeding 94%, providing a calorific value comparable to that of conventional diese l[40,41].

Additional derivatives include bio-based lubricants, surfactants for detergents and soaps, and specialty chemicals for the textile and cosmetic industries. The shift toward renewable, sustainable raw materials is driving the increasing adoption of palm-based oleochemicals as petrochemical alternatives.

Conclusion

Synthesis of Key Findings

This qualitative literature review of recent research (2020-2025) provides compelling evidence that palm oil and its derivatives offer significant health benefits when consumed appropriately. The bioactive constituents—particularly tocotrienols, carotenoids, phytosterols, and coenzyme Q10—confer cardiovascular protection through multiple mechanisms, including cholesterol reduction, anti-inflammatory effects, and atherosclerosis prevention. Tocotrienol-rich fractions demonstrate 7-38% reductions in LDL cholesterol through HMG-CoA reductase inhibition, while increasing HDL cholesterol and improving overall lipid profiles.

Antioxidant and anti-inflammatory properties stem from activation of Nrf2-mediated antioxidant pathways and suppression of NF-κβ inflammatory signaling. These mechanisms protect against oxidative stress across multiple organ systems, including the liver, brain, and cardiovascular tissues. Neuroprotective effects include cognitive improvement in Alzheimer's disease models, β-amyloid reduction, and maintenance of glutathione levels in neural tissue.

Dermatological applications demonstrate wound healing, anti-aging, and anti-inflammatory benefits, while palm-derived oleochemicals dominate the USD 28.51 billion personal care market. Anti-cancer properties manifest through selective inhibition of cancer cell growth, enhanced apoptosis, and immunomodulation, without harming healthy cells. Critically, red palm oil addresses vitamin A deficiency, which affects 190 million children globally, by providing highly bioavailable provitamin A to populations at the highest risk.

Resolution of Misconceptions

Scientific evidence contradicts persistent misconceptions regarding palm oil and health. Palm oil contains no cholesterol, as cholesterol is produced exclusively by animals; plants produce phytosterols, which actually reduce cholesterol absorption. The palmitic acid in palm oil differs metabolically from animal-sourced palmitic acid because it is stereospecifically located in triacylglycerols, with unsaturated fatty acids preferentially placed at the sn-2 position, enhancing digestibility.

Meta-analyses confirm that palm oil consumption does not increase cardiovascular risk compared with other cooking oils when consumed as part of a balanced diet. While palm oil increases LDL cholesterol slightly compared to highly polyunsaturated fats, it significantly increases HDL cholesterol and does not adversely affect triglycerides. The balanced saturated: unsaturated fatty acid ratio (approximately 50:50), combined with antioxidant content, mitigates potential lipid concerns.

Importantly, processing methods significantly affect metabolic outcomes. Natural, non-interesterified palm oil demonstrates neutral to beneficial metabolic effects, while chemically interesterified palm oil impairs insulin signaling and promotes insulin resistance. These findings emphasize the importance of consuming minimally processed, natural palm oil products.

Implications for Clinical Practice and Public Health

Evidence supports incorporating palm oil, particularly red palm oil, into balanced nutritional frameworks for populations at risk of vitamin A deficiency. Healthcare providers should update practice guidelines to reflect current evidence, recognizing palm oil's cardiovascular safety when consumed appropriately and its therapeutic potential through bioactive compounds.

Tocotrienol-rich fractions demonstrate clinical efficacy as adjunctive therapy in hyperlipidemia, chronic kidney disease, and potentially neurodegenerative diseases. Standardized extracts may serve targeted health applications, though optimal dosing, formulation, and duration require further investigation.

Public health initiatives in regions with a high prevalence of vitamin A deficiency should consider red palm oil fortification programs. Studies from Africa, Asia, and Latin America demonstrate effectiveness, bioavailability, and cultural acceptability. Cost-effectiveness analyses support expanded use in areas where red palm oil remains underutilized despite local palm cultivation.

Research Gaps and Future Directions

Despite substantial evidence, research gaps remain. Long-term longitudinal clinical trials examining dose-response relationships, population-specific metabolic responses, and gene-nutrient interactions would enhance precision nutrition applications. Mechanistic research elucidating synergistic effects among bioactive compounds—tocotrienols, carotenoids, phytosterols, and coenzyme Q10—may reveal greater therapeutic potential than their isolated constituents.

Emerging research areas include personalized nutrition approaches based on genetic polymorphisms affecting lipid metabolism, optimal tocotrienol dosing for specific health outcomes, and the development of bioavailability-enhanced formulations. Product development opportunities exist for standardized palm phytonutrient concentrates, targeted dermatological applications, and sustainable derivative innovations addressing both health and environmental considerations.

Effectiveness trials and cost-benefit analyses examining red palm oil interventions in vitamin A-deficient populations would inform scale-up strategies. Operational research addressing supply chain logistics, cultural acceptability, and integration with existing supplementation programs could accelerate implementation.

Final Recommendations and Conclusions

Scientific evidence accumulated over the past five years establishes that palm oil and its derivatives provide substantial health benefits through bioactive constituents operating via multiple mechanisms. When consumed as part of balanced diets, palm oil does not pose incremental cardiovascular risk and offers measurable protective effects across cardiovascular, neurological, metabolic, and immunological systems.

Stakeholder recommendations include: **Researchers** should continue mechanistic investigations, conduct long-term clinical trials, and explore personalized nutrition applications. **Healthcare providers** must update clinical guidelines to reflect an evidence-based understanding rather than outdated perceptions, and consider tocotrienol supplementation for specific patient populations. **Industry** should emphasize sustainable sourcing through certified supply chains, maintain transparent labeling distinguishing natural from interesterified products, and invest in bioavailable phytonutrient formulations. **Consumers** should make informed decisions based on scientific evidence, incorporate appropriate amounts into balanced diets, and preferentially select sustainably sourced products.

Policymakers in regions with vitamin A deficiency should prioritize red palm oil fortification programs as cost-effective public health interventions. The paradigm shift from perception-based to evidence-based understanding recognizes palm oil's nutritional value when responsibly sourced and appropriately consumed, balancing health benefits with environmental sustainability considerations. Future research integrating nutritional science, sustainable agriculture, and public health will optimize palm oil's contribution to global nutrition while addressing legitimate environmental concerns through certified sustainable production practices.

Acknowledgement

There is no acknowledgement statement.

References

[1] S. Alkandari *et al.*, "Pilot scale production of functional foods using red palm olein: Antioxidant, vitamins' stability and sensory quality during storage," *Saudi J. Biol. Sci.*, vol. 28, no. 10, pp. 5547–5554, Oct. 2021, doi: 10.1016/j.sjbs.2021.06.032.

[2] D. Sari *et al.*, "The Effects of Red Palm Oil, Koja Bay Leaves, and Passion Fruit Seeds Formulation on Antioxidant Activity, Antihyperlipidemia, BDNF, and Lipase Enzyme Activity on Sprague-Dawley Rats," *J. Exp. Pharmacol.*, vol. Volume 16, pp. 271–284, Sep. 2024, doi: 10.2147/JEP.S466494.

[3] GAPKI, "Palm Oil: Case of Cholesterol Or Phytosterol?," GAPKI Publication Recent News. Accessed: Nov. 16, 2025. [Online]. Available: https://gapki.id/en/news/2024/09/23/palm-oil-case-of-cholesterol-or-phytosterol/

[4] L. Judijanto, "Benefits of Palm Oil Consumption on Human Health: A Review," *Eur. J. Med. Heal. Res.*, vol. 3, no. 5, pp. 126–137, Sep. 2025, doi: 10.59324/ejmhr.2025.3(5).18.

[5] M. Ruscica, W. J. Loh, C. R. Sirtori, and G. F. Watts, "Phytosterols and phytostanols in context: From physiology and pathophysiology to food supplementation and clinical practice," *Pharmacol. Res.*, vol. 214, p. 107681, Apr. 2025, doi: 10.1016/j.phrs.2025.107681.

[6] S. Zio, B. Tarnagda, F. Tapsoba, C. Zongo, and A. Savadogo, "Health interest of cholesterol and phytosterols and their contribution to one health approach: Review," *Heliyon*, vol. 10, no. 21, p. e40132, Nov. 2024, doi: 10.1016/j.heliyon.2024.e40132.

[7] Y. Zhang *et al.*, "Effects of phytosterol-rich foods on lipid profile and inflammatory markers in patients with hyperlipidemia: a systematic review and meta-analysis," *Front. Pharmacol.*, vol. 16, Jul. 2025, doi: 10.3389/fphar.2025.1619922.

[8] H. Snyder, "Literature review as a research methodology: An overview and guidelines," *J. Bus. Res.*, vol. 104, pp. 333–339, Nov. 2019, doi: 10.1016/j.jbusres.2019.07.039.

[9] G. Pare and S. Kitsiou, "Methods for Literature Reviews," Handbook of e-Health Evaluation: An Evidence-based Approach. Accessed: Apr. 04, 2025. [Online]. Available: https://www.ncbi.nlm.nih.gov/books/NBK481583/

[10] A. D. Looi, U. D. Palanisamy, M. Moorthy, and A. K. Radhakrishnan, "Health Benefits of Palm Tocotrienol-Rich Fraction: A Systematic Review of Randomized Controlled Trials," *Nutr. Rev.*, vol. 83, no. 2, pp. 307–328, 2024, doi: 10.1093/nutrit/nuae061.

[11] W. L. Hall *et al.*, "Postprandial lipid and vascular responses following consumption of a commercially-relevant interesterified palmitic acid-rich spread in comparison to functionally-equivalent non-interesterified spread and spreadable butter: a randomised controlled trial i," *Food Funct.*, vol. 15, no. 5, pp. 2733–2750, 2024, doi: 10.1039/D3FO05324E.

[12] Z. Zainal, H. Khaza'ai, A. Kutty Radhakrishnan, and S. K. Chang, "Therapeutic potential of palm oil vitamin E-derived tocotrienols in inflammation and chronic diseases: Evidence from preclinical and clinical studies," *Food Res. Int.*, vol. 156, p. 111175, 2022, doi: https://doi.org/10.1016/j.foodres.2022.111175.

[13] C. Y. May, N. M. Han, M. A. Ngan, and Y. Basiron, "Production of Coenzyme Q10 from Crude Palm Oil," *MPOB Inf. Ser.*, vol. MPOB TT No, no. 239, pp. 1–2, 2024.

[14] M. Dewi, D. Martianto, N. Andarwulan, R. Kazimierczak, and D. Średnicka-Tober, "Plant Sterol-Enriched Palm Oil Intervention to Improve Lipid Profile and Inflammation Status in Hyperlipidemic Individuals," *Nutrients*, vol. 16, no. 19, p. 3370, Oct. 2024, doi: 10.3390/nu16193370.

[15] B. Mishra, Z. Badade, J. Lingidi, S. Jaiswal, and B. Anand, "Tocotrienols and its impact in cardiovascular health," *Int. J. Clin. Trials*, vol. 1, no. 3, p. 87, 2014, doi: 10.5455/2349-3259.ijct20141102.

[16] L. Trugilho *et al.*, "Effects of Tocotrienol on Cardiovascular Risk Markers in Patients With Chronic Kidney Disease: A Randomized

Controlled Trial," *J. Nutr. Metab.*, vol. 2025, no. 1, Jan. 2025, doi: 10.1155/jnme/8482883.
[17] Y. Sun, N. Neelakantan, Y. Wu, and R. M. van Dam, "Abstract P065: Effects of Palm Oil Consumption on Blood Lipids: A Meta-Analysis of Clinical Trials," *Circulation*, vol. 129, no. suppl_1, Mar. 2014, doi: 10.1161/circ.129.suppl_1.p065.
[18] S. Abd Muid *et al.*, "Effects of Palm Oil Derived Tocotrienol Rich Fraction and Vitamin E Isomers on Biomarkers of Early Atherogenesis in Stimulated Human Umblilical Vein Endothelial Cells," *Malaysian Appl. Biol.*, vol. 51, no. 4, pp. 145–152, Oct. 2022, doi: 10.55230/mabjournal.v51i4.23.
[19] O. R. Ajuwon, J. L. Marnewick, O. O. Oguntibeju, and L. M. Davids, "Red Palm Oil Ameliorates Oxidative Challenge and Inflammatory Responses Associated with Lipopolysaccharide-Induced Hepatic Injury by Modulating NF-κβ and Nrf2/GCL/HO-1 Signaling Pathways in Rats," *Antioxidants*, vol. 11, no. 8, p. 1629, Aug. 2022, doi: 10.3390/antiox11081629.
[20] N. Prangdimurti, E. Zakaria, and F. Rungkat, "Crude Palm Oil Consumption Was Accepted and Improved Lipid Profile and Reduced C-Reactive Protein Content in Plasma of Childbearing Age Women In Sub District Dramaga Bogor," *IPB Univ. Sci. Repos.*, 2020, [Online]. Available: https://repository.ipb.ac.id/handle/123456789/58327
[21] M. T. Golmakani, A. Soltani, and S. Sahraeian, "Investigating the impact of palm oil refining on the oxidative stability of its blend with linseed oil: a kinetic approach," *Grasas y Aceites*, vol. 75, no. 4, p. 2212, Dec. 2024, doi: 10.3989/gya.1200232.2212.
[22] M. E. Mulyono, I. F. Rizki, F. R. Panjaitan, and B. D. Bajra, "Storage stability of hydrolyzed palm kernel oil and red palm super olein blend and its softgel capsule," *E-Journal Menara Perkeb.*, vol. 93, no. 1, pp. 1–11, Apr. 2025, doi: 10.22302/iribb.jur.mp.v93i1.607.
[23] A. R. Siregar, M. Ichwan, J. Siregar, D. Anggraini, and H. Hasibuan, "Effective ness of red palm oil supplementation on motor impairment in a paraquat-induced parkinson's disease rat mode," *Int. J. Trop. Vet. Biomed. Res.*, vol. 9, no. 1, pp. 11–20, 2024.
[24] A. do E. S. Macedo *et al.*, "Neuroprotective effects of essential oils in animal models of Alzheimer's and Parkinson's disease: a systematic review," *PeerJ*, vol. 13, p. e19643, Jul. 2025, doi: 10.7717/peerj.19643.
[25] GAPKI, "Tracing True Relations Between Palm Oil and Diabetes," GAPKI Publication Recent News. Accessed: Nov. 18, 2025. [Online]. Available: https://gapki.id/en/news/2024/09/23/tracing-true-relations-between-palm-oil-and-diabetes/
[26] D. A. Berkoh, E. T. Donkoh, A. Asamoah, R. A. Sakibu, and R. A. Ngala, "Effect of Tropical Vegetable Oil Consumption on Lipid Profile and Glycaemic Control in Type 2 Diabetics," *Trends Med. Res.*, vol. 19, no. 1, pp. 247–284, Jul. 2024, doi: 10.3923/tmr.2024.274.284.
[27] B. C. Martins *et al.*, "Consumption of interesterified palm oil leads inflammation of white adipose tissue and triggers metabolic disturbances in mice on a high-fat diet," *Sci. Rep.*, vol. 14, no. 1, p. 12530, May 2024, doi: 10.1038/s41598-024-63488-9.
[28] New Straits Times, "Sustainable palm oil provides exceptional benefits for your skin and body," New Straits Times. Accessed: Nov. 18, 2025. [Online]. Available: https://www.astra-agro.co.id/2024/06/06/sustainable-palm-oil-provides-exceptional-benefits-for-your-skin-and-body/
[29] Asian Agri, "Is Palm Oil Good for Skin & Hair?," Asian Agri. Accessed: Nov. 18, 2025. [Online]. Available: https://www.asianagri.com/en/media-publications/faqs/is-palm-oil-good-for-skin-hair/
[30] A. Rada–Bula *et al.*, "Compositional characteristics and shelf life of interspecific hybrid palm oil (E. oleifera×E. guineensis)," *Food Packag. Shelf Life*, vol. 42, p. 101240, Mar. 2024, doi: 10.1016/j.fpsl.2024.101240.
[31] GAPKI, "Is Palm Oil Beneficial or Detrimental for Human Skin?," GAPKI Publication Recent News. Accessed: Nov. 18, 2025. [Online]. Available: https://gapki.id/en/news/2024/11/27/is-palm-oil-beneficial-or-detrimental-for-human-skin/
[32] H. M. L. Lubis, E. Purwoningsih, A. A. Nasution, and Q. M. Salim, "Mechanism of Action of Tumorigenesis of Anticancer Molecules of Palm Oil Tocotrienols (Elaeis Guieensis Jacq.): A Systematic Review," *Eduvest - J. Univers. Stud.*, vol. 2, no. 2, pp. 431–440, Feb. 2022, doi: 10.59188/eduvest.v2i2.379.
[33] M. Younes, G. Loubnane, C. Sleiman, and S. Rizk, "Tocotrienol isoforms: The molecular mechanisms underlying their effects in cancer therapy and their implementation in clinical trials," *J. Integr. Med.*, vol. 22, no. 1, pp. 1–11, 2024, doi: https://doi.org/10.1016/j.joim.2024.01.002.
[34] N. R. Morgan, S. Bhuvanendran, P. Krishnappa, and A. K. Radhakrishnan, "Daily Supplementation of High Doses of Tocotrienol-Rich Fraction From Palm Oil Produced No Toxic Effects in Healthy Mice," *J. Toxicol.*, vol. 2025, no. 1, Jan. 2025, doi: 10.1155/jt/9464952.
[35] R. Ranasinghe, M. Mathai, and A. Zulli, "Revisiting the therapeutic potential of tocotrienol," *BioFactors*, vol. 48, no. 4, pp. 813–856, Jul. 2022, doi: 10.1002/biof.1873.
[36] A. L. Rice and J. B. Burns, "Moving from Efficacy to Effectiveness: Red Palm Oil's Role in Preventing Vitamin A Deficiency," *J. Am. Coll. Nutr.*, vol. 29, no. sup3, pp. 302S-313S, Jun. 2010, doi: 10.1080/07315724.2010.10719845.
[37] TradeAsia, "Palm Oil-Based Oleochemicals: A Rising Force in Global Manufacturing Supply Chains," TradeAsia Article. Accessed: Nov. 18, 2025. [Online]. Available: https://www.oleochemicalsasia.com/market-insights/palm-oil-based-oleochemicals-global-supply-chains-2025
[38] E. D. Daryono and D. A. Anggorowati, "Identification of FAME (Fatty Acid Methyl Esters) Products from the Interesterification Reaction of Palm Oil with Methyl Ester Co-solvent," *G-Tech J. Teknol. Terap.*, vol. 9, no. 4, pp. 2175–2186, Oct. 2025, doi: 10.70609/g-tech.v9i4.8243.
[39] S. Peranginangin, H. B. Sembiring, Y. V. M. Nainggolan, N. S. Sihotang, and H. Wijoyo, "Fatty acid methyl ester (FAME) production from sludge solid waste as K2O/SiO2 catalyst," *Case Stud. Chem. Environ. Eng.*, vol. 10, p. 100933, Dec. 2024, doi: 10.1016/j.cscee.2024.100933.
[40] B. Y. Lamba, S. Jain, and S. Jha, "Biodiesel from Palm Vegetable Oil," *Nat. Resour.*, vol. 15, no. 02, pp. 51–60, 2024, doi: 10.4236/nr.2024.152004.
[41] Y. M. Oo, P. Juera-Ong, and K. Somnuk, "Methyl ester production process from palm fatty acid distillate using hydrodynamic cavitation reactors in series with solid acid catalyst," *Sci. Rep.*, vol. 14, no. 1, p. 27732, Nov. 2024, doi: 10.1038/s41598-024-78974-3.