

The attenuating effect of the aqueous extract of the Flower of *Opuntia ficus indica* against lead-induced reprotoxicité in the wistar rat

Naouel Boudjema (✉ naouelboudj23@gmail.com)

Université Badji Mokhtar Annaba

Zyneb Ladouali

Université Badji Mokhtar Annaba

Research Article

Keywords: *Opuntia ficus indica*, CASA sperm, lead, DNA, testosterone

Posted Date: February 8th, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-3925409/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

The objective of this study is to investigate the beneficial effect of *Opuntia ficus indica* flower aqueous extract against lead (Pb)-induced reprotoxicité in male *Wistar* rats administered orally for 4 weeks. These are divided into 6 groups: the control group (C), 2 positive controls treated with the aqueous extract of the flower of *Opuntia ficus indica* F1 and F2 respectively (250 mg/kg, 500 mg/kg of body weight/day) ;1 group treated with lead (50 mg/kg bodyweight/day) and 2 groups of combined treatment treated with lead and aqueous extract of *Opuntia ficus indica* flowers (F1 + Pb), (F2 + Pb). Extracts of *Opuntia ficus indica* flowers are the subject of a phytochemical study. After 30 days of oral administration, the absolute weight of the testicles and of the epididymis, the plasma levels of testosterone are evaluated; in addition to sperm biology parameters (concentration, motility, vitality, velocity (VCL, VSL and VAP), cross-level (BCF) and DNA fragmentation test are evaluated by the CASA system. Testicular glutathione and epididymal (GSH), Malondialdehyde (MDA) and glutathione peroxidase (GPX) are also evaluated. The results obtained show a significant reduction in testosterone levels, absolute testicular and epididymal weights, live sperm, VCL, VSL, VAP, BCF, GSH and GPX as well as an increase in dead spermatozoa and MDA in the group treated with lead (Pb) compared to the control group (C) and the positive controls, the administration of the aqueous extract of the flowers of *Opuntia ficus indica* in the combination groups (F1 + Pb and F2 + Pb) caused an attenuating effect against lead-induced toxicity. Even the phytochemical study confirms that the methanolic extract of *Opuntia ficus indica* extract is rich in tannins and flavonoids.

INTRODUCTION

Environmental pollution by heavy metals is a global problem that reduces the quality of life and has negative consequences on living organisms (James Daniel et al., 2018). These persistent environmental pollutants have the ability to accumulate in the tissues of living organisms and cause more or less significant damage (INRS, 2018). Among these pollutants, lead is a very abundant compound with a fairly wide distribution in all types of ecosystems, with a bioconcentrating potential in the different body systems (Drif et al., 2019). Lead is not biodegradable (Seregin and Ivanov, 2001), and therefore can induce a wide range of central, peripheral, and hematopoietic nervous system dysfunctions (Burroughs and Rollins, 2017; Derouiche et al., 2018). Lead enters the body mainly by ingestion or inhalation, it can also pass through the skin (Anomaa et al., 2017). Once absorbed, lead passes into the bloodstream (Andjelkovic et al., 2019). And is then distributed to several tissues such as bones, brain, kidneys and liver. It is also found in the male reproductive system (epididymis, seminal vesicles, testicles and prostate.) (Picot, 2010; Andjelkovic et al., 2019), leading to disruption of male reproduction (Alouche et al; 2009). This negatively affects the number of sperm, their movements and delays their activity (Berredjem et al., 2014). The mechanism of lead-induced oxidative stress involves a balance between the generation and elimination of ROS (reactive oxygen species) in tissues and cellular components causing damage to the membrane, DNA and proteins (Patra et al., 2011; Berrahal et al. 2011; Flora et al. 2012; Abdel-Moneim et al. 2015; Wu et al. 2016; BaSalamah et al. 2018). Lead significantly reduces the antioxidant system). To remedy the poisoning caused by heavy metals and lead in particular, many studies have focused on the therapeutic virtues of the flowers of the prickly pear (*Opuntia ficus indica*), which is a plant native to Mexico, but cultivated in different countries of the world (Toure, 2017). This plant is characterized by rapid growth and great adaptation to difficult abiotic conditions (Pérez-Torrero et al. 2017). The flowers of *Opuntia ficus indica* are known for their antioxidant properties, helping to protect cells against oxidative damage caused by radicals (Toure, 2017) They are also used as an antihemorrhagic, antibacterial, antifungal, antiseptic and diuretic (Ammar et al., 2012; Aragona et al., 2017; Keller et al., 2019). Among the indications of *Opuntia ficus indica* flowers relieve kidney pain and effectively fight against cholesterol, type II

diabetes, prostate adenoma, burns, peptic ulcers and even certain types of cancers whose progression it would slow down (ref). Analysis of the physicochemical properties of broths and infusions prepared from *Opuntia ficus-indica* flowers reveals that these preparations are rich in minerals (potassium and calcium), polyphenols, flavonoids and tannins (Chahdoura et al., 2014; Ammarel et al., 2015). The main active compounds in *Opuntia ficus indica* flowers are polysaccharides which accelerate the repair process throughout the tissue. Glycoproteins, aromatic compounds, in particular betacyanins, betaxanthins, polyphenols, fatty acids, sterols, β -carotene, due to their antioxidant activity and polyphenolic flavonoids are metabolites which have numerous pharmacological properties (Panico et al., 2007). Other studies have shown the beneficial effects of *Opuntia ficus indica* flowers on sperm DNA fragmentation after sperm cryopreservation in humans (Meamar et al., 2012). Other researchers have also observed a significant reduction in sperm DNA fragmentation as well as improved sperm quality, motility and viability (Allai et al., 2016).

MATERIALS AND METHODS

Ethical approval

The Ethical Committee of Animal Sciences at the University of Badji Mokhtar-Annaba has given the authorization to realize the Ph.D. research program.

Plant material

The flowers of *Opuntia. Ficus indica* were collected in May 2019, in the region of Ben Azouz, an area located to the North East of Algeria about a hundred kilometers from the city of Sikkda. The dried flowers were then ground into powder and filtered through a fine mesh fabric and kept in paper bags away from light to prevent the photo-oxidation of the active substances contained in the powder. Distilled water was added to the powder to obtain a homogeneous suspension at 250 mg and 500 mg/kg body weight) according to the protocol of (Alimi et al., 2011).

The dosage of phenolic compounds

Preparation of the methanolic extract

The preparation of this extract consists in macerating 1g of vegetable powder in 20ml of absolute methanol for 24 hours at room temperature, the extracts are then filtered and then evaporated under reduced pressure to dryness by a rotary evaporator at $T = 65^{\circ}\text{C}$. The dry residue is taken up in a few milliliters of methanol and stored in an amber flask at $+4^{\circ}\text{C}$.

Determination of phenolic compounds:

The determination of the total phenolic compounds was carried out by the method using the Folin-Ciocalteu reagent. This method was described in 1965 by Singleton and Rossi.

This assay is based on the quantification of the total concentration of hydroxyl groups present in the extract. A volume of 200 μl of the stock solution of gallic acid and of the methanolic extract of the species studied is introduced into test tubes, 1 ml of the Folin-Ciocalteu reagent diluted 10 times is added there to. After standing for 5 min, 800 μl of 7.5% sodium carbonate are introduced. The tubes are shaken and stored for 30 minutes at room temperature and protected from light. Absorbance is measured at 765 nm against a blank using a SHIMADZU UV-1202 spectrophotometer. A calibration curve is performed in parallel under the same operating conditions using gallic

acid as a positive control. The results are expressed in milligram (mg) equivalent of gallic acid per gram of dry plant matter (mg EAG / g of dry matter).

Dosage of tannins:

The total tannin content was evaluated according to the colorimetric method described by **Polshettiware et al. (2007)**, based on the reduction, in an alkaline medium, of phosphomolybdic and tungstic acids contained in the Folin-Denis reagent (RFD). The reaction gives a blue color, the intensity of which is measured by spectrophotometry at 755 nm.

0.5 ml of RFD and 1 ml of sodium carbonate (0.5% Na₂CO₃) were introduced into volumetric flasks. 100 µl of the dilution of the methanolic extract was added. There to, the volume of the solution was adjusted to 5 mL with distilled water. After a 30 min incubation, the absorbance was measured by a SHIMADZU UV-1202 spectrophotometer. The total tannin content was determined by extrapolation on a calibration curve obtained from a series of dilutions of tannic acid (MS 10 mg / ml). The results were expressed in milligrams of tannic acid equivalent per gram of dry plant material (mg EAT / g DM).

Determination of flavonoids:

Principle: The quantification of the flavonoids was carried out by a method adapted by **Zhishen et al. (1999)**, based on the formation of complexes between flavonoids and aluminum trichloride. The complexes produced, yellow in color, absorb in the visible range at 510 nm.

500 µl of the crude extract is mixed with 2 ml of distilled water, and added with 150 µl of 5% sodium nitrite (NaNO₂). After 5 min of incubation, 100 µl of 10% aluminum trichloride is added to the mixture. After a further 6 min incubation, add 1 ml of 1M sodium carbonate. The mixture is completely stirred in order to homogenize the content and the absorbance of the solution is determined at 510 nm against a blank. A calibration curve is carried out in parallel under the same operating conditions using quercetin as positive control. The total flavonoid content is expressed in milligram (mg) equivalent of quercetin per gram of dry plant material (mg EQ / g DM).

Evaluation of the anti-free radical activity by the DPPH test:

The anti-free radical activity of the extract of *Inula viscosa* L. was evaluated in vitro, by the DPPH test. This spectrophotometric method uses the violet-colored DPPH (2,2'-diphenyl-1-picrylhydrazyl) radical as a reagent, which turns yellow, in the presence of free radical scavengers, and is reduced to 2,2'-diphenyl-1-picrylhydrazine (**Cuendet et al., 1997; Buritsand Bucar, 2000**). This makes it possible to follow the discoloration kinetics at 517 nm.

Evaluation of anti-inflammatory activity:

The anti-inflammatory activity of methanolic extract from the aerial part of *Inula viscosa* L. is evaluated by the protein denaturation method described by **Alhakmani et al. (2013)**, with some modifications. By this method, acetylsalicylic acid (aspirin) is an anti-inflammatory drug used as a benchmark anti-inflammatory.

The test consists of two experiments. In the first experiment, the reaction mixture consists of 2 ml of extract of *Opuntia Ficus* Flower. (methanolic) at different concentrations (2-1.6-1.2 mg / ml) and 2.8 ml of distilled water adjusted to pH = 6.4 with HCl, then mix with 2 ml of egg albumin. Incubate at 37 ° C for 10 minutes. In the second experiment, the extract of *Opuntia ficus* Flower. is replaced by acetylsalicylic acid for each of the experiments, a control is prepared. It consists of 2 ml of distilled water in place of the inula extract and acetylsalicylic acid, 2.8 ml of

distilled water and 2ml of egg albumin. Denaturation is induced by placing the reaction mixture in a water bath heated to 72 ° C for 5 minutes. After cooling, the absorbance is measured at 660nm using distilled water as a white.

Protocol HPLC

HPLC (high performance liquid chromatography) is an analytical separation technique. first, we prepared the methanolic extract as we saw in the study of the components of *opuntia ficus indica* flowers, then we put 5 mg of *opuntia ficus indica* flower extract in 100 ml of distilled/directed water and 900 µl of methanol (MeOH). Chromatographic analysis was performed using Shimadzu Prominence-I LC-2030C (Japan) HPLC liquid chromatography with UV detector, C18 (4.6 x 25 mm, 5 µm) chromatography column (USA). column is maintained at 45°C.

Chemical Product

In this study, we followed the protocol of (Hussein AL-Azawi, 2015). Lead acetate trihydrate (Géorgie, USA) was dissolved daily in distilled water at a rate of 50mg/kg body weight.

Animals

Fifty-five (55) rats from the Pasteur Institute in Algiers. each weighing 196±8 g. were raised in the pet store of the Badji Mokhtar-Annaba University. They were subjected 15 days period of adaptation to the conditions of the pet store (T°22C°+2C°, 40% humidity and a clear and dark photoperiod of 12 hours/12 hours). Fifty-five rats were divided into 06 equal groups.

Group 1 (C), which is the control group, each rat receives 1 ml of tap water by gavage, Group 2 (F1), which is the positive control group, each rat receives 1.5 ml of the extract aqueous extract of *opuntia ficus indica* flowers by gavage at 250 mg/kg body weight, group 3 (F2), which is the positive control group, each rat received 1.5 ml of the aqueous extract of *opuntia ficus indica* flowers by gavage at a dose of 500 mg/kg of body weight, group 4 (Pb) which is the group treated with lead, each rat received 1 ml of Pb acetate solution (50 mg/Kg of body weight) by gavage. , Group 5 (F1+Pb), each rat receives the combination of Pb acetate (50mg/kg/body weight) and the aqueous extract of *Opuntia ficus indica* flowers (250mg/kg bw by gavage. Group 6 (F2+Pb), each rat receives the combination of Pb acetate (50mg/kg/body weight) and the aqueous extract of *Opuntia ficus indica* flowers (5mg/kg bw) by gavage. were treated for four weeks. The animals were weighed once a week throughout the experimental period. At the end of the treatment period, the rats were sacrificed by decapitation. The organs (testis and epididymis) were removed, weighed and a part was frozen at -20°C for the assay of markers linked to oxidative stress and the other part was fixed in formalin in order to make the histological sections.

Studied Parameters

At the time of the sacrifice, the blood was collected in heparinized tubes and then centrifuged at 3000 rpm for 10min. The plasma obtained was stored at -20 °C along with the which had already been weighed, till further analysis. Animals' treatments were authorized by the Ethical Committee of Animal Sciences at the University of Badji Mokhtar-Annaba, before starting the experimental work Semen analysis.

The sperm analysis is carried out immediately after the sacrifice, by taking a sample of the epididymis, estimated at 1 µl, which is diluted with a physiological solution of NaCl at 0.09%, then 5 ml of the mixture is deposited in an empty chamber slide (Gold Cyto model) The results are read and analyzed under an electron microscope on a Nikon

Eclipse microscope (Nikon E200-LED) with phase objective (x4), Connected to a computer (CASA) at the using a group analysis of spermatozoa (SCA®, Microptic, Barcelona, Spain), automatic sperm analysis allowed the calculation of concentration, velocities, vitality, velocity parameters (VCL, VSL, VAP), and cross-beat frequency (BCF), were automatically calculated. For the DNA fragmentation test, a diluent semen sample at a concentration of 5 to 10 million per millilitre was used (n = 4). The test was performed using the Gold Cyto DNA kit (Gold Cyto Biotechcrop. Room 1602, 6F, NO, 16) as illustrated in the instruction manual. The spermatozoa were recognized and automatically classified into fragmented spermatozoa. According to established analysis criteria.

Hypo - osmotic swelling (HOS) test

The study of sperm vitality is carried out by carrying out the Bright Vit Protocol. This test was developed to measure two functional aspects of sperm, namely living sperm in a simple way and secondarily the quality of cell membrane integrity and the ability of the cell membrane to resist osmotic swelling without bursting. Bright vit, a neosine négrosine solution stains dead cells pink, living cells will remain white. This analysis is done by exposing a drop of sperm, extracted from the tail of the epididymis, to a hypo osmotic solution consisting of a négrosine neosine.

Analysis of Oxidative Stress Markers

100 mg of testis samples and epididymis were taken and placed in test tubes for measurement. For the determination of glutathione (GSH) using the method of Cory and Weckbecker. The principle of this assay is based on the measurement of the optical absorbance of 2-nitro-5-mercaptopuric acid. 200 mg of each tissue was placed in the presence of 8 ml of a solution of ethylene diamine tetra acetic acid (EDTA) at 0.02 M, then cold ground using an ultra-Sound mill (4 °C) to obtain a homogenate. Read optical densities at 412 nm against white. The testicular and epididyme total proteins were quantified according to the colorimetric method of Bradford. by using CoomassieBlueG-250 (Sigma, St. Louis, USA) Malondialdehyde (MDA) was estimated by using the method of **Ohkawa et al., 1979**. The determination is based on the formation in an acidic and hot medium (100 ° C) between MDA and thiobarbituric acid (TBA) of an absorbent-colored pigment at 530 nm, extractable by organic solvents such as butanol. the enzymatic activity of glutathione peroxidase (Gpx) was measured by the method of **Flohé and Günzler (1984)**. This method is based on the reduction of hydrogen peroxide (H₂O₂) in the presence of reduced glutathione (GSH), the latter is transformed into (GSSG) under the influence of GPX. read optical densities at 412 nm against white

Histological examination of samples

Tissue samples are immediately fixed in buffered formaldehyde (10%), the samples are then dehydrated with a series of ethanol, cleaned with xylene, embedded in paraffin, cut using a 3 µm microtome. The tissue sections are mounted on gelatinized slides, then stained with eosin hematoxylin, then examined under a light microscope (Optika B-350). A second toluidine blue stain is used (Appendix I). The microscopic photos are taken with a digital camera (Optikam B5) according to the recommendations (**Aeffner et al., 2018**). Tissue samples are immediately fixed in buffered formaldehyde (10%), the samples are then dehydrated with a series of ethanol, cleaned with xylene, included in paraffin, cut using a microtome at 3 µm. The fabric sections are mounted on gelatinized blades and then stained with hematoxylin eosin then examined under an optical microscope (). A second color of toluidine blue is used (Appendix I). Microscopic photos are taken with a digital camera (Optikam B5) according to the recommendations (**Aeffner et al., 2018**).

Statistical analysis

Statistical analysis of all results was analyzed using the Microsoft Excel program (2013). Data analysis was performed by student's test by comparing the means of each parameter analyzed in pairs between groups, using the software prism7.

RESULTS

Quantitative evaluation of the phenolic compounds of *Opuntia ficus indica* flowers (OFI)

The quantitative analysis of the phenolic compounds of the methanolic extract of the flowers of *Opuntia ficus indica* affirms the richness of this species in Total polyphenols, the Tannins as well as the flavonoids.

Table 1: the quantitative analysis of the phenolic compounds of the methanolic extract of the flowers of *Opuntia ficus indica* affirms the richness of this species in Total polyphenols, the Tannins as well as the flavonoids

Phenolic compounds	Total polyphénols (mg EAG/g DM)	Flavonoïdes (mg EAG/g DM)	Tannins (mg EAG/g DM)
Teneur (OMFE)	970,4503	16,85013	883,835

EAG/g DM): Milligram equivalent of gallic acid per gram of dry matter

Evaluation of anti-free radical activity by the DPPH test (Diphenyl-picryl-hydrazyl)

The results obtained show that the methanolic extract of OFI flowers has an exceptional antioxidant effect against the DPPH radical with an IC50 of the order of 0.843mg/ml compared to: Inhibitory concentration of 50% of the ascorbic acid IC50 = 0.290mg/ml.

Table 2: the quantitative analysis of the phenolic compounds of the methanolic extract of the flowers of *Opuntia ficus indica* affirms the richness of this species in Total polyphenols, the Tannins as well as the flavonoids

50% inhibitory concentration	IC50 (mg/ml)
Ascorbic Acid	0,290 mg / ml
Flower extract concentration	0,843mg /ml

Evaluation of anti-inflammatory activity

The results obtained show that the methanolic extract of OFI Flowers has an anti-inflammatory activity in vitro. The addition of increasing concentration of the methanolic extract of the OFI flowers of the plant to the albumin solution prevents the precipitation of this protein.

Table 3: the quantitative analysis of the phenolic compounds of the methanolic extract of the flowers of *Opuntia ficus indica* affirms the richness of this species in Total polyphenols, the Tannins as well as the flavonoids

Concentration of flower extract mg/ml	% inhibition methanolic extract
100	13,39
200	16.12
300	32.79
500	55.74
750	84.43

Qualitative analysis of flavonoids

The analytical separation technique allows the identification of the phenolic compounds of the aqueous extract of the flowers of *Opuntia ficus indica*. Chromatographic profiles of aqueous extracts of prickly fig flowers analysed by HPLC show several spikes which appear to contain several types of phenolic compounds. The comparison between the reference molecules (standards) and the phenolic compounds of the aqueous extract of the flowers of *Opuntia ficus indica* is based on retention time (TR), that is, the height of the spike. So, we identified phenolic compounds such as flavonoids, and Iso quercetin, the other phenolic compounds are not identified because of a lack of standards. According to chromatographic profiles the aqueous extracts of *Opuntia ficus indica* flowers are rich in phenolic compound.

Chromatogram

The analytical separation technique allows the identification of the phenolic compounds of the aqueous extract of the flowers of *Opuntia ficus indica*. Chromatographic profiles of aqueous extracts of prickly fig flowers analysed by HPLC show several spikes which appear to contain several types of phenolic compounds. The comparison between the reference molecules (standards) and the phenolic compounds of the aqueous extract of the flowers of *Opuntia ficus indica* is based on retention time (TR), that is, the height of the spike. So, we identified phenolic compounds such as flavonoids, and Iso quercetin, the other phenolic compounds are not identified because of a lack of standards. According to chromatographic profiles the aqueous extracts of *Opuntia ficus indica* flowers are rich in phenolic compound.

General Assessment

General assessment During the experimental period, the rats of the control group (C) and the positive groups (F1, F2) were in very good condition unlike the group treated with lead (Pb) which presented a set of changes in the behavior of the animals such as rigor, aggressiveness, and exhaustion, as well as a decrease in body gain and the appearance of lymph nodes in some rats.

Organs and total body Weights

the treatment of rats with lead at 50 mg/kg of body weight caused a non-significant decrease in the body weight of the rats compared to the control group (C) and the positive control groups (F1, F2). On the other hand, a non-significant increase in the weight of the rats in the group treated with the Pb association and the aqueous extract of *OFI* was observed compared to the Pb group, observed and administration of the aqueous extract of *OFI* flowers at 250 mg/kg and 500 mg/kg by gavage did not cause any significance in body mass in the different groups of rats.

The treatment of rats with lead (Pb) also induces a significant reduction in the absolute weight of the testes and epididymides compared to the control group (C) and the positive control groups (F1, F2). Co-administration of the aqueous extract of *OFI* flowers with lead acetate caused a significant increase in the absolute weight of the testicles and epididymides in the combination groups, especially (F2+Pb) compared to the (Pb) group.

Table 4: The total body weight (g) and the absolute weight (g) of reproductive organs (mean±SEM) of Wistar rats exposed to lead acetate and co-administrated with the aqueous extract of *OFI* for 4 consecutive weeks.

Groups	Body weight	Testis	Epididymis
C	246,6 ± 9,495	1,514 ± 0,035	0,459 ± 0,013
F1	250,9 ± 10,73	1,524 ± 0,033	0,504 ± 0,023
F2	253,4 ± 11,24	1,564 ± 0,025	0,565 ± 0,020 ^{a: **}
Pb	229,8 ± 3,693	1,32 ± 0,030 ^{a: **, bc: ***}	0,329 ± 0,009 ^{abc: ***}
F1+Pb	235,1 ± 5,238	1,501 ± 0,029 ^{d: **}	0,499 ± 0,014 ^{d: ***}
F2+Pb	241,7 ± 7,655	1,557 ± 0,030 ^{d: ***}	0,526 ± 0,020 ^{d: ***}

Spermatozoa Concentration and Motility

The results indicate a significant decrease ($p < 0.05$) in the concentration and motility of spermatozoa in the group (Pb) compared to the control group (C) and the positive control groups (F1, F2). However, administration of water extract from *Opuntia ficus indica* flowers in all groups combined was significantly higher than in lead-poisoned rats.

Sperm Velocity

The results show that the fast, medium and slow sperm and velocity parameters. The sperm velocity and velocity parameters (VCL, VSL, VAP and BCF) of the Pb group sperm were significantly lower than those of the control groups. The velocities and VCL, VSL, VAP and BCF of the sperm of the combined groups were significantly higher than those of the (Pb) group especially of the combined (F2 + Pb) group and were not statistically different from the control group (Figure).

Sperm vitality

The results indicate that the percentage of dead sperm in the (Pb) group was significantly higher compared to living sperm in the control group (C) and the positive control groups (F1, F2). However, the results reveal that the treatment of rats with the aqueous extract of the flowers of *Opuntia ficus indica* in the combined groups (F1+Pb), (F2+Pb) succeeded in preventing lead-induced alterations, which we observed a significant increase in the percentage of living spermatozoa and the percentage of dead spermatozoa was remarkably low, especially in the group (F2+Pb).

Testosterone levels

The results are shown in the figure (). indicate a significant decrease in the level of testosterone in the group (Pb) compared to the control group (T) and the positive control groups (F1, F2). On the other hand the supplementation of the aqueous extract of the flowers (*OFI*) in the groups of the combination (F1+Pb), (F2+Pb) improved the results, especially in the group (F2+Pb).

DNA Fragmentation

The results show a significant increase in DNA fragmentation in group (Pb) by contribution to control group (C) and groups (F1, F2). at the same time, a significant decrease in DNA fragmentation in the combined groups (F1+Pb) and especially in the group (F2+Pb). while F1 had levels close to those of the control group. while F2 was better than group (C) and positive control groups

Malondialdehyde Level

The results obtained in our experiment show that gavage of lead acetate for 4 weeks caused a significant increase in the concentration of MDA, in the testes and epididymides compared to the control group (C) and the positive control groups (F1, F2).. However, treatment of rats with *OFI* flower suspension in the combined (F1+Pb) and (F2+Pb) groups successfully prevented the deleterious effect of lead acetate. Exhibited a significant decrease in MDA concentration compared with (pb)-treated group.

Glutathione Concentration

The enzymatic activity of GSH in the testes and epididymis of rats was significantly reduced in the lead-treated group compared to the control group (C) and the positive control groups (F1, F2). However, the activity of GSH in the combined treatments (F1+Pb, F2+Pb) was significantly higher than in the (Pb) group.

Glutathione Peroxidase Activity

We observed that testicular and epididymal glutathione peroxidase activity was significantly reduced in rats exposed to lead acetate compared to control group (C) and positive control groups (F1, F2). However, the treatment of rats with *Opuntia ficus indica* flower suspension in the combined groups (F1+Pb) and (F2+Pb) successfully prevented the toxic effect of lead acetate. Showed a significant increase, especially in the (F2 + Pb) groups.

Table 5: Testicular and epididymal GSH concentration, GPx activity, and MDA level (mean±SEM) of Wistar rats exposed to lead acetate and coadministered with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.

Groups	umol GSH/mg proteins	GSH	Umol Gpx/mg proteins	GPX	nmol/100 mg tissues	MDA
C Testis	49,06 ± 1,592		41,43 ± 0,7		5,386 ± 0,223	
Epididymis	53,06 ± 0,951		7435,2 ± 0,451		4,977 ± 0,015	
F1 Testis	53,2 ± 0,730 ^{a : *}		43,88 ± 0,731		5,177 ± 0,181	
Epididymis	56,35 ± 0,275 ^{a : *}		35,9 ± 0,511		4,898 ± 0,023	
F2 Testis	59,93±0,456 ^{a :***, b :*}		41,43 ± 0,774 ^{a :**}		4,926 ± 0,041	
Epididymis	57,95 ± 0,463 ^{a : ***}		42,28 ± 0,746 ^{ab :**}		4,833 ± 0,019	
Pb Testis	33,2 ± 1,691 ^{abc :***}	34,32 ± 0,554 ^{abc}	43,88 ± 0,731 ^{abc :***}	17,53 ± 0,807 ^{abc :***}	10,78 ± 0,063 ^{abc :***}	
Epididymis					10,28 ± 0,032 ^{abc :***}	
F1+Pb Testis	48,31±2,329 ^{b :**, cd :***}	52,17 ± 1,038 ^{b :**, cd :***}	38,89±0,874 ^{b :**, cd :***}	29,9 ± 0,809 ^{abd :***}	6,162 ± 0,069 ^{a :*, b :**, cd :***}	
Epididymis					5,634 ± 0,23 ^{ab :**, cd :***}	
F2+Pb Testis	55,32 ± 1,503 ^{a :***, c :*, d :***}		41,55 ± 0,493 ^{c :*, d :***}		5,442 ± 0,209 ^{d :***, e *}	
Epididymis	56,01 ± 0,762 ^{d :***, e :*}		29,9 ± 0,809 ^{abd :***}		5,136±0,145 ^{d :***, e :*}	

Histological study

The estimated histological examination of the testis of different groups of rats after four (4) weeks of treatment is mentioned in the figures. The histological study of the testicle of the control group, showed several seminiferous tubes, each tube consists of a central lumen bordered by a seminiferous epithelium containing germ cells, the testicles of the positive control (F1, F2) present preserved histological aspects like healthy tissue (C). Rats exposed to lead acetate (50mg/kg) by gavage, showed some changes which are noted according to their severity, we noticed dystrophic parenchyma, widening of interstitial spaces. The lumens of the seminiferous tubules are empty or few spermatozoa, the cell layers in the seminiferous tubules are scarce thus vascular congestion. On the other hand, the rats treated with the (F1+Pb) and (F2+Pb) combinations demonstrated a normal structural organization of the testicles in the two combination groups (F1+Pb, F2+Pb), a well-defined epithelium. The protective effect of aqueous extract of *OFI* flowers revealed recovery of spermatogenesis, seminiferous tubules become more normal, and seminiferous lumen is enriched with spermatozoa. The epididymis is a very coiled canal where the maturation of the spermatozoa takes place (acquisition of their forward mobility essential to their fertilization functions). The epididymis of the control group shows a well-structured tissue architecture, the central lumen presents a density of spermatozoa, same observation in the histological sections of the positive control groups (F1 and F2). Exposure to

lead acetate caused alterations, we observed architectural disorganization (photo No. 5), fibrosis of the wall of the epididymal tubes, with narrowing of light and absence of germ cells compared to the group control and positive control groups (F1, F2). However, co-administration of aqueous extracts of *OFI* flowers in groups (F1+Pb and F2+Pb) at different doses improved the results, which was revealed by well-defined tissue architectures, normal cell distribution, increased density of sperm in the epididymal lumen.

The control testicle (C) composed of well-defined seminiferous tubules (Photo N°01). These seminiferous tubules consisting of a central lumen bordered by a seminiferous epithelium (Photo N°2). Those of the positive controls (F1, F2) have a histological appearance like healthy tissue (Photos N°03 and N°04). The testicles treated with 50 mg/kg of lead acetate present disorganized seminiferous tubules, empty interstitial space (L.I.V), lumen of the seminiferous tubules are empty (LTS.V) and enlargement of the interstitial spaces (EEI) (photo N°5, 6, 7, 8). The testicle reveals an atrophic parenchyma (AP) and dystrophic parenchyma (PD) (Photo N°9). Presence of vessel congestion (CVx).

DISCUSSION

Fertility physiology in mammals is very sensitive to disturbances in the body by exogenous agents. Numerous studies indicate a decrease in the number and quality of male sex cells in recent years induced by several pollutants (Jegou, 1996, debouche et al., 2018). Heavy metals that pose serious ecological problems, because of their presence in the biosphere, and have the ability to accumulate in the tissues of living organisms (Drif et al., 2019). Lead is a cumulative toxic element that causes changes in rat sperm quality and is an inducer of oxidative stress (WHO, 2017). Disrupts spermatogenesis (Anomaa et al., 2017; Wani et al., 2015). Lead acetate induces in male rats functional and morphological abnormalities on spermatozoa, disrupts spermatogenesis, steroidogenesis, acrosome function, neuroendocrine system, this causes male infertility (Anomaa et al., 2017; Wani et al., 2015; INRS, 2018). As well as the number, motility of sperm and increased fragmentation of DNA (Akcha et al., 2020; Li et al., 2018; Wani et al., 2015). Perhaps explained by tissue atrophy induced by inhibition of spermatogenesis, so could be related to the harmful effects of lead on the hypothalamo-pituitary-testicular axis (Abdel-Emam, 2021). Other studies also reported that ingestion of lead acetate reduced the diameter or thickness of seminiferous tubes, a neuroendocrine disruption that leads to testicular atrophy (Didolkar et al., 1988; Chanel et al., 2017). On the other hand, a mitigating effect is observed on body weight and sexual organs, in groups treated with the combination of aqueous extract of *OFI* and lead especially in the lot (F2+Pb). Our results are consistent with studies by Ncibi et al., 2008; Leo et al., 2010; Gouthamchandra et al., 2010. Explained this protection by the antioxidant capacity of the *OFI* flower extract, these bioactive components act as traps and lipid peroxidation inhibitors of ROS. According to phytochemical screening, the aqueous extracts of *OFI* flowers present its richness in phenolic compound. In the present study we observed a positive correlation between testicular weight, epididymis and testosterone concentration in the experimental groups. Our results are consistent with studies by Mokhtari et al., 2011 confirmed that the experimental groups that received 50 and 100 mg/kg of lead acetate showed a significant decrease in testosterone levels. Other recent studies have shown that the toxicity of lead in rats treated with 50 mg/kg lead acetate results in reductions in serum FSH, LH, testosterone, body weight and testis, as well as 17 β -Steroid hydroxy dehydrogenase was significantly reduced (Abdel-Emam, 2021). Other studies have shown that the decrease in testosterone concentration may be mainly due to the toxic effect of lead on Leydig cells (Anomaa et al., 2017). So, the alteration of spermatogenesis during lead exposure can be attributed to hormonal imbalance at the hypothalamic-pituitary axis, this translates into the reduction of follicular-stimulating hormone-luteinizing (FSH), the gonadotropin-liberating hormone (Gn, and testosterone concentration (Doumouchsis et al., 2009, Rezaei et al., 2018; Ahmed and Abdel-Emam, 2019). In contrast, the combination of flower extract (*OFI*) with lead significantly improved serum

testosterone levels. According to (Sakly et al., 2014). Showed that the administration of *OFI* flower extract (250 and 500 mg/ kg) in rats against nickel-induced toxicity in the testes of rats resulted in a significant recovery of testosterone, follicle-stimulating hormone (FSH) levels and luteinizing hormone (LH). The therapeutic power of *OFI* flower extract is validated and could be explained by several cumulative factors, including phenolic compounds, flavonoids and polysaccharides of plant origin, that act as traps and inhibitors of lipid peroxidation of ROS via hydrogen (Leo et al., 2010; Gouthamchandra et al., 2010). In the present study the administration of lead acetate to rats showed a clear toxic alteration in the parameters of sperm biology, the results of which indicate that the concentration, vitality and different types of sperm velocity significantly reduced after oral gavage of lead acetate and velocity parameters (VCL, VSL, VAP, BFC) were decreased. Analysis of sperm mobility informs us about their essential kinetic qualities: speed, vitality and progressivity in a straight line. These factors are essential for fertilization of the oocyte. Our results are in perfect synchronization with those of Pizent et al., 2012; Abdel-Emam, 2021. Numerous studies have reported that lead exposure disrupts the parameters of sperm biology leading to male infertility (Kumar, 2018; Martin et al., 2017; Wahab et al., 2019). According to Naha et al., 2005, Moreover, Li et al., 2018; Wani et al., 2015 demonstrated that high concentrations of lead in drinking water cause decreased sperm motility, density, and viability. Also, they lead to significant morphological abnormalities, and alteration of endocrine function of reproductive organs (Chowdhury, 2009). Lead-induced spermatogenesis (Abdel-Emam, 2021). Metal ions induce Sperm concentration is reduced in our study can be attributed to dysfunction apoptosis of Leydig cells so the decrease of all sperm cell populations, in particular, the sperm and sperm forms (Shubina et al., 2016; Corpas et al., 2002). The deleterious effects of lead on different types of sperm velocity can be attributed to changes in the intermediate part, particularly in mitochondria, axoneme and flagella function (Castellini et al., 2009). By also inducing phosphorylation of axonem proteins or by reducing the production of ATP in sperm cells that are necessary for the wavy movement of sperm (Aitken and Baker, 2006). In addition, the decrease in sperm concentration and viability in our experiment could be reasonably assumed also on lead-induced oxidative stress. Which, it was noted that lead causes excessive production of ROS could also be related to mitochondrial dysfunction leading to lipid peroxidation whose consequences, decreased sperm motility, concentrations, viability, increased DNA fragmentation and apoptosis (Sainath et al., 2011; Chowdhury, 2009). Lead also affects sperm membrane morphology (Shubina et al., 2016). The exploration of male infertility in first line is based on the analysis of the spermogram and spermocytogram, these two examinations give an overall idea of the fertilizing potential of the spermatozoon, but does not inform us on the quality of sperm DNA. Lead acetate in our research caused increased fragmentation of DNA. DNA damage causes disruption of cell functions including somatic and germ cells (Akcha et al., 2020). These results are certainly explained as reported by several studies, by the deleterious effect of ROS when present in high concentrations, which induces peroxidation of the lipids of the spermatid membrane and leads to the deterioration of the axonal structures of spermatozoa, thereby reducing their mobility (asthenospermia (Saleh et al., 2002). On the other hand; by the Caspases, mediators of apoptosis that are found in high concentration in the sperms of individuals characterized by asthenospermia (Hamamah et al., 2000). On the other hand, oral administration of the aqueous extract of *Opuntia ficus indica* with lead in the groups (F1+Pb, F2+Pb) attenuated the biological parameters of sperm against lead toxicity whose results, leading to an improvement in fertility parameters (concentration, viability, velocity, kinetic parameters (VCL, VSL, VAP, BFC) were significantly increased compared to the toxic group (Pb) with no significant change in comparison with the control group, plus decreased DNA fragmentation. Indeed, the antioxidant property of the aqueous extract of *OFI* improved the biological parameters of sperm in combination groups (F1+Pb, F2+Pb). Its results are supported by (Sakly et al., 2014). Also showed that the addition of *OFI* flowers improved sperm parameters, testosterone levels, sperm concentration and mobility in sheep. Studies by (Allai et al., 2016) have reported that prickly pear flowers have beneficial and mitigating effects against DNA fragmentation in men's sperm. They observed a significant reduction

in sperm DNA fragmentation as well as an improvement in sperm quality, mobility and viability. Indeed, our phytochemical results confirm that the methanolic extract of *OFI* flowers exerts an exceptional antioxidant and anti-inflammatory function. Lead is an exogenous poison that can induce ROS production, disrupt and inhibit the activity of free radical scavenging enzymes, improve lipid peroxidation and cause tissue damage (WHO, 2017). In our experiment sub chronic exposure of rats to lead acetate for 4 weeks resulted in a significant increase in MDA thus significantly reducing the level of GSH and Gpx enzyme activity in the testes and epididymis. Several literature shows that lead induces a significant decrease in the activities of antioxidant enzymes GSH, GPX and CAT in animals, and a significant increase in the level of expression of Malondialdehyde (MDA) (Chen et al., 2019, Han et al., 2017). Our results are consistent with (Piasecka et al., 1997; Gavrić et al., 2017). Explained that lead has a high affinity for sulfhydryl groups and direct interaction with biological membranes inducing lipid peroxidation, and decreased free radical trapping enzymes and glutathione. Lead ions also lead to lipid peroxidation of the sperm membrane due to its richness in polyunsaturated fatty acids that could cause serious damage to the membrane, this can change the structure of the sperm and the integrity of the membrane, on the other hand changes in the DNA (Aitken et al., 1987; Chen et al., 2004; Hou et al., 2018). A recent study has shown that oxidative stress plays an important role in Pb-induced cell damage that not only influences the antioxidant defense system, but also leads to tissue damage (Ademuyiwa et al., 2009). Our histological study confirms tissue damage induced by lead. Nrf2 is an essential regulator between the cellular antioxidant response and the xenobiotic metabolism of heavy metals that are generally multifactorial, with ROS being the main contributor to heavy metal toxicity (Rehman et al., 2018). However the aqueous extract of *OFI* managed to protect the organs to study against oxidative stress induced by lead in groups (F1+Pb, F2+Pb) and bring them back to normal levels. We noted the decrease in the concentration of MDA as well as the increase in the concentration of GSH, and the activity of Gpx in the testicles and epididymis. These results are consistent with other studies that confirm that the administration of the aqueous extract of *Opuntia ficus indica* flowers to protect organs from oxidative stress induced by several xenobiotics (Hamed et al., 2014; Chougui et al., 2013; Ammar et al., 2015). Showed that *OFI* flower extract had antioxidant effect because of these bioactive compounds. According to Sakly et al., 2014; Allai et al., 2016 in turn demonstrated that they managed to overcome oxidative damage induced by nickel at the testicular level and It was sported that *OFI* flowers managed to prevent lead-induced stress (BENKAHOUL, 2018). lead leads to depletion of glutathione and proteins, leading to increased production of reactive oxygen species such as peroxide ion, hydroxyl radical and H O ROS increase lipid peroxidation by oxidation of polyunsaturated fatty acids of membrane phospholipids. These alterations modify the fluidity of membranes (Liu et al., 2012; Laamech et al., 2017). In testicles and epididymis our results agree with (Piasecka et al., 1997; Gavrić et al., 2017). Nrf2 is an essential regulator between the cellular antioxidant response and the xenobiotic metabolism of heavy metals that are generally multifactorial, with ROS being the main contributor to heavy metal toxicity (Rehman et al., 2018). Our histological study confirms tissue damage induced by lead. However, the aqueous extract of *OFI* flowers managed to protect the organs from oxidative stress induced by lead in the groups (F1+Pb, F2+Pb) and bring them back to normal levels. We noticed the decrease in the concentration of MDA as well as the increase in the concentration of GSH, and the activity of Gpx in the testicles and epididymis. These results are consistent with other studies that confirm that the administration of aqueous *OFI* flower extract to protect organs from oxidative stress induced by several xenobiotics (Hamed et al., 2014; Chougui et al., 2013; Ammar et al., 2015) showed that *OFI* flower extract had antioxidant effect because of these bioactive compounds. According to Sakly et al., 2014; Allai et al., 2016 in turn demonstrated that they managed to overcome oxidative nickel-induced damage to the testicle. It has been planted that *OFI* flowers have successfully prevented oxidative stress induced by lead (Chahdoura et al., 2014; Rabhi et al., 2015; BENKAHOUL, 2018).

Conclusion

In conclusion, lead is a toxic element that disrupts sperm quality parameters, induces oxidative stress and affects testicular MDA, GSH and Gpx. Co-administration of aqueous extracts of *Opuntia ficus indica* with lead succeeded in maintaining levels of oxidative stress markers close to their normal physiological values and improved sperm quality. The extract of *OFI* flowers has an exceptional antioxidant effect, attenuating the toxicity of lead.

Declarations

Author Contribution

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us. We understand that the Corresponding Author is the sole contact for the Editorial process. She is responsible for communicating with the other authors about progress.

References

A.

1. **Abdel- Emam, Esraa, A., Ahmed ., Rania A.(2021).** Ameliorative effect of L- carnitine on chronic lead- induced reproductive toxicity in male rats
2. **Abdel Moneim, A. (2015).** Indigofera oblongifolia Prevents Lead Acetate-Induced Hepatotoxicity, Oxidative Stress, Fibrosis and Apoptosis in Rats. PLOS ONE, 11: (7) 1-18
3. **Ademuyiwa, O., Agarwal, R., Chandra, R., Behari, J. R. (2009).** Lead-induced phospholipidosis and cholesterogenesis in rat tissues. Chem Biol Interact, 179:(2) 314-320.
4. **Ahmed, E. A., & Abdel- Emam, R. A. (2019).** The potential impact of 1st and 2nd generation antihistamines on male fertility. Comparative Clinical Pathology, 28, 1465– 1470. <https://doi.org/10.1007/s00580-019-02993-0>
5. **Aitken, R.J., Clarkson, J.S. (1987).** Cellular basis of defective sperm function and its association with the genesis of reactive oxygen species by human spermatozoa. J Reprod Fertil;81:459–469.
6. **Aitken, R. J., & Baker, M. A. (2006).** Oxidative stress, sperm survival and fertility control. Molecular and Cellular Endocrinology, 250, 66– 69. <https://doi.org/10.1016/j.mce.2005.12.026>
7. **Akacha, A., Badraoui, R., Rebai, T., & Zourgui, L. (2020).** Effect of *Opuntia ficus indica* extract on methotrexate-induced testicular injury: a biochemical, docking and histological study, Journal of Biomolecular Structure and Dynamics, DOI: 10.1080/07391102.2020.1856187 To link to this article: <https://doi.org/10.1080/07391102.2020.1856187>
8. **Alhakmani, F., Sokindra., Kumar., Shahalam., Khan.(2013).** Estimation of total phenolic content, in-vitro antioxidant and anti-inflammatory activity of flowers of *Moringa oleifera*. Asian Pacific journal of tropical biomedicine. DOI:10.1016/S2221-1691(13)60126-4
9. **Alimi, H., Hfaiedh, N., Bouoni, Z., Sakly, M., Ben Rhouma, K. (2011).** Evaluation of antioxidant and antiulcerogenic activities of *Opuntia ficus indica* f. inermis flowers extract in rats. Environ. Toxicol. Pharmacol. 32, 406–416.
10. **Allai, L., Druart, X., Öztürk, M., BenMoula, A., Nasser, B., & El Amiri, B. (2016).** Protective effects of *Opuntia ficus-indica* extract on ram sperm quality, lipid peroxidation and DNA fragmentation during liquid storage.

11. **Allouche, L., Hamadouche, M., Touabti, A. (2009).** Chronic effects of low lead levels on sperm quality, gonadotropins and testosterone in albinos rats. *Exp Toxicol Pathol* [Epub ahead of print].
12. **Ammar, I., Ennouri, M., Khemakhem, B., Yangui, T., Attia, H. (2012).** Variation in chemical composition and biological activities of two species of *Opuntia* flowers at four stages of flowering. *Industrial Crops and Products*, 37 : (1) 34–40
13. **Ammar, I., Ennouri, M., Attia, H. (2015).** Phenolic content and antioxidant activity of cactus (*Opuntia ficus-indica* L.) flowers are modified according to the extraction method. *Ind. Crops Prod.* 64, 97–104.
14. **Andjelkovic, M., Buha Djordjevic, A., Antonijevic, E., Antonijevic, B., Stanic, M., Kotur-Stevuljevic, J., Bulat, Z. (2019).** Toxic Effect of Acute Cadmium and Lead Exposure in Rat Blood, Liver, and Kidney. *International Journal of Environmental Research and Public Health*, 16 : (2) 274
15. **Anomaa, G., Barek-Deligny, N., Daneta, S., Bouard, B. (2017).** Occupational exposure to lead: practical situation in a shooting range. *medicine and armies*, 4: (3) 357-363
16. **Aragona, M., Lauriano, E. R., Pergolizzi, S., Faggio, C. (2017).** *Opuntia ficus-indica* (L.) Miller as a source of bioactivity compounds for health and nutrition. *Natural Product Research*, 32: (17), 2037–2049.
17. **Azawi, N.H., Berg, K.D., Thamsborg, A.K.M., Dahl, C., Jepsen, J.V., Kroman-Andersen B., et al. (2015).** Laparoscopic and robotic nephroureterectomy: does lymphadenectomy have an impact on the clinical outcome? *Int Urol Nephrol.* 2015;49:1785–92.

B

18. **BaSalamah, M.A., Abdelghany, A.H., El-Boshy, M., Ahmad, J., Idris, S., Refaat, B. (2018).** Vitamin D alleviates lead induced renal and testicular injuries by immunomodulatory and antioxidant mechanisms in rats. *Scientific Reports* volume 8, Article number: 48530 ORCID: orcid.org/0000-0003-4267-1016³
19. **Berrahal, A., Lasram, M., El Elj, N., Kerkeni, A., Gharbi, N., El-Fazâa, S. (2011).** Effect of age-dependent exposure to lead on hepatotoxicity and nephrotoxicity in male rats. *Environmental Toxicology*, 26: (1) 68–78.
20. **Berredjem, R., Mansouri, O., Abdennour, C., Boulakoud, M.K., Khelili, K. (2014).** What is the benefit of garlic supplementation to Pb intoxicated rats? *Adv Environ Biol* 8:268–275
21. **Bruits, M and Bucar, F. (2000).** Antioxidant activity of *Nigella sativa* essential oil. *Phytother. Res.*, 14, 323-328.
22. **Burroughs Peña, M.B., Rollins Allman, T. (2017).** Environmental Exposures and Cardiovascular Disease: A Challenge for Health and Development in Low- and Middle-Income Countries, *Medicine, Cardiology clinics*.

C

23. **Castellini, C., Mourvaki, E., Sartini, B., Cardinali, R., Moretti, E., Collodel, G., Fortanero, S., Sabbioni, E., Renieri, T. (2009).** In vitro toxic effects of metal compounds on kinetic traits and ultra structure of rabbit spermatozoa. *Reprod Toxicol* 27:46–54. <https://doi.org/10.1016/j.reprotox.2008.12.003>
24. **Chahdoura, H., João, C. M., Barros, L., Santos-Buelga, C., Ferreira, I. C., & Achour, L., Chia, S.E., Yap, E., and Chia, K.S. (2014).** Delta-aminolevulinic acid dehydratase (ALAD) polymorphism and susceptibility of work-exposed to inorganic lead and its effects on neurobehavioral functions. *Neurotoxicol* 2014; 25: 1041-1047.
25. **Chanel, O., Dollfus, C., Haguenoer, J.M., Hartemann, P., Béatrice Larroque, G.H., Lison, D., Marret, S., Pinon-Lataillade, G., Prémont, J., et al. (2017).** Lead in the environment: what health risks. HAL ID: hal-01571950. <https://hal-lara.archives-ouvertes.fr/hal-01571950>.
26. **Chen, S.S. (2004).** Flavonoids in Food and their health benefits. *Plant Food Human Nutrition*; 59: p113-122

27. **Chen, L., Shi, G.R., Huang, D.D., Li, Y., Ma, C.C., Shia, M., Su, B.X., Shi, G.J. (2019).** Male sexual dysfunction: A review of literature on its pathological mechanisms, potential risk factors, and herbal drug intervention. *Biomedicine and Pharmacotherapy*, 112, 108-585.
28. **Chougui, N., Louaileche, H., Mohedeb, S., Mouloudj, Y., Hammoui, Y., Tamendjari, A. (2013).** Physico-chemical characterization and antioxidant activity of some *Opuntia ficusindica* varieties grown in North Algeria. *African Journal of Biotechnology*. 12 : 299-307.
29. **Chowdhury, A.R. (2009).** Recent advances in heavy metals induced effect on male reproductive function-a retrospective. *Al Ameen J Med Sci*2:37–42.
30. **Corpas, I., Castillo, M., Marquina, D., Benito, M.J. (2002).** Lead intoxication in gestational and lactation periods alters the development of male reproductive organs. *Ecotox Environ Safe*53:259–266. <https://doi.org/10.1006/eesa.2002.2230>
31. **Cuendet M., H.K., Dyatmiko W. &Potterat O.(1997).** Iridoid Glucosides with Free Radical Scavenging Properties from *Fagvaea blumei* *Helvetica Chimica Acta*, 80, 1144-1152.

D

32. **Debouche, A., Moreau, L., Bernard, N., Bracikowski, N. (2018).** Study of the optimal multiplication ratio of a high torque magnetic planetary transmission. *Electrical Engineering Symposium, University of Lorraine [UL]*, Jul 2018, Nancy, France. ffhal-02981859f
33. **Derouiche, S., Zeghib, k., Gharbi, S., Khelef, Y. (2018).** Beneficial Effects of *Aristolochia Longa* and *Aquilaria Malaccensis* on Lead-Induced Hematological Alterations and Heart Oxidative Stress in Rats. *Journal of Chemical and Pharmaceutical Research*, 10 : (9) 8-15.
34. **Didolkar, A.K., Keizer-Zucker, A., Sundaram, K., Bardin, C.W., Agback, H., et Johansson, E.D.B. (1988).** Effect of sulfasalazine and its analogs on fertility in male rats, *Contraception* 37(5), p.539-548.
35. **Doumouchtsis, K. K., Doumouchtsis, S. K., Doumouchtsis, E. K., & Perrea, D. N. (2009).** The effect of lead intoxication on endocrine functions. *Journal of Endocrinological Investigation*, 32, 175– 183. <https://doi.org/10.1007/BF03345710>
36. **Drif, F., Abdenmour, C., Ciğerci, İH., Muddassir, A.M., Mansouri, O., Messarah, M. (2019).** Preliminary assessment of stress and genotoxicity biomarkers in bivalve molluscs from the Gulf of Annaba, Algeria. *Bull Environ Contam Toxicol* 102:555– 559. <https://doi.org/10.1007/s00128-019-02583-4>

F

37. **Flohé, L., and Günzler, W.A. (1984).** Assays of Glutathione Peroxidase. *Methods in Enzymology*, 105, 114-120.
38. **Flora, G., Gupta, D., Tiwari, A. (2012).** Toxicity of lead: A review with recent updates. *Interdiscip Toxicol*. 2012 Jun. 5 (2):47-58.

G

39. **Gavrić, J., Anđelković, M., Tomović, L., Prokić, M., Despotović, S., & Gavrilović Bet Saičić, Z. (2017).** Oxidative stress biomarkers, cholinesterase activity and biotransformation enzymes in the liver of dice snake (*Natrix tessellata* Laurenti) during pre-hibernation and posthibernation: A possible correlation with heavy metals in the environment. *Ecotoxicology and Environmental Safety*. 138: 154-162
40. **Gouthamchandra., Bose. (2010),** modern marketing, eastern economy.

H

41. **Hamamah, S., Corcia, L., Entezami, G., El Wией, H. (2000).** "ART techniques and their indications in male subfertility". *Andrology*, 10, 03, 279-283.

42. **Hamed, N.A., Trnka, J. (2014).** The Amplituhedron. *J. High Energ. Phys.* 30 (2014).
<https://doi.org/10.1007/JHEP10>
 43. **Han, L., et al. (2017).** S. cerevisiae Trm140 has two recognition modes for 3-methylcytidine modification of the anticodon loop of tRNA substrates. *RNA* 23(3):406-419
 44. **Hou, G., Surhio, M. M., Ye, H., Gao, X., Ye, Z., Li, J., Ye, M. (2018).** Protective effects of a Lachnum polysaccharide against liver and kidney injury induced by lead exposure in mice. *International Journal of Biological Macromolecules*.
- I
45. **INRS (Institut national de recherche scientifiques). (2018).** Lead and mineral compounds. Toxicological data sheets, n°59. www.inrs.fr/fichetox
- J
46. **Jamesdaniel, S., Rosati, R., Westrick, J., & Ruden, D. M. (2018).** Chronic lead exposure induces cochlear oxidative stress and potentiates noise-induced hearing loss. *Toxicology Letters*, 292, 175–180.
[doi:10.1016/j.toxlet.2018.05.004](https://doi.org/10.1016/j.toxlet.2018.05.004)
 47. **JEGOU (Bernard). (1996).** Do men become less fertile Source: THE RESEARCH. No. 288, 1996/06, pages 60-65. Document type: ARTICLE.
- K
48. **Keller, A.L., Girard, C., Chaumont, J. P. (2019).** Opuntia ficus indica L. Mill, the prickly pear or nopal, a plant with multiple uses. plant sheet.
 49. **Kumar, S. (2018).** Occupational and environmental exposure to lead and reproductive health impairment: An overview. *Indian Journal of Occupational and Environmental Medicine*, 22(3), 128– 137.
- L
50. **Laamech, J., El-Hilaly, J., Fetoui, H., Chtourou, Y., Gouitaa, H., Tahraoui, A., Lyoussi, B. (2017).** Berberis vulgaris L. effects on oxidative stress and liver injury in lead-intoxicated mice. *Journal of Complementary and Integrative Medicine*, 14: (1) 1-11
 51. **Leo et al. (2010).** Profiling the chemical content of Opuntia ficus-indica flowers by HPLC-PDA-ESI-MS and GC/EIMS analyses. *Phytochem. Latvian*.
 52. **Li, C., Zhao, K., Zhang, H., Liu, L., Xiong, F., Wang, K., & Chen, B. (2018).** Lead exposure reduces sperm quality and DNA integrity in mice. *Environmental Toxicology*, 33(5), 594– 602. <https://doi.org/10.1002/tox.22545>
 53. **Liu, C.M., Ma, J.Q., Sun, Y.Z. (2012).** Puerarin protects the rat liver against oxidative stress-mediated DNA damage and apoptosis induced by lead. *Exp. Toxicol. Pathol.* 64:575-582.
- M
54. **Martin, K. K., Arsene, A. M., Melaine, M. G., Ernest, Z. N., Joseph, D. A., Mireille, D., & David, N. J. (2017).** Effects of chronic lead exposure on zinc concentration and spermatoc parameters in Wistar rats. *AMBS*, 3(2), 51– 58.
 55. **Meamar, M., Zribi, N., Cambi, M., Tamburrino, L., Marchiani, S., Filimberti, E., Fino, M.G., Biggeri, A., Menezo, Y., Forti, G., et al. (2012).** Sperm DNA fragmentation induced by cryopreservation: new insights and effect of a natural extract from Opuntia ficus-indica. *Fertility and Sterility* 98 326–333.
([doi:10.1016/j.fertnstert.2012.05.001](https://doi.org/10.1016/j.fertnstert.2012.05.001))
 56. **Mokhtari, M., Zanboori, M. (2011).** The Effects of Lead Acetate on Sexual Behavior and the Level of Testosterone in Adult Male Rats. *Int J FertilSteril.* 1:13–20.
- N

57. **Naha, N., Bhar, R. B., Mukherjee, A., & Chowdhury, A. R. (2005).** Structural alteration of spermatozoa in the persons employed in lead acid battery factory. *Indian Journal of Physiology and Pharmacology*, 49, 153– 162.
 58. **Ncibi, S., Ben Othman, M., Akacha, A., Krifi, M., Zourgui, L. (2008).** Opuntia ficus indica extract protects against chlorpyrifos-induced damage on mice liver. *Food and Chemical Toxicology*, 46: (2) 797–802.
- O**
59. **Ohkawa, H., Ohishi, N., Yagi, K. (1979).** Assay for lipid peroxidation in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry*, 95, 351-358.
- P**
60. **Panico, M., Cardile, V., Garufi, F., Puglia, C., Bonina, F., Ronsisvalle, S. (2007).** Effect of hyaluronic acid and polysaccharides from *Opuntia ficus-indica* (L.) cladodes on the metabolism of human chondrocyte cultures .*J. Ethnopharmacol*, 111 :315-321.
 61. **Patra, R.C., Rautray, A.K., Swarup, D. (2011).** Oxidative stress in lead and cadmium toxicity and its amelioration. *J. Vet. Intern. Med.*, 2011: 457327.
 62. **Pérez-Torrero, E., Garcia-Tovar, SE., Luna-Rodriguez, LE., Rodríguez-Garcia, M.E. (2017).** Chemical composition of prickly pads from (*Opuntia ficus-indica* (L.) Miller related to maturity stage and environment. *Int J Plant Biol Res.* 5(1):1061.
 63. **Piasecka, M., Wenda-Rozewicka, L., Kram, A. (1997).** Computerized analysis of cytochemical reactions of spermatozoa in rats chronically treated with lead acetate [Pb(II)]. *Folia Histochemica et Cytobiologica*. 35(2): p.129-131
 64. **Picot, A. (2010).** Heavy metals: major toxins / The trio mercury, lead, cadmium. A ssociation of toxicology and chemistry (ATC) Paris (2010). CNRS France.
 65. **Pizent, A., Tariba, B., & Živković, T. (2012).** Reproductive toxicity of metals in men. *Archives of Industrial Hygiene and Toxicology*, 63(Suppl 1), 35– 46. <https://doi.org/10.2478/10004-1254-63-2012-2151>
 66. **Polshettiwar, S.A., Ganjiwale, R.O., Wadher, S.J., Yeole, P.G. (2007).** "Spectrophotometric estimation of total tannins in some ayurvedic eye drops". *Indian J Pharm Sci*, 69:574-6.
- R**
67. **Rabhi, A., Falleh, H., Limam, F., Ksouri, R., Abdelly, C., Raies, A. (2015).** Upshot of the ripening time on biological activities, phenol content and fatty acid composition of Tunisian *Opuntia ficus-indica* fruit. *Afr J Biotechnol.* 12 : 5875–5885
 68. **Rehman, M.A., et al. (2018).** The Nem1/Spo7-Pah1/lipin axis is required for autophagy induction after TORC1 inactivation. *FEBS J* 285(10):1840-1860
 69. **Rezaei, N., Mardanshahi, T., Shafaroudi, M. M., Abedian, S., Mohammadi, H., & Zare, Z. (2018).** Effects of L-carnitine on the follicle- stimulating hormone, luteinizing hormone, testosterone, and testicular tissue oxidative stress levels in streptozotocin- induced diabetic rats. *Journal of Evidence- Based Integrative Medicine*, 23, 1– 10. <https://doi.org/10.1177/2515690X18796053>
- S**
70. **Sainath, S.B., Meena, R., Supriya, C., Reddy, K.P., Reddy, P.S. (2011).** Protective role of *Centella asiatica* on lead-induced oxidative stress and suppressed reproductive health in male rats. *Environ Toxicol Pharmacol* 32:146– 154. <https://doi.org/10.1016/j.etap.2011.04.005>
 71. **Sakly, C., Rekik, M., Ben Salem, I., Lassoued, N., Gonzalez-Bulnes, A., & Ben Salem, H. (2014).** Reproductive response of fat-tailed Barbarine ewes subjected to short-term nutritional treatments including spineless cactus (*Opuntia ficus-indica* f. *inermis*) cladodes. *Journal of Animal*.

72. **Saleh, R.A.,Agarwal,A, (2002).** Oxidative stress and male infertility: from research bench to clinical practice.Nov-Dec;23(6):737-52.
 73. **Shubina, O.S., Dudenkova, N.A. (2016).**The effect of lead on the process of spermatogenesis in sex glands of male albino rats. Vet World 9:1129–1134. <https://doi.org/10.14202/vetworld.2016.1129-1134>
 74. **Seregin, I.V and Ivanov, V.B. (2001).** Physiological Aspects of Cadmium and Lead Toxic Effects on Higher Plants. Russian Journal of Plant Physiology, 48, 523-544.
- T**
75. **Toure, H.A. (2017).** Contribution to the Valorization of Opuntia ficus indica in Nutraceuticals and Phytotherapy. Doctoral thesis. Mohammed V University Rabat, Morocco
- W**
76. **Wahab, O. A., Princely, A. C., Oluwadamilare, A. A., Oore- oluwapo, D. O., Blessing, A. O., & Alfred, E. F. (2019).** Clomiphene citrate ameliorated lead acetate- induced reproductive toxicity in male Wistar rats. JBRA Assisted Reproduction, 23(4), 336– 343.
 77. **Wani, A. L., Ara, A., Usmani, J. A. (2015).** Lead toxicity: a review. Interdisciplinary Toxicology, 8 :(2) 55–64.
 78. **World Health Organization (WHO). (2017).** Lead poisoning and health. Available from: <http://www.who.int/mediacentre/factsheets/fs379/en/>[Accessed 2017 28/09].
- Z**
79. **Zhishen, J., Mengcheng, T., et Jianming, W. (1999).** The determination of flavonoid contentsin mulberry and their scavenging effects on superoxide radicals. *Food chemistry.*, 64(4), 555559.

Appendix

Appendix I is not available with this version.

Figures

Datafile Name:Extrait Flavo NFO 2025.lcd
Sample Name:Extrait Flavo 2022
Sample ID:001





Figure 1

Chromatogram of the aqueous extract of *Opuntia ficus indica* flower.



the concentration of spermatozoa in rats exposed to lead and combined with the aqueousextract of *Opuntia ficus indica* flowers for 4 weeks of treatment.



Sperm velocity (fast, medium, slow and motionless) of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.




Figure 4

VCL (curvilinear velocity), VSL (straight-line velocity), VAP (velocity average path), levels of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.




Figure 5

Trajectory crossover frequency (BCF) of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.




Figure 6

Percentage of live and dead spermatozoa (%) of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment




Figure 7

Testosterone levels of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.




Figure 8

Percentage of DNA fragmentation (%) of rats exposed to lead and combined with the aqueous extract of *Opuntia ficus indica* flowers for 4 weeks of treatment.




Figure 9

Characteristic of the testicles of rats exposed to lead acetate and co-administered with aqueous extract of *OFI* flowers for 4 weeks of treatment.(G×40)




Figure 10

Characteristic of the epididymis of rats exposed to lead acetate and co-administered with aqueous extract of *OFI* flowers for 4 weeks of treatment.(G×40) (C, control; PB, lead acetates; F1, positive control (aqueous extract of *Opuntia ficus indica* flowers at 250 mg/kg dose); F2; positive control (aqueous extract of *Opuntia ficus indica* flowers at 500mg/kg dose); F1+b combination of lead and (aqueous extract of the flowers of *Opuntia ficus indica* at 250 mg/kg); F2+Pb; combination between lead and (aqueous extract of the flowers of *Opuntia ficus indica* at 500 mg/kg dose). L.C. E; light from efferent ducts; R. L; Narrowing of light; F. R; wall fibrosis; L. V; empty light; S Z; sperm.