Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Physicochemical quality properties of peach (Prunus persica L.) varieties at Holetta, Ethiopia

Tajebe Mosie

tajebemu@gmail.com

Ethiopian Institute of Agricultural Research

Habtam Setu

Ethiopian Institute of Agricultural Research

Getaneh Seleshi

Ethiopian Institute of Agricultural Research

Research Article

Keywords: fruit, physicochemical, quality, varieties, TSS, TA, ripening index

Posted Date: November 25th, 2023

DOI: https://doi.org/10.21203/rs.3.rs-3589445/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Abstract

Holetta has different peach fruit varieties that could be used for various purposes. However, their fruit quality characteristics were not fully identified. Studies have indicated that the physicochemical qualities of peach fruits are influenced by a number of factors, with the varietal factor being one of the most important. Limited information regarding the factors that affect these qualities is available in the country, specifically at Holetta. Therefore, this study was initiated to evaluate the physicochemical quality properties of peach fruit varieties. Their physical quality traits such as fruit length, fruit diameter, fruit shape index, and average fruit weight, as well as chemical quality parameters like TSS, specific gravity, TA, ripening index, ascorbic acid content, and pH, were evaluated. The result revealed that both physical and chemical quality parameters were significantly affected by varietal factors. Amongst, Bonny Gold variety had the longest fruit length and Florida down had the largest fruit diameter. As far as average fruit weight is concerned, 88 – 18 W had the heaviest weight with 111.98 g, while Transvalia had the highest TSS and specific gravity. However, the Summersun and 9A-35C varieties recorded the highest TA and ripening index, respectively. Thus, the fruit's physicochemical quality properties were highly altered with peach varieties.

1. Introduction

Peaches [*Prunus persica* (L.) Batsch] are the world's most widely produced temperate tree fruits after apples and pears [5]. Their increased consumption can make them a significant part of the diet, economically and nutritionally significant [36]. Peach fruits are rich in carbohydrates, vitamins A and C, and minerals content. Peaches can be eaten as fresh fruit and also be processed into canned fruit, jams, jellies, juices, pulp for yogurts, and liquors [32]. Studies have demonstrated that consuming fruits reduces the risk of developing cancer [24], diabetes, cardiovascular diseases, and obesity [2]; [6]. In addition to improving energy balance and weight management [3], they also treat neurodegenerative diseases [39]. Their flavor and aroma are produced due to the balance between sugar, phenolic compounds, organic acids, carotenoids, and volatile compounds [41]; [43]. These attributes influence consumer preference and consumption [17]. Many of these traits are quantitatively inherited, but their genetic control remains unclear [20].

Numerous studies have reported significant variations among peach varieties that could be attributed to climatic conditions, geographical zones, crop genetics, and cultural practices [4]; [18]; [43]. Organoleptic qualities such as sweetness, juiciness, and flavor vary among peach varieties [10]. Individual fruits should also be monitored for changes during ripening, since the ripening pattern of one cultivar may not be applicable to other varieties within the same species [27]. Due to their possible beneficial effects on health, the biochemical constituents of peaches and other fruits have recently received increased attention [35]. Sucrose is the utmost available soluble sugar in peaches, subsequently glucose and fructose, with lower levels of sorbitol [8]. These sugars account for approximately 60% of the soluble sugar content (SSC) in ripe fruit [11]. The concentration of glucose and fructose increases steadily during fruit development, but sucrose accumulates primarily during maturation [29].

It is crucial to study the information on the overall fruit quality to understand the product's performance throughout the supply chain, even if climatic conditions, cultural practices, variety differences, and maturity stages at harvesting, transportation, and storage conditions influence the physicochemical characteristics of peaches. In relation to varietal diversity in Ethiopian environments, there is, however, limited evidence on fruit quality. The highlands of Ethiopia now produce temperate fruits, like peaches, and demand for them has grown over time. In the meantime, agro-processing businesses are springing all throughout the nation, giving peach growers the chance to produce premium fruits that meet consumer demand. The Holetta Agricultural Research Center has introduced various peach fruit varieties from abroad that perform better in terms of productivity and quality. Although the physicochemical properties of fruits can vary with climate, cultivation practices, and varieties, extensive information on fruit quality with various attributes of relevance is sparse. Assessing the physicochemical characteristics of peach fruits is crucial to understanding product behavior across the value chain and utilization for the intended purpose. Studying this high-quality information also assists breeders in setting up new breeding programs and identifying the difference between varieties in terms of their suitability for fresh or processed produce. Therefore, the study was designed to evaluate the physicochemical properties of peach fruit varieties.

2. Materials and Methods

Fruits with uniform maturity stages were collected from different peach varieties at the experimental fields of the Holetta Agricultural Research Center (HARC) which is located in (N9°00', E 38°30'; 2400 m elevation above sea level). The average annual rainfall in the area was 1041.4 mm and the relative humidity of 58.7 percent. The average annual minimum and maximum temperatures were 6.7 and 21.7°C, respectively [21].

Fruits were hand-harvested and taken to the Temperate Fruits Research Laboratory at HARC. The fruits were then washed to cool down the field heat and fifteen fruits with more or less similar maturity were selected and subjected to physicochemical quality assessment. The trial was arranged using a completely randomized design (CRD) and replicated three times. Five fruits were assigned for each plot. Physical parameters such as fruit length, fruit diameter, and fruit weight were first measured for each individual fruit and averaged thereafter. The fruit shape index was also calculated as the ratio of the fruit length to the fruit width [42].

The chemical quality parameters were evaluated using extracted and cleared fruit juices of five fruits per treatment. The titratable acidity was determined using the method of [25], and the vitamin C content was also measured by the method of [7]. Whereas TSS was measured by a digital refractometer (HANNA HI96801) and juice pH was measured by a pH meter (Orion star A211). Finally, specific gravity (SG) was extrapolated from the Brix reading [14] while the TSS/TA ratio was calculated by dividing the value of the Brix reading by titratable acidity.

Data analysis

An analysis of variance (ANOVA) was performed using SAS software version 9.3 [37]. To identify the varieties with substantial differences, the least significant difference (LSD) was utilized.

3. Results and Discussion

3.1. Physical quality traits of peach fruit varieties

The peach varieties were assessed via the fruit length, fruit diameter, fruit shape, and average fruit weight (Table 1). Peach varieties showed highly significant variation in all of the tested physical quality characteristics. Out of the tested thirteen peach varieties that were evaluated, 46% had fruits that were longer and wider fruit diameter than the average (5.29 and 5.31 cm, respectively), while the rest varieties had fruits that were shorter and less wide. The fruit lengths of these varieties lay between 4.85 and 5.81 cm. The Bonny Gold variety had the longest fruit, followed by Transvalia and Early Grand, which had 9.80%, 6.77%, and 4.51% over the mean, respectively, while the 9A-35C variety had the shortest fruits, which were 8.34% shorter than the grand mean. Peach varieties varied in length at both immature and mature stages since the fruits varied in the amount of accumulated assimilates through the growing season [48]. The length and breadth of the fruit were also found to be significantly influenced by variety [40]. With regard to the fruit diameter, peach varieties had a fruit diameter ranging from 4.41 to 6.62 cm. Florida Down variety had the largest fruit diameter, followed by Florida Star and 88-18W which had 24.67%, 16.01%, and 8.47% wider fruit diameters, respectively, as compared to the grand mean. In comparison, the May Crust variety recorded the smallest fruit diameter, 16.95% less wide than the mean diameter. According to [22] the diameter of peach varieties varied from 4.81cm to 6.40 cm.

Table 1
Fruit length, fruit diameter, fruit shape index, and average fruit weight of peach fruit varieties at Holetta

Variety	Fruit length (cm)	Fruit diameter (cm)	Fruit shape index	Average Fruit weight (g)
McRed	5.11fg	4.62gh	1.11b	71.70fgh
88 – 18 W	5.33de	5.76c	0.92e	111.98a
90-19H	5.18def	5.19def	1.00d	63.00hi
9A-35C	4.85h	4.87fg	1.00d	75.37efg
Tropic Beauty	5.36cd	5.46cd	0.98de	68.57gh
88-22C	5.15ef	4.98ef	1.04bcd	58.03i
Early Grand	5.53bc	5.16def	1.07bc	88.41c
Florida Down	5.33de	6.62a	0.81f	76.07efg
Florida Star	4.95gh	6.16b	0.81f	63.32hi
May Crust	5.29def	4.41h	1.20a	78.96def
Summer Sun	5.25def	5.03ef	1.04bcd	85.50cd
Transvalia	5.65ab	5.32de	1.07bc	81.90cde
Bonny Gold	5.81a	5.45cd	1.07bc	97.74b
Mean	5.29	5.31	1.01	78.50
LSD (5%)	0.18	0.32	0.06	8.53
CV (%)	1.98	3.56	3.56	6.48

Means indicated by the same letters within the column have no statistically significant differences ($P \leq 0.05$).

Both the length and diameter of the fruit contribute to its shape. Over 53.85% of the varieties tested varieties at HARC had an ovate shape (Table 1). When a fruit's shape index value is less than one, it is more likely to have a round shape; when is more, it will have an ovate shape. Fruit size is one potential indicator of fruit maturity; however, it can also be influenced by crop load, climatic conditions, and cultural practices. Fruit shape and/or cheek fullness indicate maturity [31]. Stone fruits are considered mature when the fruits have well-developed shoulders and sutures. This criterion, however, must be combined with other indicators such as skin color to be reliable [31].

With regard to the average fruit weight, the variation. The88-18 W variety had the heaviest fruit weight of 111.98 g, while the 88-22C variety had the lowest (58.03 g). Apart from this, in several peach hybrids, fruit weight variations were observed [22]. Fruit weight is a significant quantitative inherited factor

influencing yield, fruit quality, and consumer acceptability [19]. Fruit weight varies greatly between varieties due to differences in tree production, and fruit numbers per tree [1].

3.2. Chemical quality traits

The result showed that there were statistically significant differences in all of the chemical quality traits among peach varieties (Table 2). About 46.15% of the varieties were recorded above the mean for all chemical quality traits except ascorbic acid which accounted for 30.77% of the varieties. The total soluble solids content (TSS) of these varieties ranged from 10.27 to 16.87 ^{o}Brix, with Transvalia receiving the highest (16.87 ^{o}Brix) and McRed the lowest (10.27 ^{o}Brix). Such variation might be attributed to the formation of different sugars such as sucrose, glucose, fructose, and sorbitol, and the rapid movement of these sugars from leaves to fruit, which are genetically controlled and regulated by source-sink relationships. However, the sugar content varied with genotype, seasonal conditions, yield, position of the fruits in the canopy, and maturity [8] similar observations in different peach varieties were reported by [38]. Soluble solids content below 10% are generally unacceptable to consumers [13]. However, this standard may vary from one country to another [23]. Similar to this, the variety Transvalia recorded the highest specific gravity (1.067), whereas the McRed variety recorded the lowest (1.041). However, the ripe peaches typically have a specific gravity of 0.99 [46]. In accordance with these results, [45] found a considerable variation in the specific gravity of ripe fruit among the different peach varieties.

Table 2
TSS, SG, TA, TSS/TA, Ascorbic acid, and pH of peach fruit varieties at HARC during the 2021 cropping season

Variety	Total soluble solid (°Brix)	Specific gravity (SG)	Titratable acidity (%)	Ripening index (TSS/TA)	Ascorbic acid (mg/100gm)	pH
McRed	10.27h	1.041h	0.80g	12.87def	9.11ef	3.41e
88 – 18 W	12.33ef	1.049ef	1.03cd	11.98efg	8.89f	3.42e
90-19H	11.40g	1.045g	0.94def	12.21efg	10.67c	3.44de
9A-35C	13.67c	1.055c	0.82fg	16.71a	10.22cd	3.57cd
Tropic Beauty	11.87fg	1.047fg	0.95de	12.46defg	9.78de	3.48de
88-22C	11.13g	1.045g	0.74g	15.11b	8.00f	3.49de
Early Grand	12.80de	1.051de	0.85efg	14.98bc	9.33ef	3.46de
Florida Down	14.73b	1.059b	1.21b	12.24efg	8.89f	3.91ab
Florida Star	14.67b	1.059b	1.09bc	13.49cde	8.00f	3.80b
May Crust	13.50cd	1.054cd	1.19b	11.42fg	9.33ef	4.03a
Summer Sun	15.43b	1.062b	1.41a	10.96g	16.67a	3.82b
Transvalia	16.87a	1.067a	1.17b	14.45bc	14.22b	3.82b
Bonny Gold	13.10cd	1.052cde	0.96de	13.84bcd	14.67b	3.66c
Mean	13.21	1.053	1.03	13.29	10.60	3.64
LSD (5%)	0.74	0.003	0.12	1.42	0.74	0.13
CV (%)	3.32	0.171	7.22	6.36	4.17	2.13

Means indicated by the same letters within the column have no statistically significant differences ($P \leq 0.05$).

With regard to the titratable acidity (TA), the varieties showed a significant variation with the range of 0.74 (88-22C) to 1.41% (Summer Sun). Titratable acidity in peaches is determined by several factors among which variety is the one [9]. Such TA variation mainly contributed from the formation of organic acids mainly malic, followed by citric, quinic, and succinic within the fruits which is mainly governed by genetics. The TA generally ranges from 0.07 to 0.14% in low-acid varieties to 1.1 to 1.45% in the acidic

ones [5] When the TA values are lower than 0.9%, it is considered as the maximum limit for low-acidity peaches [30]. On the other hand, consumer satisfaction is mainly attributed to fruit firmness, soluble solid content, and TA [15] which mainly vary with genotype, cultural practices, climatic conditions, and ripening stages [5].

The highest value of ripening index, which is directly associated with total soluble solid and titratable acidity, was obtained from variety 9A-35C (16.71) and the lowest was from summer sun (10.96). According to [44], the relationship between total soluble solids and total acidity is a very important parameter in determining fruit quality, because it provides information on the sugar/acids balance in the fruits. Naturally higher total soluble solids and lower titratable acidity of a variety in comparison with the other varieties, caused the highest TSS/TA ratio [28]. The ripening index is a major organoleptic quality trait of the mature fruit and is commonly used as a quality index in peaches [4]. The relationship between TA and TSS has an important role in consumer acceptance of some apricots, peaches, nectarine, and plum varieties. Consumer acceptance is controlled by the interaction between TA and TSS rather than TSS alone [15].

The maximum ascorbic acid value of 16.67 mg/100 gm was observed on variety Summer Sun and the minimum values of 8 mg/100 gm were obtained on both varieties 88-22C and Florida Star. This result in general revealed that there was a considerable difference in ascorbic acid content among peach varieties (Table 2). The synthesis of ascorbic acid content in fruit depends upon a sufficient supply of hexose Sugar, which declines at the ripening stage and might be due to a diminishing in acidity that would be attributed to the oxidation of ascorbic acid [33]. Comparably, [26] reported statistically significant variations in the vitamin C content of peach varieties. Similarly, [28] revealed that, significant differences in vitamin C among studied varieties.

Peach varieties grown at HARC had pH values ranging from 3.41 to 4.03. Amongst, the highest pH value of 4.03 was obtained from a variety of May crust and the lowest (3.41) was from Mc-red. This indicated that most of the varieties can more or less be grouped as high-acidity peaches. Peaches have a natural acidity of around 3–4 [12], and the studied varieties fall within this range. The report of [47] and [34], revealed that the pH of low-acidity peaches is greater than 4.0. Fruit pH is an important quality attribute for peaches and others in general.

3.3. Cluster analysis

Complete linkage hierarchical cluster analysis was used to measure the dissimilarity of observations using Euclidean distance measures. Hierarchical cluster analysis (HCA) of the thirteen peach varieties was performed using the ten physicochemical characteristics from the previous literature, as listed in Tables 1 and 2. The Euclidean method was used for the distance calculation for the cluster analysis. Thirteen peach varieties were grouped into four different clusters (Fig. 1) Clusters 1 and 2 were made from two varieties each. Whereas, clusters 3 and 4 consisted of six and three varieties respectively. In most of the chemical properties Florida Down and Florida Star which were in the same cluster were not

significantly different. However, the rest of the varieties in the same cluster had significantly different physicochemical properties.

3.4 Principal Component Analysis

Principal component analysis (PCA) was used to visualize the differences in physicochemical qualities among the thirteen peach varieties. The percentage of the cumulative contribution of variance of the two PCs was 63.1%; apart, PC1 presented 41.1% and PC2 presented 22.0% (Fig. 2). Fruit length, fruit weight, fruit width, pH, specific gravity, total soluble solid, titratable acidity, and ascorbic acid content were the main contributors to PC1 (positive side), Still, the ripening index and fruit shape index were on the negative side. On the other hand, the main contributors to PC2 (positive side) were fruit shape index, fruit length, fruit weight, and ascorbic acid content whilst ripening index, titratable acidity, total soluble solids, pH, specific gravity, and fruit width were the key contributors to the negative side.

Thus, it was demonstrated that the high ascorbic acid content and heavy and long fruit were all shared by the varieties May Crust, Summer Sun, and Transvalia. The physical and chemical composition of peach fruit varies significantly depending on the variety, as confirmed by the PCA analysis conducted for this study.

3.5 Correlation among physicochemical parameters

The correlation among physicochemical quality parameters of the evaluated peach varieties has been presented in Fig. 3. Total soluble solids and specific gravity showed a strongly positive correlation. While fruit shape index was highly negatively correlated with fruit width. However, pH has not correlated with fruit shape index and fruit weight. Generally, most of the parameters were correlated with each other even if the correlation coefficient varied considerably.

4. Conclusions

In general, peach fruit physicochemical quality characteristics were highly influenced by varieties. Hence, this study indicated that these varieties can be considered for various purposes and uses of interest according to their quality profile. However, other factors like pre-harvest management practices, environment, and climatic conditions may need to be assessed to understand their contribution to physicochemical quality characteristics.

Declarations

Author Contribution

Every author took part in the physical research activity, which included fieldwork and laboratory analysis. The corresponding author then authored the drafted manuscript following the data collection, and the other writers corrected it. The second author also designed the figures.

Data availability statements

The data that support the findings of this study are available from the corresponding author upon reasonable request. Data are located in controlled access data storage at the Ethiopian Agricultural Research Institute.

Acknowledgment

The authors extended their heartfelt gratitude to the ex-researchers who started the collection of the varieties, the field workers who managed the fruit trees, and the Ethiopian Institute of Agricultural Research for its financial support.

References

1. Abid W, Jiminez S, Moreno MA, and Gogorcena Y. Evaluation of antioxidant compounds and total sugar content in a nectarine [*Prunus persica* (L.) Batsch] progeny. International journal of molecular sciences, 2011. 12(10), pp.6919–6935.
2. Amiot MJ, and Lairon D. Fruit and vegetables, cardiovascular disease, diabetes and obesity. 2008. pp. 95–118. *In: F.A. Tomás-Barberán and M.I. Gil (eds.), Improving the Health-Promoting Properties of Fruit and Vegetable Products*. *Woodhead Publishing Limited*, *Cambridge, England*.
3. Arguin H, and Tremblay A. Fruit and vegetables, energy balance and weight management. 2008. pp. 182–200. *In: F.A. Tomás-Barberán and M.I. Gil (eds.), Improving the Health-Promoting Properties of Fruit and Vegetable Products*. *Woodhead Publishing Limited*, *Cambridge, England*.
4. Bassi D, and Selli R. Evaluation of fruit quality in peach and apricot. Advances in Horticulture Science 1990. 4: 107–112.
5. Bassi D, Mignani I, Spinardi A, and Tura D. Peach (Prunus persica (L.) Batsch). In *Nutritional composition of fruit cultivars*.2016. (pp. 535–571). Academic Press.
6. Bazzano LA. Epidemiologic evidence for the effect of fruit and vegetables on cardiovascular diseases, diabetes, and obesity. 2008. pp. 119–144. *In: F.A. Tomás-Barberán and M.I. Gil (eds.), Improving the Health-Promoting Properties of Fruit and Vegetable Products. Woodhead Publishing Limited, Cambridge, England*.
7. Bessey OA, and King C. The distribution of vitamin C in plant and animal tissues, and its determination, Journal of Biological Chemistry, vol. 103, 1933. pp. 687–698.
8. Brooks SJ, Moore JN, and Murphy JB. Quantitative and qualitative changes in sugar content of peach genotypes [*Prunus persica* (L.) Batsch.]. Journal of the American Society for Horticultural Science, *118*(1), 1993. pp.97–100.

9. Byrne D. Breeding peaches and nectarines for mild-winter climate areas: state of the art and future directions. *In Proceedings of the First Mediterranean Peach Symposium, Eds. F. Marra and F. Sottile, Agrigento, Italy*, 10 September 2003. pp. 102–109.
10. Cano-Salazar J, Lopez M, Crisosto C, Echeverri, G. Volatile compound emissions and sensory attributes of 'Big Top' nectarine and 'Early Rich' peach fruit in response to a pre-storage treatment before cold storage and subsequent shelf-life. Postharvest Biology and Technology 2013.76: 152–162.
11. Cantin C, Moreno MA, Gogorcena Y. Evaluation of the antioxidant capacity, phenolic compounds, and vitamin C content of different peach and nectarine [*Prunus persica* (L.) Batsch] breeding progenies. Journal of Agricultural and Food Chemistry 2009. 57: 4586–4592.
12. CFSAN (Center for Food Safety and Applied Nutrition). Approximate pH of Foods and Food Products Department of Health and Human Services. 2008.
13. Clareton M. Peach and nectarine production in France: Trends, consumption and perspectives. *In Prunus Breeders Meeting* 2000. (pp. 83–91).
14. Colin K. "Refractometers," 2017. Newsletter.
15. Crisosto C, Crisosto G, and Neri F. Understanding tree fruit quality based on consumer acceptance. *In: (Eds). Conference*. 2006. pp 183–190.
16. Crisosto CH, and Crisosto GM. Relationship between ripe soluble solids concentration (RSSC) and consumer acceptance of high and low acid melting flesh peach and nectarine [*Prunus persica* (L.) Batsch] cultivars. Postharvest Biology and Technology. 2005. 38, 239–246.
17. Crisosto CH. How do we increase peach consumption? Acta Hort.592: 601–605. 2002. https://doi.org/10.17660/ActaHortic.2002.592.82
18. Di Vaio C, Marallo N, Graziani G, Ritieni A, and Di Matteo A. Evaluation of fruit quality, bioactive compounds, and total antioxidant activity of flat peach cultivars. Journal of the Science of Food and Agriculture, 95 (10), 2124–2131. 2014. doi: 10.1002 / jsfa.6929
19. Dirlewanger E, Moing A, Rothan C, Svanella L, Pronier V, Guye A, Plomion C, Monet R. Mapping QTLs controlling fruit quality in peach [*Prunus persica* (L.) Batsch]. Theoretical and Applied Genetics, 1999. 98, 18–31.
20. Eduardo I, Pacheco I, Chietera G, Bassi D, Pozzi C, Vecchietti A, and Rossini L. QTL analysis of fruit quality traits in two peach intraspecific populations and importance of maturity date pleiotropic effect. Tree Genetics and Genomes; 2011. 7:323–35. https://doi.org/10.1007/s11295-010-0334-6
21. EIAR (Ethiopian Institute of Agricultural Research).2017. *Holeta Agricultural Research Center progress report*. Holeta, Ethiopia.
22. El-Morshedy FA, Salama M I, Hamdia M, Ayaad, and Abdelmonem AE. Evaluation of Peach Hybrids Using Some Pomological Characters and Rapd Markers. *Journal of Agricultural Research Kafr El-Sheikh University* 2016. pp: 554–570, Vol. 42(4).
23. Farina V, Lo Bianco R, and Mazzaglia A. Evaluation of late-maturing peach and nectarine fruit quality by chemical, physical, and sensory determinations. *Agriculture*, 9(9), 2019. p.189.

24. Franco B, and Anna V. Fruit and vegetables and cancer. pp. 75–94. In: F.A. Tomás-Barberán and M.I. Gil (eds.), Improving the Health-Promoting Properties of Fruit and Vegetable Products. *Woodhead Publishing Limited*, *Cambridge, England* 2008.
25. Garner D, Crisosto C, Wiley P, and Crisosto G. Measurement of pH and titratable acidity. Quality evaluation methodology. USA: Kearney Agricultural Center2005.
26. Gecer MK. Biochemical content in fruits of peach and nectarine cultivars. Turkish Journal of Agriculture and Forestry, 44(5), 2020. PP.500–505.
27. Goulao LF, Oliveira CM. Cell wall modifications during fruit ripening: when the fruit is not the fruit, Trends Food Science and Technology 2008. 19: 4–25.
28. Hajilou J, and Fakhimrezaei S. Evaluation of fruit physicochemical properties in some peach cultivars. Research in Plant Biology, *1*(5), 2011. PP.16–21.
29. Hancock JF. Strawberries. CABI, Wallingford, 1999.P 77.
30. Hilaire C. The peach industry in France: state of art, research, and development. *In: (Eds). Conference*. 2003. pp 27–34.
31. Kader AA, Mitchell FG. Postharvest physiology. *In: La Rue, J.H.; Johnson, R.S.; (eds) Peaches, Plums and Nectarines: Growing and Handling for Fresh Market. University of California Department of Agriculture and Natural Resources* 1989. *pp*. 158–164.
32. Kelley KM, Primrose R, Crassweller R, Hayes JE, Marini R. Consumer peach preferences and purchasing behavior: A mixed methods study. Journal of Science of Food and Agriculture, 96(7), 2451–2461. 2016. https://doi.org/10.1002/jsfa.7365.
33. Kumari M, Karanjeev K, Sanjay S, and Feza A. Assessment of Chemical Characteristics of Low Chilling Peach Cultivars under Agro-climatic Conditions of Bihar. International Journal of Current Microbiology and Applied Sciences 2020. ISSN: 2319–7706 (9)2.
34. Moing A, Svanella L, Monet R, Rothan C, Diakou P, Gaudillere JP, and Rolin D. Organic acid metabolism during the fruit development of two peach cultivars. Acta Horticulturae, 1998. 465, 425–432.
35. Prior RL, and Cao GH. Antioxidant phytochemicals in fruits and vegetables: diet and health implications. HortScience 2000; 35:588–592.
36. Remorini D, Tavarini S, DeglInnocenti E, Loreti F, Massai R, Guidi L. Effect of rootstocks and harvesting time on the nutritional quality of peel and flesh of peach fruits. Food Chemistry 2008.110: 361–367.
37. SAS, 2017. *Proc. 24th International Symposium*. New York, USA.
38. Singh G, Sharma KK, and Jawanda JS. Physico-chemical characteristics of some peach cultivars under Ludhiana conditions. The Punjab Horticultural Journal, 1984. 24(1–4): 92–95.
39. Singh M. and Ramassamy C. Beneficial effects of phenolic compounds from fruit and vegetables in neurodegenerative diseases. 2008. pp. 145–181. *In: F.A. Tomás-Barberán and M.I. Gil (eds.),*

Improving the Health-Promoting Properties of Fruit and Vegetable Products. Woodhead Publishing Limited, Cambridge, England.

40. Singh O, Kumar A, Rai R, and Kohli K. Quality evaluation of low chill peach cultivars for preparation of ready-to-serve 'Nectar'drink. Asian Journal of Dairy and Food Research, *35*(4), 2016. Pp.327–330.
41. Toralles RP, Vendruscolo JL, Tondovendruscolo FBD Pino, PL Autunes. Determination of reaction rate constants for ascorbic acid degradation in the peach effect of temperature and concentration 2008. 28(1): 18–23.
42. UPOV (International Union for the protection of new varieties of plants). 2012. Pomegranate. Guidelines for the conduct of tests for distinctness, uniformity and stability.
43. Veerappan K, Natarajan S, Chung H and Park J. Molecular Insights of Fruit Quality Traits in Peaches, Review. MDPI, Plants 10, 2191. 2021. https://doi.org/10.3390/plants10102191
44. Voca S, Dobricevic N, Dragovic-Uzelac V, Duralija B, Druzic J. Fruit quality of new early ripening strawberry cultivars in Croatia. Food Technology and Biotechnology, 2008. 46:292–298.
45. Wen IC, Koch KE, and Sherman WB. Comparing fruit and tree characteristics of two peaches and their nectarine mutants. Journal of the American Society for Horticultural Science, 120(1), 1995. Pp.101–106.
46. Westwood MN. Seasonal changes in specific gravity and shape of apple, pear, and peach fruits. *Proceedings of the American Society for Horticultural Science*, 1962. 80:90–96.
47. Yoshida M. Genetical studies on the fruit quality of peach varieties. 1. Acidity. Bulletin of the Tree Research Station Series A; 1970.9, 1–15.
48. Zohrabi S, Seiiedlou S, and Alipasandi A. Study some physical and mechanical properties of three cultivars of peach in maturation stages. World of Sciences Journal, 4, 2013. Pp.108–117.

Figures




Figure 1

Complete hierarchical cluster analysis based on similarity matrix using Euclidean method of thirteen peach varieties




Figure 2

PCA biplot map illustrating the relationship among the physicochemical properties of thirteen peach fruit varieties; Note: 1=88-18 W, 2=88-22C, 19-90H, 4=9A-35C, 5=Bonny Gold, 6=Early Grand, 7=Florida Down, 8=Florida Star, 9=May Crust, 10=McRed, 11=Summer Sun, 12=Transvalia, 13=Tropic Beauty




Figure 3

Correlation among physicochemical properties