



Draft Genome Sequence of *Enterobacter asburiae* NCR1, a Plant Growth-Promoting Rhizobacterium Isolated from a Cadmium-Contaminated Environment

 Charlotte J. François,^a Steven Batinovic,^b Steve Petrovski,^b  Anthony R. Gendall^a

^aDepartment of Animal, Plant and Soil Sciences, La Trobe University, Bundoora, Victoria, Australia

^bDepartment of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, Victoria, Australia

ABSTRACT *Enterobacter asburiae* NCR1 is a plant growth-promoting rhizobacterium isolated from the rhizosphere of *Carpobrotus rossii*. We report the draft genome sequence of *E. asburiae* strain NCR1, which revealed many genes facilitating beneficial interactions with plant hosts.

The genus *Enterobacter* comprises more than 20 species that are widely distributed through various environments. Some species of *Enterobacter* can enhance plant growth by increasing the availability of nutrients and water and synthesizing plant growth-promoting hormones (1, 2). *Enterobacter asburiae* NCR1 was previously isolated from the rhizosphere of *Carpobrotus rossii* grown in a cadmium-contaminated environment (3).

E. asburiae NCR1 was grown at 27°C for 16 h in Luria-Bertani broth, and total genomic DNA was extracted using the Bioline Isolate II genomic DNA kit and converted to sequencing libraries using the Nextera XT DNA library prep kit (Illumina) with 1 ng of input DNA, according to the manufacturer's instructions. The quality and concentration of the genomic DNA were assessed using an Invitrogen Qubit 3.0 fluorometer. The libraries were sequenced on an Illumina MiSeq instrument using a MiSeq v.3 reagent kit (600 cycles), which generated 919,623 reads of 300-bp paired-end reads. The sequence reads were trimmed to a Phred threshold quality score of 20 using Trimmomatic v.0.36.6 (4), and assembly was performed using SPAdes v.3.12.0 (5), both on Galaxy Australia (6). The total length of the genome sequence of *E. asburiae* NCR1 was 4,546,475 bp, with a G+C content of 56.08%. The final number of contigs was 15, with an N_{50} value of 2,597,842 bp, and the final genome coverage was 50×. Default parameters were used unless otherwise stated.

The annotation of the assembled genomic sequence was performed using the Prokaryotic Genome Annotation Pipeline (PGAP) v.4.10 of the National Center for Biotechnology Information (NCBI) (7). The genome sequence of *E. asburiae* NCR1 was closely related to that of *E. asburiae* GN04222 with an average nucleotide identity (ANI) of 98.06%, as calculated by a JSpeciesWS v.3.8.2 analysis (8). A total of 4,237 genes were identified, with 4,240 coding sequences (CDS), including 87 RNA genes. The RNA genes comprised 4 5S rRNAs, 1 partial 16S rRNA, 1 partial 23S rRNA, and 73 tRNA genes.

The annotated genome sequence revealed several genes pertinent to plant-microbe interactions, including those involved in auxin synthesis (*iaa* and *ipdC*) and phosphorus metabolism (*phoP*). Bacterially derived auxin can enhance the growth of plants, while the catabolism of phosphates may improve phosphate availability to the host plants (9).

Heavy metal resistance to Zn, Cd, and Pb is conferred through *zntA*, which acts as a P-type ATPase pump (10) and is suggestive of this strain's ability to survive in challenging environments. This draft genome sequence of *E. asburiae* NCR1 could provide

Citation François CJ, Batinovic S, Petrovski S, Gendall AR. 2021. Draft genome sequence of *Enterobacter asburiae* NCR1, a plant growth-promoting rhizobacterium isolated from a cadmium-contaminated environment. Microbiol Resour Announc 10:e00478-21. <https://doi.org/10.1128/MRA.00478-21>.

Editor Vincent Bruno, University of Maryland School of Medicine

Copyright © 2021 François et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Charlotte J. François, charlotte.francois@latrobe.edu.au.

Received 13 May 2021

Accepted 10 August 2021

Published 2 September 2021

valuable information regarding the plant growth-promotive properties and heavy metal resistance and tolerance of this strain.

Data availability. The *E. asburiae* NCR1 whole-genome sequencing project has been deposited at GenBank under accession number [JAAAJX000000000](#), BioProject accession number [PRJNA600757](#), and BioSample accession number [SAMN13831070](#). The raw sequences were deposited in the Sequence Read Archive under accession number [SRR14460571](#).

REFERENCES

1. Ramesh A, Sharma SK, Sharma MP, Yadav N, Joshi OP. 2014. Plant growth-promoting traits in *Enterobacter cloacae* subsp. *dissolvens* MDSR9 isolated from soybean rhizosphere and its impact on growth and nutrition of soybean and wheat upon inoculation. *Agric Res* 3:53–66. <https://doi.org/10.1007/s40003-014-0100-3>.
2. Jha CK, Aeron A, Patel BV, Maheshwari DK, Saraf M. 2011. Enterobacter: role in plant growth promotion, p 159–182. In Maheshwari DK (ed), *Bacteria in agrobiolgy: plant growth responses*. Springer, Berlin, Germany. https://doi.org/10.1007/978-3-642-20332-9_8.
3. Liu W, Wang Q, Wang B, Hou J, Luo Y, Tang C, Franks AE. 2015. Plant growth-promoting rhizobacteria enhance the growth and Cd uptake of *Sedum plumbizincicola* in a Cd-contaminated soil. *J Soils Sediments* 15: 1191–1199. <https://doi.org/10.1007/s11368-015-1067-9>.
4. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Afgan E, Sloggett C, Goonasekera N, Makunin I, Benson D, Crowe M, Gladman S, Kowsar Y, Pheasant M, Horst R, Lonie A. 2015. Genomics virtual laboratory: a practical bioinformatics workbench for the cloud. *PLoS One* 10:e0140829. <https://doi.org/10.1371/journal.pone.0140829>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
8. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
9. Spaepen S, Vanderleyden J. 2011. Auxin and plant-microbe interactions. *Cold Spring Harb Perspect Biol* 3:a001438. <https://doi.org/10.1101/cshperspect.a001438>.
10. Naz N, Young HK, Ahmed N, Gadd GM. 2005. Cadmium accumulation and DNA homology with metal resistance genes in sulfate-reducing bacteria. *Appl Environ Microbiol* 71:4610–4618. <https://doi.org/10.1128/AEM.71.8.4610-4618.2005>.