

Melatonin alleviates acid-induced stress in peanut (*Arachis hypogaea* L.) through physiological, antioxidative regulation, and molecular defense mechanisms

Muhammad Hafeez Ullah Khan¹, Ruoyi Fu¹, Ali Muhammad², Shuaichao Zheng¹, Daijing Zhang², Zhiyong Zhang^{1*}, Quanyong Liu^{1*}

¹Xinxiang Key Laboratory of Crop Root Biology and Green Efficient Production/Henan Key Laboratory of Quality and Stress-Resilient Bio-breeding and Safe Production of Cotton and Oil Crops, School of Life Sciences, Henan Institute of Science and Technology, Xinxiang, Henan, 453000, China

²College of Life Sciences, Henan Normal University, Xinxiang 453007, Henan, China.

Correspondence Author:

Zhiyong Zhang*

(Email: z_zy123@126.com)

Muhammad Hafeez Ullah Khan; hafeezkhan_88@yahoo.com, Ruoyi Fu; fu_ruoyi0123@163.com, Ali Muhammad; alikhaksar87@gmail.com, Shuaichao Zheng; yndxzsc@126.com, Daijing Zhang; zdjdai@163.com, Zhiyong Zhang; z_zy123@126.com, Quanyong Liu; liuquanyong@hist.edu.cn.

35 Abstract

36 Acidic stress severely restricts crop growth by disrupting nutrient uptake, redox
 37 homeostasis, and membrane stability, yet mitigation strategies remain limited. Here,
 38 we investigated the role of melatonin (MT) in modulating growth, photosynthesis,
 39 oxidative stress, antioxidant defense, and proton transport in peanut seedlings grown
 40 under acidic (pH 4.0) and near-optimal (pH 6.5) conditions. Acid stress markedly
 41 reduced biomass accumulation, chlorophyll content, and redox balance, while
 42 enhancing ROS (H₂O₂) and lipid peroxidation (MDA). Exogenous MT application,
 43 particularly at 50-100 μM, significantly improved shoot and root fresh and dry
 44 weights, restored chlorophyll a, b, total chlorophyll, and carotenoids, and reduced
 45 H₂O₂ and MDA accumulation, with more pronounced effects under pH 4.0 than pH
 46 6.5. MT strongly activated antioxidant enzymes (SOD, CAT, APX), while POD
 47 activity declined, reflecting melatonin's dual role as both a direct ROS scavenger and
 48 a regulator of enzymatic redox networks. Notably, MT induced strong, dose-
 49 dependent upregulation of H⁺-ATPase genes (AH1 and AH2) in both leaves and roots
 50 under acidic conditions, suggesting enhanced proton extrusion, intracellular pH
 51 homeostasis, and stress adaptation. Although MT also conferred benefits at pH 6.5,
 52 responses were generally moderate compared with acid stress. Collectively, these
 53 results demonstrate that MT confers robust acid stress tolerance in peanut by
 54 integrating growth regulation, antioxidant defense, and proton homeostasis. This
 55 study provides the first integrated evidence that melatonin enhances acid stress
 56 tolerance by coordinating antioxidant defense with H⁺-ATPase-mediated proton
 57 homeostasis, revealing a novel mechanism for crop adaptation to low-pH soils.

58 **Keywords:** Melatonin supplementation, Peanut seedling resilience, Acidic soil
 59 adaptation, ROS-pH coordination, Low-pH crop physiology

60

61

62

63

64

65

66

67

68 Introduction

69 Soil acidification has become an escalating global concern, now affecting nearly 30–
70 40% of the world's arable lands and posing a serious threat to agricultural
71 productivity and food security ([Huang et al., 2023](#)). Soils with a pH below 5.5 are
72 considered acidic, and such conditions limit the availability of essential
73 macronutrients like phosphorus (P), calcium (Ca^{2+}), and magnesium (Mg^{2+}). At the
74 same time, they increase the solubility of toxic metal ions, especially aluminium
75 (Al^{3+}) and manganese (Mn^{2+}), which can be highly detrimental to plant growth
76 ([Delfim et al., 2025](#)). This imbalance in nutrient and ion dynamics often results in
77 inhibited root elongation, reduced nutrient uptake, and disrupted photosynthetic
78 processes. Furthermore, an excess of hydrogen ions (H^+) in the rhizosphere disturbs
79 pH stability and alters microbial community structure, compounding physiological
80 stress in plants ([Naz et al., 2022](#); [Thapa et al., 2025](#)).

81 Peanut (*Arachis hypogaea* L.), a major legume widely cultivated in tropical
82 and subtropical regions, is particularly vulnerable to acid soil stress ([Li et al., 2024b](#);
83 [Jamir et al., 2025](#)). During early seedling stages, low pH and other abiotic stress
84 conditions restrict root development, damage cell membranes, and interfere with ion
85 transport mechanisms, ultimately impairing water and nutrient absorption. These
86 physiological disruptions often lead to reduced chlorophyll levels, decreased
87 photosynthetic efficiency, and poor biomass accumulation ([Liu et al., 2018](#);
88 [Borhannuddin Bhuyan et al., 2019](#); [Li et al., 2021](#)). On a cellular scale, acid stress
89 intensifies the generation of reactive oxygen species (ROS), including superoxide
90 anions ($\text{O}_2^{\cdot-}$), hydrogen peroxide (H_2O_2), and hydroxyl radicals ($\cdot\text{OH}$), which can
91 damage lipids, proteins, and nucleic acids ([Ahmad et al., 2023b](#); [Hasan et al., 2025](#)).
92 Although plants possess both enzymatic antioxidants (such as SOD, POD, APX, and
93 CAT) and non-enzymatic compounds (like ascorbate, glutathione, and flavonoids).
94 These defense mechanisms are often inadequate to neutralize the rapid oxidative surge
95 induced by acidic environments, particularly in sensitive crops like peanut.

96 In recent years, melatonin (N-acetyl-5-methoxytryptamine) has been identified
97 as an effective plant growth regulator and stress alleviator ([Zheng et al., 2025](#);
98 [Muhammad et al., 2026](#)). Melatonin was first found in animals, and is now widely
99 recognized in plants to modulate plant growth, development and defence responses
100 ([Ahmad et al., 2023a](#)). Exogenous melatonin has been reported to alleviate various

abiotic stresses such as drought, salinity, acid, extreme temperature, nutrient deficiency and heavy metal toxicity by enhancing photosynthetic efficiency, maintaining redox balance and elevating antioxidant capacity (Basharat *et al.*, 2025; Dou *et al.*, 2025; Kong *et al.*, 2025). Melatonin is also involved in alleviating aluminium toxicity under suppressed soil pH, maintaining the integrity of cellular membrane, stabilizing chloroplast structure and activating the antioxidant system (Wang *et al.*, 2025a). In addition to its direct ROS-scavenging ability, melatonin indirectly regulates redox homeostasis by modulating the expression of redox-responsive genes (Hasan *et al.*, 2025; Jiang *et al.*, 2025a).

Proton pumps, particularly plasma membrane and vacuolar H⁺-ATPases, are fundamental molecular systems that enable plants to sustain cytoplasmic pH homeostasis under an acidic environment (Cosse & Seidel, 2021; Zeng *et al.*, 2024; MacDonald *et al.*, 2025). These ATPases actively transport protons from the cytoplasm into the apoplast or vacuole, thereby establishing electrochemical gradients that drive essential physiological processes such as nutrient uptake, ion transport, stomatal regulation and stress adaptation (MacDonald *et al.*, 2025; Sun *et al.*, 2025). In peanut, various ATPase genes have been characterized, among which AH1 and AH2 are key regulators of intracellular pH balance and ion homeostasis under acid stress. The upregulation of H⁺-ATPase genes has been reported in several crops as an adaptive mechanism to low pH, facilitating proton extrusion and protecting cellular metabolism (Li *et al.*, 2022; Li *et al.*, 2024a). However, the regulatory dynamics of these genes under melatonin treatment remain largely unexplored in the peanut seedling.

Melatonin has been shown in other crop species to influence proton pump activity and ion transport. For instance, melatonin-induced activation of H⁺-ATPases can enhance nutrient absorption, stabilize membrane potential, and promote root development under stress conditions (Huang *et al.*, 2022; Tahjib-Ul-Arif *et al.*, 2025). Therefore, integrating physiological, biochemical and transcriptional analysis, particularly through expression profiling of AH1 and AH2, can provide valuable mechanistic insight into how melatonin supports pH regulation and overall homeostasis in peanut seedlings under acidic conditions.

The present study aims to elucidate the integrated physiological, antioxidative, and molecular mechanisms underlying melatonin-mediated acid tolerance in peanut seedlings, with a particular focus on AH1 and AH2 gene expression. We demonstrate

that melatonin enhances peanut tolerance to acid stress by improving growth and photosynthetic traits, strengthening antioxidant defense, and upregulating key H⁺-ATPase genes to stabilize cellular pH and ion homeostasis. Unravelling these mechanisms will contribute to the development of sustainable agronomic strategies and genetic improvement of peanut for acid soil environments.

Materials and methods

Plant material and experimental design

The experiment was conducted in the greenhouse (25±2°C, 60-70% RH, 16/8 h light/dark, 200 µmol m⁻² s⁻¹) at the Henan Institute of Science and Technology, Xinxiang City, Henan Province, China, under controlled environmental conditions. A paper roll culture system was used in a hydroponic setup to evaluate the effects of melatonin on peanut growth under acidic stress conditions. The experiment followed a randomized complete design (RCD) with eight biological replications to ensure statistical reliability.

Uniform, mature, and healthy seeds of peanut (*Arachis hypogaea* L. cv. Xinbailua 25) were selected as experimental material. Seeds were surface-sterilized with 1% sodium hypochlorite solution for 5 min and rinsed several times with distilled water to remove any residual disinfectant. To investigate the effects of melatonin on plant growth, photosynthetic pigments, antioxidant responses, and gene expression, the seeds were soaked for 10 hours in melatonin solutions of 0.5 µM, 5 µM, 50 µM, and 100 µM. Seeds soaked in distilled water under identical conditions served as the control (CK).

For each replicate, seven seeds were placed vertically on moistened germination paper to ensure uniform orientation and root emergence. The papers were then carefully rolled and secured with rubber bands to maintain contact and prevent unrolling during incubation. Following treatment, the paper rolls containing the seeds were placed vertically in plastic containers filled with half-strength Hoagland nutrient solution prepared according to the modified composition described by Cai et al. (2021) (Cai et al., 2021). Two nutrient pH regimes were maintained: a normal condition with pH 6.5 and an acidic condition with pH 4.0. The pH of each solution was carefully adjusted using 1 M HCl or NaOH and monitored daily to ensure consistent stability throughout the experiment. The lower ends of the paper rolls were immersed approximately 2-3 cm into the solution to facilitate the upward capillary movement of

nutrients and water while maintaining adequate aeration around the root zone. To prevent nutrient depletion and pH fluctuations, the nutrient solutions were replaced every two days. During the early germination stage, the paper rolls were kept under dark conditions until radicle emergence to ensure uniform germination. After emergence, seedlings were exposed to normal light conditions for continued growth. After 15 days of cultivation, seedlings were harvested for measurements of growth parameters, photosynthetic pigments, antioxidant enzyme activities, and gene expression analyses.

Plant Germination and Growth Parameters

Seed germination was monitored daily until radicle emergence, which was defined as a visible protrusion of at least 2 mm from the seed coat, and the germination percentage (GP) was calculated at day 7. After 15 days of cultivation, seedlings were carefully removed from the paper rolls and gently rinsed with distilled water to remove residual nutrient solution. Shoot length was measured from the base to the tip of the longest leaf, and root length from the base to the tip of the longest root, using a ruler to the nearest millimeter. Fresh weights of shoots and roots were determined separately after blotting with filter paper to remove surface moisture, and samples were then oven-dried at 70 °C for 72 h or until a constant weight was achieved to determine dry weights. All measurements were performed for each seedling in every replicate, and the mean values of the eight replicates were used for statistical analysis. Data are presented as mean $\pm$ standard error (SE) and used for subsequent physiological, biochemical, and molecular analyses.

Photosynthetic Pigment Measurement

Photosynthetic pigments, including chlorophyll a, chlorophyll b, and carotenoids, were extracted from 0.2 g of fresh peanut leaves using 80% (v/v) acetone. The leaf samples were homogenized in chilled acetone, kept on ice for 10-15 min in the dark, and centrifuged at $10,000 \times g$ for 10 min at 4 °C. The clear supernatant was collected, and absorbance was recorded at 663, 645, and 470 nm using a UV–visible spectrophotometer with 80% acetone as blank. Pigment contents were calculated using standard equations and expressed as mg g⁻¹ fresh weight. All extractions were performed under low-light conditions to minimize pigment degradation, and six independent replicates were analyzed per treatment.

Antioxidant Enzyme Assays

To assess the antioxidant defense response in peanut seedlings, the activities of catalase (CAT), superoxide dismutase (SOD), peroxidase (POD), and ascorbate peroxidase (APX), content, were determined from fresh leaf tissues. Approximately 0.5 g of fresh leaves was homogenized in 5 mL of ice-cold 50 mM phosphate buffer (pH 7.0) containing 1 mM EDTA and 1% (w/v) polyvinylpyrrolidone (PVP). The homogenate was centrifuged at $12,000 \times g$ for 15 min at 4 °C, and the resulting supernatant was used for all enzyme assays.

Superoxide dismutase (SOD; EC 1.15.1.1) activity was assayed by the inhibition of nitroblue tetrazolium (NBT) reduction under light, with one unit of SOD activity defined as the amount of enzyme required to cause 50% inhibition of NBT photoreduction. Catalase (CAT; EC 1.11.1.6) activity was determined by monitoring the decrease in absorbance at 240 nm due to H_2O_2 decomposition. Peroxidase (POD; EC 1.11.1.7) activity was measured using guaiacol as a substrate, recording the increase in absorbance at 470 nm as guaiacol was oxidized. Ascorbate peroxidase (APX; EC 1.11.1.11) activity was estimated by monitoring the decrease in absorbance at 290 nm as ascorbate was oxidized in the presence of H_2O_2 .

Quantification of Malondialdehyde (MDA) and Hydrogen Peroxide (H_2O_2)

The levels of malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) were determined to evaluate lipid peroxidation and oxidative stress in peanut leaves and roots. Lipid peroxidation was determined by quantifying malondialdehyde (MDA) using the thiobarbituric acid (TBA) reaction. For MDA estimation, 1 mL of enzyme extract was mixed with 2 mL of 0.6% (w/v) TBA in 10% (w/v) trichloroacetic acid (TCA) and heated at 95 °C for 15 min, then rapidly cooled and centrifuged at $10,000 \times g$ for 10 min. The absorbance of the supernatant was measured at 532 nm and corrected for nonspecific turbidity by subtracting absorbance at 600 nm. MDA content was calculated using the extinction coefficient ($\epsilon = 155 \text{ mM}^{-1} \text{ cm}^{-1}$) and expressed as nmol g^{-1} fresh weight (FW).

Hydrogen peroxide (H_2O_2) content was determined spectrophotometrically using a colorimetric KI reaction with quantification against a standard curve. Fresh leaf tissue (0.5 g) was ground in 5 mL of cold 0.1% (w/v) TCA and centrifuged at $12,000 \times g$ for 15 min at 4 °C. A reaction mixture was prepared by mixing 0.5 mL of the extract with 0.5 mL of 10 mM potassium phosphate buffer (pH 7.0) and 1.0 mL of 1 M potassium iodide (KI) (all kept on ice and protected from light). The reaction was allowed to

proceed for 1 min at room temperature in the dark, and the absorbance was read at 390 nm. A standard calibration curve was prepared in parallel using known concentrations of H₂O₂ (for example 0, 10, 25, 50, 100, 200 μM) prepared in 0.1% TCA; sample H₂O₂ content was interpolated from this curve and expressed as μmol H₂O₂ g⁻¹ FW (adjusting for extract and assay volumes and any dilutions). All reactions and standards were run in technical triplicate and blank reactions (reagents without plant extract) were included to correct baseline absorbance.

Gene Expression Analysis

Total RNA was extracted from peanut roots and leaves using the RNeasy Pure Plant Plus Kit (Qiagen Biotech, Beijing, China) according to the manufacturer's instructions. The quality and quantity of RNA were determined using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, USA) by measuring the absorbance ratios at 260/280 nm and 260/230 nm. RNA integrity was confirmed by 1.0% agarose gel electrophoresis, ensuring that samples displayed sharp and intact 28S and 18S rRNA bands without visible degradation. Only high-quality RNA (A260/A280 ratio between 1.9 and 2.1) was used for cDNA synthesis.

First-strand cDNA was synthesized from 50 μg of total RNA using the PrimeScript™ RT reagent kit with gDNA Eraser (Takara, Japan) to eliminate potential genomic DNA contamination. The synthesized cDNA was diluted 10-fold with nuclease-free water and stored at -20 °C until further analysis.

Quantitative real-time PCR (qRT-PCR) was conducted using TB Green® Premix Ex Taq™ II (Takara, Japan) on a QuantStudio™ 5 Real-Time PCR System (Applied Biosystems, USA). Each 20 μL reaction contained 10 μL of 2× SYBR Green master mix, 0.4 μL each of forward and reverse primers (10 μM), 2 μL of diluted cDNA, and 7.2 μL of nuclease-free water. The PCR cycling conditions were as follows: initial denaturation at 95 °C for 30 s, followed by 40 amplification cycles of 95 °C for 5 s and 60 °C for 30 s. A melting curve analysis (60-95 °C) was performed at the end of each run to confirm the specificity of amplification.

Each sample was analyzed in triplicate, and the mean values of three technical replicates from three independent biological replicates were used for statistical analysis. The relative transcript abundance of target genes was calculated using the 2^{-ΔΔCt} method (Livak and Schmittgen, 2001). The AhGAPDH gene was used as an

internal reference due to its stable expression across different tissues and treatments.

Primer sequences used in the study are listed in Supplementary Table 1.

Statistical Analysis

All data were analyzed using SPSS v26.0. Results are presented as mean $\pm$ SE. Two-way ANOVA was used to compare treatments, and significant differences were determined using Tukey's test ($p < 0.05$). For qRT-PCR, the average Ct values of technical replicates were used, and fold changes were calculated using the $2^{-\Delta\Delta Ct}$ method.

Table 1. Primer sequences used for qRT-PCR analysis of plasma membrane H⁺-ATPase genes in peanut.

Gene Name	Primer Sequence
AH1 H ⁺ - ATPase 1-F	GCCTCTGACATAGTTCTAAC
AH1 H ⁺ - ATPase 1-R	TAAAGCCAACCACAATACGG
AH2 H ⁺ - ATPase 2-F	GAAAAGAATGGGCGTGGTTG
AH2 H ⁺ - ATPase 2-R	TCAAGGGACACAATCTCACC

The table lists forward (F) and reverse (R) primer sequences designed for the amplification of *AhAH1* and *AhAH2* genes.

Results

Effect of melatonin on peanut seed germination under acid condition

Seed germination of peanut was significantly affected by pH and melatonin treatment (Fig. 1a and b). At acidic pH (4), control seeds exhibited the lowest germination (~70%). Application of MT markedly improved germination in a dose-dependent manner: 0.5 μ M MT increased germination to ~88% (+18%), 5 μ M MT to ~90% (+20%), 50 μ M MT to ~93% (+23%), and 100 μ M MT to ~98% (+28%) with related to control, indicating a strong protective effect of MT under acid stress. Under near-neutral conditions (pH 6.5), germination was consistently high across treatments, with control seeds reaching ~85% and MT-treated seeds ranging from 90% to nearly 100%. The relative improvement of germination by MT was more pronounced under acidic stress, indicating that MT alleviates pH-induced inhibition. These results demonstrate that melatonin significantly mitigates the inhibitory effects of low pH on peanut seed germination, with the largest relative benefit observed at 100 μ M MT under acidic

conditions, and that moderate MT concentrations are sufficient to maintain near-maximal germination at near-neutral pH.

Melatonin supplementation enhances peanut plant growth and biomass accumulation under acid stress

To evaluate the role of melatonin in modulating peanut seedling growth under contrasting pH conditions, root and shoot development as well as biomass accumulation were assessed under acidic (pH 4) and near-neutral (pH 6.5) environments (Figs. 1c–h). Acidic stress (pH 4) markedly suppressed seedling growth, as evidenced by substantial reductions in root and shoot elongation in control plants, indicating strong inhibition of early developmental processes. Melatonin treatment markedly alleviated this inhibition, resulting in a dose-dependent enhancement of root length, shoot length, and biomass. Root length under pH 4 increased progressively from approximately 6 cm in control plants to nearly 20 cm at 100 μ M MT, while shoot length increased from about 2.5 cm in the control to over 6 cm at the highest MT concentration, demonstrating a strong stimulatory effect of MT on seedling elongation under acid stress (Fig. 1c and d). Under neutral conditions (pH 6.5), root and shoot lengths were generally higher than those at pH 4 in control plants; however, the growth-promoting effect of MT was less pronounced. Low MT concentration (0.5 μ M) enhanced root and shoot length, while higher MT doses showed either moderate improvement or stabilization of growth, suggesting that MT exerts a stronger relative effect under stress conditions than under optimal pH (Fig. 1c and d).

Consistent with growth responses, root fresh weight and shoot fresh weight under pH 4 increased substantially with increasing MT concentration. Root fresh weight rose from 1.09 g in control plants to 3.45 g at 100 μ M MT, while shoot fresh weight increased from 1.09 g to 2.86 g, indicating enhanced biomass accumulation (Fig. 1e and f). This trend was further supported by dry weight measurements, where root dry weight increased from 0.07 g in control plants to 0.17 g at 100 μ M MT, and shoot dry weight increased from 0.06 g to 0.15 g, confirming that MT-induced growth was associated with true biomass gain rather than water accumulation alone (Fig. 1g and h).

Under pH 6.5, overall growth and biomass parameters were higher than those observed under acidic conditions in the control plants; however, the response pattern

326 to MT differed from that at pH 4. Low MT concentration (0.5 μ M) promoted root and
 327 shoot growth and fresh weight accumulation, whereas higher MT concentrations (5-
 328 100 μ M) resulted in a gradual decline in root and shoot length and biomass (Fig. 1e-
 329 h). Notably, root dry weight and shoot dry weight at pH 6.5 showed little or no
 330 improvement with increasing MT concentration and tended to decrease at higher
 331 doses, suggesting a concentration-dependent shift in MT responsiveness under non-
 332 stress conditions.

333 Overall, these results demonstrate that melatonin strongly enhances root and
 334 shoot growth and biomass accumulation under acidic stress (pH 4), with maximal
 335 effects observed at higher MT concentrations, whereas under near-neutral conditions
 336 (pH 6.5), low MT levels are sufficient to promote growth and excessive MT may exert
 337 inhibitory effects. This indicates that the growth-promoting role of melatonin is highly
 338 dependent on environmental pH and stress intensity.




Figure 1. Effect of melatonin (MT) on seed germination and seedling growth under normal (pH 6.5) and acidic (pH 4.0) conditions on peanut. (a) seed germination and seedling growth phenotypes, (b) seed germination (%), (c) shoot length (cm), (d) root length (cm), (e) shoot fresh weight (g), (f) root fresh weight (g), (g) root dry weight (g), (h) shoot dry weight. Statistical differences among treatments were analyzed using appropriate multiple comparison tests. ns indicates not significant, while *, **, *, and **** denote significant**

differences at $P < 0.05$, $P < 0.01$, $P < 0.001$, and $P < 0.0001$, respectively. Data in the figures are means of eight replications ($n = 8$) $\pm$ standard deviation (SD). Bar = 2 cm.


Melatonin modulates photosynthetic pigments in response to acid stress in peanut

Melatonin application significantly influenced the accumulation of photosynthetic pigments in peanut seedlings under both acidic (pH 4) and near-neutral (pH 6.5) conditions (Figs. 2a-d). Under pH 4, control plants exhibited relatively low levels of chlorophyll a, chlorophyll b, total chlorophyll, and carotenoids, indicating impaired pigment biosynthesis under acid stress. However, MT treatments progressively enhanced pigment accumulation, with a clear dose-dependent response.

Chlorophyll a content under pH 4 increased gradually with increasing MT concentration, showing moderate enhancement at 5 μ M MT and a pronounced increase at 50 and 100 μ M MT, where values reached their maximum (Fig. 2a). A similar trend was observed for chlorophyll b, resulting in a substantial rise in total chlorophyll content, particularly at 50 and 100 μ M MT. These increases were significantly higher than the control ($P < 0.05$), demonstrating that MT effectively mitigates acid-induced inhibition of chlorophyll synthesis (Fig. 2b and c). Under pH 6.5, pigment contents were generally higher than those under pH 4 in control plants, reflecting more favorable growth conditions. Notably, low MT concentration (0.5 μ M) caused a marked increase in chlorophyll a, chlorophyll b, and total chlorophyll, reaching peak values under this pH. However, higher MT concentrations (50-100 μ M) resulted in a gradual decline in pigment levels, suggesting that excessive MT under optimal pH may suppress chlorophyll accumulation (Figs. 2a-c).

Carotenoid content showed a distinct but complementary pattern. Under pH 4, carotenoid levels increased steadily with rising MT concentration, reaching the highest values at 100 μ M MT, which were significantly greater than the control ($P < 0.05$). This indicates enhanced antioxidant capacity under acid stress (Fig. 2d). In contrast, under pH 6.5, carotenoid content peaked sharply at 0.5 μ M MT and declined at higher concentrations, paralleling the response observed for chlorophyll pigments (Fig. 2d). Overall, these results demonstrate that melatonin enhances photosynthetic pigment accumulation in peanut seedlings in a pH-dependent and concentration-dependent manner. Low MT concentrations are optimal for pigment enhancement under near-neutral conditions, whereas higher MT doses are more effective in

379 protecting pigment biosynthesis under acidic stress, likely through improved stress
380 tolerance and antioxidant regulation.



381 **Figure 2. Effect of melatonin (MT) on photosynthetic pigments of peanut under normal**
382 **(pH 6.5) and acidic (pH 4.0) conditions. (a) Chlorophyll a (mg/g FW), (b) chlorophyll b**
383 **(mg/g FW), (c) total chlorophyll content (mg/g FW), (d) carotenoids (mg/g FW).** Statistical
384 differences among treatments were analyzed using appropriate multiple comparison tests. ns
385 indicates not significant, while ***, and **** denote significant differences at $P < 0.001$, and
386 $P < 0.0001$, respectively. Data in the figures are means of eight replications ($n = 8$) $\pm$ standard
387 deviation (SD).

388 **Melatonin differentially regulates antioxidant defense system in response** 389 **to acid stress in peanut**

390 We evaluated the enzymatic antioxidant activity in acid-stressed peanut plants treated
391 with melatonin to understand how exogenous melatonin mitigates acid stress. Our
392 findings revealed that the radical scavenging activities of SOD, POD, CAT, and APX,
393 were markedly affected by both external pH and melatonin treatment in peanut leaves,
394 reflecting dynamic regulation of reactive oxygen species (ROS) homeostasis under
395 stress and non-stress conditions (Figs. 3a-h).

396
397

Superoxide Dismutase (SOD) Activity

SOD activity showed a strong induction under acidic conditions (pH 4) compared with pH 6.5 across all treatments, indicating enhanced superoxide radical generation under acid stress. In control plants, SOD activity at pH 4 was substantially higher (60.79 U mg⁻¹ FW) than at pH 6.5 (18.70 U mg⁻¹ FW), reflecting a basal stress response. Application of melatonin significantly increased SOD activity in a dose-dependent manner, with 100 µM MT reaching 137.1 U mg⁻¹ FW, representing a ~128% increase over control, suggesting that MT promotes superoxide detoxification capacity under severe acidic stress (Fig. 3a). In contrast, under pH 6.5, SOD activity was relatively low and showed only moderate induction by MT, with a peak at 0.5 µM MT followed by a decline at higher concentrations (Fig. 3a). This differential response indicates that melatonin enhances ROS-scavenging systems primarily under stress conditions, while under near-optimal pH it avoids excessive activation of antioxidant machinery, thereby maintaining redox balance.

In roots, SOD activity exhibited a similar but more pronounced response. At pH 4, melatonin increased SOD activity from 54.34 U mg⁻¹ FW (control) up to 119.39 U mg⁻¹ FW at 100 µM MT (~126% increase). This strong induction indicates that MT substantially enhances superoxide detoxification in roots exposed to acidic stress (Fig. 3b). At pH 6.5, the increase was smaller but still significant (18.70 U mg⁻¹ FW control to 42.29 U mg⁻¹ FW at 0.5 µM MT, ~126% increase), however, MT application still increased SOD relative to control, though to a lesser extent than under acidic conditions (Fig. 3b). These results suggest that MT-mediated regulation of SOD is stress-responsive and tissue-specific, with a stronger effect in roots facing direct exposure to low-pH environments.

Peroxidase (POD) Activity

POD activity exhibited an opposite trend compared with SOD, particularly under acidic conditions. At pH 4, control plants showed the highest POD activity (175.53 U mg⁻¹ FW), reflecting stress-induced activation of cell wall-associated and vacuolar peroxidases, which are often involved in stress signaling, lignification, and ROS-mediated defense responses. Melatonin application led to a progressive decrease in POD activity, with 100 µM MT reducing activity to 29.24 U mg⁻¹ FW (~83% reduction compared to control). This decline suggests that melatonin alleviates oxidative stress, thereby reducing the requirement for POD-mediated hydrogen

peroxide consumption (Fig. 3c). Importantly, this reduction highlights the dual role of melatonin: while it enhances ROS scavenging through SOD, CAT, and APX, it simultaneously suppresses excessive POD activity that is often associated with oxidative damage, rigidification of cell walls, and growth inhibition. Under pH 6.5, POD activity was generally lower than at pH 4 and displayed a distinct pattern, with moderate increases at higher MT concentrations (Fig. 3c). This suggests that under non-stress conditions, POD may participate more in fine-tuning redox signaling rather than bulk ROS detoxification.

In roots, POD activity displayed a similar pattern. At pH 4, POD activity also decreased from 168.59 U mg⁻¹ FW in control to 35.63 U mg⁻¹ FW at 100 μM MT (~79% reduction), mirroring the trend observed in leaves and supporting the role of melatonin in alleviating oxidative stress by reducing H₂O₂ accumulation (Fig. 3d). However, at pH 6.5, POD activity increased progressively with melatonin concentration, reaching near-control or higher levels at 100 μM MT (Fig. 3d). This contrasting behavior highlights the dual role of melatonin: under stress, it suppresses excessive ROS-scavenging enzyme activity by lowering ROS production, whereas under non-stress conditions, it maintains or enhances POD activity to support redox homeostasis and signaling.

Catalase (CAT) Activity

CAT activity was strongly stimulated by melatonin under both pH conditions, though the magnitude and pattern of induction differed. Under acidic pH, CAT activity increased from 139.4 U mg⁻¹ FW in control to 273.2 U mg⁻¹ FW at 100 μM MT (~96% increase over control) (Fig. 3e). This indicates that melatonin enhances the rapid decomposition of hydrogen peroxide into water and oxygen, a crucial mechanism for preventing oxidative damage under acid-induced stress. At pH 6.5, CAT activity was generally higher than control and showed a pronounced induction at low to moderate MT concentrations, with the highest activity observed at 0.5 μM MT, followed by a gradual decline at higher doses (Fig. 3e). This suggests that under near-neutral pH, lower MT concentrations are sufficient to optimize hydrogen peroxide homeostasis, while excessive MT may reduce the need for CAT due to improved redox equilibrium.

In roots, CAT activity similarly increased under acidic stress, from 167.6 U mg⁻¹ FW (control) to 331.6 U mg⁻¹ FW at 100 μM MT (~98% increase) (Fig. 3f).

At near-neutral pH, control activity was 147.9 U mg⁻¹ FW, and melatonin slightly increased CAT at low concentrations (0.5-5 μM MT) but decreased at higher doses (100 μM MT, 147.2 U mg⁻¹ FW) (Fig. 3f). These results indicate that melatonin effectively boosts CAT-mediated H₂O₂ scavenging, especially under acid stress, enhancing antioxidant defense in both leaves and roots.

Ascorbate Peroxidase (APX) Activity

APX activity increased consistently with melatonin concentration under pH 4, indicating a strong MT-mediated activation of the ascorbate-glutathione cycle under acidic stress. At pH 4, control leaves had 28.8 U mg⁻¹ FW, which rose to 66.9 U mg⁻¹ FW at 100 μM MT (~132% increase) (Fig. 3g). The highest APX activity at 100 μM MT suggests that APX plays a central role in fine-scale hydrogen peroxide detoxification in chloroplasts and cytosol when plants experience acid-induced oxidative pressure. Under pH 6.5, APX activity showed a different pattern, increased from 26.8 U mg⁻¹ FW (control) to 62.9 U mg⁻¹ FW at 0.5 μM MT, while higher MT concentrations showed slightly reduced activity (~34.7–51.8 U mg⁻¹ FW), indicating a dose-dependent modulation (Fig. 3g). This response implies that under optimal conditions, melatonin primarily enhances APX activity at low doses to maintain signaling-compatible ROS levels, while higher concentrations may downregulate APX as oxidative stress is already minimal.

APX activity in roots increased significantly with MT under acidic conditions, from 18.3 U mg⁻¹ FW in control to 63.3 U mg⁻¹ FW at 100 μM MT (Fig. 3h). At pH 6.5, APX activity peaked at moderate MT doses (66.2 U mg⁻¹ FW at 0.5 μM), while higher MT doses slightly reduced activity (Fig. 3h), showing that APX is finely regulated by both MT concentration and pH to balance ROS detoxification with normal metabolism.

Collectively, these results demonstrate that melatonin orchestrates a coordinated antioxidant response in peanut leaves by enhancing ROS-scavenging enzymes (SOD, CAT, APX) while downregulating stress-associated POD activity, particularly under acidic conditions. This dual regulatory role allows melatonin to mitigate oxidative damage while preserving ROS signaling necessary for growth and adaptation. The contrasting responses observed between pH 4 and pH 6.5 further highlight that melatonin acts as a stress-dependent modulator of redox homeostasis

rather than a constitutive antioxidant, enabling peanut plants to adapt efficiently to acidic stress environments.




Figure 3. Effect of melatonin (MT) on antioxidant contents of peanut under normal (pH 6.5) and acidic (pH 4.0) conditions. (a) leaf SOD (U/g FW), (b) leaf POD ((U/g FW), (c) leaf CAT (U/g FW), (d) leaf APX (U/g FW), (e) root SOD (U/g FW), (f) root POD ((U/g FW), (f) root CAT (U/g FW), (g) root APX (U/g FW). Statistical differences among treatments were analyzed using appropriate multiple comparison tests. ns indicates not significant, while **,

, and * denote significant differences at $P < 0.01$, $P < 0.001$, and $P < 0.0001$, respectively. Data in the figures are means of eight replications ($n = 8$) $\pm$ standard deviation (SD).

Melatonin reduces hydrogen peroxide and lipid peroxidation contents in peanut under acidic condition

Melatonin is widely recognized for its role in mitigating oxidative stress across various biological systems, including plants. We conducted a comprehensive study to assess how melatonin influences peanut seedling growth under acidic condition. Our findings revealed that acid-treated plants exhibited significantly elevated levels of H_2O_2 and MDA compared to MT treated plants (Figs. 4a-d)

Hydrogen Peroxide (H_2O_2) Content

Under control conditions (CK), H_2O_2 levels were high at pH 4 (79.40 $\mu\text{mol g}^{-1}$ FW) and pH 6.5 (71.06 $\mu\text{mol g}^{-1}$ FW), indicating substantial oxidative stress, particularly under acidic conditions. Melatonin application significantly reduced H_2O_2 accumulation at pH 4 in a dose-dependent manner. Compared with the control, 0.5 μM MT reduced H_2O_2 to 57.95 $\mu\text{mol g}^{-1}$ FW ($\approx 28\%$ decrease), 5 μM MT to 49.28 $\mu\text{mol g}^{-1}$ FW ($\approx 38\%$ decrease), 50 μM MT to 46.31 $\mu\text{mol g}^{-1}$ FW ($\approx 43\%$ decrease), and 100 μM MT to 39.99 $\mu\text{mol g}^{-1}$ FW ($\approx 50\%$ decrease) (Fig. 4a). In contrast, at pH 6.5, low MT concentrations initially decreased H_2O_2 , with 0.5 μM MT reducing levels to 44.38 $\mu\text{mol g}^{-1}$ FW ($\approx 36\%$ decrease) and 5 μM MT to 53.03 $\mu\text{mol g}^{-1}$ FW ($\approx 26\%$ decrease). However, higher MT doses led to a reversal of this trend, with H_2O_2 increasing to 62.34 $\mu\text{mol g}^{-1}$ FW at 50 μM MT ($\approx 11\%$ decrease) and 68.71 $\mu\text{mol g}^{-1}$ FW at 100 μM MT (nearly comparable to control) (Fig. 4a). This suggests a dose and pH-dependent dual role of melatonin, acting as a strong antioxidant under acidic stress but allowing moderate ROS signaling at optimal pH.

Root tissues showed a similar but more pronounced response. Under pH 4, control plants accumulated 85.04 $\mu\text{mol g}^{-1}$ FW H_2O_2 . Melatonin significantly lowered H_2O_2 to 64.25 $\mu\text{mol g}^{-1}$ FW at 0.5 μM ($\approx 24\%$ reduction), 58.54 $\mu\text{mol g}^{-1}$ FW at 5 μM ($\approx 32\%$ reduction), 48.76 $\mu\text{mol g}^{-1}$ FW at 50 μM ($\approx 44\%$ reduction), and 43.81 $\mu\text{mol g}^{-1}$ FW at 100 μM MT ($\approx 51\%$ reduction) (Fig. 4b). At pH 6.5, H_2O_2 content in roots declined at low MT doses (44.25 $\mu\text{mol g}^{-1}$ FW at 0.5 μM ; $\approx 46\%$ decrease) but increased at higher concentrations, reaching 74.78 $\mu\text{mol g}^{-1}$ FW at 100 μM MT ($\approx 9\%$ decrease compared with control) (Fig. 4b). These results indicate that

melatonin efficiently scavenges excessive ROS under acid stress, while at neutral pH high MT may permit ROS accumulation for signaling functions.

Malondialdehyde (MDA) Content

MDA content, an indicator of lipid peroxidation, was markedly higher under acidic conditions. In leaves at pH 4, control plants exhibited 12.35 nmol g⁻¹ FW MDA. Melatonin treatments progressively reduced MDA levels to 8.29 nmol g⁻¹ FW at 0.5 μM (≈32% reduction), 6.38 nmol g⁻¹ FW at 5 μM (≈48% reduction), 4.94 nmol g⁻¹ FW at 50 μM (≈60% reduction), and 3.89 nmol g⁻¹ FW at 100 μM MT (≈69% reduction) (Fig. 4c). At pH 6.5, control MDA content was lower (9.0 nmol g⁻¹ FW). Low MT concentrations further decreased MDA to 3.92 nmol g⁻¹ FW at 0.5 μM (≈56% reduction) and 5.52 nmol g⁻¹ FW at 5 μM (≈39% reduction). However, higher MT doses increased MDA to 6.66 nmol g⁻¹ FW at 50 μM (≈26% reduction) and 7.83 nmol g⁻¹ FW at 100 μM MT (≈12% reduction) (Fig. 4c), again reflecting melatonin's concentration-dependent dual behavior.

Root MDA followed trends similar to leaves but with overall lower values. At pH 4, control roots had 11.16 nmol g⁻¹ FW MDA. This decreased to 9.04 nmol g⁻¹ FW at 0.5 μM (≈19% reduction), 7.15 nmol g⁻¹ FW at 5 μM (≈35% reduction), 5.93 nmol g⁻¹ FW at 50 μM (≈46% reduction), and 3.65 nmol g⁻¹ FW at 100 μM MT (≈68% reduction) (Fig. 4d). At pH 6.5, MDA declined sharply at low MT doses (3.83 nmol g⁻¹ FW at 0.5 μM; ≈64% reduction) but increased again at higher concentrations, reaching 8.25 nmol g⁻¹ FW at 100 μM MT (≈20% reduction compared with control) (Fig. 4d). This indicates that melatonin effectively protects membrane integrity under acidic stress, while excessive doses at optimal pH may attenuate this protective effect.

Collectively, these results demonstrate that melatonin markedly alleviates oxidative stress and lipid peroxidation in peanut leaves and roots, particularly under acidic (pH 4) conditions. The strong reduction in H₂O₂ and MDA at pH 4 indicates that melatonin primarily functions as an antioxidant under stress. In contrast, under pH 6.5, low melatonin concentrations enhance oxidative protection, whereas higher doses partially restore ROS and lipid peroxidation levels, suggesting a regulated balance between ROS scavenging and signaling. This dual role of melatonin

highlights its importance in fine-tuning redox homeostasis in peanut plants under contrasting pH environments.




Figure 4. Effect of melatonin (MT) on H₂O₂ and MDA contents of peanut under normal (pH 6.5) and acidic (pH 4.0) conditions. (a) leaf H₂O₂ (μmol/g FW), (b) leaf MDA (nmol/g FW), (c) root H₂O₂ (μmol/g FW), (d) root MDA (nmol/g FW). Statistical differences among treatments were analyzed using appropriate multiple comparison tests. ns indicates not significant, while *, **, and **** denote significant differences at P < 0.05, P < 0.01, and P < 0.0001, respectively. Data in the figures are means of eight replications (n = 8) ± standard deviation (SD).

Melatonin induces dose-dependent upregulation of H⁺-ATPase genes under acid stress

To investigate the transcriptional response of plasma membrane H⁺-ATPase genes to melatonin (MT) under acidic conditions, the relative expression levels of *AhAH1* and *AhAH2* were analyzed in peanut leaves and roots under acid stress (Figs. 5a and b). Both genes are central to proton extrusion and pH homeostasis during environmental stress.


Melatonin treatment significantly enhanced the expression of plasma membrane H⁺-ATPase genes *AhAH1* and *AhAH2* in peanut seedlings, with a pronounced induction under acidic conditions. The basal expression of *AhAH1* was

low in control plants, with minimal transcript accumulation observed in both leaves and roots. MT supplementation resulted in a consistent and progressive increase in *AhAH1* expression at all concentrations tested. At 0.5 and 5 μ M MT, *AhAH1* transcript levels increased by approximately 2.4, and 2.6-fold in leaves and 1.5, and 2.5-fold in roots relative to the control (Fig. 5a). This induction intensified with increasing MT concentration, reaching ~3.7-fold in leaves and ~3.2-fold in roots at 50 μ M MT. The strongest response was observed at 100 μ M MT, where *AhAH1* expression increased by ~7.0-fold in leaves and ~5.1-fold in roots compared with control plants, indicating robust, dose-dependent transcriptional activation (Fig. 5a). The stronger induction in roots highlights the pivotal role of root-localized H⁺-ATPase activity in coping with acidic stress.

A similar positive response was observed for *AhAH2*, which exhibited higher basal expression than *AhAH1* in control plants, particularly in roots. Nevertheless, MT treatment further enhanced *AhAH2* expression at all concentrations under acid stress. At low MT concentrations (0.5 and 5 μ M), *AhAH2* transcript abundance increased by approximately 1.7- and 3.1-fold in leaves and 2.6- and 4.5-fold in roots, respectively, relative to the control (Fig. 5b). This upregulation became progressively stronger with increasing MT dosage, reaching approximately 5.1-fold in leaves and 8.2-fold in roots at 50 μ M MT. The maximum induction occurred at 100 μ M MT, where *AhAH2* expression was elevated by approximately 6.6-fold in leaves and 11.6-fold in roots compared with control plants (Fig. 5b), demonstrating a robust and concentration-dependent transcriptional response. Importantly, All MT treatments significantly increased *AhAH2* transcript levels relative to the control, with the highest induction observed at 100 μ M MT. This consistent upregulation under acidic stress suggests that *AhAH2* is positively responsive to melatonin and likely contributes to enhanced proton extrusion and pH homeostasis. The pronounced upregulation of *AhAH2* at higher melatonin concentrations suggests that this gene encodes a stress-adaptive H⁺-ATPase that remains transcriptionally active under strong melatonin stimulation, thereby contributing to the maintenance of cytosolic pH homeostasis and sustained root function under acidic stress conditions.

Overall, these results demonstrate that melatonin induces a consistent, concentration-dependent increase in H⁺-ATPase gene expression at all MT treatments relative to the control, with the strongest fold induction occurring under acidic conditions. This transcriptional enhancement provides molecular evidence for

623 melatonin-mediated reinforcement of proton transport capacity, supporting improved
624 acid tolerance in peanut seedlings.



625 **Figure 5. Melatonin-induced expression of plasma membrane H⁺-ATPase genes in**
626 **peanut leaves and roots.** (a) Relative expression of *AhAH1* and (b) relative expression of
627 *AhAH2* in leaves and roots under different melatonin (MT) concentrations (0.5, 5, 50, and 100
628 μM) compared with the control. Bars represent mean ± SD of biological replicates. Statistical
629 differences among treatments were analyzed using appropriate multiple comparison tests. ns
630 indicates not significant, while *, **, ***, and **** denote significant differences at P < 0.05,
631 P < 0.01, P < 0.001, and P < 0.0001, respectively.

632 Discussion

633 Melatonin alleviates acid stress-induced growth inhibition by promoting 634 seed germination and seedling growth in peanut

635 Acidic stress conditions severely constrain seed germination and early seedling
636 establishment by disrupting metabolic activation, reserve mobilization, and cell
637 expansion (Msimbira & Smith, 2020; Hasan et al., 2025). Seed germination and early
638 seedling establishment are among the most pH-sensitive developmental stages in
639 crops, and acidic stress is known to impair metabolic activation, reserve mobilization,
640 and cell expansion (Shavrukov & Hirai, 2016; Ali et al., 2023; Wang et al., 2025b). In
641 this study, peanut seeds exposed to acidic pH (4.0) showed a clear reduction in
642 germination percentage and seedling growth compared with near-neutral pH (6.5),
643 confirming the high pH sensitivity of early developmental stages. Exogenous
644 melatonin (MT) markedly alleviated this inhibition, significantly improving
645 germination and promoting vigorous seedling growth, with a much stronger relative
646 effect under acidic stress than under near-neutral conditions (Figs.1a-h). The enhanced
647 germination under acidic pH likely reflects improved metabolic activation during

imbibition, as MT is known to stimulate hydrolytic enzymes such as α -amylases and proteases, thereby facilitating efficient mobilization of stored reserves under stress (Liu *et al.*, 2023; Fu *et al.*, 2025). In addition, MT interacts with key phytohormones, including gibberellins and auxins, which regulate radicle protrusion and early seedling vigor, further supporting rapid germination under low-pH conditions (Arnao & Hernández-Ruiz, 2019; Fu *et al.*, 2025). Consistent with improved germination, MT substantially enhanced root and shoot growth and biomass accumulation under acidic conditions, indicating improved seedling establishment (Figs.1a-h). Similar MT-mediated promotion of early growth under unfavourable soil conditions has been reported in other crops, where MT improved root development and stress resilience by maintaining cellular homeostasis and nutrient acquisition (Muhammad *et al.*, 2024; Chung *et al.*, 2025; Hasan *et al.*, 2025). In contrast, under near-neutral pH, low MT doses were sufficient to slightly enhance seedling performance, whereas higher concentrations often reduced growth and physiological indices, highlighting the role of MT as a stress-responsive regulator rather than a constitutive growth promoter. This inhibitory effect likely results from the disruption of finely tuned redox and hormonal signaling, as melatonin also functions as a regulatory molecule and excessive levels can suppress seedling growth under non-stress conditions (Arnao & Hernández-Ruiz, 2019; Ameen *et al.*, 2024; Hasan *et al.*, 2025). Collectively, these results demonstrate that melatonin effectively mitigates acid-induced inhibition of peanut seed germination and early seedling growth, underscoring its potential for improving crop establishment in acidic soils.

Melatonin enhances photosynthetic pigment stability and efficiency under acid stress in peanut

Acidic stress markedly impaired photosynthetic pigment accumulation in peanut seedlings, as evidenced by reduced chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents under pH 4, indicating disrupted chloroplast function and pigment biosynthesis (Figs. 2a-d). Such reductions are well documented consequences of low pH-induced proton toxicity, metal ion imbalance, and oxidative damage, which collectively destabilize chlorophyll synthesis and accelerate pigment degradation during early vegetative growth (Sharma *et al.*, 2020; Yang *et al.*, 2023; Jiang *et al.*, 2025b). Our results showed that exogenous melatonin application substantially alleviated this inhibition, resulting in a dose-dependent restoration of chlorophyll and

carotenoid levels under acidic conditions, with maximal pigment accumulation observed at higher MT concentrations (Figs. 2a-d). This response suggests that melatonin preserves pigment integrity under acid stress, likely by protecting chloroplast membranes and photosynthetic complexes from oxidative injury through its strong antioxidant and redox-buffering capacity (Dou et al., 2025; Hasan et al., 2025). Carotenoid accumulation followed a similar pattern, increasing progressively with MT concentration under pH 4, suggesting enhanced photoprotection and reactive oxygen species scavenging (Figs. 2a-d). The associated increase in carotenoid content under acidic conditions further supports a protective role of melatonin, as carotenoids are essential for quenching excess excitation energy and preventing photo-oxidative damage under stress. Similar MT-induced stabilization of photosynthetic pigments has been reported in crops exposed to salinity, drought, and metal toxicity, where melatonin maintained chlorophyll biosynthesis and delayed stress-induced senescence (An et al., 2025; Basharat et al., 2025; Chen et al., 2025; Hasan et al., 2025; Li et al., 2025d). In contrast, under near-neutral pH (6.5), pigment levels were inherently high in control plants, and low MT concentrations were sufficient to enhance chlorophyll accumulation, whereas higher doses led to pigment stabilization or decline. This concentration-dependent response indicates that melatonin primarily functions as a stress-responsive regulator, exerting its strongest effects when pigment homeostasis is challenged rather than under optimal growth conditions (Ali et al., 2023; Yang et al., 2023). Overall, these results demonstrate that melatonin enhances photosynthetic pigment stability in peanut in a pH and dose-dependent manner, with higher MT concentrations effectively counteracting acid-induced pigment loss. This protective role of melatonin on chlorophyll and carotenoid metabolism likely contributes to improved photosynthetic competence and stress resilience of peanut seedlings under acidic environments, a condition that remains underexplored in crop systems.

Melatonin-regulated antioxidant redox homeostasis confers acid stress tolerance in peanut

Acidic stress markedly disrupts redox homeostasis in plants by enhancing reactive oxygen species (ROS) production and impairing antioxidant efficiency, thereby constraining early growth and physiological stability (Hlihor et al., 2022; Jahan et al., 2022; Hasan et al., 2025). In the present study, peanut seedlings exposed to acidic pH (4) exhibited strong oxidative pressure, as evidenced by differentially modulated basal

activities of stress-responsive antioxidant enzymes, such as SOD, POD, CAT and APX. Exogenous MT application significantly reprogrammed the antioxidant defense system, resulting in a coordinated and stress-adaptive response. Under acidic conditions, MT markedly enhanced SOD, CAT, and APX activities in both leaves and roots, indicating improved superoxide dismutation and hydrogen peroxide detoxification, while simultaneously suppressing excessive POD activity (Figs. 3a-h). The marked reduction in POD activity observed in melatonin-treated peanut leaves and roots under acidic conditions likely reflects melatonin-mediated alleviation of oxidative stress rather than suppression of antioxidant capacity. Exogenous melatonin significantly reduced POD activity in a dose-dependent manner, indicating that MT effectively lowers ROS production and H₂O₂ availability, thereby reducing the requirement for POD-dependent scavenging (Figs. 3b and c). It has been reported that melatonin has been shown to modulate antioxidant enzyme activities and reduce ROS accumulation under various stress conditions, contributing to redox homeostasis (meta-analysis in rice and other plants) (Yang *et al.*, 2021; Muhammad *et al.*, 2025). Moreover, in other systems melatonin reduced ROS accumulation while lowering peroxidase activity, supporting the interpretation that decreased POD activity can accompany improved oxidative balance (Yang *et al.*, 2021; Hasan *et al.*, 2025). Thus, the reduced POD activity in MT-treated peanut seedlings reflects a functional shift from stress-induced peroxidase activation to a more energy-efficient antioxidant strategy, highlighting melatonin's dual role as both a ROS scavenger and a redox regulator under acidic stress (Figs. 3b and c). This pattern suggests that MT not only strengthens ROS-scavenging capacity but also limits stress-associated peroxidative reactions linked to cell wall rigidification and growth inhibition. Similar MT-mediated enhancement of enzymatic antioxidants and redox balance under acidic or oxidative stress has been reported in many crops, where MT improved stress tolerance by activating ROS-detoxifying enzymes and their associated genes (Ahmed *et al.*, 2020; Ahmed *et al.*, 2024; Chakraborty & Raychaudhuri, 2025; Chung *et al.*, 2025; Li *et al.*, 2025c). Notably, the contrasting responses observed at pH 6.5 further support the stress-dependent role of melatonin. Under near-neutral conditions, MT induced only moderate or dose-limited changes in antioxidant activity, indicating that MT fine-tunes redox signaling rather than overstimulating defense pathways when oxidative stress is minimal (Figs. 3a-h). This dual behavior, strong activation under stress and restrained modulation under optimal conditions, has been recognized as a

hallmark of melatonin function in plants (Moustafa-Farag *et al.*, 2020; Menhas *et al.*, 2025; Shehzadi *et al.*, 2025). Collectively, these findings demonstrate that melatonin enhances acid-stress tolerance in peanut by orchestrating a balanced antioxidant response, promoting efficient ROS detoxification while preserving redox signaling, thereby enabling physiological stability and growth under low-pH environments.

Melatonin fine-tunes redox homeostasis by limiting hydrogen peroxide accumulation and membrane lipid peroxidation under acidic stress

The pronounced accumulation of hydrogen peroxide (H₂O₂) and malondialdehyde (MDA) in peanut seedlings under acidic conditions confirms that low pH imposes severe oxidative stress, leading to excessive ROS generation and membrane lipid peroxidation (Hlihor *et al.*, 2022; Hasan *et al.*, 2025). Under pH 4, control plants exhibited markedly elevated H₂O₂ and MDA levels in both leaves and roots, reflecting impaired redox balance and enhanced oxidative damage. Exogenous melatonin significantly and dose-dependently reduced H₂O₂ accumulation under acidic conditions, with the highest MT concentration lowering H₂O₂ by ~50% in leaves and ~51% in roots relative to the control (Figs. 4a and b). This strong suppression of ROS indicates that melatonin acts primarily as an efficient antioxidant under acid stress, either through direct free-radical scavenging or by reinforcing enzymatic detoxification pathways, thereby reducing cellular H₂O₂ availability (Arnao & Hernández-Ruiz, 2019; Ahammed *et al.*, 2020; Ahammed *et al.*, 2024; Hasan *et al.*, 2025; Muhammad *et al.*, 2025).

Consistent with reduced ROS levels, MDA content declined sharply in MT-treated seedlings under pH 4, with up to ~69% and ~68% reductions in leaves and roots, respectively, indicating effective protection of membrane integrity (Figs. 4c and d). The tight coupling between decreased H₂O₂ and lower lipid peroxidation suggests that melatonin limits oxidative damage by stabilizing cellular membranes and preventing ROS-induced fatty acid oxidation, a response widely associated with improved stress tolerance (Ali *et al.*, 2023; Ahammed *et al.*, 2024; Hasan *et al.*, 2025). In contrast, under near-neutral pH (6.5), low MT concentrations enhanced oxidative protection, whereas higher doses partially restored H₂O₂ and MDA levels (Figs. 4a-d), supporting the concept that melatonin permits controlled ROS accumulation for signaling under non-stress conditions (Hasan *et al.*, 2025; Li *et al.*, 2025a). Collectively, these results demonstrate that melatonin functions as a redox modulator in peanut, acting

predominantly as an antioxidant under acidic stress while maintaining ROS-dependent signaling at optimal pH, thereby ensuring balanced cellular homeostasis.

Melatonin enhances plasma membrane H⁺-ATPase transcription to maintain pH homeostasis under acidic stress

Plasma membrane H⁺-ATPases are key regulators of proton extrusion, cytosolic pH stability, and nutrient uptake, and their function becomes especially critical under acidic stress, where excessive proton influx disrupts cellular metabolism and root growth (Chao & Chao, 2025; Dong *et al.*, 2025). In the present study, melatonin (MT) significantly and dose-dependently upregulated the expression of the H⁺-ATPase genes *AhAH1* and *AhAH2* in peanut seedlings under acidic conditions, with markedly stronger induction in roots than in leaves. This transcriptional response provides molecular evidence that MT reinforces proton transport to the apoplast to improve acid tolerance (Figs. 5a and b).

The strong MT-induced expression of *AhAH1* and *AhAH2* is consistent with previous reports showing that melatonin enhances H⁺-ATPase activity and expression under various abiotic stresses (Wang *et al.*, 2025c). For example, MT has been shown to stimulate plasma membrane H⁺-ATPases in cucumber and tomato under salinity and nutrient stress, thereby stabilizing intracellular pH and improving ion homeostasis (Li *et al.*, 2022; Zeng *et al.*, 2024; Li *et al.*, 2025b). These findings align closely with our observation that *AhAH1* and *AhAH2* expression was most strongly induced at higher melatonin concentrations under pH 4, highlighting a conserved role of melatonin in activating proton transport machinery under acidic stress (Figs. 5a and b). Under acidic or aluminium-stress conditions, enhanced H⁺-ATPase activity promotes proton efflux and mitigates rhizotoxic effects, supporting root elongation and function (Zhang *et al.*, 2017; Munyaneza *et al.*, 2024). The robust upregulation of *AhAH2* at higher melatonin doses indicates that this isoform may function as a stress-adaptive proton pump, supporting sustained root activity, membrane potential stability, and nutrient acquisition under acidic conditions. The pronounced induction of *AhAH2*, particularly in roots, suggests isoform-specific stress responsiveness, in line with studies in *Arabidopsis* and rice, where certain H⁺-ATPase genes are preferentially activated under low-pH stress to maintain root

apoplasmic and cytosolic pH balance (Gómez-Arjona *et al.*, 2022; Chakraborty *et al.*, 2024; Jain & Schmidt, 2024).

Mechanistically, melatonin-mediated upregulation of H⁺-ATPase genes likely reflects its integrative role in redox and hormonal signaling. Melatonin modulates ROS levels and interacts with auxin and calcium-dependent pathways, both of which are known regulators of H⁺-ATPase transcription and activation (Huang *et al.*, 2022; Yang *et al.*, 2024; Chung *et al.*, 2025). By alleviating oxidative stress while maintaining controlled ROS signaling, MT may create favorable cellular conditions for sustained proton pump expression and activity under acidic environments. Overall, these findings demonstrate that melatonin enhances acid stress tolerance in peanut by transcriptionally activating plasma membrane H⁺-ATPase genes, particularly *AhAH2* in roots. This response likely strengthens proton extrusion, stabilizes intracellular pH, and supports root performance under low-pH conditions, highlighting H⁺-ATPase regulation as a key molecular component of melatonin-mediated adaptation to acidic stress in crops.

Interaction between melatonin concentration and pH regulates peanut physiological responses


Across all measured parameters, including germination, seedling growth, biomass accumulation, photosynthetic pigments, oxidative stress markers, antioxidant enzyme activities, and H⁺-ATPase gene expression, a clear interaction between melatonin concentration and pH level was observed. Acidic conditions (pH 4) consistently amplified the positive effects of melatonin, whereas near-neutral pH (6.5) markedly reduced or altered MT responsiveness. This interaction indicates that melatonin does not act as a simple growth promoter, but rather functions as a stress-dependent regulator whose efficacy is tightly linked to environmental pH (Fig. 6).

Under acidic stress, increasing MT concentrations progressively improved germination, root and shoot development, pigment accumulation, redox homeostasis, and proton pump gene expression. These responses reflect a coordinated mitigation of proton toxicity, oxidative damage, and metabolic disruption, suggesting that higher MT availability is required to counteract the elevated ROS burden and pH imbalance typical of acid stress (Fig. 6). Similar pH or stress-intensity-dependent melatonin responses have been reported under salinity, metal toxicity, and drought, where stronger stress necessitated higher MT doses to restore cellular homeostasis (Arnao &

846 Hernández-Ruiz, 2019; Ahmad et al., 2023b; Ahammed et al., 2024; An et al., 2025;
847 Basharat et al., 2025).

848 In contrast, under near-neutral pH, most physiological parameters peaked at
849 low MT concentrations and declined at higher doses, indicating that excessive
850 melatonin can disrupt optimal redox and hormonal signaling when stress pressure is
851 minimal (Fig. 6). Basal ROS are essential signaling molecules for normal growth,
852 photosynthesis, and development, and over-scavenging by high MT levels may
853 suppress these signaling pathways, leading to reduced growth and metabolic
854 efficiency (Back, 2021; Liu et al., 2023; Dzinyela et al., 2025). Thus, the MT and pH
855 interaction demonstrates that melatonin acts within an optimal concentration window
856 that shifts upward with increasing stress severity.

857 Collectively, this interaction highlights melatonin as a conditional bioregulator
858 whose beneficial range depends on environmental pH, emphasizing the importance of
859 dose optimization when applying MT in agronomic systems.



860 **Figure 6.** Melatonin and pH interaction shows that a higher melatonin level is required to
861 alleviate seedling growth and yield, oxidative damage, photosynthetic pigments, ROS and
862 H^+ -ATPase genes expression and pH imbalance under acidic stress, whereas excessive
863 melatonin can disrupt normal growth under optimal conditions.

864 **Conclusion**

The present study provides comprehensive evidence that exogenous melatonin application markedly enhances peanut tolerance to acidic stress by integrating redox regulation, growth maintenance, and pH homeostasis. Acidic conditions severely restricted seed germination, seedling growth, biomass accumulation, pigment synthesis, and differentially affect antioxidants, while inducing excessive H₂O₂ accumulation and membrane lipid peroxidation. Melatonin application significantly reversed these effects, particularly at pH 4, by suppressing oxidative damage, protecting the photosynthetic apparatus, and sustaining biomass production. This protection was associated with a coordinated reprogramming of the antioxidant system, characterized by enhanced SOD, CAT and APX activities and reduced POD activity, indicating alleviation of oxidative pressure and a shift toward more efficient H₂O₂ detoxification. Importantly, melatonin strongly induced the plasma membrane H⁺-ATPase genes *AhAH1* and *AhAH2*, especially in roots, providing molecular evidence that melatonin reinforces proton extrusion capacity and cellular pH stability under acid stress.

Overall, the strong melatonin and pH interaction observed across all parameters demonstrates that higher melatonin concentrations are required to confer effective protection under acidic stress, whereas excessive melatonin under near-neutral conditions can impair normal physiological regulation. Collectively, these findings establish melatonin as a multifunctional regulator of acid stress adaptation in peanut and reveal a previously underexplored role of melatonin in mechanistically linking redox control with proton transport to support early crop establishment on acidic soils.

Author Contributions

Muhammad Hafeez Ullah Khan: conceptualization, investigation, writing – original draft. **Fu Ruoyi:** methodology and formal analysis. **Ali Muhammad:** visualization and formal analysis. **Shuaichao Zheng:** data curation and formal analysis. **Daijing Zhang:** writing – review and editing. **Zhiyong Zhang:** supervision, funding acquisition, writing – review and editing. **Quanyong Liu:** supervision, funding acquisition, writing – review and editing.

Funding and Acknowledgement

This research was partially supported by the National Natural Science Foundation of China (31571600, 31271648), the Key R&D and Promotion Projects of Henan Province (222102110303, 232102110221, 242102111094, 252102111072), the International Science and Technology Cooperation Project of Henan Province

(242102521050), and the Henan Center of Outstanding Overseas Scientists (GZS2024018). We gratefully acknowledge Professor Xiaoping Pan from East Carolina University for her great input, comments, and proofreading of this paper.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

- Ahammed, G. J., Li, Z., Chen, J., Dong, Y., Qu, K., Guo, T., Wang, F., Liu, A., Chen, S. & Li, X. 2024. Reactive oxygen species signaling in melatonin-mediated plant stress response. *Plant Physiology and Biochemistry*, **207**, 108398.
- Ahammed, G. J., Wu, M., Wang, Y., Yan, Y., Mao, Q., Ren, J., Ma, R., Liu, A. & Chen, S. 2020. Melatonin alleviates iron stress by improving iron homeostasis, antioxidant defense and secondary metabolism in cucumber. *Scientia Horticulturae*, **265**, 109205.
- Ahmad, I., Song, X., Hussein Ibrahim, M. E., Jamal, Y., Younas, M. U., Zhu, G., Zhou, G. & Adam Ali, A. Y. 2023a. The role of melatonin in plant growth and metabolism, and its interplay with nitric oxide and auxin in plants under different types of abiotic stress. *Front Plant Sci*, **14**, 1108507.
- Ahmad, R., Alsahli, A. A., Alansi, S. & Altaf, M. A. 2023b. Exogenous melatonin confers drought stress by promoting plant growth, photosynthetic efficiency and antioxidant defense system of pea (*Pisum sativum* L.). *Scientia Horticulturae*, **322**, 112431.
- Ali, S., Gill, R. A., Ulhassan, Z., Zhang, N., Hussain, S., Zhang, K., Huang, Q., Sagir, M., Tahir, M. B., Gill, M. B., Mwamba, T. M., Ali, B. & Zhou, W. 2023. Exogenously applied melatonin enhanced the tolerance of *Brassica napus* against cobalt toxicity by modulating antioxidant defense, osmotic adjustment, and expression of stress response genes. *Ecotoxicology and Environmental Safety*, **252**, 114624.
- Ameen, M., Zafar, A., Mahmood, A., Zia, M. A., Kamran, K., Javaid, M. M., Yasin, M. & Khan, B. A. 2024. Melatonin as a master regulatory hormone for genetic responses to biotic and abiotic stresses in model plant *Arabidopsis thaliana*: a comprehensive review. *Functional Plant Biology*, **51**.
- An, W., Wang, G., Dou, J., Zhang, Y., Yang, Q., He, Y., Tang, Z. & Yu, J. 2025. Protective mechanisms of exogenous melatonin on chlorophyll metabolism and photosynthesis in tomato seedlings under heat stress. *Frontiers in Plant Science*, **16**, 1519950.
- Arnao, M. B. & Hernández-Ruiz, J. 2019. Melatonin: a new plant hormone and/or a plant master regulator? *Trends in Plant Science*, **24**, 38–48.
- Back, K. 2021. Melatonin metabolism, signaling and possible roles in plants. *The Plant Journal*, **105**, 376–391.

941 Basharat, S., Saeed, W., Mubeen, S., Khan, L. U., Zhang, S., Liu, P. & Waseem, M. 2025.
 942 Exogenous Melatonin Regulates Hormone Signalling and
 943 Photosynthesis-Related Genes to Enhance Brassica napus. Yield: A
 944 Transcriptomic Perspective. *Journal of Pineal Research*, **77**, e70077.
 945 Borhannuddin Bhuyan, M., Hasanuzzaman, M., Nahar, K., Mahmud, J. A., Parvin, K.,
 946 Bhuiyan, T. F. & Fujita, M. 2019. Plants behavior under soil acidity stress:
 947 Insight into morphophysiological, biochemical, and molecular responses.
 948 *Plant abiotic stress tolerance: Agronomic, molecular and biotechnological*
 949 *approaches*, 35–82.
 950 Cai, B., Vancov, T., Si, H., Yang, W., Tong, K., Chen, W. & Fang, Y. 2021. Isolation and
 951 Characterization of Endomycorrhizal Fungi Associated with Growth Promotion
 952 of Blueberry Plants. *J Fungi (Basel)*, **7**.
 953 Chakraborty, N., Das, A., Pal, S., Roy, S., Sil, S. K., Adak, M. K. & Hasanuzzaman, M.
 954 2024. Exploring Aluminum tolerance mechanisms in plants with reference to
 955 rice and Arabidopsis: A comprehensive review of genetic, metabolic, and
 956 physiological adaptations in acidic soils. *Plants*, **13**, 1760.
 957 Chakraborty, S. & Raychaudhuri, S. S. 2025. Melatonin improves the lead tolerance in
 958 Plantago ovata by modulating ROS homeostasis, phytohormone status and
 959 expression of stress-responsive genes. *Plant Cell Reports*, **44**, 39.
 960 Chao, Z.-F. & Chao, D.-Y. 2025. Barriers and carriers for transition metal homeostasis
 961 in plants. *Plant Communications*, **6**.
 962 Chen, T., Xu, L., Yang, P., Tong, J., Liu, Y., Su, Y. & Sun, S. 2025. Plant melatonin: roles
 963 and regulatory mechanisms in plant growth, development, and responses to
 964 biotic and abiotic stresses. *Plant Signaling & Behavior*, **20**, 2585629.
 965 Chung, I.-M., Thiruvengadam, M., Samynathan, R., Lee, S.-B., Choi, H.-J., Jung, B.-S.,
 966 Park, Y., Jeon, D., Venkidasamy, B. & Kim, S.-H. 2025. Molecular insights into
 967 melatonin-mediated stress tolerance in rice. *Plant Cell Reports*, **44**, 271.
 968 Cosse, M. & Seidel, T. 2021. Plant proton pumps and cytosolic pH-homeostasis.
 969 *Frontiers in Plant Science*, **12**, 672873.
 970 Delfim, J., Moreira, A. & Moraes, L. A. 2025. Wheat growth, yield, nutrient
 971 concentration and soil chemical properties influenced by liming in an acid soil.
 972 *Journal of Plant Nutrition*, **48**, 87–100.
 973 Dong, T., Yan, L., Wang, J., Niu, Y. & Wang, L. 2025. Salt Tolerance in Soybean (Glycine
 974 max L.): A Comprehensive Review of Molecular Mechanisms, Key Regulators,
 975 and Future Perspectives for Saline Soil Utilization. *Plants*, **14**, 3668.
 976 Dou, J., Tang, Z., Yu, J., Wang, G., An, W., Zhang, Y. & Yang, Q. 2025. Effects of
 977 exogenous melatonin on the growth and photosynthetic characteristics of
 978 tomato seedlings under saline-alkali stress. *Scientific Reports*, **15**, 5172.
 979 Dzinyela, R., Manda, T., Hwarari, D., Ramakrishnan, M., Ahmad, Z., Agassin, R. H.,
 980 Yang, L. & Movahedi, A. 2025. Melatonin-mediated phytohormonal crosstalk
 981 improves salt stress tolerance in plants. *Planta*, **262**, 86.
 982 Fu, D., Wu, W., Mustafa, G., Yang, Y. & Yang, P. 2025. Molecular mechanisms of rice
 983 seed germination. *New Crops*, **2**, 100051.
 984 Gámez-Arjona, F. M., Sánchez-Rodríguez, C. & Montesinos, J. C. 2022. The root
 985 apoplastic pH as an integrator of plant signaling. *Frontiers in Plant Science*, **13**,
 986 931979.

987 Hasan, M. K., Imon, M. A. A.-H., Jahan, I., Das, T. K., Mishu, M. S., Shopan, J., Saha, M.,
988 Khan, M. S. I., Islam, M. S. & Ahammed, G. J. 2025. Melatonin-mediated
989 adaptation of spinach (*Spinacia oleracea* L.) in acidic soil. *Plant Physiology and*
990 *Biochemistry*, **223**, 109816.

991 Hlihor, R. M., Roşca, M., Hagiuz-Zaleschi, L., Simion, I. M., Daraban, G. M. & Stoleru, V.
992 2022. Medicinal plant growth in heavy metals contaminated soils: Responses
993 to metal stress and induced risks to human health. *Toxics*, **10**, 499.

994 Huang, K., Li, M., Li, R., Rasul, F., Shahzad, S., Wu, C., Shao, J., Huang, G., Li, R., Almari,
995 S., Hashem, M. & Aamer, M. 2023. Soil acidification and salinity: the
996 importance of biochar application to agricultural soils. *Frontiers in Plant*
997 *Science*, **Volume 14 - 2023**.

998 Huang, X., Tanveer, M., Min, Y. & Shabala, S. 2022. Melatonin as a regulator of plant
999 ionic homeostasis: implications for abiotic stress tolerance. *Journal of*
1000 *Experimental Botany*, **73**, 5886–5902.

1001 Jahan, I., Shopan, J., Rahman, M. M., Sarkar, A., Baset, M. A., Zhang, Z., Li, X.,
1002 Ahammed, G. J. & Hasan, M. K. 2022. Long-term traditional fertilization alters
1003 tea garden soil properties and tea leaf quality in Bangladesh. *Agronomy*, **12**,
1004 2128.

1005 Jain, D. & Schmidt, W. 2024. Protein phosphorylation orchestrates acclimations of
1006 arabidopsis plants to environmental pH. *Molecular & Cellular Proteomics*, **23**.

1007 Jamir, Z., Tzudir, L., Kumari, S., Solo, V., Dolie, S., Nongmaithem, D., Yadav, R., Kikon, N.
1008 & Mollier, M. 2025. Productivity of Groundnut (*Arachis hypogaea* L.) under
1009 Different Sowing Times, Utrient Supply Levels and Planting Geometry in Sub-
1010 Humid Tropical Environment. *Current World Environment*, **20**, 121.

1011 Jiang, D., Ou, Y., Jiang, G., Dai, G., Liu, S. & Chen, G. 2025a. Melatonin-priming
1012 ameliorates aluminum accumulation and toxicity in rice through enhancing
1013 aluminum exclusion and maintaining redox homeostasis. *Plant Physiology and*
1014 *Biochemistry*, **219**, 109433.

1015 Jiang, Z., van Zanten, M. & Sasidharan, R. 2025b. Mechanisms of plant acclimation to
1016 multiple abiotic stresses. *Communications Biology*, **8**, 655.

1017 Kong, X., Wang, R., Jia, P., Li, H., Khan, A., Muhammad, A., Fiaz, S., Xing, Q. & Zhang, Z.
1018 2025. Physio-biochemical and molecular mechanisms of low nitrogen stress
1019 tolerance in peanut (*Arachis hypogaea* L.). *Plant Molecular Biology*, **115**, 19.

1020 Li, J., Yang, J., Liang, X., Zhan, S., Bai, Y. & Ruan, L. 2025a. Melatonin-Mediated
1021 Regulation of Antioxidant Defense Enhances the Resistance of Tea Plants
1022 (*Camellia sinensis* L.) to Lead-Induced Stress. *Plants*, **14**, 1417.

1023 Li, J., Zhang, Z. & Shi, G. 2024a. Genome-wide identification and expression profiling
1024 of heavy metal ATPase (HMA) genes in peanut: Potential roles in heavy metal
1025 transport. *International Journal of Molecular Sciences*, **25**, 613.

1026 Li, L., Cheng, X., Zhang, Y., Kohtz, D., Wang, X., Zhang, X., Kong, X., Xue, H., Jia, P., Bai,
1027 N., Li, Z., Xiao, P., Pan, X., Zhang, Z. & Zhang, B. 2024b. Exogenous melatonin
1028 improves peanut field productivity and quality at reduced nitrogen
1029 application. *Field Crops Research*, **319**, 109650.

1030 Li, L., Li, Q., Davis, K. E., Patterson, C., Oo, S., Liu, W., Liu, J., Wang, G., Fontana, J. E.,
1031 Thornburg, T. E., Pratt, I. S., Li, F., Zhang, Z., Zhou, Y., Pan, X. & Zhang, B. 2021.
1032 Response of Root Growth and Development to Nitrogen and Potassium

1033 Deficiency as well as microRNA-Mediated Mechanism in Peanut (*Arachis*
1034 *hypogaea* L.). *Frontiers in Plant Science*, **Volume 12** - 2021.

1035 Li, Q., Zhang, Y., Liu, Y., Li, T., Xu, H., Wei, Q., Zeng, H., Ni, H. & Li, S. 2025b. Multiple
1036 functions of exogenous melatonin in cucumber seed germination, seedling
1037 establishment, and alkali stress resistance. *BMC Plant Biology*, **25**, 359.

1038 Li, W.-Q., Li, J.-Y., Bi, S.-J., Jin, J.-Y., Fan, Z.-L., Shang, Z.-L., Zhang, Y.-F. & Wang, Y.-J.
1039 2025c. Melatonin enhances maize germination, growth, and salt tolerance by
1040 regulating reactive oxygen species accumulation and antioxidant systems.
1041 *Plants*, **14**, 296.

1042 Li, Y., Zeng, H., Xu, F., Yan, F. & Xu, W. 2022. H⁺-ATPases in plant growth and stress
1043 responses. *Annual Review of Plant Biology*, **73**, 495–521.

1044 Li, Z., Tang, Z., Song, J., Li, J., Wang, X. & Zhang, X. 2025d. Melatonin stabilizes
1045 photosynthesis and scavenges ROS to enhance salt tolerance in cucumber.
1046 *BMC Plant Biology*, **25**, 1433.

1047 Liu, H., Ren, X., Zhu, J., Wu, X. & Liang, C. 2018. Effect of exogenous abscisic acid on
1048 morphology, growth and nutrient uptake of rice (*Oryza sativa*) roots under
1049 simulated acid rain stress. *Planta*, **248**, 647–659.

1050 Liu, Z., Dai, H., Hao, J., Li, R., Pu, X., Guan, M. & Chen, Q. 2023. Current research and
1051 future directions of melatonin's role in seed germination. *Stress Biology*, **3**, 53.

1052 MacDonald, E., Johannes, L. & Wunder, C. 2025. Acidification on the plasma
1053 membrane. *Current Opinion in Cell Biology*, **95**, 102531.

1054 Menhas, S., Chen, M., Jin, H., Xu, J., Zhu, S. & Lin, D. 2025. Plant growth stage and
1055 melatonin concentration dependency together drive the metal-nutrient
1056 dynamics of rice in paddy soil. *International Journal of Phytoremediation*, **27**,
1057 958–971.

1058 Moustafa-Farag, M., Elkelish, A., Dafea, M., Khan, M., Arnao, M. B., Abdelhamid, M.
1059 T., El-Ezz, A. A., Almoneafy, A., Mahmoud, A., Awad, M., Li, L., Wang, Y.,
1060 Hasanuzzaman, M. & Ai, S. 2020. Role of Melatonin in Plant Tolerance to Soil
1061 Stressors: Salinity, pH and Heavy Metals. *Molecules*, **25**, 5359.

1062 Msimbira, L. A. & Smith, D. L. 2020. The roles of plant growth promoting microbes in
1063 enhancing plant tolerance to acidity and alkalinity stresses. *Frontiers in*
1064 *Sustainable Food Systems*, **4**, 106.

1065 Muhammad, A., Kong, X., Li, L., Khan, M. H. U., Jia, P., Miao, C. & Zhang, Z. 2026.
1066 Melatonin Enhances Peanut Productivity by Enriching Root-Associated
1067 Nitrogen-Fixing Bacteria. *Journal of Pineal Research*, **78**, e70105.

1068 Muhammad, I., Ahmad, S. & Shen, W. 2024. Melatonin-mediated molecular
1069 responses in plants: Enhancing stress tolerance and mitigating environmental
1070 challenges in cereal crop production. *International Journal of Molecular*
1071 *Sciences*, **25**, 4551.

1072 Muhammad, I., Fahad, S., Khalofah, A., Zheng, B. & Shen, W. 2025. Melatonin
1073 Enhances Antioxidant Defense Systems and Stress Tolerance in Plants under
1074 Variable Environmental Conditions. *Rice (N Y)*, **18**, 70.

1075 Munyaneza, V., Zhang, W., Haider, S., Xu, F., Wang, C. & Ding, G. 2024. Strategies for
1076 alleviating aluminum toxicity in soils and plants. *Plant and Soil*, **504**, 167–190.

1077 Naz, M., Dai, Z., Hussain, S., Tariq, M., Danish, S., Khan, I. U., Qi, S. & Du, D. 2022. The
1078 soil pH and heavy metals revealed their impact on soil microbial community.
1079 *Journal of Environmental Management*, **321**, 115770.

1080 Sharma, A., Kumar, V., Shahzad, B., Ramakrishnan, M., Singh Sidhu, G. P., Bali, A. S.,
1081 Handa, N., Kapoor, D., Yadav, P. & Khanna, K. 2020. Photosynthetic response
1082 of plants under different abiotic stresses: a review. *Journal of Plant Growth*
1083 *Regulation*, **39**, 509–531.

1084 Shavrukov, Y. & Hirai, Y. 2016. Good and bad protons: genetic aspects of acidity stress
1085 responses in plants. *Journal of Experimental Botany*, **67**, 15–30.

1086 Shehzadi, K., Maqsood, M. F., Kanwal, R., Shahbaz, M., Naqve, M., Zulfiqar, U., Jamil,
1087 M., Khalid, N., Ali, M. F. & Soufan, W. 2025. Enhancing cadmium stress
1088 resilience in chickpea (*Cicer arietinum* L.) via exogenous melatonin application.
1089 *International Journal of Phytoremediation*, **27**, 794–809.

1090 Sun, G., Luan, M., Xue, S., Yan, J. & Lan, W. 2025. Vacuolar H⁺-ATPase Is Required for
1091 Efficient Vacuolar Phosphate Storage and Systemic Pi Homeostasis in
1092 *Arabidopsis*. *Plant, Cell & Environment*, **48**, 587–598.

1093 Tahjib-Ul-Arif, M., Zahan, I., Hossain, M. S., Imran, S., Hasanuzzaman, M., Dawood, M.
1094 F., Dawood, A. F., Asaduzzaman, M., Rhaman, M. S. & Sour, Z. 2025.
1095 Melatonin-mediated ionic homeostasis in plants: Mitigating nutrient
1096 deficiency and salinity stress. *Discover Plants*, **2**, 143.

1097 Thapa, A., Hasan, M. R. & Kabir, A. H. 2025. Transcriptional reprogramming and
1098 microbiome dynamics in garden pea exposed to high pH stress during
1099 vegetative stage. *Planta*, **261**, 83.

1100 Wang, F., Li, X., Chen, C., Zhao, L. & Wei, Y. 2025a. Mechanism of Melatonin in
1101 Alleviating Aluminum Toxicity in Plants: A Review. *Biology*, **14**, 1316.

1102 Wang, J., Jin, D., Deng, Z., Zheng, L., Guo, P., Ji, Y., Song, Z., Zeng, H. Y., Kinoshita, T. &
1103 Liao, Z. 2025b. The apoplastic pH is a key determinant in the hypocotyl
1104 growth response to auxin dosage and light. *Nature Plants*, 1–16.

1105 Wang, L., Ju, C., Han, C., Yu, Z., Bai, M. Y. & Wang, C. 2025c. The interaction of
1106 nutrient uptake with biotic and abiotic stresses in plants. *Journal of*
1107 *Integrative Plant Biology*, **67**, 455–487.

1108 Yang, S.-J., Huang, B., Zhao, Y.-Q., Hu, D., Chen, T., Ding, C.-B., Chen, Y.-E., Yuan, S. &
1109 Yuan, M. 2021. Melatonin enhanced the tolerance of *Arabidopsis thaliana* to
1110 high light through improving anti-oxidative system and photosynthesis.
1111 *Frontiers in Plant Science*, **12**, 752584.

1112 Yang, X., Liu, D., Liu, C., Li, M., Yan, Z., Zhang, Y. & Feng, G. 2024. Possible melatonin-
1113 induced salt stress tolerance pathway in *Phaseolus vulgaris* L. using
1114 transcriptomic and metabolomic analyses. *BMC Plant Biology*, **24**, 72.

1115 Yang, Y., Guan, S., Jiang, X., Li, M., Wei, S. & Diao, M. 2023. Exogenous melatonin
1116 alleviates the inhibitory effect of NaHCO₃ on tomato growth by regulating the
1117 root pH value and promoting plant photosynthesis. *Agronomy*, **13**, 2777.

1118 Zeng, H., Chen, H., Zhang, M., Ding, M., Xu, F., Yan, F., Kinoshita, T. & Zhu, Y. 2024.
1119 Plasma membrane H⁺-ATPases in mineral nutrition and crop improvement.
1120 *Trends in Plant Science*, **29**, 978–994.

1121 Zhang, J., Wei, J., Li, D., Kong, X., Rengel, Z., Chen, L., Yang, Y., Cui, X. & Chen, Q. 2017.
1122 The role of the plasma membrane H⁺-ATPase in plant responses to aluminum
1123 toxicity. *Frontiers in Plant Science*, **8**, 1757.

1124 Zheng, S., Tang, Z., Li, L., Jia, P., Kong, X., Xue, H., Cheng, X., Wang, X., Zhang, Y. & Bai,
1125 N. 2025. Melatonin reprograms soil microbial community, creates friendly soil

1126 environments, and promotes peanut growth. *Plant Physiology and*
1127 *Biochemistry*, **218**, 109307.
1128