

Growth, abundance, and diversity of black mangrove (*Avicennia germinans*) pneumatophores in deforested and sand-filled mangrove forest at Eagle Island, Niger Delta Nigeria

Aroloye Numbere (✉ aroloyen@yahoo.com)

University of Port Harcourt

Gerardo Camilo

Saint Louis University

Research Article

Keywords: pneumatophore, branching, muddy soil, climate change, oxygen, black mangrove

Posted Date: August 22nd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1810526/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Pneumatophore is a major channel of oxygen circulation in mangrove forest. It is hypothesized that soil condition, canopy cover and stagnant pool of water can influence pneumatophore growth. Pneumatophore abundance, diversity indices, microbial and chemical composition in sand-filled mangrove forest was studied. Six plots were established within a rectangular area measuring 4152.24m² in a random block design to investigate the effect of soil conditions on pneumatophore growth. A total of 9,586 pneumatophores were physically counted and differentiated into four types namely 1-branch, 2-branch, 3-branch, and 4-branch pneumatophores. The ANOVA results shows that there is significant difference in the abundance of pneumatophore types ($F_{3,20} = 7.61, p < .001$). The most abundant pneumatophore type is the one branch pneumatophore ($n = 4747$) while the least is the four branched pneumatophores. The stagnant pool site with silty and muddy soil has the most abundant pneumatophore growth whereas plots in the seashore site with sandy soil has the least abundant pneumatophores. In contrast, the seashore site has the highest diversity ($H = 1.367$) while the stagnant pool has the lowest diversity ($H = 0.956$). Metal concentration was higher in the stagnant pool site while microbial count is higher in the seashore site. The study shows that soil condition, canopy cover, stagnant pool plus tidal action influenced pneumatophore growth.

Introduction

Mangroves are coastal halophytic plants that can survive in marine, freshwater, and estuarine environments (Spencer et al. 2022). They are aquatic and semi-aquatic because of their growth in the interface between the land and the sea. Mangroves are reported to be land forming by using their adventitious roots to trap sediments (Carlton 1974), which metamorphose into muddy soil and later a hard land (Krause et al. 2014). Mangroves can survive the difficult, turbulent and salty coastal terrain because of their unique root system that is used for respiration (Pratolongo 2022) and excretion of heavy metals (Hilmi et al. 2022) and salts (Munir et al. 2022). The adventitious root of mangroves is called “breathing root” because it is used for gaseous exchange through the transmission of atmospheric oxygen in and out of the plant (Numbere, 2018). The breathing root thus, serves as the lungs of the mangrove trees (Ke et al. 2022; Hoogeveen 2020). This is the reason they can survive when submerged in an aquatic environment or impacted by tidal currents.

The pneumatophore is part of the root system of the black mangroves (*Avicennia germinans*), which are vertical finger-like projections that protrude from the forest ground beneath the trees (Eisenberg 2002). The pneumatophore grows out of the soil into the air to facilitate atmospheric gaseous exchange (Clough 2013). The atmospheric oxygen complements the underwater oxygen for submerged roots. There are thousands of these pneumatophores at the base of black mangrove trees forming radial circumference around the forest canopy (Ong and Gong, 2013). The pneumatophores also have the biological significance of regulating salinity exchange between the plant and the marine environment (Munir et al. 2022). They act as a shutdown system that prevent the excessive intake of salt solutions into the plant, which helps the mangrove to survive high concentration of sodium chloride and prevent the osmotic

collapse of its cell from excess salt crystals. The pneumatophores also serve as an erosion break (Stewart and Fairfull, 2013), conduit pipe for the transmission of nutrient minerals and water into the plant and the excretion of salt crystals and other harmful waste substances from the plant. The pneumatophore is one of the organs of the mangroves that has made it to be successful as an aquatic plant. Pneumatophores also play other vital roles in the mangrove ecosystem by providing soil nutrients themselves through the decomposition of their death parts. They increase the soil nutrients when their litter attract microbes that carry out the decomposition of the organic materials.

Pneumatophores are good spawning ground for fingerlings of fish (e.g., tilapia) and other crustaceans e.g., fiddler crabs that reside in the mangrove forest (Mandal et al. 2013). The finger like projections of the pneumatophores also traps other organisms during low tides, which serve as food for other organism in the food chain. Crabs and other organisms feed on the soft inner part of the pneumatophores (Roy, 2011). Because of the numerous physical, chemical, and biological roles played by the pneumatophores, it is thus, important to study the factors that affect their growth, distribution, abundance, and diversity in the deforested sand-filled area. The chemical composition of the pneumatophore is studied because they don't only serve as a sanctuary for spawning organisms, but also serve as a trap for harmful waste such as plastics, which decompose to produce harmful substances consumed by the fish, which is later transferred to human thereby leading to public health disaster (Fig. 3). The black mangrove was selected because it is the most dominant mangrove species in this study areas and the only mangroves with that produce pneumatophores. Other mangroves have giant adventitious roots system that protrude from the ground and form long chains and knee joins but not like that of the black mangroves e.g., red mangroves (e.g., *Rhizophora* species).

The goal of this study, therefore, is to determine the pneumatophore abundance and diversity, and the chemical, and microbial composition of soil around the pneumatophore in the sand-filled area. The following questions were thus addressed: (1) Are there differences in pneumatophore abundance and diversity within plots and sites? If so which part of the plot or site will have higher pneumatophore abundance and diversity, that is, the stagnant pool that is muddy or seashore that is sandy? (2) Are there differences in chemical and microbial composition of the soil at the stagnant pool and seashore sites of the deforested and sand filled mangrove forest? (3) Are there correlations between living versus dead or length versus weight, length versus branch and weight versus branch of pneumatophores?

Methods

Description of study area

The study was conducted in a deforested and sand filled mangrove forest directly behind the back gate of Rivers State university (Fig. 1). Several years ago, this area was covered by a luxuriant mangrove forest with some scattered nypa palm trees. But now part of the forest is gone because it was cut down for the establishment of a sand mine, which has also been abandoned. The area is currently in an early stage of succession by young mangrove seedlings with other non-mangrove plants such as grasses and aquatic

weed. There are four mangrove species present in the area namely red (*Rhizophora racemosa*), black (*Avicennia germinans*), grey (*Acrosticum aureum*) and white (*Laguncularia racemosa*). However, the black mangrove is the most dominant species in this ecosystem (Numbere, 2021). During high tides the plant trap plastic and other types of waste within the sand filled area. The soil is light to dark brown in color and has low to medium plasticity because of the muddy nature of the soil and has a fine to coarse sand particles. The temperature is between 28 to 34°C and the pH is from 6 to 7 while the salinity ranges from 1.45–1.62‰. The muddy area is brownish soft and silty while the sandy area is whitish, coarse, and rough in texture. The area experiences a 6-hourly tidal cycle with water rushing into the sand-filled area during high tide and flowing back into the sea during the low tide.

Experimental design

The study used a random block design in an area measuring 94.78 m × 43.76 m (4152.24m²), which was further sub divided into six plots (habitat types) (Fig. 2). Each plot was delineated equally using a standard tape measure at an accuracy of 0.1m. Two key areas of the plots are the seashore that is sandy and coarse and close to the river (Plots 1–3) and stagnant pool (Plots 4–6) that is muddy and silty and away from the river. Furthermore, the seashore site is always dry during the low tide and covered with water during the high tide while the stagnant pool is constantly wet because it is supplied by river water during high tide and rainwater. The wetness of the stagnant pool has caused the growth of algae (spirogyra) and the standing pool of water serves as a death trap for organisms ranging from insects, fish, and crabs (Fig. 3). In each plot all the pneumatophores that were visible were painstakingly counted physically to get the abundance using line transects thrown across the ground in series. Pictures and videos were also used to get a view of the number of pneumatophores. The pneumatophores were grouped according to the number of branches that was identified for the purpose of this study as follows: 1-branch, 2-branch, 3-branch, and 4-branch. They refer to the pneumatophores that have one, two, three and four branches growing between 5 cm of the sub surface and surface soil. The six plots were georeferenced with a Garmin GPS (USA) (Table 1).

Table 1
Soil characteristics in different plots at Eagle Island, Niger Delta, Nigeria

Plots	Coordinates	Elevation (m)	Moisture content (%)	Total organic content (TOC)	Pore water salinity (%)	Soil compaction (kg/cm ²)	pH	Temp (°C)
Plot 1	N04°47.289; E006°58.558	02.00	20.00	1.44 ± 0.01	1.16 ± 0.02	0.25 ± 0.01	6.8 ± 0.1	26.1 ± 0.1
Plot 2	N04°47.305; E006°58.553	03.00	20.00	1.40 ± 0.01	1.45 ± 0.01	0.21 ± 0.02	6.6 ± 0.1	26.2 ± 0.1
Plot 3	N04°47.315; E006°58.561	03.00	30.00	1.13 ± 0.01	1.48 ± 0.02	0.26 ± 0.01	6.8 ± 0.2	26.4 ± 0.1
Plot 4	N04°47.296; E006°58.521	54.00	60.00	2.03 ± 0.01	1.53 ± 0.02	0.20 ± 0.01	5.4 ± 0.1	29.1 ± 0.1
Plot 5	N04°47.299; E006°58.519	22.00	38.00	1.79 ± 0.01	1.55 ± 0.01	0.23 ± 0.03	6.1 ± 0.1	27.7 ± 0.1
Plot 6	N04°47.293; E006°58.552	04.00	55.00	2.22 ± 0.01	1.55 ± 0.02	0.21 ± 0.02	5.3 ± 0.1	28.5 ± 0.1

Pneumatophore growth and development

Pneumatophores are dominant in this study area because of the large population of black mangroves (Numbere, 2021). They grow in large or small population underneath the black mangrove trees. They are made up of a soft dark outer and a light inner coat. The outer coat is slippery to the touch and can easily be pulled away. The pneumatophores prevent the growth of other plant species at ~ 1.8 m circumference around the black mangrove tree.

Pneumatophore abundance and diversity

All pneumatophores in sight were counted in situ, and their abundance recorded in a data sheet. The Shannon diversity index (H) was used to determine the pneumatophore diversity. It is the natural logarithm which considers low and high species diversities based on abundances. High diversity index means high biodiversity while low diversity index means low biodiversity (Shannon & Weiner, 1949).

Soil sand pneumatophore samples collection

A hand-held soil augur was used to randomly collect soil samples from each plot 5cm below the soil surface within each transect ($n = 10$) in each plot. The samples were placed in a well-labelled

polyethylene bags and sent to the laboratory for physico-chemical and microbial analysis. The seashore and stagnant pool sites were studied in detail to determine their influence on the growth and proliferation of the pneumatophore around the coastal mangrove ecosystem. The soil texture and composition were also studied. Ten pneumatophore samples from each plot were randomly pulled by hand, bagged, and sent to the laboratory for measurements of length (cm) and weight (g) in m at the level of accuracy of 0.1m and 0.1 g respectively and physico-chemically analyzed.

Physico-chemical analysis

Nitrate, Iron, Cadmium, Lead, Zinc, Copper, and total hydrocarbon content (THC) were determined using the method of Aigberua & Tarawou, (2018) where aliquots of 0.25g of air-dried sediment samples were weighed into a Teflon inset of a microwave digestion vessel, and 2 ml concentrated (90%) nitric acid (Sigma-Aldrich, Dorset, UK) were added. The metals were extracted using a microwave accelerated reaction system (MARS Xpress, CEM Corporation, Matthews, North Carolina) at 1500 W power (100%), ramped to 175°C in 5.5 min, held for 4.5 min, and allowed to cool down for 1 h. The cool digest solution was filtered through the Whatman 42 filter paper and made up to 100 ml in a volumetric flask by adding de-ionized water. The detection limit for the metals analyzed was 0.001 mg/l.

Soil pH was determined with a Kelway soil tester while the soil compaction was determined with a pocket penetrometer. Soil temperature was determined with a digital dual sensor thermometer to a detection unit of $\pm 1^\circ\text{C}$. Salinity of the pore water soil was determined with a salinity meter (OAKTON Salt 6 Acorn Series). The salinity meter probe was used to test standing water in dug out holes during low tide.

Total organic content (TOC) was determined using Walkey-Black titrimetric method (Table 1). The TOC was used to determine the nutrients in the soil. The TOC was determined because soil organic content influences soil texture and composition, which in turn influences mangrove growth (Alongi, 2009).

Microbial analysis

Total heterotrophic bacteria (THB) and total heterotrophic fungi (THF) were analyzed from soils collected randomly from each plot. Firstly, the laboratory procedure for the THB was determined as follows: 1g of soil sample was weighed into 9ml sterile diluents (0.85% NaCl) under aseptic condition. It was then shaken vigorously to homogenize and serially diluted. Then 0.1ml aliquot of the inoculums was collected using a sterile pipette, inoculated on nutrient agar surface. The inoculums were spread evenly with a sterile rod. Plates were incubated at 37°C for 24 hours. Thereafter, colonies were counted to obtain colony forming unit (cfu) value per ml of the soil sample. Distinct colonies were picked and streaked on freshly prepared nutrient agar medium to obtain pure culture after 24 hours incubation at 37°C. The pure culture was gram stained for microscopic examination. Secondly, the laboratory procedure for the THF was determined as follows: 1g of soil sample was weighed into 9ml sterile diluents (0.85% NaCl) under aseptic condition. It was then shaken vigorously to homogenize and serially diluted. 0.1ml aliquot of inoculums is inoculated on Potato Dextrose Agar (PDA) acidified with 0.1% lactic acid to inhibit growth of bacteria and allowed for only the growth of fungi. Inoculated plates were incubated at ambient temperature for 3–

5 days. Cultural characteristics of isolates were observed and sub-cultured for purification. Microscopic examination was done using lacto phenol cotton blue stain with $\times 400$ magnifications.

Statistical analysis

An analysis of variance (ANOVA) was conducted since there were multiple samples per block ($n = 60$) to test whether there was any significant difference in pneumatophore abundance within plots and habitat types following the example of Quinn & Keough (2002). Similarly, an ANOVA was used to determine if there were any significant differences in metal concentration and microbial population between plots and sites. Logarithmic transformation of the data was performed to meet assumptions of normality and homoscedasticity (Logan, 2010). Similarly, a post-hoc Tukey's HSD test was done to investigate pair wise mean differences between groups. Pearson's product-moment correlation was done to compare whether there was any significant difference between living vs. dead, pneumatophore and length vs. weigh, length vs. branch, and weight vs. branch of pneumatophores. Regression analysis was also done to compare pneumatophore abundance and siltiness of the soil. All analyses were performed in *R* statistical environment, 3.0.1 (*R* Core Team, 2013).

Results

Pneumatophore abundance

The ANOVA result shows that there is a significant difference in the abundance of pneumatophore types ($F_{3,20} = 7.61, p < .001$, Table 2. Figure 4). Similarly, there was significant difference between pneumatophore abundance and sites ($F_{1,22} = 6.82, p < .02$, Table 2. Figure 3). However, there was no significant difference in the abundance of pneumatophore between plots ($F_{5,18} = 1.33, p = 0.30$, Table 2. Figure 4). A post hoc Tukey's test showed that 1-branch and 4-branch pneumatophores differed most significantly at $P < .05$ (Fig. 3). Moreover 1-branch pneumatophores were the most abundant ($n = 4747$) followed by 2-branch ($n = 3620$) and 3-branch pneumatophores ($n = 1099$) in all plots. In contrast the 4-branch pneumatophores were the least. Plots 4–6 in the stagnant pool site has the most abundant pneumatophore growth whereas plots 1–3 in the seashore site have the least abundant pneumatophores. In the overall plot 6 has the highest number of pneumatophores while plot 1 has the least number of pneumatophores (Table 2).

Pneumatophore diversity

The pneumatophore diversity was estimated from the pneumatophore abundance within each plot. The result (Table 2) indicates that plot 3, along the seashore has the highest diversity ($H = 1.367$) followed by plot 2 ($H = 1.190$) and plot 1 ($H = 1.062$), which are all at the seashore and made of sandy soil.; while plot 5, at the stagnant pool made of muddy soil has the lowest diversity ($H = 0.956$).

Table 2

Pneumatophore abundance and diversity indices in different plots at Eagle Island, Niger Delta, Nigeria. It shows that the seashore site (Plots 1–3) has higher biodiversity than the stagnant pool site (Plots 4–6).

Site	Plot	Pneumatophore type	Abundance	Proportion (Pi)	Ln (Pi)	Pi* Ln (Pi)	H
	Plot 1	1-branch	502	0.4883	-0.717	-0.350	
		2-branch	390	0.3794	-0.969	-0.368	
		3-branch	100	0.0973	-2.330	-0.227	
		4-branch	36	0.0350	-3.352	-0.117	
		Total	1028				-1.062
	Plot 2	1-branch	146	0.3842	-0.957	-0.368	
		2-branch	155	0.4079	-0.897	-0.366	
Seashore		3-branch	54	0.1421	-1.951	-0.277	
		4-branch	25	0.0658	-2.721	-0.179	
		Total	380				-1.190
	Plot 3	1-branch	412	0.4568	-0.784	-0.358	
		2-branch	362	0.4013	-0.913	-0.768	
		3-branch	98	0.1087	-2.219	-0.241	
		4-branch	30	0.0333	-3.402	-0.113	
		Total	902				-1.367
	Plot 4	1-branch	1440	0.5012	-0.691	-0.346	
		2-branch	1119	0.3895	-0.943	-0.367	
		3-branch	302	0.1051	-2.253	-0.237	
		4-branch	12	0.0042	-5.473	-0.023	
		Total	2873				-0.973
	Plot 5	1-branch	1025	0.4971	-0.699	-0.348	
		2-branch	835	0.4050	-0.904	-0.366	
		3-branch	195	0.0946	-2.358	-0.223	

Site	Plot	Pneumatophore type	Abundance	Proportion (Pi)	Ln (Pi)	Pi* Ln (Pi)	H
Stagnant pool	Plot 6	4-branch	7	0.0034	-5.684	-0.019	
		Total	2062				-0.950
		1-branch	1220	0.5220	-0.650	-0.339	
		2-branch	759	0.3242	-1.126	-0.365	
		3-branch	350	0.1495	1.901	0.284	
		4-branch	10	0.0043	-5.449	-0.023	
		Total	2341				-1.011
		Gross Total	9586				

Physico-chemical analysis

The ANOVA result showed that there is a significant difference between concentration of metals ($F_{3,20} = 197.1$, $P < 0.001$, Table 3, Fig. 5). But there was no significant difference in metal concentration between sites ($F_{1,22} = 0.15$, $P > 0.05$, Table 3). However, there was higher Nitrate concentration in plots 1 (1.48 mg/kg) and 3 (1.50 mg/kg) (Table 3). Similarly, there was higher iron concentration in the seashore plots (1–3) than the stagnant pool plots (4–6).

Table 3

Soil metal composition of different study plots at Eagle Island, Niger Delta, Nigeria. Seashore plots are 1–3 and stagnant pool plots are 4–6.

Metals (mg/kg)						
Metal	Plot 1	Plot 2	Plot 3	Plot 4	Plot 5	Plot6
THC	1.22 ± 0.01	0.01 ± 0.00	0.20 ± 0.01	0.40 ± 0.01	2.00 ± 0.01	5.50 ± 1.10
Cadmium	0.21 ± 0.02	0.001 ± 0.00	0.12 ± 0.00	0.001 ± 0.00	0.001 ± 0.00	0.001 ± 0.00
Lead	0.36 ± 0.02	4.63 ± 1.20	0.001 ± 0.00	1.05 ± 0.01	0.46 ± 0.00	1.62 ± 0.1
Zinc	0.001 ± 0.00	0.001 ± 0.01	0.001 ± 0.00	10.06 ± 0.11	2.47 ± 0.20	2.43 ± 0.2
Iron	1631.48 ± 3.22	888.93 ± 7.34	2310.87 ± 2.55	158.38 ± 1.13	156.61 ± 4.66	619.63 ±
Nitrate	1.48 ± 0.10	1.24 ± 0.01	1.50 ± 0.01	1.42 ± 0.01	1.30 ± 0.01	1.24 ± 0.01
Copper	2.01 ± 0.20	0.85 ± 0.01	2.02 ± 0.01	0.21 ± 0.00	0.10 ± 0.00	1.33 ± 0.00

Microbial analysis

The result indicates that there is significant difference between bacterial and fungal populations ($F_{1,10} = 7.40$, $P = 0.02$, Fig. 6). Similarly, there was significant difference in microbial population across plots ($F_{1,10} = 5.50$, $P < 0.04$).

Growth of pneumatophores

The growth of pneumatophore at different sites were recorded by measuring the length (Fig. 7) and weight. The ANOVA result shows that there is a significant difference in the length of the pneumatophores growing in the seashore and stagnant pool sites ($P_{1,49} = 35.7$, $P < 0.001$, Fig. 7a). In contrast there is no significant difference in the weight of pneumatophore ($P_{1,48} = 0.02$, $P = 0.897$, Fig. 7b),

Correlation between living and dead pneumatophores and their length, branch, and weight

Furthermore, the correlation results indicate that there was little or no correlations in the number of dead vs. live ($t = 0.033$, $df = 37$, $p\text{-value} = 0.974$, $cor = 0.005402276$, Fig. 8), the length vs. branch ($t = -0.43$, $df = 48$, $p\text{-value} = 0.664$, $cor = 0.06289677$); and the weight vs. branch ($t = 0.108$, $df = 48$, $p\text{-value} = 0.914$, $cor = 0.0156338$) of pneumatophores at both sites. However, there is slight correlation between length and weight of pneumatophores ($t = 3.427$, $df = 48$, $p\text{-value} = 0.00126$, $cor = 0.4433579$). This means the taller the pneumatophore the heavier it is.

Discussion

Pneumatophores play unique role in oxygen circulation by acting as the breathing apparatus of mangrove forest globally (Kathiresan and Bingham, 2001, Hogarth, 2015). Pneumatophores don't only circulate oxygen in mangroves but also play key role in climate change by helping to sequester carbon through the absorption of atmospheric carbon dioxide by the network of their root systems (Zhang et al. 2022; Romero-Urbe et al. 2022; Ouyang et al. 2011). Therefore, this study is one of the first in the Niger Delta to report the growth of different types of pneumatophore branches and the factors that influence their proliferation and survival in coastal areas. The result of this study shows that soil chemical and microbial composition influence the growth of pneumatophores (e.g., Fusi et al. 2022). Areas with silt and muddy soils have taller (Fig. 6a) and more abundant (Table 2) pneumatophores compared to areas with sandy soil (E.g., Sarker et al. 2022; Best et al. 2022). Pneumatophores with one branch (i.e., 1-branch) is the most abundant and has high population in the stagnant pool (Fig. 3). In contrast the sandy seashore site has more diversity by having more pneumatophore types (1-branch, 2-branch, 3-branch, and 4-branch). Nonetheless, the length and weight of the pneumatophores have little or no correlations to the number of branches (Fig. 7). However, the presence of more pneumatophore types is a survival tendency for the *A. germinans* tree to survive in an nutrient-depleted environment such as the seashore where the coast is constantly battered by harsh tidal currents. In other words, the more the branches of the pneumatophore the better it becomes in the utilization of oxygen in an anaerobic environment. (Al-khayat and Alatalo, 2022). On the other hand, the stagnant pool being a more stable environment supports more single branch pneumatophores (1-branch), which grow taller and transmit more oxygen into the mangrove environment and surrounding sediment (Krauss et al. 2003). Although, the rich supply of pneumatophores attracts spawning organisms (Wickramasinghe et al. 2022; Deng et al. 2022), it also serves as a trap for plastic pollutants (Cesarini and Scalici, 2022) that contaminates organisms (Fig. 3) (Portz et al. 2022).

This study showed that three key factors influence the growth of pneumatophores in the deforested terrain, namely: stagnant pool of water, soil composition, and tree canopy cover. The stagnant pool has the largest population of pneumatophores because the standing water act as a trap for organisms (Fig. 2), which die and decompose to increase the total organic content of the soil (Table 1). In addition, when sun heats the pond, chemical reactions do occur (i.e., hydration and hydrolysis), which erodes the sub soil leading to an increase in the acidity and heavy metal concentration of the water (Fig. 4). The acidic water breaks down any plant and animal that enter the stagnant pool (Das et al. 2022). The disintegrated organic matter increases the TOC, which in turn accelerates the growth of pneumatophores. In contrast, the seashore site is made of sandy soil that has little organic content mostly because of the “flushing action” of the tidal force that sweeps the surface clean of any plant and animal materials during high tide. Nevertheless, the high microbial action at the seashore site (Fig. 5) does not translate to high decomposition rate, leading to low pneumatophore growth. Furthermore, the presence of the black mangrove canopy cover at the stagnant pool sites influences the local climate, which regulates the temperature of the soil and water beneath (Mariano et al. 2022). The warming of the stagnant pool influences its chemistry and microbiology. Similarly, increased litter fall from nearby trees lead to the

increase in decomposition rate of the fallen leaves. The decomposed leaves form organic matter that increases the fertility of the soil leading to the growth of more pneumatophores.

The presence of a stagnant pool is also a strong factor that facilitates the growth of the pneumatophores by changing the chemical and microbial dynamics of the soil beneath the pool of water. The water helps to hydrolyze and liquify dead plant and animal matter and convert them to manure for the micro-organisms (e.g., bacteria, and fungi) to act upon. Pneumatophores help the black mangroves to survive the harsh swampy environmental condition by acting as the channel for oxygen transmission into the plants. Their presence in the mangrove forest also contribute to climate stabilization through their action of carbon sequestration. Soil condition, forest canopy and stagnant pool facilitate the growth and development of pneumatophores. The more the population of the pneumatophores the better the environmental quality, which can lead to the proliferation of spawning aquatic organisms useful to humans. Therefore, the abundance and growth in length of the pneumatophores can be used as biological indicator to determine the level of pollution of the mangrove forest, which is significant in pollution studies, restoration ecology and fish population studies.

Declarations

Acknowledgements

I thank my research team comprising of Mr. Chimezie Brown and my graduate students Mr. Emoyoma Udi and Miss Peace Ohia for assistance in sample collection.

Funding Not applicable

Conflict of Interest

The author claims no competing interests.

Author Contribution

Aroloye O. Numbere: Conceptualization (lead); Data curation (lead); Formal analysis (lead); Investigation (lead); Resources (lead); Supervision (lead); Writing-original draft (lead); Writing-review & editing (lead)

Gerardo R. Camilo: Conceptualization, Data curation, Supervision, review, and editing

References

1. Al-Khayat, J. A., & Alatalo, J. M. (2021). Relationship between tree size, sediment mud content, oxygen levels, and pneumatophore abundance in the mangrove tree species *Avicennia marina* (Forssk.) Vierh. *Journal of Marine Science and Engineering*, 9(1), 100.
2. Best, Ü. S., van der Wegen, M., Dijkstra, J., Reyns, J., van Prooijen, B. C., & Roelvink, D. (2022). Wave attenuation potential, sediment properties and mangrove growth dynamics data over Guyana's

- intertidal mudflats: assessing the potential of mangrove restoration works. *Earth System Science Data*, 14(5), 2445–2462.
3. Carlton, J. M. (1974). Land-building and stabilization by mangroves. *Environmental conservation*, 1(4), 285–294.
 4. Cesarini, G., & Scalici, M. (2022). Riparian vegetation as a trap for plastic litter. *Environmental Pollution*, 292, 118410.
 5. Clough, B. (2013). Continuing the journey amongst mangroves. *ISME mangrove educational book series*, (1), 86.
 6. Das, N., Mondal, A., & Mandal, S. (2022). Polluted waters of the reclaimed islands of Indian Sundarban promote more greenhouse gas emissions from mangrove ecosystem. *Stochastic Environmental Research and Risk Assessment*, 36(5), 1277–1288.
 7. Deng, H., Fu, Q., Zhang, Y., Li, D., He, J., Feng, D., ... & Ge, C. (2022). Bacterial communities on polyethylene microplastics in mangrove ecosystems as a function of exposure sites: compositions and ecological functions. *Journal of Environmental Chemical Engineering*, 107924.
 8. Eisenberg, E. (2002). A Field Guide to the Southeast Coast and Gulf of Mexico. In *A Field Guide to the Southeast Coast and Gulf of Mexico*. Yale University Press.
 9. Fusi, M., Booth, J. M., Marasco, R., Merlino, G., Garcias-Bonet, N., Barozzi, A., ... & Daffonchio, D. (2022). Bioturbation Intensity Modifies the Sediment Microbiome and Biochemistry and Supports Plant Growth in an Arid Mangrove System. *Microbiology Spectrum*, e01117-22.
 10. Hilmi, E., Dewi, R., Sudiana, E., Mahdiana, A., Sari, L. K., & Cahyo, T. N. (2022). The Clustering and Distribution of Heavymetal Accumulation and Translocation as an Ability of Mangrove Vegetation to Reduce Impact of Heavymetal (Hg, Cd and Zn) Pollution.
 11. Hogarth, P. J. (2015). *The biology of mangroves and seagrasses*. Oxford University Press.
 12. Hoogeveen, S. J. (2020). *Mangrove dynamics in the Richmond River's estuary* (Master's thesis, University of Twente).
 13. Kathiresan, K., & Bingham, B. L. (2001). Biology of mangroves and mangrove ecosystems. 81–251.
 14. Ke, Y., Li, J., Yuan, W., Chen, Y., Zhao, B., Tang, Z., ... & Tang, Y. (2022). Mangrove Root-Inspired Carbon Nanotube Film for Micro-Direct Methanol Fuel Cells. *ACS Applied Materials & Interfaces*.
 15. Krauss, K. W., Allen, J. A., & Cahoon, D. R. (2003). Differential rates of vertical accretion and elevation change among aerial root types in Micronesian mangrove forests. *Estuarine, Coastal and Shelf Science*, 56(2), 251–259.
 16. Krauss, K. W., McKee, K. L., Lovelock, C. E., Cahoon, D. R., Saintilan, N., Reef, R., & Chen, L. (2014). How mangrove forests adjust to rising sea level. *New Phytologist*, 202(1), 19–34.
 17. Mariano, H., Aguilos, M., Dagoc, F. L., Sumalinab, B., & Amparado, R. (2022). Abandoned Fishpond Reversal to Mangrove Forest: Will the Carbon Storage Potential Match the Natural Stand 30 Years after Reforestation?. *Forests*, 13(6), 847.

18. Munir, N., Hasnain, M., Roessner, U., & Abideen, Z. (2022). Strategies in improving plant salinity resistance and use of salinity resistant plants for economic sustainability. *Critical Reviews in Environmental Science and Technology*, 52(12), 2150–2196.
19. Numbere, A. O. (2018). Mangrove species distribution and composition, adaptive strategies and ecosystem services in the Niger River Delta, Nigeria. *Mangrove ecosystem ecology and function*, 7, 17.
20. Ong, J. E., & Gong, W. K. (2013). *Structure, function, and management of mangrove ecosystems*. International Society for Mangrove Ecosystems.
21. Ouyang, X., Lai, D. Y., Marchand, C., & Lee, S. Y. (2022). Carbon storage and mineralization in coastal wetlands. In *Carbon Mineralization in Coastal Wetlands* (pp. 295–310). Elsevier.
22. Portz, L., Manzolli, R. P., Villate-Daza, D. A., & Fontán-Bouzas, Á. (2022). Where does marine litter hide? The Providencia and Santa Catalina island problem, seaflower reserve (Colombia). *Science of the Total Environment*, 813, 151878.
23. Pratolongo, P. D. (2022). Salt Marshes and Mangroves: Tidal Saline Wetlands Dominated by Vascular Plants. *Marine Biology: A Functional Approach to the Oceans and their Organisms*, 211.
24. Romero-Urbe, H. M., López-Portillo, J., Reverchon, F., & Hernández, M. E. (2022). Effect of degradation of a black mangrove forest on seasonal greenhouse gas emissions. *Environmental Science and Pollution Research*, 29(8), 11951–11965.
25. Roy, S. (2011, December). Seasonally and spatially coordinated strategy of detritus conservation and use in the world's largest mangrove ecosystem. In *Proceedings of the Zoological Society* (Vol. 64, No. 2, pp. 63–71). Springer-Verlag.
26. Sarker, S., Masud-Ul-Alam, M., Hossain, M. S., Rahman Chowdhury, S., & Sharifuzzaman, S. M. (2021). A review of bioturbation and sediment organic geochemistry in mangroves. *Geological Journal*, 56(5), 2439–2450.
27. Spencer, T., Möller, I., & Reef, R. (2022). Mangrove systems and environments. *Journal: Treatise on Geomorphology*, 675–712.
28. Stewart, M., & Fairfull, S. (2008). Mangroves. In *Industries, Ndop (Ed.)*. Sydney, NSW Government.
29. Wickramasinghe, S., Wijayasinghe, M., & Sarathchandra, C. (2022). Sri Lankan Mangroves: Biodiversity, Livelihoods, and Conservation. In *Mangroves: Biodiversity, Livelihoods and Conservation* (pp. 297–329). Springer, Singapore.
30. Zhang, C., Zhang, Y., Luo, M., Tan, J., Chen, X., Tan, F., & Huang, J. (2022). Massive methane emission from tree stems and pneumatophores in a subtropical mangrove wetland. *Plant and Soil*, 473(1), 489–505.

Figures




Figure 1

Map of study area at Eagle Island, Niger Delta, Nigeria




Figure 2

Experimental design of pneumatophore growth, abundance and diversity study carried out in a 94.78 m × 43.76 m (4152.24m²) plot at Eagle Island, Niger delta. The picture indicates (a) the stagnant pool and (b) the seashore sites where the pneumatophores were studied. The drawing of the study area shows the points of sample collection.




Figure 3

Slides showing (a) fish brought in by tidal current into the sand fill area (b) fish on grass (*longibrateatus*), (c) accumulated plastic waste trapped in grasses, (d) dead fish dissected to show decaying internal organs few minutes after death, which might be caused by chemical poisoning from accumulated waste.




Figure 4

Mean values ($\pm$ SE) of abundance of different types of pneumatophores of black mangroves (*A. germinans*) in different sites at Eagle Island, Niger Delta. It shows that stagnant pool area with muddy soil has high pneumatophore abundance than the seashore area with sandy soil. There is also an increase in siltiness (muddy soil condition) from the seashore to the stagnant pool.




Figure 5

Physico-chemical analysis of soil at some locations at Eagle Island, Niger Delta, Nigeria. It shows that seashore has more Cd and THC while the stagnant pool has more concentration of Pb, and Zn.




Figure 6

Microbial population at different sites at Eagle Island, Niger Delta, Nigeria. It shows that bacterial population is more dominant than the fungi population. Similarly, seashore site has more microbial population than the stagnant pool site.




Figure 7

length and weight comparison of pneumatophores growing in different sites at Eagle Island, Niger Delta, Nigeria




Figure 8

Regression analysis of (a) no. of dead vs, no of alive and (b) length versus weight, (c) length versus branch and (d) weight versus branch of pneumatophores at Eagle Island.